









20. Harbige LS, Sharief MK. Polyunsaturated fatty acids in the pathogenesis and treatment of multiple sclerosis. *Br J Nutr.* 2007;98:S46–S53.
21. Shiraishi K, Matsuzaki S, Ishida H, et al. Impaired erythrocyte deformability and membrane fluidity in alcoholic liver disease: participation in disturbed hepatic microcirculation. *Alcohol Alcohol Suppl.* 1993;1:59–64.
22. Tempelhoff GF, Ullrich S, Heilmann L. Erythrocyte deformability in pregnancy-induced hypertension. *Zentralbl Gynakol.* 1994;116:160–163.
23. Patel KV, Mohanty JG, Kanapuru B, et al. Association of the red cell distribution width with red blood cell deformability. *Adv Exp Med Biol.* 2013;765:211–216.
24. Brück W. Clinical implications of neuropathological findings in multiple sclerosis. *J Neurol.* 2005;252:10–14.
25. Stinissen P, Raus J, Zhang J. Autoimmune pathogenesis of multiple sclerosis: role of autoreactive T lymphocytes and new immunotherapeutic strategies. *Crit Rev Immunol.* 1997;17:33–75.
26. Ghaffari S. Oxidative stress in the regulation of normal and neoplastic hematopoiesis. *Antioxid Redox Signal.* 2008;10:1923–1940.
27. Fukuta H, Ohte N, Mukai S, et al. Elevated plasma levels of B-type natriuretic Peptide but not C-reactive protein are associated with higher red cell distribution width in patients with coronary artery disease. *Internat Heart J.* 2009;50:301–312.
28. Song CS, Park DI, Yoon MY, et al. Association between red cell distribution width and disease activity in patients with inflammatory bowel disease. *Dig Dis Sci.* 2012;57:1033–1038.