

The impact of weight gain during adjuvant chemotherapy on survival in breast cancer

Can Atalay¹, Ali İmran Küçük²

ABSTRACT

Objective: Weight gain can be detected during adjuvant chemotherapy in breast cancer patients, leading to administration of lower drug doses than planned and a decrease in long-term survival. In this study, the effect of weight gain on survival in breast cancer patients was investigated.

Material and Methods: Breast cancer patients treated with adjuvant chemotherapy between May 2002 and May 2003 were prospectively included in the study. Patients treated with neoadjuvant chemotherapy or with chemotherapy for metastatic disease were excluded from the study. Data regarding patients' demographic, clinical and pathologic characteristics and chemotherapy protocols were recorded. Patients were grouped as those with weight gain less than or more than 3 kg, and those with a body mass index of less than or more than 30. The impact of weight gain on patients' disease-free and overall survival was investigated. Log-rank test and Cox regression analyses were utilized for survival analyses. $P < 0.05$ value was accepted as statistically significant.

Results: Eighty-eight consecutive female patients with a median age of 46 (29-71) were included in the study. Patients received anthracycline based chemotherapy protocols. Weight gain was detected in 79 patients (89.8%), with more than 3 kg weight gain detected in 38 patients (43.2%). In a median follow-up time of 98 months (62-120), distant metastases were detected in 21 patients (23.9%), and 11 patients (12.5%) died. Mean disease-free survival of patients with a weight gain less than and more than 3 kg during chemotherapy was 89.1 ± 3.9 and 84.7 ± 4.2 months ($p = 0.007$), whereas mean overall survival was 95.6 ± 2.2 and 92.5 ± 2.1 months ($p = 0.01$), respectively. Mean disease-free survival of patients with a body mass index less than and more than 30 was 87.3 ± 2.3 and 85.1 ± 3.6 months ($p = 0.4$), whereas mean overall survival was 94.2 ± 2.3 and 92.1 ± 1.1 months ($p = 0.35$), respectively.

Conclusion: Weight gain during adjuvant chemotherapy has a negative effect on both disease-free and overall survival in patients with breast cancer.

Keywords: Breast cancer, chemotherapy, weight gain, survival

INTRODUCTION

Modern treatment options for breast cancer have prolonged survival, especially in patients diagnosed at early stages. Weight gain is observed during or following adjuvant chemotherapy in breast cancer patients who were followed-up for a long period. The cause of weight gain has not been revealed in studies conducted so far. Previous studies have reported that having a high body mass index on diagnosis is a poor prognostic factor (1, 2). However, studies on the effect of weight changes in patients receiving adjuvant chemotherapy on survival reported conflicting results. Clarification of controllable factors that have an effect on survival, such as weight gain, will be beneficial in order to improve outcome in breast cancer treatment.

Doses of the drugs used in adjuvant chemotherapy regimens are determined based on extensive studies. It is aimed to achieve maximum efficiency by giving a minimum amount of drug in the calculation of these doses. In addition, the systemic side effects of the drug are intended to be minimized. Since drug doses are calculated according to body surface area, weight gain during chemotherapy may result in lower drug doses than planned. This in turn may lead to shorter disease-free and overall survival since these patients do not receive the required drug dose. In this study, we aimed to investigate the effect of weight gain during chemotherapy on survival in breast cancer patients treated in our clinic.

MATERIAL AND METHODS

Patients who were diagnosed with breast cancer and received surgical and adjuvant chemotherapy in our clinic between May 2002 to May 2003 were enrolled in this prospective study. Patients who underwent neoadjuvant chemotherapy or chemotherapy for metastatic breast cancer were excluded. Demographic data (age, gender), and clinic-pathological characteristics (menopausal status, TNM stage, tumor estrogen, progesterone, and HER2 receptor status, grade) were recorded along with the chemotherapy regimens applied. Patient weight and height were measured before and at the end of chemotherapy. Patient's body mass index (kg/m^2) was calculated before treatment. Patients were grouped according to weight gain of more or less than 3 kg, and pre-treatment body mass index more or less than 30, based on measurements before and after chemotherapy. Effects of menopausal status, initial body mass in-

¹Clinic of General Surgery, Ankara Oncology Training and Research Hospital, Ankara, Turkey

²Clinic of General Surgery, Derince Training and Research Hospital, Kocaeli, Turkey

Address for Correspondence

Can Atalay

Ankara Onkoloji Eğitim ve Araştırma Hastanesi, Genel Cerrahi Kliniği, Ankara, Türkiye
Phone: +90 532 263 55 14
e-mail: atalay_can@hotmail.com

Received: 24.02.2015

Accepted: 21.03.2015

©Copyright 2015
by Turkish Surgical Association
Available online at
www.uluscerrahidergisi.org

dex and the amount of weight gain during chemotherapy on disease-free and overall survival were evaluated.

Statistical Analysis

Statistical Package for the Social Sciences 15.0 (SPSS Inc., Chicago, Illinois, USA) was used for analysis. Descriptive statistics were presented as mean ± standard deviation for continuous variables, and as frequencies and percentages for categorical variables. For evaluation of disease-free and overall survival, log-rank test was used for univariate analysis and Cox regression analysis for multivariate analysis. P<0.05 was considered statistically significant.

RESULTS

The study enrolled 88 consecutive female patients. The median age of patients was 46 years (29-71). Demographic characteristics of patients and tumor histopathology are outlined in Table 1. There were 53 (60%) premenopausal patients. Most patients (66%) had stage II breast cancer. As surgical treatment, 65 patients (73.9%) underwent modified radical mastectomy while the remaining received breast-conserving therapy and axillary dissection. Axillary lymph node metastases were detected in 55 patients (62.5%). Twenty-eight patients had 1-3, 14 patients 4-9 and 13 patients had >9 lymph node metastases. Estrogen and progesterone receptors were positive in 50 (57%) and 59 (67%) patients, respectively. HER2 positivity was detected in 38 patients (43%). Patients mostly had grade 3 tumors (49%). The median body mass index before treatment was 27 (17-40). This value was higher than 30 in 26 patients (29.5%).

The anthracycline-containing chemotherapy regimens were applied in patients with proper indications according to the stage of breast cancer after surgical treatment. Chemotherapy was started within 3-6 weeks after surgery. Chemotherapy protocols included CAF (cyclophosphamide 600 mg/m², doxorubicin 60 mg/m², 5-fluorouracil 600 mg/m²) and CEF (cyclophosphamide 600 mg/m², epirubicin 50 mg/m², 5-fluorouracil 600 mg/m²). The anthracycline used in the chemotherapy protocol was epirubicin in 57 patients (65%), and doxorubicin in 31 patients. All patients received 6 cycles of chemotherapy in the planned doses. In addition, hormone receptor-positive patients were treated with tamoxifen or an aromatase inhibitor after chemotherapy according to their menopausal status.

1-12 kg weight gain was detected in seventy-nine patients (89.8%) during chemotherapy, with a median weight gain of 3 kg. Weight gain was greater than 3 kg in 38 patients (43.2%). Distant metastases were detected in 21 patients (23.9%) during a median of 98 months (62-120) follow-up and 11 patients (12.5%) died. Local-regional recurrence was not detected in any of these patients. The mean disease-free survival and overall survival was 89.1±3.9 and 84.7±4.2 months (p=0.007), and 95.6±2.2 and 92.5±2.1 months (p=0.01) in patients with less than or more than 3 kg weight gain during chemotherapy, respectively. Five-year disease-free survival and 5-year overall survival was 85.2% and 78.0%, and was 93.7% and 90.0% in patients with less than or more than 3 kg weight gain, respectively.

However, disease-free survival and overall survival was 87.3±2.3 and 85.1±3.6 months (p=0.4), and 94.2±2.3 and 92.1±1.1 months (p=0.35) in patients with a body mass in-

dex less or more than 30 at the time of diagnosis. The 5-year disease-free survival and 5-year overall survival was 84% and 78.3%, and was 90.8% and 89.8% in patients with a body mass index less or more than 30, respectively. Similarly, there was no statistically significant difference in terms of disease-free survival (p=0.6) and overall survival (p=0.23) between premenopausal and postmenopausal patients.

DISCUSSION

Previous studies have reported high body mass index at the time of breast cancer diagnosis as a negative prognostic factor (1, 2). However, conflicting results have been published regarding the effect of weight gain during chemotherapy on prognosis. The difference in results may be due to reasons such as retrospective data analysis, administration of anthracycline-free chemotherapy protocols, lack of data if patients have received the planned doses or not, monitoring weight gain at different time intervals, and short follow-up period. According to results of our study, weight gain (>3 kg) during adjuvant chemotherapy in patients with breast cancer has negative effects on both disease-free and overall survival. This

Table 1. Patient demographics and tumor histopathology characteristics

	n	%
Age	Median 46	Range 29-71
Body mass index		
<30	62	70.5
≥30	26	29.5
Menopausal status		
Premenopausal	53	60.0
Postmenopausal	35	40.0
Pathologic stage*		
I	11	12.5
II A	31	35.0
II B	27	31.0
III A	19	21.5
Axillary status		
Positive	55	52.5
Negative	33	37.5
Estrogen receptor		
Positive	50	57.0
Negative	38	43.0
Progesterone receptor		
Positive	59	67.0
Negative	29	33.0
Grade		
I	10	11.0
II	35	40.0
III	43	49.0

*According to American Joint Committee on Cancer 2002

finding may be related to the prospective study plan, administration of standard chemotherapy protocol containing anthracycline, and a median follow-up period longer than 8 years. In contrast, it was determined that high body mass index at the time of diagnosis and menopausal status did not have an effect on either disease-free or overall survival.

There are studies on the effects of weight gain or obesity prior to the diagnosis of breast cancer on survival; however, there are only limited number of studies evaluating the progress during chemotherapy. In a recent meta-analysis of 43 studies, general and breast cancer specific mortality rates have been reported to be higher in obese women as compared to non-obese (respectively, HR=1.33, 95% CI: 1.21 to 1.47 and HR=1.33, 95% CI: 1.19 to 1.50). Nevertheless, when interpreting these results it should be considered that studies spanning over a wide time range (1963-2005) and with varying number of patients (100 to 424.168) were included in this meta-analysis (3). While weight gain before diagnosis increases the risk of breast cancer in postmenopausal women, both premenopausal and postmenopausal patients gain weight after diagnosis. Initial studies on weight gain in breast cancer detected weight gain in one fourth, with more than 10 kg increase in weight (3, 4). As it has been shown that weight gain may adversely affect prognosis of breast cancer, physicians began warning their patients about this effect thus resulting in a decrease in weight gain. In contrast to these, there are reports stating that patients who lose weight after diagnosis of breast cancer have a poor prognosis. In this case, whether the poor prognosis was related to weight loss or disease progression could not be clearly determined.

Weight gain during chemotherapy can have a direct effect on survival as well as an indirect effect by causing patients to receive lower doses of chemotherapy drugs. Weight gain due to hormonal changes can be expected in patients undergoing menopause as a result of disruption of ovarian function during chemotherapy. In accordance, weight gain during chemotherapy is observed more in premenopausal women. Moreover, chemotherapy leads to weight gain in patients with reduced physical activity (5, 6). There are reports stating that resting energy consumption by the body is decreased in patients receiving chemotherapy (5, 7). The reasons for weight gain during chemotherapy include the reduction in energy consumption (slowing down of the basal metabolic rate, decreased physical activity, thermogenesis) and an increase in food intake (8).

The results of studies on the effects of weight gain after diagnosis of breast cancer on prognosis are contradictory. Some studies report increased disease recurrence and mortality with weight gain (4, 8-10), while other studies did not reveal this relationship (11-13). Similar to our study, Thivat et al. (8) retrospectively investigated the effect of weight change on prognosis of patients receiving anthracycline-based chemotherapy protocol. Within a follow-up period of more than twenty-years, more than 5% change in weight during chemotherapy increased the risk of recurrence and mortality more than two-fold. In their study including 13000 patients, Caan et al. (14) reported that preservation of the same weight for a couple of years after diagnosis of breast cancer has a positive impact on the survival. In contrast, weight loss and concomitant diseases reduce survival rate. Bradshaw et al. (15) conducted a study on

1,436 patients, and reported poor prognosis in patients with more than 10% weight gain within 5 years of chemotherapy (HR=2.67, 95% CI: 1.37, 5.05). This negative effect was more prominent in those who gained weight within the first two years.

Similar to our results, in their series with more than 7 years follow-up Camoriano et al (4) reported that breast cancer patients with weight gain more than 5.9 kg had a 1.5 times higher recurrence rate and 1.6-fold increase in mortality. Similarly, in the Nurses' Health Study with 5,000 patients, weight gain was reported as a factor that increased both recurrence and mortality in breast cancer (9). In contrast to these, studies not reporting weight gain in patients treated with anthracycline-containing chemotherapy regimens are also available (8).

Different mechanisms have been proposed to explain the negative prognostic effects of weight gain in breast cancer patients. The increase in fat tissue in the body with weight gain leads to increased estrogen production particularly in postmenopausal women, and reduced sex hormone binding protein. As a result of these changes, the increased free estrogen in the blood may play a role in local-regional recurrence of breast cancer, formation of a new cancer in the opposite breast and distant metastases. Additionally, an increase in insulin, insulin resistance and C-peptide levels is also detected in this patient group, leading to hyperglycemia. Similarly, an increase in inflammatory markers such as C-reactive protein and interleukins has been reported in breast cancer patients with weight gain (1, 16). The resulting metabolic scenario has a negative impact on the prognosis of breast cancer. In addition to these, it has been reported that insulin and insulin-like growth factors lead to the development of resistance to chemotherapy and radiotherapy in tumor cells (17).

Patients usually implement high carbohydrate diet during adjuvant chemotherapy due to fear of losing weight. In addition, they consume foods low in fat and rich in fruits and vegetables. The weight loss during adjuvant therapy or follow-up of breast cancer patients is not due to the inability to feed but due to aggressive tumor biology and accelerated metabolism. Patients with early breast cancer should not be eating more than usual, since these alterations are unlikely. The nutrition strategy in breast cancer patients should be directed at reduction of excess weight while restricting carbohydrates, and increasing fat and protein intake (18-20). To improve prognosis in breast cancer patients, it is suggested that physical activity and a diet rich in fruits, vegetables and grains that are low in fat and refined carbohydrates should be increased (21). Weight control methods should be individualized for patients with breast cancer, as is the case in adjuvant therapy. Exercise and dietary recommendations should be made in accordance with the patient's body mass index, the muscle-fat ratio and existing co-morbidities prior to diagnosis. Studies have shown high rate of patient compliance with exercise and diet programs initiated during chemotherapy (22, 23).

Study Limitations

Study limitations include the relatively small number of patients, and lack of data on distribution of excess weight in the tissues. Weight measurements were made during chemotherapy, and if weight gain was monitored further in the follow-up

of patients after chemotherapy, it is likely that patients with ongoing weight gain would have a shorter survival.

CONCLUSION

According to our study results, weight gain (> 3 kg) during adjuvant chemotherapy in patients with breast cancer negatively affects both disease-free and overall survival. In contrast, high body mass index on diagnosis and menopausal status were not found to influence either disease-free or overall survival.

Ethics Committee Approval: Ethics committee approval was not received for this study since it only involved weight measurement.

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - C.A.; Design - C.A.; Supervision - C.A.; Funding - C.A.; Materials - C.A., A.I.K.; Data Collection and/or Processing - C.A., A.I.K.; Analysis and/or Interpretation - C.A., A.I.K.; Literature Review - C.A.; Writer - C.A.; Critical Review - C.A.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Rock CL, Demark-Wahnefried W. Nutrition and survival after the diagnosis of breast cancer: a review of the evidence. *J Clin Oncol* 2002; 20: 3302-3316. [\[CrossRef\]](#)
2. Kumar NB, Cantor A, Allen K, Cox CE. Android obesity at diagnosis and breast carcinoma survival. *Cancer* 2000; 88: 2751-2757. [\[CrossRef\]](#)
3. Demark-Wahnefried W, Winer EP, Rimer BK. Why women gain weight with adjuvant chemotherapy for breast cancer. *J Clin Oncol* 1993; 11: 1418-1429.
4. Camoriano JK, Loprinzi CL, Ingle JN, Therneau TM, Krook JE, Veeder MH. Weight change in women treated with adjuvant therapy or observed following mastectomy for node-positive breast cancer. *J Clin Oncol* 1990; 8: 1327-1334.
5. Demark-Wahnefried W, Hars V, Conaway MR, Havlin K, Rimer BK, McElveen G, et al. Reduced rates of metabolism and decreased physical activity in breast cancer patients receiving adjuvant chemotherapy. *Am J Clin Nutr* 1997; 65: 1495-1501.
6. Demark-Wahnefried W, Peterson BL, Winer EP, Marks L, Aziz N, Marcom PK, et al. Changes in weight, body composition, and factors influencing energy balance among premenopausal breast cancer patients receiving adjuvant chemotherapy. *J Clin Oncol* 2001; 19: 2381-2389.
7. Harvie MN, Campbell IT, Baildam A, Howell A. Energy balance in early breast cancer patients receiving adjuvant chemotherapy. *Breast Cancer Res Treat* 2004; 83: 201-210. [\[CrossRef\]](#)
8. Thivat E, Therondel S, Lapirot O, Abrial C, Gimbergues P, Gadea E, et al. Weight change during chemotherapy changes the prognosis in non metastatic breast cancer for the worse. *BMC Cancer* 2010; 10: 648. [\[CrossRef\]](#)
9. Kroenke CH, Chen WY, Rosner B, Holmes MD. Weight, weight gain, and survival after breast cancer diagnosis. *J Clin Oncol* 2005; 23: 1370-1378. [\[CrossRef\]](#)
10. Nichols HB, Trentham-Dietz A, Egan KM, Titus-Ernstoff L, Holmes MD, Bersch AJ, et al. Body mass index before and after breast cancer diagnosis: associations with all-cause, breast cancer, and cardiovascular disease mortality. *Cancer Epidemiol Biomarkers Prev* 2009; 18: 1403-1409. [\[CrossRef\]](#)
11. Caan BJ, Kwan ML, Hartzell G, Castillo A, Slattery ML, Sternfeld B, et al. Pre-diagnosis body mass index, post-diagnosis weight change, and prognosis among women with early stage breast cancer. *Cancer Causes Control* 2008; 19: 1319-1328. [\[CrossRef\]](#)
12. Caan BJ, Emond JA, Natarajan L, Castillo A, Gunderson EP, Habel L, et al. Post-diagnosis weight gain and breast cancer recurrence in women with early stage breast cancer. *Breast Cancer Res Treat* 2006; 99: 47-57. [\[CrossRef\]](#)
13. Makari-Judson G, Judson CH, Mertens WC. Longitudinal patterns of weight gain after breast cancer diagnosis: observations beyond the first year. *Breast J* 2007; 13: 258-265. [\[CrossRef\]](#)
14. Caan BJ, Kwan ML, Shu XO, Pierce JP, Patterson RE, Nechuta SJ, et al. Weight change and survival after breast cancer in the after breast cancer pooling project. *Cancer Epidemiol Biomarkers Prev* 2012; 21: 1260-1271. [\[CrossRef\]](#)
15. Bradshaw PT, Ibrahim JG, Stevens J, Cleveland R, Abrahamson PE, Satia JA, et al. Post-diagnosis change in bodyweight and survival after breast cancer diagnosis. *Epidemiology* 2012; 23: 320-327. [\[CrossRef\]](#)
16. Siiteri PK. Adipose tissue as a source of hormones. *Am J Clin Nutr* 1987; 45 (Suppl 1): 277-282.
17. Turner BC, Haffty BG, Narayanan L, Yuan J, Havre PA, Gumbs AA, et al. Insulin-like growth factor-1 receptor overexpression mediates cellular radioresistance and local breast cancer recurrence after lumpectomy and radiation. *Cancer Res* 1997; 57: 3079-3083.
18. Forsythe CE, Phinney SD, Fernandez ML, Quann EE, Wood RJ, Bibus DM, et al. Comparison of low fat and low carbohydrate diets on circulating fatty acid composition and markers of inflammation. *Lipids* 2008; 43: 65-77. [\[CrossRef\]](#)
19. Volek JS, Phinney SD, Forsythe CE, Quann EE, Wood RJ, Puglisi MJ, et al. Carbohydrate restriction has a more favorable impact on the metabolic syndrome than a low fat diet. *Lipids* 2009; 44: 297-309. [\[CrossRef\]](#)
20. Seshadri P, Iqbal N, Stern L, Williams M, Chicano KL, Daily DA, et al. A randomized study comparing the effects of a low-carbohydrate diet and a conventional diet on lipoprotein subfractions and C-reactive protein levels in patients with severe obesity. *Am J Med* 2004; 117: 398-405. [\[CrossRef\]](#)
21. McTiernan A, Irwin M, VonGruenigen V. Weight, physical activity, diet, and prognosis in breast and gynecologic cancers. *J Clin Oncol* 2010; 28: 4074-4080. [\[CrossRef\]](#)
22. Djuric Z, Ellsworth JS, Weldon AL, Ren J, Richardson CR, Resnicow K, et al. A diet and exercise intervention during chemotherapy for breast cancer. *Open Obes J* 2011; 3: 87-97. [\[CrossRef\]](#)
23. Champ CE, Volek JS, Siglin J, Jin L, Simone NL. Weight gain, metabolic syndrome, and breast cancer recurrence: Are dietary recommendations supported by the data? *Int J Breast Cancer* 2012; 506868. [\[CrossRef\]](#)