


LETTER TO THE EDITOR

Open Access



Severity of hypoxia modulates effect of CPAP on myocardial stress as measured by highly sensitive troponin T

Amir Sharafkhaneh^{1,3*} , Jennifer Katigbak⁴, Max Hirshkowitz^{1,3}, Hossein Sharafkhaneh³, Saba P. Sharafkhaneh⁵, Christie M. Ballantyne^{3,6}, Biykem Bozkurt^{2,3} and Vijay Nambi^{2,3,6}

Dear editor

Obstructive sleep apnea (OSA) is associated with increased risk for cardio- and cerebrovascular diseases [1, 2]. Animal studies propose intermittent hypoxia and the resulting inappropriate activation of sympathetic nervous system [3, 4] as the major link between OSA and cardio- and cerebrovascular comorbidities. However, the OSA severity level at which continuous positive airway pressure (CPAP) therapy offers cardio- and cerebrovascular benefit (reduction in indices of myocardial stress/injury) is a matter of debate. Sub-clinical injury to the myocardium is considered to be a precursor for the development of incident cardiovascular disease including heart failure [5]. Troponin T measured with a high sensitivity assay (Hs-TnT) has been shown to predict incident heart failure and cardiovascular death [6]. Further, increase or decrease in Hs-TnT over time predicted increased or decreased incidence of heart failure and cardiovascular death [7, 8]. Hs-TnT has also shown to be increased with presence and severity of OSA [9]. Stronger correlations were reported with nadir SpO₂ [10]. Roca and colleagues, in a cohort of 1645 subjects free of coronary artery disease and heart failure with a median follow up of 12.4 years, reported that hs-TnT was associated with risk of death or incident heart failure across various categories of OSA after adjusting for 17 potential confounders [11]. Surprisingly, a recent study showed increased Hs-TnT after 12 months of CPAP therapy in patients with OSA. However, the utilization of CPAP was not reported [12]. We report changes in Hs-TnT level as a

marker of myocardial stress/injury in a cohort of CPAP compliant patients with OSA.

Methods

This study was approved by institutional review board of Baylor College of Medicine and Research & Development Committee of Michael E. DeBakey VA Medical Center. All participants signed an informed consent form. We enrolled adult subjects with confirmed OSA (apnea + hypopnea index (AHI) of ≥ 15 obstructive and/or mixed events/h) using attended polysomnography (PSG). Subjects qualifying for the PSG underwent blood sampling and completed an Epworth Sleepiness Scale (ESS). The participants underwent attended CPAP titration and were placed on therapy. The best pressure was the one associated with the lowest AHI while the patient slept 20 min, or more. After titration, subjects received a CPAP machine and related accessories (Respironics, REMStar Pro) with card reader to monitor the compliance of CPAP and were followed for 6 months. Subjects were seen 2–3 times during the study and CPAP compliance was checked during the visit. CPAP efficacy was rechecked with overnight pulse oximetry at the end of the study. Blood was collected in the morning between 7 and 8 AM in EDTA-containing tubes and kept at 4 °C during processing at baseline and after 6 months of CPAP use. Aprotinin (100 μ L containing 0.6 TIU per mL of blood) was added to one of the tubes and the samples were then centrifuged at 3000 rpm for 30 min. Hs-TnT concentrations were measured with a novel high sensitivity assay, Elecsys Troponin T (Roche Diagnostics[®]), on an automated Cobas e411 analyzer with a limit of measurement of 3 ng/L [13].

* Correspondence: amirs@bcm.edu

¹Section of Pulmonary, Critical Care and Sleep Medicine, Medical Care Line, Michael E. DeBakey Veterans Affairs Medical Center, MED VA Medical Center, Bldg. 100 (111i), 2002 Holcombe Blvd, Houston, TX 77025, USA

³Department of Medicine, One Baylor Plaza, Houston, TX 77030, USA

Full list of author information is available at the end of the article

Results

Twenty-three subjects were enrolled and 20 subjects completed the study. One subject died unexpectedly at home, from unknown causes. Two subjects were lost to follow up. Baseline and follow samples were available for analysis in 13 and 10 subjects, respectively. The subjects were all male, mean age was 59.7 (SD ± 2) years, body mass index (BMI) 36.5 (SD ± 1.8) kg/m². Baseline polysomnogram showed a total sleep time of 264 min (SD ±121.9), sleep efficiency of 80 % (SD ±10.9 %), latency to sleep onset of 14.3 min (SD ±15.77), wake after sleep onset of 48.8 min (SD ±34.7); with stage N1 of 11 % (SD ±6.5), N2 of 70 % (SD ±14.1), N3 of 11.8 (SD ±10.7) and REM of 11.8 (SD ±10.7). Respiratory parameters at baseline were AHI of 50 (SD ± 6) per hour of sleep, nadir SpO₂ of 77 (SD ± 3), and mean time below SpO₂ of 90 % of 59.37 min (SD 98.77). Hs-TnT at baseline was elevated and correlated with baseline nadir SpO₂ (SpO₂) (Correlation Coefficient of 0.889, *p* <0.001), mean time below SpO₂ of 90 % (Correlation of Coefficient of 0.832, *p* <0.001) but not with AHI or arousal index (Fig. 1). Subjects used CPAP for 165 ± 17 days and 5.3 ± 0.35 h/night. ESS decreased with CPAP therapy (14.6 vs 9.1, *p* <0.05). Systolic and diastolic blood pressures and heart rate remained unchanged throughout the study period. CPAP effectively reversed hypoxia in all subjects (nadir SpO₂ 77 ± 3 % at baseline and 89.3 ± 3 post CPAP, *p* = 0.005) (Table 1). Hs-TnT decreased only in subjects with baseline nadir SpO₂ of 80 % or lower (*n* = 4) (Fig. 2). The absolute and percent changes in Hs-TnT compared to baseline strongly correlated with baseline SpO₂ (Correlation Coefficients of 0.835 and 0.6419 respectively, both *p* <0.05). The absolute and percent

Table 1 Sleep and metabolic parameters before and after CPAP use

Pre and Post Treatment Data		
	Baseline	Post-CPAP
AHI (episodes/h)	50 ± 6	-
CPAP pressure (cm H ₂ O)	-	10 ± 3.2
CPAP use (days)	-	165 ± 17
CPAP use (Hrs/day)	-	5.3 ± 0.35
ESS	14.6 ± 1	9.5 ± 1*
Nadir SpO ₂ (%)	77 ± 3	89.3 ± 3*
Mean SpO ₂ (%)	93.2 ± 0.7	93.8 ± 0.62
Systolic blood pressure (mmHg)	124 ± 3	129 ± 4
Diastolic blood pressure (mmHg)	76 ± 2	76 ± 2
Heart rate (bpm)	77 ± 3	72 ± 3
Body weight (Kg)	108 ± 5.3	109.6 ± 5.4*
BUN (mg/dl)	14.1 ± 1.19	14.26 ± 1.41
Creatinine (mg/dl)	0.96 ± 0.09	1.0 ± 30.06
Hs-TnT (micrograms/l)	0.012 ± 0.0085	0.011 ± 0.0035*

AHI Apnea + hypopnea index, ESS Epworth sleepiness scale, Hs-TnT Highly sensitive troponin T

*: Significant differences compared to baseline (*p* ≤ 0.05) appear in bold

changes in Hs-TnT did not correlate with baseline AHI.

Discussion

Our study, consistent with others, suggests that Hs-TnT as a biomarker of myocardial stress/injury is elevated in patients with OSA [11]. In contrast, the increased Hs-TnT was associated with severity of hypoxia but not AHI or sleep interruption as measured by arousal index. The CPAP effectively reversed the pathophysiology in





Fig. 1 Baseline Hs-TnT correlates strongly with baseline nadir O₂ saturation. Y axis presents the nadir SpO₂ during the baseline sleep study



the study subjects but only was associated with decline in Hs-TnT in subjects with more severe hypoxia at baseline. Thus, the data suggests that cardio- and cerebrovascular adverse outcomes related to OSA and the benefits seen with CPAP therapy may greatest to those with pronounced hypoxia as assessed by nadir SpO2 versus other parameters used to assess sleep apnea. A major short coming of our study is the small sample size. While our results are provocative, they must be viewed with caution due to the limited sample size and should be further studied and characterized in larger studies.

Competing interests

Dr. C.M. Ballantyne and Dr. V. Nambi are co-investigators on a provisional patent filed along with Roche diagnostics for the use of biomarkers in the prediction of heart failure. Dr. C.M. Ballantyne has received research support from Roche diagnostics. Other authors do not report any conflict of interest.

Authors' contribution

AS designed the study, analyzed the data, conducted statistical analysis and prepared the manuscript. JK analysed the data and prepared the manuscript. MH designed the study and analysed the data and prepared the manuscript. HS designed the study and analysed the data and prepared the manuscript. SPS analyzed the data and prepared the manuscript. CMB analyzed the data and prepared the manuscript. BB analyzed the data and prepared the manuscript. VN analyzed the data and prepared the manuscript. All authors read and approved the final manuscript.

Acknowledgement

Roche diagnostics provided reagents and loan of an instrument to conduct the highly sensitive troponin T assay

Author details

¹Section of Pulmonary, Critical Care and Sleep Medicine, Medical Care Line, Michael E. DeBakey Veterans Affairs Medical Center, MED VA Medical Center, Bldg. 100 (111i), 2002 Holcombe Blvd, Houston, TX 77025, USA. ²Section of Cardiology, Michael E. DeBakey Veterans Affairs Medical Center, 2002 Holcombe Blvd, Houston, TX 77025, USA. ³Department of Medicine, One

Baylor Plaza, Houston, TX 77030, USA. ⁴University of Houston, Main campus, Houston, TX, USA. ⁵University of Saint Thomas, Houston, TX 77007, USA. ⁶Center for Cardiovascular Prevention, Houston Methodist DeBakey Heart and Vascular Center, Houston, TX, USA.

Received: 18 June 2015 Accepted: 3 October 2015

Published online: 16 October 2015

References

- Shivalkar B, Van de Heyning C, Kerremans M, Rinkevich D, Verbraecken J, De BW, et al. Obstructive sleep apnea syndrome: more insights on structural and functional cardiac alterations, and the effects of treatment with continuous positive airway pressure. *J Am Coll Cardiol.* 2006;47:1433–9.
- Lavie P, Lavie L. Cardiovascular morbidity and mortality in obstructive sleep apnea. *Curr Pharm Des.* 2008;14:3466–73.
- Fletcher EC. Sympathetic over activity in the etiology of hypertension of obstructive sleep apnea. *Sleep.* 2003;26:15–9.
- Lavie L. Oxidative stress in obstructive sleep apnea and intermittent hypoxia ΓÇô Revisited ΓÇô The bad ugly and good: Implications to the heart and brain. *Sleep Med Rev.* 2015;20:27–45.
- Neeland IJ, Drazner MH, Berry JD, Ayers CR, DeFilippi C, Seliger SL, et al. Biomarkers of chronic cardiac injury and hemodynamic stress identify a malignant phenotype of left ventricular hypertrophy in the general population. *J Am Coll Cardiol.* 2013;61:187–95.
- Saunders JT, Nambi V, de Lemos JA, Chambless LE, Virani SS, Boerwinkle E, et al. Cardiac troponin T measured by a highly sensitive assay predicts coronary heart disease, heart failure, and mortality in the Atherosclerosis risk in communities study. *Circulation.* 2011;123:1367–76.
- DeFilippi CR, De Lemos JA, Christenson RH. Association of serial measures of cardiac troponin t using a sensitive assay with incident heart failure and cardiovascular mortality in older adults. *JAMA.* 2010;304:2494–502.
- Glick D, DeFilippi CR, Christenson R, Gottdiener JS, Seliger SL. Long-term trajectory of two unique cardiac biomarkers and subsequent left ventricular structural pathology and risk of incident heart failure in community-dwelling older adults at low baseline risk. *JACC Heart Fail.* 2013;1:353–60.
- Randby A, Namtvedt SK, Einvik G, Hrubos-Strom H, Hagve TA, Somers VK, et al. Obstructive sleep apnea is associated with increased high-sensitivity cardiac troponin T levels. *Chest.* 2012;142:639–46.

10. Hall TS, Herrscher T, Jarolim P, Fagerland MW, Jensen T, Hallen J, et al. Obstructive sleep apnea: no independent association to troponins. *Sleep Breath*. 2014;18:351–8.
11. Roca GQ, Redline S, Punjabi N, Claggett B, Ballantyne CM, Solomon SD, et al. Sleep apnea is associated with subclinical myocardial injury in the community. The ARIC-SHHS study. *Am J Respir Crit Care Med*. 2013;188:1460–5.
12. Barcelo A, Esquinas C, Bauca JM, Pierola J, de la Pena M, Arque M, et al. Effect of CPAP treatment on plasma high sensitivity troponin levels in patients with obstructive sleep apnea. *Respir Med*. 2014;108:1060–3.
13. Agarwal SK, Avery CL, Ballantyne CM, Catellier D, Nambi V, Saunders J, et al. Sources of variability in measurements of cardiac troponin T in a community-based sample: the atherosclerosis risk in communities study. *Clin Chem*. 2011;57:891–7.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

