

# A new species of *Symplocos* (Symplocaceae) from southern Ecuador

Carmen Ulloa Ulloa<sup>1,2</sup>, Bertil Ståhl<sup>3</sup>, Danilo Minga<sup>4</sup>, Raffaella Ansaloni<sup>4</sup>

**1** Investigador Prometeo, Facultad de Ciencia y Tecnología, Universidad del Azuay, Cuenca, Ecuador **2** Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166, U.S.A. **3** Department of Organismal Biology, Uppsala University Campus Gotland, SE62167 Visby, Sweden **4** Facultad de Ciencia y Tecnología, Universidad del Azuay, Av. 24 de Mayo 7-77, Cuenca, Ecuador

Corresponding author: Carmen Ulloa Ulloa (carmen.ulloa@mobot.org)

---

Academic editor: Peter de Lange | Received 14 May 2015 | Accepted 4 September 2015 | Published 24 September 2015

**Citation:** Ulloa Ulloa C, Ståhl B, Minga D, Ansaloni R (2015) A new species of *Symplocos* (Symplocaceae) from southern Ecuador. *PhytoKeys* 56: 41–46. doi: 10.3897/phytokeys.56.5306

---

## Abstract

A new species from Ecuador, *Symplocos limonensis*, is here described and illustrated. It resembles *S. clethrifolia* but differs by having larger leaves with evident (i.e., not concealed) areoles on lower surface, sessile inflorescences, smaller white corollas, and fewer stamens. The species is only known from three collections in the Andean forests of Morona-Santiago Province in southern Ecuador.

## Resumen

Se describe e ilustra una nueva especie, *Symplocos limonensis*, de Ecuador. Se parece a *S. clethrifolia* pero se diferencia por tener las hojas más grandes, con las aréolas evidentes (no ocultas) en el envés, las inflorescencias sésiles y las flores más pequeñas, blancas y con menos estambres. La especie solo se conoce de tres ejemplares de los bosques andinos de la provincia de Morona-Santiago en el sur del país.

## Keywords

*Symplocos*, Ecuador, Andes

## Introduction

The genus *Symplocos* Jacq. is represented in Ecuador by ca. 33 species (Ståhl 2010a), the majority of which occur in Andean forests at 2500–3500 m elevation (Ståhl 1995). Thirteen of the species are country endemics (Barriga 2011) and have been described in the last 25 years, significantly increasing the knowledge of this still poorly known genus. In his treatment of the Symplocaceae for the Flora of Ecuador series, Ståhl (1991) described several new species of *Symplocos*, especially from southern Ecuador. One of these, *S. clethrifolia* Ståhl, was based on five collections, one of which (*Ulloa* 487) was indicated to deviate from the remaining four in floral and leaf features. This collection was provisionally treated under *S. clethrifolia* until more material could help elucidate its taxonomic status.

## Materials and methods

Recent examination of the *Symplocos* collections in Ecuadorian herbaria revealed additional gatherings from the same general area that have led us to conclude that the *Ulloa* collection, along with some new ones, belong to a previously undescribed species, which is described here. Acronyms of the herbaria follow Thiers (2015).

## Description of the new species

### *Symplocos limonensis* Ståhl, C. Ulloa & Minga, sp. nov.


urn:lsid:ipni.org:names:77150152-1

Fig. 1

**Diagnosis.** *Symplocos limonensis* differs from *S. clethrifolia* Ståhl by having larger leaves (up to 13.5 vs. 9.5 cm) with evident (i.e., not concealed) venation areoles abaxially (vs. densely ferruginous-velutinous with matted hairs concealing areoles), sessile inflorescences (vs. pedunculate), somewhat smaller, white (vs. red to pink) corollas, and fewer stamens (40–50 vs. 60–70).

Type: ECUADOR. Morona-Santiago: road Cuenca-Macas (road Gualaceo-Limón East of the pass), 2700–3400 m, 19 Aug 1987, C. *Ulloa* 487 (Holotype: QCA-209874!; isotypes AAU!, GB!, MO-6500819!).

Tree or treelet to 12 m tall; branchlets reddish brown, sparsely villous to glabrescent, apical buds densely long-villous. Leaves alternate, petiolate; blade oblong to widely elliptic, 8–13.5 × 4–9 cm, the blades exceptionally almost circular and ca. 3 × 3 cm, coriaceous, light green and long-villous beneath, olive green and glabrous above, base rounded or truncate, slightly decurrent on petiole, apex rounded to widely obtuse, mucronulate at the tip, margins denticulate, teeth 3–4 mm apart, lateral veins 8–10 per side, midvein, lateral veins, and veinlets on lower side prominently raised and



**Figure 1.** *Symplocos limonensis*. **A** Flowering branch **B** Detail of venation on leaf lower surface **C** Corolla and stamens whorls **D** Young flower in longitudinal section. (**A**, **B** from Jørgensen 92876; **C**, **D** from Ulloa 487). Illustration by A.L. Arbeláez.

verrucose, impressed above; petiole 1–1.5 cm long, rounded and long-villous beneath, flattened and glabrous above. Inflorescences fasciculate, sessile, borne in axils of extant leaves and along branchlets beneath the foliage; flowers 8–10 per inflorescence; bracts 4, very broadly ovate, 2–3 × 2–2.5 mm, strigulose on midvein toward apex, otherwise glabrous, margins ciliate. Flowers with the calyx synsepalous, tube ca. 1.5 mm long, lobes 5, very broadly ovate, 1.5–2.5 × 2.2–2.8 mm, margins ciliate; corolla sympetalous, glabrous, white, 5–6 mm long, lobes 5 to 7, broadly oblong, 1.8–4 mm

wide, margins entire; stamens 40–45 in 3(4) rows, filaments smooth, fused for ca. 2 mm to the corolla tube, those of the inner whorl fused most of their length leaving ca. 0.5 mm of the filaments free, stamens of the outer whorls with free filaments 1–2.8 mm long, more or less flattened and constricted at apex, anthers ca. 0.5 mm long; disc intrastaminal, dome-shaped, densely villous; style ca. 2.5 mm long, glabrous; stigma capitate and irregularly 3-lobed; ovary inferior, 3-locular with 3 or 4 ovules per locule. Mature fruits not seen.

**Specimens examined (Paratypes).** ECUADOR. **Morona-Santiago:** road Gualaceo-Limón, km 33.3, 03°02'S, 78°35'W, 3010 m, 27 Dec 1990 (fl), *P.M. Jørgensen, C. Ulloa & B. Øllgaard* 92876 (AAU!, QCA!, QCNE!); Collay, Maylas, on road to Limón 03°00.36'S, 78°38.00'W, 3150 m, 14 Jun 2000 (bud), *F. Serrano, D. Minga & A. Verdugo* 1459 (HA!).

**Etymology.** The name of the species refers to the town of Limón, officially known as General Leonidas Plaza Gutiérrez. The road leading from the Andean town of Gualaceo to Limón, situated at 1400 m altitude on the east Andean slopes, has long been considered an important locality for botanical exploration where many new species have been found.

**Distribution and conservation status.** The species is known solely from three collections made in disturbed upper Andean forests and scrub páramo on the highest point of the Gualaceo-Limón-Macas road in Morona-Santiago Province of southern Ecuador. The area of occupancy (AOO) of the species is less than 20 km<sup>2</sup> and falls within the “Área de Conservación Municipal Tinajillas-Río Gualaceño,” a locally managed reserve; however, the category of protection is of lesser status than that of a National Park. Since 2000, we have searched for additional plants but could not locate any, finding only one population of *Symplocos quitensis* Brand (1901: 76), a very different species with a wide Andean distribution. The road was poorly maintained and treacherous until recently, but has now been considerably widened in the process of being paved, and consequently the natural vegetation alongside is heavily destroyed. Ongoing wood extraction for charcoal production, expansion of areas under cultivation, and mining activities for clay, gravel, and metals are threats to the natural habitats in this region. Given current knowledge, we assign a provisional IUCN conservation status of Endangered (IUCN 2014) to this species.

**Discussion.** *Symplocos limonensis* resembles *S. clethrifolia* Ståhl and *S. golondrinae* Ståhl by having leaves with conspicuous venation, the veins of the lower side being prominent and verrucose and those on the upper side impressed. It differs from *S. clethrifolia*, which also is restricted to southern Ecuador (but from other localities), by having larger leaves (up to 13.5 vs. 9.5 cm) with evident (i.e., not concealed) venation areoles abaxially (vs. densely ferruginous-velutinous with matted hairs concealing areoles), sessile inflorescences (vs. pedunculate), smaller (5–6 vs. c. 8 mm long), white (vs. red to pink) corollas, and fewer stamens (40–50 vs. 60–70). From *S. golondrinae* it differs by the larger size of leaves and flowers, with the leaves being coriaceous (vs. cartilaginous), long-villous abaxially (vs. sparsely strigose) and having longer (1–1.5 vs.

< 1 cm) petioles, and by having a densely villous flower disk (vs. glabrous). Moreover, *S. golondrinae* occurs in northwestern Ecuador, on the opposite side of the Andes.

*Symplocos quitensis* has been collected in the same area of the new species, but it is readily distinguished from both *S. limonensis* and *S. clethrifolia* by the densely hispid branchlets, smaller (up to 6.5 × 4 cm), membranaceous leaves, and pinkish flowers borne in short racemes.

In the most recent key to Andean *Symplocos* (Ståhl 2010a), *S. limonensis* keys to *S. clethrifolia* (albeit with the inflorescence pedunculate) or to *S. robusta* Ståhl. The latter species is known only from Bolivia and differs from *S. limonensis* in, e.g., its larger (to 14.5 × 8.5 cm), longer-petiolate (to 2 cm) leaves, more numerous flowers (up to 20) per inflorescence, and strigose corollas (vs. glabrous).

*Symplocos limonensis* has flowers with 5 to 7 petals, but being notoriously unstable in many species of *Symplocos*, and often not studied, the number of petals may show to be of little taxonomic significance.

In the key to infrageneric taxa of the genus (Fritsch et al. 2008), *Symplocos limonensis* falls into the tropical American clade *Symplocos* subg. *Symplocos* sect. *Symplocos* having exerted, monodelphous stamens adnate to the corolla beyond its base, and the filaments flattened and constricted apically.

## Acknowledgments

We thank the Ministerio del Ambiente del Ecuador for research permits 059-IC-DFP, 003-15 IC-FLO-DNB/MA. Ulloa's research at Universidad del Azuay was supported with a Prometeo fellowship from Secretaría de Educación Superior, Ciencia, Tecnología e Innovación de la República del Ecuador; she thanks the authorities at the Universidad del Azuay for support and facilities during her stay. Field work was supported in part by the National Geographic Society (6327-98 to P. Jørgensen and Ulloa) and by C. Davidson. We thank the curators at AAU, GB, HA, MO, QCA, and QCNE for access to their collections, B. Bassüner for the analysis of conservation status, A.L. Arbeláez for the illustration, and M. Conger for language editing. We thank P. Fritsch for his valuable comments on the manuscript.

## References

- Barriga P (2011) Symplocaceae. In: León-Yáñez S, Valencia R, Pitman N, Endara L, Ulloa Ulloa C, Navarrete H (Eds) Libro Rojo de las plantas endémicas del Ecuador, 2da. Edición. Publicaciones del Herbario QCA, Pontificia Universidad Católica del Ecuador, Quito, 795–797.
- Brand A (1901) IV. 242. Symplocaceae. In: Engler A von (Ed.) Das Pflanzenreich regni vegetabilis conspectus. Heft 6. W. Engelmann, Leipzig, 100 pp.

- Fritsch P, Kelly LM, Wang Y, Almeda F, Kriebel R (2008) Revised infrafamilial classification of Symplocaceae based on phylogenetic data from DNA sequences and morphology. *Taxon* 57: 823–852.
- Jacquin NJ von (1760) *Symplocos*. Enumeratio Systematica Plantarum. Theodorum Haak, Lugduni Batavorum [Leiden]. 5 pp.
- IUCN (2014) Guidelines for Using the IUCN Red List Categories and Criteria. Version 11. Prepared by the Standards and Petitions Subcommittee. <http://jr.iucnredlist.org/documents/RedListGuidelines.pdf> [accessed: 27 Feb 2015]
- Ståhl B (1991) Symplocaceae. In: Harling G, Andersson L (Eds) *Flora of Ecuador* 43. University of Göteborg, 44 pp.
- Ståhl B (1994) The genus *Symplocos* (Symplocaceae) in Bolivia. *Candollea* 49: 369–388.
- Ståhl B (1995) New or noteworthy Andean species of the genus *Symplocos* (Symplocaceae). *Candollea* 50: 445–452.
- Ståhl B (2010a) Four new species and records of *Symplocos* (Symplocaceae) from Peru and Bolivia, and a key to all species of *Symplocos* known to occur in Ecuador, Peru and Bolivia. *Nordic Journal of Botany* 28: 79–87. doi: 10.1111/j.1756-1051.2009.00445.x
- Ståhl B (2010b) Additions to the knowledge of the genus *Symplocos* (Symplocaceae) in Ecuador and Peru. *Novon* 20: 84–94. doi: 10.3417/2008079
- Thiers B (2015) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [accessed: 27 Feb 2015]