

Variability in splanchnic tissue oxygenation during preterm red blood cell transfusion given for symptomatic anaemia may reveal a potential mechanism of transfusion-related acute gut injury

Sean M. Bailey¹, Karen D. Hendricks-Muñoz², Pradeep V. Mally¹

¹Department of Pediatrics, Division of Neonatology, New York University School of Medicine, New York, NY;

²Department of Pediatrics, Division of Neonatal Medicine, Virginia Commonwealth University School of Medicine, Richmond, VA, United States of America

Background. There is increasing evidence indicating an association between red blood cell (RBC) transfusions and necrotising enterocolitis (NEC) in preterm infants, especially late-onset NEC. This phenomenon is referred to as transfusion-related acute gut injury (TRAGI). One theory as to a pathophysiological mechanism is that transfusion may result in an ischemia-reperfusion injury to intestinal tissue. We tested the hypothesis that there is significantly greater variability during transfusion in splanchnic tissue oxygen saturation (SrSO₂) than in cerebral tissue oxygen saturation (CrSO₂).

Materials and methods. This was a prospective, observational study using near-infrared spectroscopy to monitor SrSO₂ and CrSO₂ in preterm neonates undergoing RBC transfusion for symptomatic anaemia. Mean, standard deviation, highest and lowest SrSO₂ and CrSO₂ values during each transfusion were determined. The greatest difference in SrSO₂ and CrSO₂ during each transfusion was calculated, along with the coefficient of variation.

Results. We studied 37 subjects. Throughout all transfusions, the mean SrSO₂ was 45.6% ±13.8 and the mean CrSO₂ was 65.4% ±6.9 (p<0.001). The variability of SrSO₂ was significantly greater than that of CrSO₂. Averaging data from all subjects, the greatest difference in SrSO₂ was 43.8% ±13.4 compared with 23.3% ±7.6 for CrSO₂ (p<0.001). The mean coefficient of variation in all transfusions was 20.5% for SrSO₂ and 6.0% for CrSO₂ (p<0.001). Increasing post-conceptual age did not affect SrSO₂ variability (R²=0.022; p=0.379), whereas CrSO₂ variability during transfusion decreased with increasing post-conceptual age (R²=0.209; p=0.004).

Discussion. In preterm infants, there is a large degree of tissue oxygenation variability in splanchnic tissue during RBC transfusion and this does not change with increasing maturity. We speculate that these findings, combined with lower average tissue oxygenation, may demonstrate susceptibility of the preterm gut to TRAGI.

Keywords: blood transfusion, necrotising enterocolitis, near-infrared spectroscopy.

Introduction

Red blood cell (RBC) transfusion remains a mainstay in the medical management of premature infants¹. A majority of very-low birth weight infants will receive at least one RBC transfusion during their time in the neonatal intensive care unit (NICU)². Necrotising enterocolitis (NEC) is a common gastrointestinal disease affecting preterm infants in the NICU and is associated with significant morbidity and mortality³. Its exact pathogenesis is unknown, but NEC is believed to be a multifactorial process that probably includes intestinal ischaemic injury combined with an infectious process and/or the activation of an inflammatory cascade⁴. Premature infants are especially vulnerable to this condition, because, in addition to having decreased gut barrier function, an immature immune system, and

decreased gut motility, they also have poor circulatory regulation of their splanchnic organ system⁵.

Near-infrared spectroscopy (NIRS) is a well-described technique that can be used to determine both the regional splanchnic tissue oxygen saturation (SrSO₂) and the regional cerebral tissue oxygen saturation (CrSO₂) of a preterm infant⁶. Not only do these values provide information about the flow of blood to the intestines and brain, they also reflect the adequacy of the balance between tissue perfusion and oxygenation vs metabolic demand⁷.

Until recently, blood transfusions and NEC were thought to be unrelated. However, there is now growing evidence demonstrating a possible association between RBC transfusions and NEC⁸. A particular concern is developing regarding anaemic, yet stable, "feeding and

growing" premature infants who develop late-onset NEC within days after receiving a transfusion^{8,9}. This transfusion-related NEC that may occur in preterm infants has been termed transfusion-related acute gut injury (TRAGI), and several theories exist as to how this can occur. One explanation is that the splanchnic circulation is less capable of vascular regulation than other organ systems, such as the cerebral circulatory system, and an immature splanchnic circulation has especially poor vascular control capabilities^{5,10}. Some, therefore, speculate that RBC transfusions can lead to such variability in perfusion and oxygen delivery to splanchnic tissue, which is already maintained in a relatively low blood flow environment to begin with, as to result in some degree of ischaemic injury⁹. This, in turn, could create a situation in which the gut becomes more prone to developing TRAGI.

In order to examine this as a possible mechanism for RBC transfusions increasing an infant's risk of NEC, we set out to test the hypothesis that not only is mean SrSO₂ lower than mean CrSO₂ during preterm infant RBC transfusions, but that there is also greater variability in SrSO₂ during the transfusion than in CrSO₂. Our primary goal was to account for the extremes of tissue oxygenation (both high and low) that intestinal tissue may experience during a RBC transfusion, as we theorised that this could be an important factor related to TRAGI. We also sought to determine whether this finding would be seen consistently throughout the neonatal period, as TRAGI is thought to occur even in older preterm infants who generally do not manifest NEC.

Materials and methods

Patients

Preterm neonates in the NICU, who were at least 5 days old and born at New York University Langone Medical Center and Bellevue Hospital Center (New York, NY, USA), were eligible for the study. The exclusion criteria were a 5-minute Apgar score of 3 or less, a known chromosomal abnormality, a congenital cardiac malformation, grade 2 or greater intraventricular haemorrhage, a current diagnosis of NEC prior to enrolment, or a need for either vasopressor support or high frequency ventilation. This study received ethical approval by the hospital institutional review board and parental consent was obtained prior to any subject's participation.

Study design

To monitor SrSO₂ and CrSO₂ we used the INVOS 5100C Cerebral/Somatic Oximeter (Somanetics, Troy, MI; now Covidien, Mansfield, MA, USA). This NIRS device uses light emitted from a skin sensor in the near-infrared wavelength (730 nm and 810 nm) to

penetrate both soft tissue and bone in order to determine the amount of oxygenated haemoglobin (oxy-Hb) and deoxygenated haemoglobin (deoxy-Hb) in the organ tissue below. The result is displayed on the monitor as a percentage [$\text{oxy-Hb}/(\text{oxy-Hb} + \text{deoxy-Hb})$] which represents the regional tissue oxygen saturation (rSO₂) of that particular tissue. This technology has been demonstrated to be capable of measuring both SrSO₂ and CrSO₂ accurately in preterm neonates⁶.

Investigators were contacted when a transfusion was ordered by the primary NICU team because a study eligible infant had developed symptomatic anaemia. NIRS sensors were then placed on the subject if parental consent was obtained and monitoring could begin prior to starting the transfusion so that adequate baseline rSO₂ values could be established. SrSO₂ data were collected from an abdominal NIRS sensor placed just left of the umbilicus and CrSO₂ data were collected from a forehead NIRS sensor. The INVOS display screen was covered to blind the NICU staff to the information collected. Single SrSO₂ and CrSO₂ data points were measured and recorded every 30 seconds. These data were then electronically stored on a hard drive for later analysis.

All monitored RBC transfusions were ordered to be 15 mL/kg with a goal transfusion time of 4 hours. All patients received either directed donor or random donor leucocyte-reduced, irradiated, Cytomegalovirus-negative blood. Feeds were withheld only during the transfusion period, as is the standard practice in our NICU. Feeds were resumed as soon as transfusions had been completed.

Data analysis

The mean SrSO₂ and CrSO₂ during the complete RBC transfusion period were calculated for each individual subject. In addition, the highest and lowest SrSO₂ and CrSO₂ values recorded from the exact time that the transfusion started until the time that the transfusion was completed were also determined for each subject. From these highest and lowest values, the range of SrSO₂ and CrSO₂ that patients were exposed to was established, and represents the greatest variability in oxygenation that each subject's gut and brain experienced during the RBC transfusion. As an additional marker of general variability found in tissue oxygenation during a transfusion, a coefficient of variation analysis was also conducted examining both SrSO₂ and CrSO₂ for each transfusion. The coefficient of variation was calculated using the ratio: standard deviation divided by the mean. We then multiplied this value by 100 to obtain a percentage [$\text{coefficient of variation} = (\text{SD}/\text{mean}) \times 100$]. This calculation was done for SrSO₂ and CrSO₂ using all data points measured by

NIRS during each individual transfusion. This has been found useful when comparing NIRS signal variability between two separate site measurements that may have very different mean values¹¹.

The Student's *t* test was used to examine for a difference in the mean SrSO₂ compared with the mean CrSO₂ that subjects experienced during their transfusion. This test was also used to determine any difference in the amount of variability in SrSO₂ vs CrSO₂ during our subjects' monitored transfusions, both looking at the greatest degree of variability as well as the coefficient of variation. The Mann-Whitney test was used to compare median SrSO₂ and CrSO₂ values. Pearson's correlation analysis was used to determine the relationship between post-conceptual age at the time of transfusion and the greatest variability in SrSO₂ and CrSO₂ occurring throughout the transfusion. All statistical calculations were performed using SPSS 17.0 Software for Windows (SPSS Inc., Chicago, IL, USA). Two-tailed tests were used to determine statistical significance, which was established by a p-value of less than 0.05.

Results

Patients' characteristics

Thirty-seven patients were included in this study. Their demographic information is displayed in Table I. Our patients had a mean gestational age of 28.4 weeks and, at the time of their monitored transfusion, had a mean post-conceptual age of 32.7 weeks. The respiratory support needs of our study population were varied. Most of the infants were receiving feeds and 59% of all subjects were being treated with caffeine.

Blood transfusions

The mean Hb value for which our subjects received a transfusion was 9.3 g/dL, as shown in Table I. Transfusions were given to subjects because of apnoea, bradycardia, or desaturation events (22/37), tachycardia (2/37), increased respiratory support requirements (10/37) or feeding intolerance (3/37), in addition to a low Hb level. The mean transfusion volume and transfusion period remained close to the targeted 15 mL/kg over 4 hours.

All transfusions were tolerated well and there were no reports of transfusion reaction. No patients had a major change in their cardiac or ventilatory status during their transfusion. In addition, there were no reported cases of NEC within 1 week after each transfusion having taken place.

Splanchnic and cerebral tissue oxygen saturation during transfusion

The mean SrSO₂ throughout all monitored transfusions was found to be significantly less than the mean CrSO₂ (mean±SD: 45.6% ±13.8 vs 65.4%

Table I - Subjects' demographics and clinical characteristics (n=37).

Gender	
Male	65% (24/37)
Female	35% (13/37)
Twin gestation	
	19% (7/37)
Ethnicity	
Caucasian	43% (16/37)
African American	14% (5/37)
Hispanic	35% (13/37)
Asian	8% (3/37)
Measurements at birth	
<i>Gestational age (weeks)</i>	
Mean±SEM	28.4±0.48
Median; IQR	27 ^{0/7} ; 26 ^{0/7} -30 ^{0/7}
<i>Birth weight (g)</i>	
Mean±SEM	1,112±69.4
Median; IQR	1,000; 805-1,202
Measurements at transfusion	
<i>Post-conceptual age at transfusion (weeks)</i>	
Mean±SEM	32.7±0.58
Median; IQR	32 ^{6/7} ; 30 ^{1/7} -35 ^{3/7}
<i>Mean weight at transfusion (g)</i>	
Mean±SEM	1,403±78.2
Median; IQR	1,370; 1,045-1,780
Ventilation support status at transfusion	
Room air	38% (14/37)
Nasal cannula	30% (11/37)
Nasal continuous positive airway pressure	16% (6/37)
Conventional ventilation	16% (6/37)
Inspired fraction of oxygen, mean±SD	0.26±0.06
Feeds and caffeine at transfusion	
Receiving enteral feeds prior to transfusion	78% (29/37)
Volume of feeds if receiving (mL/kg/day), mean±SD	114±53.3
Receiving caffeine at time of transfusion	59% (22/37)
Transfusion information	
Pre-transfusion Hb level (g/dL), mean±SD	9.3±1.2
Post-transfusion Hb level (g/dL), mean±SD	13.0±3.3
Volume of transfusion (mL/kg), mean±SD	15.3±1.4
Transfusion duration (h), mean±SD	3.8±0.5

SEM: standard error of the mean; IQR: interquartile range; SD: standard deviation; Hb: haemoglobin.

±6.9; p<0.001). In every case, but one, the mean SrSO₂ throughout the transfusion period was always lower than the mean CrSO₂ that each individual subject experienced. In the one exception, the mean SrSO₂ was 57.8% and the mean CrSO₂ was 56.7%. The median SrSO₂ during transfusions was 49.5% compared to 66.0% for CrSO₂ (p<0.01).

Variability in splanchnic and cerebral tissue oxygen saturation during transfusion

Figure 1 displays the highest and lowest SrSO₂ and CrSO₂ values recorded during each transfusion for every subject who was monitored. It is apparent that, during the transfusion period, there is a much wider range of SrSO₂ values than CrSO₂ values. From these data points, the greatest variability that each patient experienced in SrSO₂ and CrSO₂ was determined. These values are shown in Figure 2, demonstrating a significantly greater variability in SrSO₂ than in CrSO₂ during the transfusion periods (mean±SD: 43.8% ±13.4 vs 23.3% ±7.6; p<0.001).

The average coefficient of variation for SrSO₂ in all individual transfusions was 20.5%, whereas that for CrSO₂ was 6.0%. When comparing these means, NIRS SrSO₂ monitoring had more than three times greater variability than NIRS CrSO₂ monitoring (p<0.001). In addition, the mean standard deviation in SrSO₂ monitoring for individual subjects was 8.8% compared with 3.9% for CrSO₂ (p<0.001).

Variability in tissue oxygen saturation based on post-conceptual age

There was no decrease in the maximum amount of SrSO₂ variability that each subject experienced during their blood transfusion as they increased in post-conceptual age (p=0.379) (Figure 3). However, a significant inverse correlation was found between increasing post-conceptual age at transfusion and the greatest degree of variability in CrSO₂ during that blood transfusion (p=0.004). Increasingly mature subjects were more likely to experience lesser variability in brain tissue oxygenation.

Discussion

We found that splanchnic tissue oxygenation was significantly lower and had significantly greater variability during a preterm infant RBC transfusion

when compared to tissue oxygenation of the brain. We also discovered that increasing maturity had no effect on the degree of splanchnic tissue oxygenation control during a transfusion, which was in contrast to the increasingly tightly controlled auto-regulation of cerebral tissue oxygen levels. These results support the theoretical possibility that RBC transfusions could lead to TRAGI secondary to intestinal mucosal damage from a combination of both hypoxia and ischaemia-induced gut injury, and that this could occur throughout the preterm neonatal period.

We decided to compare relative SrSO₂ values to CrSO₂ values for two reasons. To begin with, blood flow to the brain is considered relatively well controlled and consistent in delivering a high amount of constant oxygen to cerebral tissue^{12,13}. In addition, to our knowledge, there is currently only limited published data suggesting a risk to the preterm brain resulting directly from a RBC transfusion and many feel that RBC transfusion is likely to be neuroprotective¹⁴. Therefore, it seemed appropriate to use CrSO₂ as a control to compare both SrSO₂ levels and variability. We chose to focus on the highest and lowest points in rSO₂ during a transfusion for our primary analysis because we theorised that it may be the extremes of tissue perfusion status that make intestinal tissue most vulnerable to perfusion-ischaemia-reperfusion injury, which has been thought of as a potential factor that can play a role in the pathogenesis of NEC.

Conditions that lead to intestinal tissue hypoxia have been shown to be risk factors for developing NEC¹⁵. Examples include maternal-foetal placental insufficiency, umbilical vessel catheterisation occluding mesenteric blood flow, and congenital cardiac disease¹⁶. In fact, NEC is one of the primary complications of cardiac lesions associated with lower baseline arterial oxygen saturation¹⁷.

In addition to simply low tissue oxygen levels, a change from a low blood flow state back to a high blood




Figure 1 - Graphical representation of the highest and lowest SrSO₂ and CrSO₂ values that each subject experienced during their RBC transfusion (n=37).




Figure 2 - Box plot demonstrating the mean and interquartile range for the amount of variability in SrSO₂ and CrSO₂ during all RBC transfusions (n=37).

flow state has been shown in animal models to cause intestinal injury that could theoretically lead to NEC¹⁸. Several different biochemical reactions that develop as a result of ischaemia followed by reperfusion have been demonstrated as possible pathways to injury. These include an increased production of reactive oxygen free radical species and the accumulation of potentially toxic metabolites¹⁹. The substantial changes in splanchnic blood flow that we demonstrated could potentially lead to these mechanisms of injury.

NIRS appears capable of monitoring for both low oxygen states and changes in blood flow, which are both potentially responsible for TRAGI. NIRS values provide information about the adequacy of tissue oxygen supply which is primarily based on the balance between organ blood flow vs the metabolic demands of that organ⁷. If a patient remains relatively stable without rapid changes in metabolic demands, then rSO₂ provides a good representation of tissue perfusion. There have been case reports of patients with cardiac lesions with low SrSO₂ values monitored with NIRS who later developed NEC¹⁷. Moreover, Fortune *et al.* demonstrated that low SrSO₂

measured with NIRS in preterm infants were strongly associated with NEC²⁰.

Ultrasound with Doppler analysis has been able to monitor the altered mesenteric blood flow velocity response in the major vessels that can occur immediately following a blood transfusion²¹. While the NIRS method that we employed does not examine the bulk flow of blood, it may provide more information about changes in blood flow, tissue perfusion, and oxygen delivery occurring in the microcirculation, which ultrasound cannot provide. In this regard, NIRS may be more advantageous than ultrasound for predicting which infants may be at risk of subsequently developing TRAGI.

There were limitations to this study. First, the lowest rSO₂ value that the commercially available NIRS device we used can detect is 15%. If lower values could have been detected, we might have seen even greater variability in SrSO₂. Secondly, to assess the amount of change in intestinal and cerebral tissue perfusion and oxygenation that occurred, our primary analysis was based on highest and lowest SrSO₂ and CrSO₂ values during the transfusion period. Although we feel this provides a good assessment of the degree of variability in tissue oxygenation, it does not describe well acceleration or deceleration of tissue oxygenation, which could play a role in transfusion-related gut injury. The analysis showing that SrSO₂ had a much higher coefficient of variation than CrSO₂ does, however, offer more insight into this as it also showed the overall relative variability in gut tissue oxygenation was much greater.

In addition, no patients monitored actually developed TRAGI (NEC in the post-transfusion period). It would have been helpful to determine whether there was either a lower baseline SrSO₂ or an increased amount of variability in SrSO₂ during transfusion in subjects who developed TRAGI. However, in this pilot study it would have been unlikely, as the incidence of transfusion-related NEC is reported to be very low⁸.

Even if changes in splanchnic tissue oxygenation do not represent the primary mechanism through which




Figure 3 - Correlation between post-conceptual age and the variability in tissue oxygen saturation experienced during a RBC transfusion (n=37).

transfusion-related NEC can develop, this study still provides insight into the vascular regulation that occurs in preterm neonates. The degree of variability in CrSO₂ was much less than that in SrSO₂, demonstrating the auto-regulatory capabilities of the cerebral circulatory system. The decreasing variability in CrSO₂ associated with increasing post-conceptual age shows that this mechanism continues to develop throughout the third trimester. This is likely in this district, in contrast to the blood vessels supplying the splanchnic circulation.

Conclusion

While further studies are needed to define the incidence and possible causes of TRAGI more clearly, the results from this study demonstrate that variability in intestinal tissue oxygen levels remains a potential pathophysiological mechanism for this condition. Larger studies performed in a prospective manner are needed to validate these results in order to establish whether SrSO₂ variability predisposes the intestines to develop TRAGI in the post-transfusion period.

Acknowledgements

This study received support from a KiDS of NYU research grant and the NYU School of Medicine Division of Neonatology. We thank Dr. John Wells for his prior intellectual support. We are also thankful to all of our patients and their parents, as well as to our nursing staff at both NYU Langone Medical Center and Bellevue Hospital.

Authorship contributions

SMB wrote the initial draft of the manuscript and contributed to the development and design of the project. In addition, he recruited subjects, performed the NIRS monitoring, and collected and analysed data. KDH-M contributed to the development of the project and assisted in writing the manuscript. PVM contributed to the development and design of the project and assisted in writing the final version of the manuscript.

Disclosure of conflicts of interest

None of the Authors has a conflict of interest to report pertaining to this manuscript. The Neonatal Intensive Care Units at NYU Langone Medical Center and Bellevue Hospital were previously Somanetics Neonatal NIRS sensor beta-test sites.

References

- Ohls RK. Transfusions in the preterm infant. *NeoReviews* 2007; **8**: e377-86.
- Bell EF, Strauss RG, Widness JA, et al. Randomized trial of liberal vs restrictive guidelines for red blood cell transfusion in preterm infants. *Pediatrics* 2005; **115**: 1685-91.
- Holman RC, Stoll BJ, Clarke MJ, Glass RI. The epidemiology of necrotizing enterocolitis infant mortality in the United States. *Am J Public Health* 1997; **87**: 2026-31.
- Lin PW, Stoll BJ. Necrotizing enterocolitis. *Lancet* 2006; **368**: 1271-83.
- Schnabl, KL, Van Aerde JE, Thomson, AB, Clandinin MT. Necrotizing enterocolitis: a multifactorial disease with no cure. *World J Gastroenterol* 2008; **14**: 2142-61.
- Dave V, Brion LP, Campbell DE, et al. Splanchnic tissue oxygenation, but not brain tissue oxygenation, increases after feeds in stable feeds in stable preterm neonates tolerating full bolus orogastric feeding. *J Perinatol* 2009; **29**: 213-8.
- Wolfberg AJ, Plessis AJ. Near-infrared spectroscopy in the fetus and neonate. *Clin Perinatol* 2006; **33**: 707-28.
- Christensen RD, Lambert DK, Henry E, et al. Is "transfusion-associated necrotizing enterocolitis" an authentic pathogenic entity? *Transfusion* 2010; **50**: 1106-12.
- Blau J, Calo JM, Dozor D, et al. Transfusion-related acute gut injury: necrotizing enterocolitis in very low birth weight neonates after packed red blood cell transfusion. *J Pediatr* 2011; **158**: 403-9.
- Nankervis CA, Giannone PJ, Reber KM. The neonatal intestinal vasculature: contributing factors to necrotizing enterocolitis. *Semin Perinatol* 2008; **32**: 83-91.
- Mintzer JP, Parvez B, Chelala M, et al. Quiescent variability of cerebral, renal, and splanchnic regional tissue oxygenation in very low birth weight neonates. *J Neonatal Perinatal Med* 2014; **7**: 199-206.
- Liem KD, Greisen G. Monitoring of cerebral haemodynamics in newborn infants. *Early Hum Dev* 2010; **86**: 155-8.
- Greisen G. Cerebral blood flow and energy metabolism in the newborn. *Clin Perinatol* 1997; **24**: 531-46.
- Andersen CC, Collins CL. Poor circulation, early brain injury, and the potential role of red cell transfusion in premature newborns. *Pediatrics* 2006; **117**: 1464-6.
- Vanderhoof JA, Zach TL, Adrian TE. Gastrointestinal disease. In: MacDonald MG, Mullet MD, Seshia MK (editors): *Avery's Neonatology: Pathophysiology and management of the newborn*. 6th edition. Philadelphia: Lippincott Williams and Wilkins; 2005. pp. 949-50.
- Stevenson DK, Blakely ML. Historical perspective: necrotizing enterocolitis: an inherited or acquired condition? *NeoReviews* 2006; **7**: e125-34.
- McElhinney DB, Hedrick HL, Bush DM, et al. Necrotizing enterocolitis in neonates with congenital heart disease: risk factors and outcomes. *Pediatrics* 2000; **106**: 1080-7.
- Papparella A. Ischemia-reperfusion injury in the intestines of newborn pigs. *Pediatr Res* 1997; **42**: 180-8.
- Hammerman C, Goldschmidt D, Caplan MS, et al. Amelioration of ischemia-reperfusion injury in rat intestine by pentoxifylline-mediated inhibition of xanthine oxidase. *J Pediatr Gastroenterol Nutr* 1999; **29**: 69-74.
- Fortune PM, Wagstaff M, Petros AJ. Cerebro-splanchnic oxygenation ratio (SCOR) using near infrared spectroscopy may be able to predict splanchnic ischaemia in neonates. *Intensive Care Med* 2001; **27**: 1401-7.
- Krimmel GA, Baker R, Yanowitz TD. Blood transfusion alters the superior mesenteric artery blood flow velocity response to feeding in premature infants. *Am J Perinatol* 2009; **26**: 99-106.

Arrived: 22 August 2014 - Revision accepted: 16 December 2014

Correspondence: Sean M. Bailey
Department of Pediatrics - Division of Neonatology
New York University School of Medicine
NYU/Bellevue Hospital Center
462 First Ave, NBV Suite 8S15
New York, NY 10016, USA
e-mail: sean.bailey@nyumc.org