

OPEN

Fructose Malabsorption in Systemic Sclerosis

Isabelle Marie, MD, PhD, Anne-Marie Leroi, MD, PhD, Guillaume Gourcerol, MD, PhD, Hervé Levesque, MD, PhD, Jean-François Ménard, MD, PhD, and Philippe Ducrotte, MD, PhD

Abstract: The deleterious effect of fructose, which is increasingly incorporated in many beverages, dairy products, and processed foods, has been described; fructose malabsorption has thus been reported in up to 2.4% of healthy subjects, leading to digestive clinical symptoms (eg, pain, distension, diarrhea). Because digestive involvement is frequent in patients with systemic sclerosis (SSc), we hypothesized that fructose malabsorption could be responsible for intestinal manifestations in these patients.

The aims of this prospective study were to: determine the prevalence of fructose malabsorption, in SSc; predict which SSc patients are at risk of developing fructose malabsorption; and assess the outcome of digestive symptoms in SSc patients after initiation of standardized low-fructose diet.

Eighty consecutive patients with SSc underwent fructose breath test. All SSc patients also completed a questionnaire on digestive symptoms, and a global symptom score (GSS) was calculated.

The prevalence of fructose malabsorption was as high as 40% in SSc patients. We also observed a marked correlation between the presence of fructose malabsorption and: higher values of GSS score of digestive symptoms ($P=0.000004$); and absence of delayed gastric emptying ($P=0.007$). Furthermore, in SSc patients with fructose malabsorption, the median value of GSS score of digestive symptoms was lower after initiation of standardized low-fructose diet (4 before vs. 1 after; $P=0.0009$).

Our study underscores that fructose malabsorption often occurs in SSc patients. Our findings are thus relevant for clinical practice, highlighting that fructose breath test is a helpful, noninvasive method by: demonstrating fructose intolerance in patients with SSc; and identifying the group of SSc patients with fructose intolerance who may benefit from low-fructose diet. Interestingly, because the present series also shows that low-fructose diet resulted in a marked decrease of gastrointestinal clinical manifestations in SSc patients with fructose malabsorption, our findings underscore that fructose malabsorption may play a significant role in the onset of gastrointestinal symptoms in these patients. Finally, we suggest that fructose malabsorption may be due to reduced fructose absorption by enterocytes, impaired enteric microbiome, and decreased intestinal permeability.

(*Medicine* 94(39):e1601)

Editor: Bulent Kantarceken.

Received: June 17, 2015; revised: August 24, 2015; accepted: August 25, 2015.

From the Department of Internal Medicine, CHU Rouen, and INSERM U 905 (IM, HL); Department of Digestive Physiology, CHU Rouen, and INSERM UMR 1073, University of Rouen IFRMP, Institute for Biochemical Research (A-ML, GG); Department of Biostatistics, CHU Rouen (J-FM); and Department of Gastroenterology, CHU Rouen, and INSERM UMR 1073, University of Rouen IFRMP, Institute for Biochemical Research, Rouen, France (PD).

Correspondence: Isabelle Marie, Department of Internal Medicine, CHU de Rouen, 76031 Rouen Cedex, France
(e-mail: isabelle.marie@chu-rouen.fr).

The authors have no funding and conflicts of interest to disclose.

Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution License 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.
ISSN: 0025-7974

DOI: 10.1097/MD.0000000000001601

Abbreviations: BMI = body mass index, dcSSc = diffuse cutaneous systemic sclerosis, ESR = erythrocyte sedimentation rate, GLUT = glucose transport, GSS score = global symptomatic score, IBS = irritable bowel syndrome, lcSSc = limited cutaneous systemic sclerosis, Low-FODMAP diet = low fermentable oligo-saccharides, disaccharides, monosaccharides and polyol diet, SSc = systemic sclerosis.

INTRODUCTION

Systemic sclerosis (SSc) is a systemic inflammatory disorder affecting the skin and other organs,^{1–3} particularly the gastrointestinal tract where lesions may lead to motor activity impairment.^{4–12} Gastrointestinal involvement occurs in 50 to 88% of patients with SSc.^{4–12} Gastric abnormalities are recognized to be associated with high morbidity, resulting in gastric hemorrhage related to bleeding from antral vascular ectasia (“watermelon stomach”) and delayed gastric emptying.^{4–6} Intestinal involvement may also be responsible for severe complications, including hemorrhage due to small intestinal ectasia, malabsorption syndrome, and intestinal pseudo-obstruction.^{7–12}

Recently, the deleterious effect of fermentable saccharides/polyols, especially fructose that is increasingly incorporated in many beverages, dairy products, and canned/baked or processed foods,^{13–16} has been described in non-SSc patients.^{17–23} Previous investigators have thus reported that 2.4% to 16% of healthy subjects exhibited fructose malabsorption.^{21,24–27} Other authors have further found that fructose malabsorption resulted in abdominal clinical manifestations (ie, pain, discomfort, bloating, distension, nausea, diarrhea),^{21,23,26,28–30} whereas decreased fructose intake led to significant improvement of gastrointestinal symptoms in these patients.^{16,21,23,26,28–30}

Currently, however, there are no studies in SSc assessing the relationship between food intake and gastrointestinal dysfunction; in particular, the link between fructose malabsorption and the occurrence of gastrointestinal impairment has not yet been documented in SSc. Indeed, the aims of the present prospective study were to: determine the prevalence of fructose malabsorption, using fructose breath test, in unselected patients with SSc; evaluate the correlation between fructose malabsorption and digestive symptoms, nutritional status, and gastrointestinal involvement; make prediction about which SSc patients exhibit fructose malabsorption; and assess the outcome of digestive symptoms in SSc patients after initiation of standardized low-fructose diet.

METHODS

Patients

From January 2011 to June 2014, 80 consecutive patients with a definite diagnosis of SSc, attending a tertiary care center, were included in the study. The criteria used for the diagnosis of

SSc were based on the American College of Rheumatology/European League against Rheumatism classification.³¹ Ethical approval was obtained from the local ethical committee (CERNI for the CCP de Haute Normandie), and informed consent was obtained from all patients.

The study cohort consisted of 14 men and 66 women with a median age of 52.5 years (range, 22–79 yr). The median duration of the disease, considered from the onset of the first non-Raynaud's phenomenon clinical manifestations, was 4 years (range, 1–25 yr). Patients were grouped according to criteria of LeRoy et al:³² 23 patients (28.8%) had diffuse cutaneous SSc (dcSSc) and 57 (71.2%) had limited cutaneous SSc (lcSSc). No patient had: other connective tissue disorders (ie, systemic lupus erythematosus, polymyositis/dermatomyositis, rheumatoid arthritis, or Sjögren's syndrome), antiphospholipid syndrome and vasculitis; and history of liver (eg, alcoholic/hepatitis B or C, cirrhosis) or digestive (eg, celiac disease, Crohn disease, ulcerative colitis) diseases, diabetes mellitus, gastric surgery, or vagotomy. Moreover, no patient received nonsteroidal anti-inflammatory drugs.

SSc patients had pulmonary involvement as follows: interstitial lung disease (ILD) ($n=25$; 31.3%); and pulmonary arterial hypertension characterized by pulmonary arterial systolic pressure >40 mm Hg at rest on echocardiography (confirmed by right heart catheterization) ($n=4$; 5%). Forty-three patients had digital ulcers/pitting scars (53.8%) and 31 patients had joint involvement (38.8%). Finally, 2 patients had previous history of renal crisis (2.5%).

All patients had routinely undergone esophageal manometry. Based on manometry findings, Hurwitz's criteria for degree of esophageal involvement was applied as follows: stage I: normal esophageal motility; stage II: uncoordinated peristalsis with normal pressure wave amplitude; stage III: uncoordinated peristalsis with normal low-pressure wave amplitude; and stage IV: both aperistalsis and decreased low esophageal sphincter pressure.^{5–12} According to Hurwitz's criteria, 48 patients (60%) had severe esophageal motor impairment.

Among the 80 SSc patients, 58 had routinely undergone ¹³C octanoic acid breath test, as described previously,⁹ 26 patients exhibited (44.8%) delayed gastric emptying.

Digestive Symptoms

In our SSc patients, the median value of body mass index (BMI) was 24.2 [range, 16.8–34.5].

Before undergoing fructose breath test, SSc patients were interviewed, using a standardized questionnaire, regarding occurrence of gastrointestinal symptoms, that is, nausea, vomiting, abdominal pain/discomfort, bloating, diarrhea, constipation, abdominal tenderness, dysuria, tenesmus, fever, and general illness; each symptom carried a score from 0 (no symptoms) to 3 (severe).^{9,10,33} A global symptomatic score (GSS score), calculated as the sum of all symptom scores, was assigned to each patient (maximum score, 33), as described and validated previously.^{9,10,33} In our patients, median GSS score of digestive manifestations was 2 [range, 0–21].

Furthermore, levels of anxiety and depression were also evaluated, using the hospital anxiety and depression scales.³⁴ In our patients, median values for levels of anxiety and depression were 8 [range, 2–16] and 3 [range, 0–16], respectively.

Biochemical Tests

SSc patients routinely underwent biochemical tests. These included measurements of: first, erythrocyte sedimentation rate (ESR; mm/h), C-reactive protein (mg/L), serum albumin and

prealbumin levels (g/L), ferritin ($\mu\text{mol/L}$), plasma folic acid (nmol/L), vitamin B12 (pmol/L), magnesium (mmol/L), zinc ($\mu\text{mol/L}$), selenium ($\mu\text{mol/L}$), and 25 (OH) vitamin D (nmol/L) levels; and second, antibody status: anti-centromere, anti-topoisomerase I (anti-Scl70), and anti-RNA polymerase III antibodies.

Breath Tests

Each patient underwent 2 breath tests. First, a glucose H₂/CH₄ breath test was performed to exclude underlying small intestinal bacterial overgrowth (SIBO), which is a condition that may promote false-positive results of sugar breath tests.¹⁸ If negative, a fructose breath test was then performed after a 25 g fructose load (10% solution). If the glucose H₂/CH₄ breath test was positive, the patient was given antibiotic therapy, and then underwent the fructose breath test if the glucose breath test was negative.

Before both breath tests, patients were instructed to follow a diet for 48 h. Patients had been instructed to avoid foods likely to generate hydrogen for the 3 days before the test; the day preceding the examination, all patients had low carbohydrate meals (ie, nonseasoned boiled rice and meat cooked on a hot plate or boiled fish and noncarbonated water). Cigarette smoking and physical exercise were not allowed for 2 h before and during the test, to avoid hyperventilation and subsequent changes in breath hydrogen.²¹ After a 12-h fasting, breath testing started after thorough mouth washing with 40 mL of 1% chlorhexidine solution to eliminate an early hydrogen peak related to action of oral bacteria on test sugars.²¹

Glucose Breath Test

SSc patients underwent glucose H₂/CH₄ breath test as previously described.^{10,18} The test was considered positive for SIBO when, at least one of the following criteria was present: H₂ and/or CH₄ increase >20 p.p.m. above basal value; H₂ and/or CH₄ increase >10 p.p.m. on 2 consecutive measurements within the 2 first hours; and H₂ and/or CH₄ increase >10 p.p.m. between minimal and maximal values after glucose ingestion.^{18,33} When the glucose breath test was positive, patients received an antibiotic treatment with quinolone or metronidazole for 10 days per month over 3 successive months. A second glucose breath test was then performed. The fructose breath test was only made in patients exhibiting negative control glucose breath test during the next following day.

Fructose Breath Test

Patients ingested 25 g of fructose dissolved in 250 cc of sterile water; then, end alveolar breath samples were collected every 30 min for 5 h, as described previously.¹⁸ Both H₂ and CH₄ levels were calculated. The test was considered positive and defined a fructose malabsorption in case of a rise of H₂ and/or CH₄ levels above 20 p.p.m.¹⁸

Moreover, during the test, gastrointestinal symptoms were closely monitored and were collected by the laboratory technician every 30 min of the fructose breath test. The occurrence of gastrointestinal symptoms was then analyzed and defined fructose intolerance.¹⁸

Comparison of Fructose Breath Test Findings and Other Manifestations of SSc

We compared various characteristics between SSc patients with and without abnormal fructose breath test.

First, digestive manifestations were compared between these groups of SSc patients. Digestive manifestations included: the values of BMI; the values of: GSS score of digestive

symptoms, anxiety and depression scales; the findings from esophageal manometry according to the Hurwitz criteria for the degree of esophageal involvement on manometry, that is, patients with and without severe esophageal motor dysfunction; and the findings from ¹³C octanoic acid breath test, that is, patients with and without delayed gastric emptying.

Second, patients with and without abnormal fructose breath test were also compared for the following characteristics: median age, median duration of SSc, distribution of SSc subsets, median modified Rodnan score, prevalence of digital ulcers/pitting scars, joint involvement, ILD, pulmonary arterial hypertension, and previous history of renal crisis; autoantibody status (anticentromere, anti-Scl70, and RNA polymerase III antibodies); and findings from biochemical tests, that is, ESR, C-reactive protein, serum albumin, and prealbumin levels, levels of ferritin, plasma folic acid, vitamin B12, magnesium/zinc/selenium, and 25 (OH) vitamin D.

Fructose Restrictive Diet

The SSc patients with fructose intolerance were referred to an experienced dietician for a standardized diet adaptation, consisting of a 1-month low-fermentable oligo-saccharides, disaccharides, monosaccharides, and polyol (FODMAP) diet.¹⁶ These SSc patients were interviewed, using a standardized questionnaire regarding occurrence of gastrointestinal symptoms, using the GSS score.^{5,9,10,33}

Dietary compliance was checked by direct interview by the dietician and the physician, 5 weeks after the initiation of dietary changes; compliance was considered adequate if patients confirmed that they adhered to the dietary guidelines during at least 55% of the meals consumed.

Statistical Analysis

For group comparison involving binary data, we used either the χ^2 test or Fisher exact test, depending on expected cell count (≥ 5 or < 5 , respectively). Comparisons involving continuous data were performed using: Student *t* test when distribution of variables was normal; and Mann–Whitney *U* test in other cases. The results were regarded as significant when the *P* value was less than 0.05.

RESULTS

Prevalence of Fructose Malabsorption

In our cohort of 80 SSc patients, the fructose breath test was positive in 32 patients (40%).

Furthermore, fructose intolerance, defined as the occurrence of gastrointestinal symptoms during fructose breath test, was found in 25 of 32 patients with SSc (78.1%).

Predictive Factors of Fructose Malabsorption

Digestive Symptoms in SSc Patients

The median value for the GSS score of digestive symptoms was significantly higher in SSc patients with fructose malabsorption than in those without (4 vs. 1; *P* = 0.000004).

There were no significant differences between SSc patients with and without fructose malabsorption for: BMI levels (*P* = 0.533), scales of anxiety (*P* = 0.267), and depression (*P* = 0.148) (Table 1).

Esophageal and Gastric Involvement in SSc Patients

As shown in Table 1, patients with normal gastric emptying on ¹³C octanoic acid breath test more often exhibited fructose malabsorption (*P* = 0.007).

We failed to show any significant difference between SSc patients with and without fructose malabsorption for severe esophageal motor disorders (stage IV) on manometry (*P* = 0.489) (Table 1).

Other Predictive Factors for Fructose Malabsorption General Clinical Characteristics

We failed to show any statistically significant differences between SSc patients with and without fructose malabsorption regarding: median age (*P* = 0.992), sex (*P* = 0.381), SSc duration (*P* = 0.487), median modified Rodnan score (*P* = 0.710); and prevalence of digital ulcers/pitting scars (*P* = 0.650), joint involvement (*P* = 0.350), ILD (*P* = 0.806), pulmonary arterial hypertension (*P* = 1) and previous history of renal crisis (*P* = 0.157) (Table 2). Additionally, fructose malabsorption tended to be more common in patients with lcSSc than in those with dcSSc, although not significantly so (*P* = 0.076). Finally, immunosuppressive therapy did not differ between SSc patients with and without fructose malabsorption for low-dose steroids (15.6% vs. 12.5%; *P* = 0.747), methotrexate (9.4% vs. 12.5%; *P* = 0.734), azathioprine (3.1% vs. 6.3%; *P* = 0.646), and mycophenolate mofetil (15.6% vs. 6.3%; *P* = 0.256).

Laboratory Findings

As seen in Table 3, regarding biochemical tests, SSc patients with and without fructose malabsorption did not exhibit different median levels of ESR (*P* = 0.579), C-reactive protein

TABLE 1. Digestive Features of SSc* Patients With Fructose Malabsorption Compared With Those Without

	Fructose Malabsorption (n = 32)	Absence of Fructose Malabsorption (n = 48)	<i>P</i>
GSS score of digestive symptoms	4 [1–21]	1 [0–9]	0.000004
Scale of anxiety	8 [2–16]	7.5 [2–15]	0.267
Scale of depression	6 [1–16]	2.5 [0–13]	0.148
Body mass index	24.9 [15.8–32]	24 [17.8–34.5]	0.533
Esophageal manometry findings			
Severe motor esophageal involvement	56.3%	64.6%	0.489
¹³ C octanoic acid breath test findings			
Absence of delayed gastric emptying	86.7%	61.9%	0.007

* GSS of digestive symptoms: global symptom score of digestive symptoms; SSc: systemic sclerosis. Except were indicated, values are median.

TABLE 2. Clinical Characteristics of SSc* Patients With Fructose Malabsorption Compared With Those Without

	Fructose Malabsorption (n = 32)	Absence of Fructose Malabsorption (n = 48)	<i>p</i>
Clinical parameters			
Age (yr)	54 [range, 27–79]	53 [range, 22–79]	0.992
Sex: male/female	12.5% males/87.5% females	29.7% males/70.3% females	0.381
SSc duration (yr)	5 [range: 1–17]	4 [range: 1–25]	0.487
SSc subset	78.1% of lcSSc*/21.9% of dcSSc*	66.7% of lcSSc*/33.3% of dcSSc*	0.076
Modified Rodnan score	4 [range: 15.8–32]	4.5 [range, 17.8–34.5]	0.710
Digital ulcers/pitting scars	50%	56.3%	0.650
Joint involvement	31.3%	43.8%	0.350
Interstitial lung disease	28.1%	33.3%	0.806
Pulmonary arterial hypertension	6.3%	4.2%	1
Previous history of renal crisis	6.3%	0%	0.157

* dcSSc: diffuse cutaneous SSc; lcSSc: limited cutaneous SSc; SSc: systemic sclerosis. Except were indicated, values are median.

($P = 0.827$), albumin ($P = 0.276$), prealbumin ($P = 0.455$), ferritin ($P = 0.941$), plasma folic acid ($P = 0.444$), vitamin B12 ($P = 0.115$), magnesium ($P = 0.835$), zinc ($P = 0.677$), selenium ($P = 0.077$), and 25 (OH) vitamin D ($P = 0.155$).

Additionally, autoantibody status did not differ between SSc patients and without fructose malabsorption for anti-centromere ($P = 1$), anti-Scl 70 ($P = 0.447$), and anti-RNA polymerase III ($P = 0.156$) antibodies (Table 3).

Outcome of Gastrointestinal Symptoms After Low-FODMAP Diet

Complete outcomes were available in all 32 SSc patients with fructose malabsorption who received low-FODMAP diet. Furthermore, 91% of these patients reported adequate dietary compliance. In overall 32 SSc patients, who received low-FODMAP diet, other specific therapy of SSc was unchanged.

As shown in Table 4, after low-FODMAP diet initiation, SSc patients with fructose malabsorption less commonly exhibited: nausea ($P = 7.09 \times 10^{-6}$), vomiting ($P = 0.024$), abdominal pain ($P = 6.62 \times 10^{-9}$), bloating ($P = 0.0009$), diarrhea ($P = 0.022$), and abdominal tenderness ($P = 0.008$). In addition, median value of GSS score of digestive symptoms was significantly lower (4 before vs. 1 after; $P = 0.00009$) after initiation of low-FODMAP diet in SSc patients.

DISCUSSION

Fructose is a 6-carbon monosaccharide molecule that is found in fruits and vegetables; furthermore, it is increasingly incorporated in processed foods.^{13–16,18,23} Fructose malabsorption has been shown to be responsible for gastrointestinal symptoms.^{16–23,26,28–30} In clinical practice, fructose breath test has been reported as a helpful and reliable method for

TABLE 3. Biochemical Characteristics of SSc* Patients With Fructose Malabsorption Compared With Those Without

	Fructose Malabsorption (n = 32)	Absence of Fructose Malabsorption (n = 48)	<i>P</i>
Biochemical parameters			
ESR* (mm/hour)	10 [range: 1–32]	8.5 [range: 1–89]	0.579
C-reactive protein (mg/L)	3 [range: 1–54]	3 [range: 1–19]	0.827
Albumin (g/L)	41 [range: 28–46]	41.25 [range: 28–50]	0.276
Prealbumin (g/L)	0.28 [range: 0.17–0.41]	0.27 [range: 0.13–0.42]	0.455
Ferritin (μmol/L)	82 [range: 8–334]	735 [range: 6–409]	0.941
Plasma folic acid (nmol/L)	17.2 [range: 3–45]	18.6 [range: 6.5–49]	0.444
Vitamin B12 (pmol/L)	357.5 [range: 126–806]	305.5 [range: 40–861]	0.115
Magnesium (mmol/L)	0.79 [range: 0.67–0.97]	0.81 [range: 0.38–0.95]	0.835
Zinc (μmol/L)	11.5 [range: 9.4–15.1]	11.9 [range: 7.8–14.4]	0.677
Selenium (μmol/L)	0.83 [range: 0.53–1.46]	0.86 [range: 0.55–1.3567]	0.077
Vitamine C (μmol/L)	27.9 [range: 2–83]	443 [range: 3–92]	0.239
25 OH vitamin D (μmol/L)	25 [range: 14–79]	43 [range: 12–104]	0.155
Anti-centromere antibody	43.8%	41.7%	1
Anti-Scl 70 antibody	42.4%	42.4%	0.447
Anti-RNA polymerase III antibody	6.3%	0%	0.156

* ESR: erythrocyte sedimentation rate; SSc: systemic sclerosis. Except were indicated, values are median.

TABLE 4. Gastrointestinal Features of the 32 SSc* Patients With Fructose Malabsorption Before and After FODMAP* Diet Initiation

	Before FODMAP* Diet	After FODMAP* Diet	P
Clinical parameters			
Nausea	46.9%	0%	7.09×10^{-6}
Vomiting	18.8%	0%	0.024
Abdominal pain	81.3%	9.4%	6.62×10^{-9}
Bloating	71.9%	28.1%	0.0009
Diarrhea	31.3%	6.3%	0.022
Constipation	12.5%	6.3%	0.672
Abdominal tenderness	53.1%	18.8%	0.008
GSS* score of digestive symptoms	4 [1–21]	1 (0–13)	0.00009

* FODMAP: low fermentable oligosaccharides, disaccharides, monosaccharides and polyol; GSS of digestive symptoms: global symptom score of digestive symptoms; SSc: systemic sclerosis. Except were indicated, values are median.

assessment of fructose malabsorption.^{17,18,20,21,35,36} However, to date, there have been no large studies using fructose breath test in SSc and consequently, the prevalence of fructose malabsorption has not yet been determined in these patients.

The present prospective study is, to the best of our knowledge, the first to evaluate fructose malabsorption in SSc. In this instance, we found a high frequency (40%) of fructose malabsorption in patients with SSc. Since we investigated 80 SSc patients without any prior selection based on clinical and digestive presentation, our sample appeared to be representative of the entire SSc population. Our findings therefore underscore that fructose malabsorption is prevalent in the whole population of SSc patients. In the current study, fructose breath test was not performed in healthy controls, since previous investigators reported that these latter subjects exhibited fructose malabsorption in only 2.4 to 16% of cases.^{21,24–27} Other authors have also observed fructose malabsorption, in 14.4 to 39% of patients with irritable bowel syndrome (IBS).^{21,37,38}

The current study interestingly underscores that fructose breath test is a reliable method for noninvasively disclosing fructose malabsorption in SSc. Our experimental protocol was carefully designed to avoid any methodological bias that could interfere with the interpretation of the test.

First, SIBO can reasonably be ruled out as a cause of increased proportion of fructose malabsorption in our SSc patients; indeed, all these patients had a negative glucose breath test making SIBO very unlikely. Indeed, because colonic bacteria population migrates proximally into the small intestine in SIBO (gaining access to sugars), this shift in fermentation has been reported to lead to abnormal fructose breath test.^{39,40} However, 1 potential limitation of our study was the use of a glucose breath test for SIBO diagnosis. In fact, the gold standard for the diagnosis of SIBO remains the culture of small bowel aspirates, even such aspirates cannot detect bacterial overgrowth in the distal jejunum and proximal ileum.

Second, we chose 25 g as the dose of fructose load for testing SSc patients, because it is usually accepted as the appropriate dose for clinical use of fructose breath testing.^{21,25,27,41–47} Although Goebel-Stengel et al^{48,49} used 50 g as the dose of fructose load, other authors have shown that healthy subjects can only absorb up to 25 g of fructose per day; indeed, while it is possible that some healthy volunteers may consume up to 50 g of fructose per day, this amount is usually not ingested all at once.^{21,25,27,41–47,50,51} In a series of 17 healthy controls, increasing fructose consumption from 15–

25 g to 50 g resulted in increased prevalence of fructose malabsorption from 5.9% to 52.9%.⁴³ Another series, which compared 3 doses of fructose load (15, 25, and 50 g), also demonstrated that 100% of healthy volunteers could absorb 15 g of fructose, 90% could absorb 25 g of fructose, but only 20 to 30% could absorb 50 g.²⁵ Altogether these findings suggest that the physiologic dose of 25 g fructose appears to be relevant to characterize clinically significant fructose malabsorption, higher doses leading to false-positive diagnosis of fructose malabsorption.⁴²

Furthermore, we used a 5-h fructose breath test duration, because previous authors have found that 10 to 16% of fructose malabsorption would be missed with a 3-h fructose breath test duration.^{18,35} In this instance, the collection of breath samples was also performed according to previous methodological recommendations to ensure end-alveolar breath samples;⁵² we measured both H₂ and CH₄ levels, as it has been shown that false-negative findings were observed in 14% of cases when CH₄ is not measured.^{53,54}

Another main finding in the present series was that fructose malabsorption was strongly associated with a greater prevalence and intensity of gastrointestinal clinical manifestations in SSc, as shown by the higher median value of the GSS score of digestive symptoms in SSc patients exhibiting fructose malabsorption. Previous investigators have indeed reported that unabsorbed fructose will reach the colon in patients, where colonic fermentation leads to gas production (including hydrogen, carbon dioxide, hydrogen sulphide, methane, and short-chain fatty acids), resulting in gastrointestinal symptoms such as abdominal pain and discomfort, bloating, distension, nausea, and diarrhea.^{18,21,23,29,48,54–57} In our experience, SSc patients with fructose malabsorption did not show a more altered psychological profile than SSc patients without fructose malabsorption, as shown by similar scales of anxiety and depression.

Taken together, our data suggest that fructose breath test should be performed in SSc patients without SIBO exhibiting unexplained clinical gastrointestinal symptoms to detect underlying fructose malabsorption.

In the current study, neither the patients' age nor SSc duration could be considered to be predictive of fructose malabsorption. Furthermore, our study shows that the prevalence of fructose malabsorption was similar in patients with dcSSc and lcSSc. In addition, we found that fructose malabsorption was not correlated with other extradigestive manifestations of SSc, including digital ulcers/pitting scars, joint

involvement, ILD, pulmonary arterial hypertension, and previous history of renal crisis. Finally, the present study also shows that autoantibody status, including anticentromere, anti-Scl 70 and anti-RNA polymerase III antibodies, could not be considered predictive factors of fructose malabsorption in patients with SSc.

Additionally, our findings raise the question of whether fructose malabsorption is a participating mechanism or an epiphenomenon in gastrointestinal involvement related to SSc.

From a physiologic standpoint, the human small intestine lacks enzyme to digest fructose; fructose is slowly absorbed across the intestinal epithelium by carrier-mediated facilitated diffusion, which is an energy-independent process.^{35,58} The fructose carrier is a member of the glucose transport (GLUT) family of genes encoding for facilitative sugar transporters.⁵⁸ Previous authors have shown that fructose is mainly absorbed through: GLUT5, which is found in the apical membrane on the luminal surface of human small intestine enterocytes,⁵² and GLUT2 (which is located in the basolateral membrane of intestinal epithelial cells), becoming relevant when high doses of fructose are ingested.^{35,58–61} Other fructose transporters have been described, including GLUT7, GLUT8, and GLUT12, allowing adaptation to high free fructose intake.^{62–64} In a GLUT5 knockout murine model, fructose absorption was decreased by 75% in the jejunum compared with wild-type mice;⁶⁵ these knockout mice, fed with high-fructose diet, developed intestinal dilation and fluid retention reminiscent of fructose malabsorption in humans.⁶⁶ Interestingly, higher prevalence of fructose malabsorption has been reported in patients exhibiting decreased expression of GLUT5;^{67,68} furthermore, in patients with noninsulin-dependent diabetes mellitus, increased expression of GLUT5 has been shown to be correlated with increased absorption of fructose.⁶⁹ By contrast, fructose malabsorption is not associated with GLUT5/GLUT2 mutation.⁶⁶ Recently, Wilder-Smith et al³⁵ have suggested that expression of the GLUT5/GLUT2 transporters may not be uniform along the small intestine. In SSc patients, we suggest that fructose malabsorption may also be, in part, explained by reduced fructose absorption by the enterocytes related to decreased expression of GLUT5/GLUT2 transporters. In essence, inflammation has been shown to decrease intestinal GLUT5 activity and mRNA expression in murine models;⁶³ similarly, in SSc patients, histological analyses of biopsy specimens have shown mononuclear inflammatory cells within intestinal wall. However, a limitation of the current study is that we did not assess the expression of GLUT5/GLUT2 transporter proteins in our SSc patients; thus, other investigations are warranted to confirm these data.

Another main finding in the present series is that fructose malabsorption was more frequent in our SSc patients with normal gastric emptying. Indeed, in healthy subjects, faster is gastric emptying, higher is the fructose load in the intestine and subsequently, higher is the osmotic load related to fructose, leading to fructose intolerance; it has been postulated that in patients with delayed gastric emptying, the postprandial small intestinal load of fructose is lower, reducing the osmotic effect of fructose, reducing the osmotic effect of fructose subsequently.^{70–72} Our findings are in accordance with this hypothesis, as SSc patients with normal gastric emptying more often exhibited fructose malabsorption.

Additionally, in SSc patients, we also suggest that fructose malabsorption may be related to: collagenous fibrosis of small intestinal wall, resulting in decreased intestinal permeability and fructose malabsorption, subsequently;⁴ and impairment of

enteric microbiome. Previous investigators have reported that fructose increases intestinal translocation of bacterial endotoxin and subsequent activation of Kupffer cells through TLR-dependent mechanisms in mice.^{72,73} However, no definite conclusion can be drawn from our findings, and further investigations are warranted to confirm these data.

Another finding in the present study was that SSc patients with fructose malabsorption more often exhibited fructose intolerance (78.1% of cases), determined by the onset of gastrointestinal clinical manifestations during the fructose breath test. We thus suggest that fructose breath test may be useful in identifying the group of SSc patients with fructose intolerance who may benefit from low-fructose diet.

In fact, fructose coexists in food with other poorly absorbed carbohydrates that are FODMAPS.¹⁹ The low-FODMAP diet specifically limits the intake of poorly absorbed short-chain carbohydrates, inducing luminal distension by promoting osmosis and colonic gas production.²³ Previous authors have reported that the low-FODMAP diet is helpful in: IBS patients with symptomatic fructose intolerance, resulting in rapid improvement (within 1 wk) of gastrointestinal manifestations;^{14,19,23,29,73–77} and patients with nonceliac gluten sensitivity, leading to improvement of gastrointestinal symptoms.⁷⁸ The hypothesized mechanism of action of low-FODMAP diet is related to decreased microbial fermentation of dietary FODMAPS, resulting in lower luminal gas generation and distension.⁷⁹ Until now, no studies have directly compared the efficacy of restricting all FODMAPS with that of restricting fructose/fructans alone in patients exhibiting fructose malabsorption. In 1 study, low-fructose/fructan diet was found to reduce digestive symptoms in patients with fructose malabsorption.⁸⁰ Altogether, to date, low-FODMAP diet has been mainly recommended in patients with symptomatic fructose malabsorption.⁸¹

The present series also suggests that low-FODMAP diet is useful in SSc patients with symptomatic fructose malabsorption, resulting in marked improvement of clinical manifestations (ie, nausea, vomiting, abdominal pain, bloating, diarrhea, and abdominal tenderness). Our findings thus underscore that fructose malabsorption may play a significant role in the onset of gastrointestinal symptoms in SSc patients without SIBO. Interestingly, we have also observed that the strategy of dietary advice, being delivered by trained dietician following fructose breath test, provided an appropriate base for SSc patients to adhere to low-FODMAP diet which is crucial for its success. Although there are limitations in our study, because this is a nonrandomized controlled study with the potential for a placebo effect from the intervention; further randomized series are therefore warranted to confirm our findings.

In conclusion, we report a high prevalence (40%) of fructose malabsorption in SSc. Our findings are thus relevant for clinical practice, highlighting that fructose breath test is a helpful, noninvasive method by: demonstrating fructose intolerance in patients with SSc; and identifying the group of SSc patients with fructose intolerance who may benefit from fructose restriction diet. Interestingly, because the present series also shows that low-FODMAP diet resulted in a marked decrease of gastrointestinal clinical manifestations in SSc patients with fructose malabsorption, our findings underscore that fructose malabsorption may play a significant role in the onset of gastrointestinal symptoms in these patients. Finally, we suggest that fructose malabsorption may be related to impairment of fructose absorption by the enterocytes and/or enteric microbiome and decrease of intestinal permeability in SSc.

REFERENCES

- Marie I, Gehanno JF. Environmental risk factors of systemic sclerosis. *Semin Immunopathol.* 2015;37:463–473.
- Marie I, Gehanno JF, Bubenheim M, et al. Prospective study to evaluate the association between systemic sclerosis and occupational exposure and review of the literature. *Autoimmun Rev.* 2014;13:151–156.
- Marie I, Menard JF, Duval-Modeste AB, et al. Association of occupational exposure with features of systemic sclerosis. *J Am Acad Dermatol.* 2015;72:456–464.
- Forbes A, Marie I. Gastrointestinal complications: the most frequent internal complications of systemic sclerosis. *Rheumatology (Oxford).* 2009;48:iii36–iii39.
- Marie I, Antonietti M, Houivet E, et al. Gastrointestinal mucosal abnormalities using videocapsule endoscopy in systemic sclerosis. *Aliment Pharmacol Ther.* 2014;40:189–199.
- Marie I, Ducrotte P, Antonietti M, et al. Watermelon stomach in systemic sclerosis: its incidence and management. *Aliment Pharmacol Ther.* 2008;28:412–421.
- Marie I, Dominique S, Levesque H, et al. Esophageal involvement and pulmonary manifestations in systemic sclerosis. *Arthritis Rheum.* 2001;45:346–354.
- Marie I, Ducrotte P, Denis P, et al. Oesophageal mucosal involvement in patients with systemic sclerosis receiving proton pump inhibitor therapy. *Aliment Pharmacol Ther.* 2006;24:1593–1601.
- Marie I, Ducrotte P, Denis P, et al. Outcome of small bowel motor impairment in systemic sclerosis: a prospective manometric 5-year follow-up. *Rheumatology (Oxford).* 2007;46:150–153.
- Marie I, Gourcerol G, Leroi AM, et al. Delayed gastric emptying determined using the ¹³C-octanoic acid breath test in patients with systemic sclerosis. *Arthritis Rheum.* 2012;64:2346–2355.
- Marie I, Leroi AM, Menard JF, et al. Fecal calprotectin in systemic sclerosis and review of the literature. *Autoimmun Rev.* 2015;14:547–554.
- Marie I, Levesque H, Ducrotte P, et al. Gastric involvement in systemic sclerosis: a prospective study. *Am J Gastroenterol.* 2001;41:1874–1883.
- Bray GA, Nielsen SJ, Popkin BM. Consumption of high-fructose corn syrup in beverages may play a role in the epidemic of obesity. *Am J Clin Nutr.* 2004;79:537–543.
- Choi YK, Kraft N, Zimmerman B, et al. Fructose intolerance in IBS and utility of fructose-restricted diet. *J Clin Gastroenterol.* 2008;42:233–238.
- Economic Research Service, USDA. Table 49-US total estimated deliveries of caloric sweeteners for domestic food and beverage use, by calendar year 2003.
- Fedewa A, Rao SS. Dietary fructose intolerance, fructan intolerance and FODMAPs. *Curr Gastroenterol Rep.* 2014;16:370.
- Riby JE, Fujisawa T, Kretschmer N. Fructose absorption. *Am J Clin Nutr.* 1993;58:748S–753S.
- Melchior C, Gourcerol G, Déchelotte P, et al. Symptomatic fructose malabsorption in irritable bowel syndrome: a prospective study. *United European Gastroenterol J.* 2014;2:131–137.
- Putkonen L, Yao CK, Gibson PR. Fructose malabsorption syndrome. *Curr Opin Clin Nutr Metab Care.* 2013;16:473–477.
- Ragnarsson G, Bodemar G. Pain is temporally related to eating but not to defaecation in the irritable bowel syndrome (IBS). Patients' description of diarrhea, constipation and symptom variation during a prospective 6-week study. *Eur J Gastroenterol Hepatol.* 1998;10:415–421.
- Sharma A, Srivastava D, Verma A, et al. Fructose malabsorption is not uncommon among patients with irritable bowel syndrome in India: a case-control study. *Indian J Gastroenterol.* 2014;33:466–470.
- Shepherd SJ, Parker FC, Muir JG, et al. Dietary triggers of abdominal symptoms in patients with irritable bowel syndrome: randomized placebo-controlled evidence. *Clin Gastroenterol Hepatol.* 2008;6:765–771.
- Turnbull JL, Adams HN, Gorard DA. The diagnosis and management of food allergy and food intolerances. *Aliment Pharmacol Ther.* 2015;41:3–25.
- Densupsoontorn N, Jirapinyo P, Thamonsiri N, et al. Fructose malabsorption in Thai adult. *Asia Pac J Clin Nutr.* 2007;16:209–212.
- Rao SS, Attaluri A, Anderson L, et al. Ability of the normal human small intestine to absorb fructose: evaluation by breath testing. *Clin Gastroenterol Hepatol.* 2007;5:959–963.
- Reyes-Huerta JU, de la Cruz-Patiño E, Ramírez-Gutiérrez de Velasco A, et al. Fructose intolerance in patients with irritable bowel syndrome: a case-control study. *Rev Gastroenterol Mex.* 2010;75:405–411.
- Truswell AS, Seach JM, Thorburn AW. Incomplete absorption of pure fructose in healthy subjects and the facilitating effect of glucose. *Am J Clin Nutr.* 1988;48:1424–1430.
- de Roest RH, Dobbs BR, Chapman BA, et al. The low FODMAP diet improves gastrointestinal symptoms in patients with irritable bowel syndrome: a prospective study. *Int J Clin Pract.* 2013;67:895–903.
- Halmos EP, Power VA, Shepherd SJ, et al. A diet low in FODMAPs reduces symptoms of irritable bowel syndrome. *Gastroenterology.* 2014;146:67–75.
- Staudacher HM, Whelan K, Irving PM, et al. Comparison of symptom response following advice for a diet low in fermentable carbohydrates (FODMAPs) versus standard dietary advice in patients with irritable bowel syndrome. *J Hum Nutr Diet.* 2011;24:487–495.
- van den Hoogen F, Khanna D, Fransen J, et al. 2013 classification criteria for systemic sclerosis: an American College of Rheumatology/European League against Rheumatism collaborative initiative. *Arthritis Rheum.* 2013;65:2737–2747.
- Leroy EC, Black C, Fleischmajer R, et al. Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *J Rheumatol.* 1988;15:202–205.
- Marie I, Ducrotte P, Denis P, et al. Small intestinal bacterial overgrowth in systemic sclerosis. *Rheumatology (Oxford).* 2009;48:1314–1319.
- Cho HS, Park JM, Lim CH, et al. Anxiety, depression and quality of life in patients with irritable bowel syndrome. *Gut Liver.* 2011;5:29–36.
- Wilder-Smith CH, Materna A, Wermelinger C, et al. Fructose and lactose intolerance and malabsorption testing: the relationship with symptoms in functional gastrointestinal disorders. *Aliment Pharmacol Ther.* 2013;37:1074–1083.
- Romagnuolo J, Schiller D, Bailey RJ. Using breath tests wisely in a gastroenterology practice: an evidence-based review of indications and pitfalls in interpretation. *Am J Gastroenterol.* 2002;97:1113–1126.
- Choi YK, Johlin FC Jr, Summers RW, et al. Fructose intolerance: an under-recognized problem. *Am J Gastroenterol.* 2003;98:1348–1353.
- Corlew-Roath M, Di Palma JA. Clinical impact of identifying lactose maldigestion or fructose malabsorption in irritable bowel syndrome or other conditions. *South Med J.* 2009;102:1010–1012.
- Montalto M, Gallo A, Ojetti V, et al. Fructose, trehalose and sorbitol malabsorption. *Eur Rev Med Pharmacol Sci.* 2013;17(Suppl 2):26–29.
- Posserud I, Stotzer PO, Björnsson ES, et al. Small intestinal bacterial overgrowth in patients with irritable bowel syndrome. *Gut.* 2007;56:802–808.

41. Berg LK, Fagerli E, Martinussen M, et al. Effect of fructose-reduced diet in patients with irritable bowel syndrome, and its correlation to a standard fructose breath test. *Scand J Gastroenterol*. 2013;48:936–943.
42. Erdogan A, Adame EC, Yu S, et al. Optimal testing for diagnosis of fructose intolerance: over-dosage leads to false positive intolerance test. *J Neurogastroenterol Motil*. 2014;20:560.
43. Frieling T, Kuhlbusch-Zicklam R, Kalde S, et al. Fructose malabsorption: how much fructose can a healthy subject tolerate? *Digestion*. 2011;84:269–272.
44. Gibson PR, Newnham E, Barrett JS, et al. Review article: fructose malabsorption and the bigger picture. *Aliment Pharmacol Ther*. 2007;25:349–363.
45. Ravich WJ, Bayless TM, Thomas M. Fructose: incomplete intestinal absorption in humans. *Gastroenterology*. 1983;84:26–29.
46. Rumessen JJ, Gudmand-Hoyer E. Absorption capacity of fructose in healthy adults. Comparison with sucrose and its constituent monosaccharides. *Gut*. 1986;27:1161–1168.
47. Sung HY, Kim YS. Fructose malabsorption in patients with irritable bowel syndrome-like symptoms: What is the role in the pathogenesis and clinical implication? *J Neurogastroenterol Motil*. 2014;20:135–137.
48. Goebel-Stengel M, Stengel A, Schmidtmann M, et al. Unclear abdominal discomfort: pivotal role of carbohydrate malabsorption. *J Neurogastroenterol Motil*. 2014;20:228–235.
49. Goebel-Stengel M, Monnikes H. Optimal testing for diagnosis of fructose malabsorption: under-dosage leads to false negative intolerance test. *J Neurogastroenterol Motil*. 2015;21:296–297.
50. Kyaw MH, Mayberry JF. Fructose malabsorption: true condition or a variance from normality. *J Clin Gastroenterol*. 2011;45:16–21.
51. Montonen J, Järvinen R, Knekt P, et al. Consumption of sweetened beverages and intakes of fructose and glucose predict type 2 diabetes occurrence. *J Nutr*. 2007;137:1447–1454.
52. Pimentel M, Chow EJ, Lin HC. Eradication of small intestinal bacterial overgrowth reduces symptoms of irritable bowel syndrome. *Am J Gastroenterol*. 2000;95:3503–3506.
53. Knudsen CD, Di Palma JA. Do you need to measure methane? *South Med J*. 2012;105:251–253.
54. de Lacy Costello BP, Ledochowski M, Ratcliffe NM. The importance of methane breath testing: a review. *J Breath Res*. 2013;7:024001.38.
55. Latulippe ME, Skoog SM. Fructose malabsorption and intolerance: effects of fructose with and without simultaneous glucose ingestion. *Crit Rev Food Sci Nutr*. 2011;51:583–592.
56. Ong DK, Mitchell SB, Barrett JS, et al. Manipulation of dietary short chain carbohydrates alters the pattern of gas production and genesis of symptoms in irritable bowel syndrome. *J Gastroenterol Hepatol*. 2010;25:1366–1373.
57. Simrén M, Stotzer PO. Use and abuse of hydrogen breath tests. *Gut*. 2006;55:297–303.
58. Biesiekierski JR. Fructose-induced symptoms beyond malabsorption in FGID. *United European Gastroenterol J*. 2014;2:10–13.
59. Burant CF, Takeda J, Brot-Laroche E, et al. Fructose transporter in human spermatozoa and small intestine is GLUT5. *J Biol Chem*. 1992;267:14523–14526.
60. Davidson NO, Hausman AM, Ifkovits CA, et al. Human intestinal glucose transporter expression and localization of GLUT5. *Am J Physiol*. 1992;262:C795–C800.
61. Gouyon F, Onesto C, Dalet V, et al. Fructose modulates GLUT5 mRNA stability in differentiated Caco-2 cells: role of cAMP-signalling pathway and PABP (polyadenylated-binding protein)-interacting protein (Paip) 2. *Biochem J*. 2003;375:167–174.
62. DeBosch BJ, Chi M, Moley KH. Glucose transporter 8 (GLUT8) regulates enterocyte fructose transport and global mammalian fructose utilization. *Endocrinology*. 2012;153:4181–4191.
63. Douard V, Ferraris RP. Regulation of the fructose transporter GLUT5 in health and disease. *Am J Physiol Endocrinol Metab*. 2008;295:E227–237.
64. Jones HF, Butler RN, Brooks DA. Intestinal fructose transport and malabsorption in humans. *Am J Physiol Gastrointest Liver Physiol*. 2011;300:G202–G206.
65. Vos MB. Nutrition, nonalcoholic fatty liver disease and the microbiome: recent progress in the field. *Curr Opin Lipidol*. 2014;25:61–66.
66. Wasserman D, Hoekstra JH, Tolia V, et al. Molecular analysis of the fructose transporter gene (GLUT5) in isolated fructose malabsorption. *J Clin Invest*. 1996;98:2398–2402.
67. Douard V, Ferraris RP. The role of fructose transporters in diseases linked to excessive fructose intake. *J Physiol*. 2013;591:401–414.
68. Jones HF, Butler RN, Moore DJ, et al. Developmental changes and fructose absorption in children: effect on malabsorption testing and dietary management. *Nutr Rev*. 2013;71:300–309.
69. Dyer J, Wood IS, Palejwala A, et al. Expression of monosaccharide transporters in intestine of diabetic humans. *Am J Physiol Gastrointest Liver Physiol*. 2002;282:G241–G248.
70. Southgate DA. Digestion and metabolism of sugars. *Am J Clin Nutr*. 1995;62:203S–210S.
71. Horowitz M, Cunningham KM, Wishart JM, et al. The effect of short-term dietary supplementation with glucose on gastric emptying of glucose and fructose and oral glucose tolerance in normal subjects. *Diabetologia*. 1996;39:481–486.
72. Moukarzel AA, Sabri MT. Gastric physiology and function: effects of fruit juices. *J Am Coll Nutr*. 1996;15(5 Suppl):18S–25S.
73. Barrett JS, Gibson PR. Fermentable oligosaccharides, disaccharides, monosaccharides and polyols (FODMAPs) and nonallergic food intolerance: FODMAPs or food chemicals? *Therap Adv Gastroenterol*. 2012;5:261–268.
74. Gibson PR, Varney J, Malakar S, et al. Food components and irritable bowel syndrome. *Gastroenterology*. 2015;148:1158–1174.
75. Pedersen N, Vegh Z, Burisch J, et al. Ehealth monitoring in irritable bowel syndrome patients treated with low fermentable oligo-, di-, mono-saccharides and polyols diet. *World J Gastroenterol*. 2014;20:6680–6684.
76. Rao SS, Yu S, Fedewa A. Systematic review: dietary fibre and FODMAP-restricted diet in the management of constipation and irritable bowel syndrome. *Aliment Pharmacol Ther*. 2015;41:1256–1270.
77. Staudacher HM, Irving PM, Lomer MC, et al. Mechanisms and efficacy of dietary FODMAP restriction in IBS. *Nat Rev Gastroenterol Hepatol*. 2014;11:256–266.
78. Biesiekierski JR, Peters SL, Newnham ED, et al. No effects of gluten in patients with self-reported non-celiac gluten sensitivity after dietary reduction of fermentable, poorly absorbed, short-chain carbohydrates. *Gastroenterology*. 2013;145:320-8.e1–320-8.e3.
79. Murray K, Wilkinson-Smith V, Hoad C, et al. Differential effects of FODMAPs (fermentable oligo-, di-, mono-saccharides and polyols) on small and large intestinal contents in healthy subjects shown by MRI. *Am J Gastroenterol*. 2014;109:110–119.
80. Shepherd SJ, Gibson PR. Fructose malabsorption and symptoms of irritable bowel syndrome: guidelines for effective dietary management. *J Am Diet Assoc*. 2006;106:1631–1639.
81. Escobar MA Jr, Lustig D, Pflugeisen BM, et al. Fructose intolerance/malabsorption and recurrent abdominal pain in children. *J Pediatr Gastroenterol Nutr*. 2014;58:498–501.