

Antigen gene transfer to human plasmacytoid dendritic cells using recombinant adenovirus and vaccinia virus vectors

Hetty J. Bontkes*, Janneke J. Ruizendaal, Marco W.J. Schreurs, Duco Kramer, Chris J.L.M. Meijer and Erik Hooijberg

Department of Pathology, VU University Medical Center, 1081 HV Amsterdam, The Netherlands

Abstract. Recombinant adenoviruses (RA_d) and recombinant vaccinia viruses (RVV) expressing tumour-associated antigens (TAA) are used as anti-tumour vaccines. It is important that these vaccines deliver the TAA to dendritic cells (DC) for the induction of a strong immune response. Infection of myeloid DC (MDC) with RA_d alone is relatively inefficient but CD40 retargeting significantly increases transduction efficiency and DC maturation. Infection with RVV is efficient without DC maturation. Plasmacytoid dendritic cells (PDC) play a role in the innate immune response to viral infections through the secretion of IFN α but may also play a role in specific T-cell induction. The aim of our study was to investigate whether PDC are better targets for RA_d and RVV based vaccines. RA_d alone hardly infected PDC (2%) while CD40 retargeting did not improve transduction efficiency, but it did increase PDC maturation (25% CD83 positive cells). Accordingly, specific CTL activation by RA_d infected PDC was limited (the number of IFN γ producing CTL was reduced by 75% compared to stimulation with peptide loaded PDC). RVV infected PDC specifically stimulated CTL but PDC were not activated. These results indicate that PDC are not ideal targets for RA_d and RVV based vaccines. However, PDC induced specific CTL activation after pulsing with recombinant protein, indicating that PDC can also cross-present antigens released from surrounding infected cells.

Keywords: Human, plasmacytoid dendritic cells, antigen presentation/processing, vaccinia, adenovirus

Abbreviations

APC, antigen presenting cell; CAR, coxsackie-adenovirus receptor; GFP, green fluorescent protein; IFN, interferon; MDC, myeloid dendritic cell; MFI, mean fluorescence index; MoDC, monocyte derived dendritic cell; PDC, plasmacytoid dendritic cell; RA_d, recombinant adenovirus; RVV, recombinant vaccinia virus; TAA, tumour associated antigen; TLR, toll-like receptor.

1. Introduction

Dendritic cells (DC) are the most potent antigen-presenting cell (APC) known to date and appear to be the only cell type capable of inducing primary T cell responses. DC are considered to be the sentinels of the

body and play a key role in delivering antigen from the periphery to the secondary lymphoid organs. Precursors, originating from the bone marrow, migrate to virtually every organ in the body where immature DC take up surrounding antigens. After antigen uptake DC mature, co-stimulatory molecules (e.g. CD40, CD80, CD86) are up-regulated, *de novo* CD83 expression is induced and DC migrate to the lymphoid tissues where they activate effector T-cells [3]. However, it has become clear that DC function in tumor patients is often suppressed, probably due to tumor derived immunosuppressive factors [30]. Furthermore, malignant cells may not lead to immune activation due to the absence of inflammation and “danger” signals [33]. Apart from immunological silence, tolerance can also occur under these non-inflammatory circumstances due to antigen presentation by non-APC (the tumor cells) or due to improperly activated, tolerance-inducing APC [29]. Activation of tumor specific CTL by vaccination with activated DC may break both immunological silence and tolerance in cancer patients [10].

* Corresponding author: Dr. Hetty Bontkes, Department of Pathology, VU University Medical Center, PO Box 7057, 1007 MB, Amsterdam, The Netherlands. Tel.: +31 20 44 44063; Fax: +31 20 44 42964; E-mail: hj.bontkes@vumc.nl.

Myeloid dendritic cells (MDC), characterized by CD11c expression are highly efficient antigen presenting cells capable of inducing primary T cell responses [3,13]. The introduction of methods to culture relatively large numbers of MDC (CD14⁺ monocyte or CD34⁺ progenitor cell derived) *in vitro*, have initiated MDC based vaccination trials for the treatment of cancer [2,28]. A second DC subset characterized by lack of CD11c and high levels of CD123 expression and by expression of the BDCA2 and BDCA4 markers [8] is the plasmacytoid dendritic cell subset (PDC) [12]. Myeloid DC and PDC differentially express toll like receptors (TLR). TLR are innate immune receptors that recognize conserved motifs on microbes and induce inflammatory signals. TLR4 is a receptor for LPS and is expressed on myeloid DC, while TLR9 is expressed on PDC [14]. Upon triggering of TLR9 on PDC by bacterial DNA high levels of IFN α are produced. Recognition of bacterial DNA is based on the presence of unmethylated CG dinucleotides in particular sequence contexts (CpG motifs). PDC are susceptible to stimulation by synthetic oligodeoxynucleotides that contain such CpG motifs (CpG ODN) through TLR 9 [19]. PDC also produce high levels of IFN α upon viral interaction, which has direct anti-viral effects and enhances cytotoxic effects of macrophages and Natural Killer cells. PDC infected by influenza virus stimulate allogeneic T-cells [16] and influenza specific T-cells [11]. PDC may not only play a role in eradicating viral infections but may also be of importance in eradicating virus induced tumors such as Human Papillomavirus induced cervical cancer [5].

Recombinant adeno (RAAd) and vaccinia viruses (RVV) encoding tumor associated antigens (TAA) [6,18,31] have been used to deliver antigens to MDCs. RAAd has been shown to have maturing effects on monocyte derived DC (MoDC) [32], but due to the lack of the adenovirus receptor CAR on MDC, high titers are necessary to obtain high transduction efficiencies *in vitro*. Titers can be reduced by using targeting approaches, which bypass the CAR and redirect the virus to antigens highly expressed on dendritic cells, such as CD40. This retargeting approach to CD40 also induces maturation of infected DCs [32]. RVV effectively infects MoDC but inhibits their maturation [9]. In this study we investigated whether PDC are better targets for these agents, and studied PDC activation/maturation as measured by IFN α production and CD83 expression as well as antigen presentation by PDC after RVV or RAAd infection and cross-presentation after recombinant protein pulsing.

2. Materials and methods

2.1. DC isolation and culture

PBMC were isolated from buffy coats of normal human volunteer donors by density centrifugation over Lymphoprep (Nycomed AS, Oslo, Norway). PDC were isolated (>90% pure as shown by BDCA2 and CD123 double staining) using the BDCA4⁺ cell MACS isolation kit (Miltenyi Biotec, Bergisch Gladbach, Germany) according to the manufacturer's recommendations. The mean yield was 6.9×10^5 /buffy coat (range $3-9 \times 10^5$). Isolated BDCA4⁺ cells were cultured at $3-5 \times 10^5$ cells per ml in Yssels medium supplement with 2% human pooled serum and 10 ng/ml IL-3 (R&D systems, Abingdon, Oxon, UK). MoDC of the same donor were generated as described previously [4].

2.2. Stimulation, infection and peptide loading of DC

BDCA4⁺ PDC were incubated for 24 hours with 3 μ g/ml CpG2216 (a generous gift of Art Krieg, Coley Pharmaceutical Group, MA) as a positive control for PDC activation. Infection with adenovirus type 5 (100 pfu/cell), encoding GFP (RAAd-GFP, kindly provided by Tanja D. de Gruijl, VUMC, Amsterdam), influenza M1 protein (RAAd128 a generous gift from Dr. Rickards and Dr. Wilkinson, University of Wales, UK) or HPV 16 E6/E7 (RAAd101, kindly provided by Dr. S. Inglis, Xenova, Cambridge, UK) all under a CMV promoter, was done using either lipofectamine or the anti-CD40/anti-fiber knob chemical conjugate (CC, kindly provided by Tanja D. de Gruijl, VUMC, Amsterdam) as previously described [4,32]. Vaccinia virus (v1114) infection: DC were washed once in serum free medium and incubated for 4 hours at 37°C with 20 pfu/cell of either the wild type vaccinia virus (RVV-wt) or vaccinia virus encoding influenza M1 protein (RVV-M1).

Peptide loading with the HLA-A2.1 binding influenza M1 derived peptide (M1₅₈₋₆₆) or the HLA-A2.1 binding HPV16 E7 derived peptide (E7₁₁₋₂₀) as a negative control (25 μ M each) was done for 2-4 hours at room temperature in serum free medium in the presence of β 2-microglobulin (3 μ g/ml). For cross-presentation experiments DC were allowed to take up recombinant influenza M1 protein (100 μ g/ml, a generous gift from Sjoerd van der Burg, LUMC, Leiden) at 4°C or 37°C. After 4 hours the cells were washed and incubated o/n at 37°C in an incubator with 5% CO₂ hu-

modified atmosphere. All infections, stimulations and peptide loading were done in the presence of 10 ng/ml IL-3 for the PDC and in the presence of 1000 U/ml IL-4 (Centraal Laboratorium Bloedbank, Amsterdam, The Netherlands) and 100 ng/ml GM-CSF (Schering-Plough, Madison, NJ) for the MoDC.

2.3. Flow cytometry

Cells were incubated at 4°C for 30 min in PBS with 0.1% BSA and 0.01% NaN₃, in the presence of appropriate dilutions of FITC- or PE-labeled isotype controls and mouse mAbs to BDCA-2 and BDCA-4 (Miltenyi Biotec, Bergisch Gladbach, Germany), CD123, CCR7 (PharMingen, San Diego, CA), CD83 (Immunotech, Marseille, France), $\alpha\nu\beta 3$ (Chemicon, Temecula, CA), $\alpha\nu\beta 5$ (Life technologies, Gaithersburg, MD) and coxsackie adenovirus receptor, CAR (a generous gift from Robert W Finberg, Harvard Medical School, Boston, MA). A second incubation step was performed for the unconjugated mAbs with FITC-labeled goat anti-mouse Abs (Centraal Laboratorium van de Bloedtransfusiedienst, Amsterdam, The Netherlands) or goat anti-mouse IgM (Southern Biotechnology Associates, Birmingham, AL). The cells were subsequently analyzed, using a FACS Calibur and CellQuest FACS analysis software (BD Biosciences, Erembodegem-Aalst, Belgium).

2.4. IFN α ELISA

IFN α production was measured in supernatants of stimulated and infected DC (5×10^5 /ml) using the human IFN α ELISA kit (Biosource, Nivelles, Belgium) according to the manufacturer's recommendations.

2.5. IFN γ ELISPOT

An HLA-A2.1 restricted influenza M1 specific (M1₅₈₋₆₆) CTL clone [27] (1000/well) with infected and stimulated DC as target cells (1000/well) were used in an IFN γ ELISPOT assay (Mabtech, Nacka, Sweden) as previously described [4].

2.6. DEX-FITC uptake

DC were incubated for 1 hour with 1 mg/ml of dextran-FITC (MW 70,000; Molecular Probes, Leiden, The Netherlands) on ice or at 37°C. Free dextran-FITC was washed away and uptake was determined by flow cytometry. The ratio of the mean fluorescence intensity of DC incubated at both temperatures was a measure of active dextran-FITC uptake.

2.7. Statistics

The amount of IFN α produced by PDC after different stimuli were compared using a two-sided Student's *T*-test.

3. Results

3.1. Low transduction efficiency of PDC by RAd

PDC and MoDC were infected with RAd-GFP with or without an anti-CD40/anti-fiber knob chemical conjugate. It has previously been shown that this conjugate, which blocks binding of the fiber knob to its natural receptor CAR, increases adenoviral transduction of MoDC. This was confirmed by our results showing an increased transduction of MoDC by the conjugate (17% versus 92% GFP positive cells). In contrast to MoDC, PDC could hardly be transduced either with or without the conjugate (1% versus 2% GFP positive cells, Fig. 1a). The fiber knob of the adenovirus binds to the CAR while either of the $\alpha\nu$ integrins; $\alpha\nu\beta 3$ or $\alpha\nu\beta 5$ is a necessary co-receptor for internalization of the virus. MoDC lack CAR expression but do express $\alpha\nu\beta 5$ [32] ensuring internalization of the virus after transduction in the presence of the anti-CD40/anti-fiber knob chemical conjugate which bypasses the CAR. However, only a low percentage of PDC express CAR (3%), $\alpha\nu\beta 3$ (3%) and $\alpha\nu\beta 5$ (7%) (Fig. 1b) explaining the low transduction efficiency even in the presence of the conjugate.

3.2. RAd and RVV do not induce phenotypic PDC maturation, but RAd induces a low level of IFN α production by PDC

RVV and RAd did not induce phenotypic maturation of PDC, illustrated by a lack of CD83 and CCR7 up-regulation. Moderate PDC maturation was only induced by the combination of the conjugate and RAd-GFP (25% CD83 positive cells). The conjugate alone and in combination with RAd-GFP induces a strong maturation of MoDC (74% and 73% CD83 positive cells, respectively). CpG2216 induced CD83 expression on about a third of the MoDC and PDC (Fig. 1c). RVV, CC, RAd or the combination of RAd and CC did not induce CCR7, which is necessary for migration to the lymph nodes, while CpG2216 did induce CCR7 expression on PDC (not shown). PDC activation was measured by IFN α production after incuba-




Fig. 1. (a) MoDC and PDC were incubated with the bi-specific chemical conjugate of anti-CD40 and anti-fiber knob mAb (CC) alone, RAd-GFP alone or the combination of CC and RAd-GFP. The numbers indicate the percentage of GFP positive cells. (b) Total PBMC were stained with BDCA2 in combination with antibodies directed against CAR, $\alpha v \beta 3$ and $\alpha v \beta 5$. The numbers indicate the proportion of BDCA2 positive cells that express these (co)-receptors (c) MoDC and PDC were incubated with medium, CpG2216, RVV, CC alone, RAd-GFP alone and CC and RAd-GFP combined. The numbers indicate the percentage of CD83 positive cells.

tion with the various agents. Low levels of IFN α production (mean 1.9 ng/ml) were induced by RAd infection alone ($p = 0.029$ compared to medium control) and in combination with the conjugate to similar levels while the conjugate alone and RVV did not induce IFN α production by PDC (Fig. 2a). However, the levels of IFN α induced by RAd were significantly lower compared to the levels induced by CpG2216 (mean 43 ng/ml; $p = 0.016$). On the other hand, MoDC did not produce IFN α upon any of the stimuli given (not shown). These results show that no or only a low level of PDC activation is induced by RVV and RAd, respectively.

3.3. PDC present endogenous antigens delivered by RAd and RVV

PDC were compared with MoDC for their capacity to process and present the influenza M1 antigen delivered by RAd and RVV to an HLA-A2.1 restricted M1₅₈₋₆₆ specific CTL clone. The DC were loaded with either a control peptide or the M1 peptide, infected with RAd encoding either HPV16 E6/E7 (control) or the M1 protein, or infected with wild type RVV or RVV encoding M1. Peptide loaded and in-

fecting DC were incubated overnight in an anti-IFN γ coated ELISPOT plate in the presence of the M1 specific CTL clone. Peptide loaded PDC induced antigen specific IFN γ release by the CTL clone (Fig. 2b). Despite the low transduction efficiency (Fig. 1a), and low (co)-receptor expression (Fig. 2b) RAd infected PDC also induced antigen specific IFN γ release by the CTL clone. MoDC induced a much stronger response, most probably due to higher transduction efficiency of MoDC by RAd. RVV infected PDC induced antigen specific IFN γ release by the CTL clone to similar levels as RVV infected MoDC (Fig. 2b). In conclusion, RVV and to a limited extent RAd deliver antigens to PDC, which are processed and presented to CD8 $^+$ T-cells.

3.4. PDC present exogenous antigens to antigen specific CD8 $^+$ CTL

Subsequently we tested whether PDC were able to take up and present exogenous antigen. PDC were capable of endocytosis as shown by their capacity to take up dextran-FITC. However, MoDC take up dextran-FITC more efficiently as illustrated by the higher MFI (Fig. 3a). PDC were capable of cross-presentation, as




Fig. 2. (a) IFN α production by PDC after incubation with medium, CpG2216, RVV and RAAd with and without CC. P-values (Student's *T*-test) indicate significant differences in IFN α production between the indicated conditions. (b) IFN γ ELISPOT assay of M1 specific CTL clone stimulated with peptide loaded and RAAd and RVV infected PDC and MoDC. The number of specific spots per 1000 CTL are shown.

shown by the antigen specific IFN γ release by the CTL clone after stimulation with PDC, which were fed with recombinant M1 protein. PDC fed at 4°C were used as negative controls (Fig. 3b). In conclusion, PDC are capable of processing and cross-presenting exogenous antigens, comparable to MoDC.

4. Discussion

Recombinant adenovirus is one of the most commonly used vectors for gene transfer into MDC, although clinical studies are limited [24]. Our results confirmed a modest transduction by RAAd alone and a high transduction and maturation of MoDC by an




Fig. 3. (a) Analysis of dextran-FITC uptake by PDC and MoDC. The ratio of the mean fluorescence intensity of cells incubated with dextran-FITC at 37°C and cells incubated on ice is given (mean fluorescence index, MFI). (b) IFN γ ELISPOT assay of M1 specific CTL clone stimulated with PDC loaded with recombinant M1 protein. The number of specific spots per 1000 CTL are shown.

anti-CD40/anti-fiber knob chemical conjugate in combination with RAAd-GFP [32]. In contrast, PDC maturation was not induced by RAAd alone, while the CD40 conjugate and in particular the conjugate in combination with RAAd had a modest effect on maturation. This indicates that RAAd increased cross-linking of the CD40 conjugate and that the maturation we observe is due to CD40 ligation rather than RAAd stimulation and/or infection. Furthermore, RAAd infects PDC with an extremely low efficiency both with and without the chemical conjugate. Since our PDC preparations were >90% pure (result not shown) we cannot exclude the presence of contaminating myeloid DC. Blood derived myeloid DC are infected, although with low efficiency, by adenovirus type 5 [22] and infected MDC may be responsible for the low level of GFP expression after RAAd-GFP infection and low level of CTL activation after RAAd-M1 infection of isolated PDC. However, Fig. 1b shows that in total PBMC a low percentage of cells positive for BDCA2, which is a specific marker for PDC and is not expressed on MDC, also express adenovirus (co)-receptors, indicating that a small proportion of PDC is susceptible to adenovirus infection. Furthermore, PDC secrete high amounts of

IFN α , whereas monocytes and MDC produce little or undetectable amounts of IFN α [17,21]. Figure 2a shows that isolated PDC produce a substantial amount of IFN α upon RAd stimulation, indicating that PDC are activated by RAd to produce IFN α . MoDC lack TLR9 expression and are therefore not expected to be activated by CpG [19], however Fig. 1c shows an up-regulation in CD83 expression after CpG stimulation. This could either be due to an indirect effect of IFN α production by contaminating PDC and/or NK cells in the MoDC cultures [23], although we never detected any IFN α production after CpG stimulation of MoDC. Another possible explanation could be trace amounts of endotoxins in the CpG preparation that can trigger TLR4, which is expressed on MoDC but not on PDC [14].

RVV is another vector used in cancer vaccination trials [6,18]. Transduction efficiency of PDC by RVV, appears to be much more efficient than RAd as suggested by the strong activation of the CTL clone by RVV infected PDC. However, our data show, as has also been shown for MoDC, that RVV does not induce PDC maturation, possible due to the ability of RVV to inhibit host cell protein synthesis, or due to the expression of proteins that interfere with the antiviral effects of IFN α [1,20]. Induction of primary tumor specific CTL depends on mature DC presenting TAA. Our data therefore indicate that PDC are not suitable APC for RVV and RAd based vaccines aiming at induction of tumor specific CTL, due to lack of transduction and/or activation.

Notably, we show here that human PDC are also capable of cross-presenting antigen, which is an important natural method of antigen presentation [7]. PDC and MoDC pulsed with antigen at 37°C induced a similar level of CTL activation. However, there was CTL activation induced by MoDC loaded with protein at 4°C. This was probably due to residual protein adhered to the cell surface, which was internalized and processed during co-culture with the CTL at 37°C. PDC loaded at 4°C did not induce CTL activation indicating differential protein adhesion and uptake by PDC and MoDC, which is also illustrated by the differential uptake of dextran-FITC. These results indicate that PDC are capable of taking up surrounding antigens and cross-presenting them to CTL, however it is not yet clear whether this holds true for all proteins. Since the up-take is mediated by (receptor)binding to the cell, different proteins may be differentially taken up by PDC.

Although the primary function of PDC will be the inhibition of viral replication [16] and induction of

plasma cell differentiation [15,25] through IFN α production, our results and those of others [11,26] indicate that stimulation of antigen specific T cells is an additional property of PDC. Whether PDC are suitable antigen presenting cells to also prime CTL, not only by converting naive MART-1 specific CTL precursors into effector CTL [26], but also by inducing tumor antigen specific CTL that are undetectable before priming, remains to be investigated.

Acknowledgements

This work was supported by Dutch Cancer Society (KWF) grants VU2002-2627 (to H.J.B and D.K.) and VU2001-2503 (to M.W.J.S.). We thank Dr. T.D. de Gruijl for reagents, helpful discussions and comments on the manuscript. We also thank, Dr. Krieg, Dr. Rickards, Dr. Wilkinson, Dr. Inglis and Dr. Finberg and Dr. van der Burg for providing us with reagents.

References

- [1] A. Alcami, J.A. Symons and G.L. Smith, The vaccinia virus soluble alpha/beta interferon (IFN) receptor binds to the cell surface and protects cells from the antiviral effects of IFN, *Journal of Virology* **74** (2000), 11230–11239.
- [2] J. Banchereau, A.K. Palucka, M. Dhodapkar, S. Burkeholder, N. Taquet, A. Rolland, S. Taquet, S. Coquery, K.M. Witkowski, N. Bhardwaj, L. Pineiro, R. Steinman and J. Fay, Immune and clinical responses in patients with metastatic melanoma to CD34(+) progenitor-derived dendritic cell vaccine, *Cancer Research* **61** (2001), 6451–6458.
- [3] J. Banchereau and R.M. Steinman, Dendritic cells and the control of immunity, *Nature* **392** (1998), 245–252.
- [4] H.J. Bontkes, T.D. de Gruijl, G.J. Schuurhuis, R.J. Scheper, C.J. Meijer and E. Hooijberg, Expansion of dendritic cell precursors from human CD34(+) progenitor cells isolated from healthy donor blood; growth factor combination determines proliferation rate and functional outcome, *Journal of Leukocyte Biology* **72** (2002), 321–329.
- [5] H.J. Bontkes, J.J. Ruizendaal, D. Kramer, C.J.L.M. Meijer and E. Hooijberg, Plasmacytoid dendritic cells are present in cervical carcinoma and become activated by Human Papillomavirus type 16 virus like particles, *Gynecologic Oncology* in press (2005).
- [6] L.K. Borysiewicz, A. Fiander, M. Nimako, S. Man, G.W. Wilkinson, D. Westmoreland, A.S. Evans, M. Adams, S.N. Stacey, M.E. Bournsnel, E. Rutherford, J.K. Hickling and S.C. Inglis, A recombinant vaccinia virus encoding human papillomavirus types 16 and 18, E6 and E7 proteins as immunotherapy for cervical cancer, *Lancet* **347** (1996), 1523–1527.
- [7] J.M. den Haan and M.J. Bevan, Antigen presentation to CD8+ T cells: cross-priming in infectious diseases, *Current Opinion in Immunology* **13** (2001), 437–441.

- [8] A. Dzionek, A. Fuchs, P. Schmidt, S. Cremer, M. Zysk, S. Miltenyi, D.W. Buck and J. Schmitz, BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood, *Journal of Immunology* **165** (2000), 6037–6046.
- [9] J. Engelmayer, M. Larsson, M. Subklewe, A. Chahroudi, W.I. Cox, R.M. Steinman and N. Bhardwaj, Vaccinia virus inhibits the maturation of human dendritic cells: a novel mechanism of immune evasion, *Journal of Immunology* **163** (1999), 6762–6768.
- [10] F.D. Finkelman, A. Lees, R. Birnbaum, W.C. Gause and S.C. Morris, Dendritic cells can present antigen in vivo in a tolerogenic or immunogenic fashion, *Journal of Immunology* **157** (1996), 1406–1414.
- [11] J.F. Fonteneau, M. Gilliet, M. Larsson, I. Dasilva, C. Munz, Y.J. Liu and N. Bhardwaj, Activation of influenza virus-specific CD4+ and CD8+ T cells: a new role for plasmacytoid dendritic cells in adaptive immunity, *Blood* **101** (2003), 3520–3526.
- [12] G. Grouard, M.C. Rissoan, L. Filgueira, I. Durand, J. Banchereau and Y.J. Liu, The enigmatic plasmacytoid T cells develop into dendritic cells with interleukin (IL)-3 and CD40-ligand, *Journal of Experimental Medicine* **185** (1997), 1101–1011.
- [13] P. Guermonprez, J. Valladeau, L. Zitvogel, C. Thery and S. Amigorena, Antigen presentation and T cell stimulation by dendritic cells, *Annual Reviews in Immunology* **20** (2002), 621–667.
- [14] D. Jarrossay, G. Napolitani, M. Colonna, F. Sallusto and A. Lanzavecchia, Specialization and complementarity in microbial molecule recognition by human myeloid and plasmacytoid dendritic cells, *Gene Therapy* **31** (2001), 3388–3393.
- [15] G. Jego, A.K. Palucka, J.P. Blanck, C. Chalouni, V. Pascual and J. Banchereau, Plasmacytoid dendritic cells induce plasma cell differentiation through type I interferon and interleukin 6, *Immunity* **19** (2003), 225–234.
- [16] N. Kadowaki, S. Antonenko, J.Y. Lau and Y.J. Liu, Natural interferon alpha/beta-producing cells link innate and adaptive immunity, *Journal of Experimental Medicine* **192** (2000), 219–226.
- [17] N. Kadowaki, Y.J. Liu, Natural type I interferon-producing cells as a link between innate and adaptive immunity, *Human Immunology* **63** (2002), 1126–1132.
- [18] A.M. Kaufmann, P.L. Stern, E.M. Rankin, H. Sommer, V. Nuessler, A. Schneider, M. Adams, T.S. Onon, T. Bauknecht, U. Wagner, K. Kroon, J. Hickling, C.M. Boswell, S.N. Stacey, H.C. Kitchener, J. Gillard, J. Wanders, J.S.C. Roberts and H. Zwierzina, Safety and immunogenicity of TA-HPV, a recombinant vaccinia virus expressing modified human papillomavirus (HPV)-16 and HPV-18 E6 and E7 genes, in women with progressive cervical cancer, *Clinical Cancer Research* **8** (2002), 3676–3685.
- [19] A. Krug, A. Towarowski, S. Britsch, S. Rothenfusser, V. Hornung, R. Bals, T. Giese, H. Engelmann, S. Endres, A.M. Krieg and G. Hartmann, Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12, *European Journal of Immunology* **31** (2001), 3026–3037.
- [20] W.X. Li, H. Li, R. Lu, F. Li, M. Dus, P. Atkinson, E.W.A. Brydon, K.L. Johnson, A. Garcia-Sastre, L.A. Ball, P. Palese and S.W. Ding, Interferon antagonist proteins of influenza and vaccinia viruses are suppressors of RNA silencing, *Proceedings of the National Academy of Sciences* **101** (2004), 1350–1355.
- [21] K. Lore, M.R. Betts, J.M. Brenchley, J. Kuruppu, S. Khojasteh, S. Peretto, M. Roederer, R.A. Seder and R.A. Koup, Toll-like receptor ligands modulate dendritic cells to augment cytomegalovirus- and HIV-1-specific T cell responses, *Journal of Immunology* **171** (2003), 4320–4328.
- [22] O.J.A.E. Ophorst, S. Kostense, J. Goudsmit, R.L. de Swart, S. Verhaagh, A. Zakhartchouk, M. van Meijer, M. Sprangers, G. van Amerongen and S. Yuksel, An adenoviral type 5 vector carrying a type 35 fiber as a vaccine vehicle: DC targeting, cross neutralization, and immunogenicity, *Vaccine* **22** (2004), 3035–3044.
- [23] A.K. Palucka, J. Gatlin, J.P. Blanck, M.W. Melkus, S. Clayton, H. Ueno, E.T. Kraus, P. Cravens, L. Bennett, A. Padgett-Thomas, F. Marches, M. Islas-Ohlmyer, J.V. Garcia and J. Banchereau, Human dendritic cell subsets in NOD/SCID mice engrafted with CD34+ hematopoietic progenitors, *Blood* **102** (2003), 3302–3310.
- [24] S. Paul, B. Calmels and R.B. Acres, Improvement of adoptive cellular immunotherapy of human cancer using ex-vivo gene transfer, *Current Gene Therapy* **2** (2002), 91–100.
- [25] H. Poeck, M. Wagner, J. Battiany, S. Rothenfusser, D. Wellisch, V. Hornung, B. Jahrsdorfer, T. Giese, S. Endres and G. Hartmann, Plasmacytoid dendritic cells, antigen, and CpG-C license human B cells for plasma cell differentiation and immunoglobulin production in the absence of T-cell help, *Blood* **103** (2004), 3058–3064.
- [26] M. Salio, M. Cella, W. Vermi, F. Facchetti, M.J. Palomowski, C.L. Smith, D. Shepherd, M. Colonna and V. Cerundolo, Plasmacytoid dendritic cells prime IFN-gamma-secreting melanoma-specific CD8 lymphocytes and are found in primary melanoma lesions, *European Journal of Immunology* **33** (2003), 1052–1062.
- [27] M.W.J. Schreurs, K.B.J. Scholten, E.W.M. Kueter, J.J. Ruizendaal, C.J. Meijer and E. Hooijberg, In vitro generation and life span extension of human papillomavirus type 16-specific, healthy donor-derived CTL clones, *Journal of Immunology* **171** (2003), 2912–2921.
- [28] B. Schuler-Thurner, E.S. Schultz, T.G. Berger, G. Weinlich, S. Ebner, P. Woerl, A. Bender, B. Feuerstein, P.O. Fritsch, N. Romani and G. Schuler, Rapid induction of tumor-specific type 1 T helper cells in metastatic melanoma patients by vaccination with mature, cryopreserved, peptide-loaded monocyte-derived dendritic cells, *Journal of Experimental Medicine* **195** (2002), 1279–1288.
- [29] M.J. Smyth, D.I. Godfrey and J.A. Trapani, A fresh look at tumor immunosurveillance and immunotherapy, *Nature Immunology* **2** (2001), 293–299.
- [30] C.C. Sombroek, A.G. Stam, A.J. Masterson, S.M. Lougheed, M.J. Schakel, C.J. Meijer, H.M. Pinedo, A.J. van den Eertwegh, R.J. Scheper and T.D. de Gruijl, Prostanoids play a major role in the primary tumor-induced inhibition of dendritic cell differentiation, *Journal of Immunology* **168** (2002), 4333–4343.
- [31] W. Song, Y. Tong, H. Carpenter, H.L. Kong and R.G. Crystal, Persistent, antigen-specific, therapeutic antitumor immunity by

- dendritic cells genetically modified with an adenoviral vector to express a model tumor antigen, *Gene Therapy* **7** (2000), 2080–2086.
- [32] B.W. Tillman, T.D. de Gruijl, S.A. Luykx-de Bakker, R.J. Scheper, H.M. Pinedo, T.J. Curiel, W.R. Gerritsen and D.T. Curiel, Maturation of dendritic cells accompanies high-efficiency gene transfer by a CD40-targeted adenoviral vector, *Journal of Immunology* **162** (1999), 6378–6383.
- [33] M. Wick, P. Dubey, H. Koeppen, C.T. Siegel, P.E. Fields, L. Chen, J.A. Bluestone and H. Schreiber, Antigenic cancer cells grow progressively in immune hosts without evidence for T cell exhaustion or systemic anergy, *Journal of Experimental Medicine* **186** (1997), 229–238.