



ORIGINAL ARTICLE

Quercetin protects human brain microvascular endothelial cells from fibrillar β -amyloid_{1–40}-induced toxicity



Yongjie Li^a, Sibai Zhou^a, Jinze Li^a, Yuhua Sun^b, Hamlati Hasimu^b,
Rui Liu^{a,b,*}, Tiantai Zhang^{a,*}

^aInstitute of Materia Medica, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100050, China

^bXinjiang Key Laboratory of Xinjiang Uygur Medicine, Xinjiang Institute of Materia Medica, Urumqi 830004, China

Received 15 October 2014; received in revised form 5 December 2014; accepted 10 December 2014

KEY WORDS

Alzheimer's disease;
Fibrillar $A\beta_{1-40}$;
Human brain microvascular
endothelial cells;
Quercetin

Abstract Amyloid beta-peptides ($A\beta$) are known to undergo active transport across the blood-brain barrier, and cerebral amyloid angiopathy has been shown to be a prominent feature in the majority of Alzheimer's disease. Quercetin is a natural flavonoid molecule and has been demonstrated to have potent neuroprotective effects, but its protective effect on endothelial cells under $A\beta$ -damaged condition is unclear. In the present study, the protective effects of quercetin on brain microvascular endothelial cells injured by fibrillar $A\beta_{1-40}$ (f $A\beta_{1-40}$) were observed. The results show that f $A\beta_{1-40}$ -induced cytotoxicity in human brain microvascular endothelial cells (hBMECs) can be relieved by quercetin treatment. Quercetin increases cell viability, reduces the release of lactate dehydrogenase, and relieves nuclear condensation. Quercetin also alleviates intracellular reactive oxygen species generation and increases superoxide dismutase activity. Moreover, it strengthens the barrier integrity through the preservation of the transendothelial electrical resistance value, the relief of aggravated permeability, and the increase of characteristic enzyme levels after being exposed to f $A\beta_{1-40}$. In conclusion, quercetin protects hBMECs from f $A\beta_{1-40}$ -induced toxicity.

© 2015 Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences. Production and hosting by Elsevier B.V. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

*Corresponding authors. Tel.: +86 1063035779.

E-mail addresses: liurui@imm.ac.cn (Rui Liu), ttzhang@imm.ac.cn (Tiantai Zhang).

Peer review under responsibility of Institute of Materia Medica, Chinese Academy of Medical Sciences and Chinese Pharmaceutical Association.

1. Introduction

In the pathogenesis of Alzheimer's disease (AD), accumulation and deposition of β -amyloid peptides ($A\beta$) in the brain plays a vital role. $A\beta$ is cleared from the brain across the blood–brain barrier (BBB)^{1,2}. In AD patients, destruction of the BBB leads to $A\beta$ clearance disruption, resulting in $A\beta$ plaques and cerebral amyloid angiopathy (CAA)^{3,4}. The pathology of microvascular CAA is closely related to the cerebrovascular dysfunction that contributes to disease progression^{2,5}. Among all the cerebral microvasculature comorbidities in AD, cerebrovascular $A\beta$ deposition is the most common pathological finding, presenting in up to 90% of AD patients. Amyloid deposition is observed at sites in the cerebrovasculature, mainly in the leptomeningeal and cortical vessels^{6,7}. Cerebrovascular amyloidosis is a central factor in AD pathogenesis and supported by increasing evidence.

Brain microvascular endothelial cells (BMECs) contribute to maintenance of brain homeostasis and the formation of the BBB⁸. In recent studies, it was found that treatment with $A\beta$ could stimulate inflammatory responses, reduce endothelial antioxidants, augment transendothelial migration, and increase the changes in morphology in BMEC monolayers^{9–11}. Progressive build-up of $A\beta$ in and around vessels induces alteration in the BBB permeability, which chronically limits blood supply and results in deprivation of oxygen and nutrients^{12,13}. Accordingly, these changes trigger a secondary cascade of metabolic events involving generation of free radicals, oxidative stress, and release of proteases, such as γ -glutamyl transpeptidase (γ -GT) and alkaline phosphatase (ALP)^{13–15}. Therefore, BMECs are considered as a potential target in AD.

Quercetin (3,3',4',5,7-pentahydroxyflavone, Fig. 1), a natural flavonoid compound, is found in common foods, such as tea, berries and apples. It exerts numerous beneficial effects on human health, including anti-inflammation, anti-ischemia, cardiovascular protection and neuroprotective effects^{16–19}. Quercetin has been reported to pass through the BBB and slow down the progression of degenerative diseases. Additionally, quercetin was shown to reduce the maturation of the amyloid precursor protein in human SH-SY5Y neuroblastoma cells^{19,20}. Quercetin significantly decreased the ratio of insoluble $A\beta$ in a mouse model of AD²¹. Furthermore, studies show that quercetin acts as a novel neuroprotectant by mitigating the increased levels of reactive oxygen species (ROS) that can accelerate the progress of AD^{16,17,19}.

Considering the protective properties of quercetin and its therapeutic potential, we speculated that quercetin might have a protective effect against $A\beta$ -induced BBB disruption. To test our hypothesis, we employed an *in vitro* BBB model of human BMECs (hBMECs). Our study assessed the effects of fibrillar




Figure 1 Chemical structure of quercetin.

$A\beta_{1-40}$ ($fA\beta_{1-40}$) exposure on hBMECs with respect to changes in cell viability, oxidative stress and the barrier function. Finally, we investigated the protective effect of quercetin on hBMECs against $fA\beta_{1-40}$ -induced toxicity.

2. Materials and methods

2.1. Cell culture and treatment

The hBMECs were purchased from ScienCell Research Laboratories (Carlsbad, USA). The hBMECs were cultured in endothelial cell complete medium (ScienCell Research Laboratories, Carlsbad, USA). The cells were incubated at 37 °C in an atmosphere consisting of 95% air and 5% CO₂ in a humidified incubator until reaching 80% confluency.

Synthetic $A\beta_{1-40}$ was purchased from Sangon Biotech Company (Shanghai, China) and dissolved in water to make a stock solution of 0.1 mmol/L to foster the fibrillization state. Quercetin (purity >98%) was purchased from Huike Botanical Development Company (Shanxi, China). It was first dissolved in DMSO at 100 mmol/L, and then diluted in endothelial cell medium at 30.0 μ mol/L, 3.0 μ mol/L and 0.3 μ mol/L. Different concentrations of quercetin were added to the cells at the start of $fA\beta_{1-40}$ injury, and incubated with/without $fA\beta_{1-40}$ for 24 h. The cells were randomly divided into eight groups: (1) control group; control groups in the presence of quercetin at (2) 0.3 μ mol/L, (3) 3.0 μ mol/L, (4) 30.0 μ mol/L for 24 h; (5) $fA\beta_{1-40}$ group treated with 20 μ mol/L $fA\beta_{1-40}$ for 24 h; $fA\beta_{1-40}$ groups treated with quercetin (6) 0.3 μ mol/L, (7) 3.0 μ mol/L, (8) 30.0 μ mol/L for 24 h.

2.2. Cell viability assay

MTS [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-tetrazolium, inner salt] assay (Promega, Madison, USA) was used for evaluating cell viability. After being treated with quercetin and $fA\beta_{1-40}$ as described above, the medium was discarded and replaced with 100 μ L of MTS solution according to the manufacturer's protocol. Then the absorbance of samples at 490 nm was read using a SpectraMax Plus microplate reader (Molecular Devices Corp, Sunnyvale, USA).

2.3. Lactate dehydrogenase (LDH) release assay

LDH, a soluble cytosolic enzyme present in the most eukaryotic cells, is released into the culture medium upon cell death due to damage to the plasma membrane. The increase in LDH activity in the culture supernatant is proportional to the number of lysed cells. The amount of LDH released by cells was determined using an LDH assay kit (Promega, Madison, USA) according to manufacturer's protocol. Briefly, after cells were exposed to $fA\beta_{1-40}$, quercetin or both for 24 h, the supernatant in each well was transferred to a new plate for LDH assay and measured in SpectraMax Plus microplate reader.

2.4. Hoechst 33342 staining assay

Nuclear changes were measured in hBMECs using Hoechst 33342 staining (Dojindo Laboratory, Kumamoto, Japan). After the cells were treated with $fA\beta_{1-40}$, quercetin, or both for 24 h, they were washed in phosphate-buffered solution (PBS), fixed in 4%

formaldehyde for 1 h, and stained with 5 $\mu\text{g/L}$ Hoechst 33342 for 30 min. Nuclear change was assessed by the value of average fluorescent intensity measured with a Cellomics ArrayScan V^{TI} HCS Reader (Cellomics Inc, Pittsburgh, USA).

2.5. Measurements of intracellular ROS and superoxide dismutase (SOD)

The intracellular ROS level was measured using 2,7-dichloro-fluorescein diacetate (DCFH₂-DA, Sigma, St. Louis, USA). After treatment, hBMECs were washed with PBS and incubated with DCFH₂-DA at a final concentration of 5 $\mu\text{mol/L}$ for 40 min at 37 °C protected from light. The cells were then washed with PBS to remove the extracellular DCFH₂-DA, and the fluorescence intensity was measured with SpectraMax Plus microplate reader at an excitation wavelength of 485 nm and an emission wavelength of 538 nm. The intracellular ROS levels were expressed as fluorescent intensity. After the exposure to $fA\beta_{1-40}$, and quercetin, hBMECs were collected and lysed by sonication (60 W with 0.5 s interval for 10 min). The lysate of hBMECs was centrifuged at 10,000 \times g for 15 min and the supernatant fraction was used to measure the SOD activity using WST-1 based SOD inhibition kit (Dojindo Laboratory, Kumamoto, Japan) according to the manufacturer's protocols. The absorbance of the endpoint reaction at 440 nm was measured. Percent inhibition of samples was calculated by the following equation:

$$\text{Inhibition (\%)} = \frac{(A_1 - A_3) - (A_5 - A_2)}{(A_1 - A_3)} \times 100\% \quad (1)$$

where A_1 , A_2 , A_3 and A_5 were the absorbances of uninhibited test, blank sample, blank reagent and sample, respectively.

2.6. Transendothelial electrical resistance (TEER)

TEER was measured using an EVOM resistance meter (World Precision Instruments, Sarasota, USA). The extracellular matrix-treated Transwell inserts (Corning Co., Corning, USA) were placed in a 12-well plate containing culture medium and used to measure the background resistance. The resistance measurements of these blank filters were then subtracted from those of filters with cells. TEER values were expressed as $\Omega \cdot \text{cm}^2$ based on culture inserts.

2.7. Transendothelial permeability for sodium fluorescein and FITC-labeled albumin

To measure transendothelial permeability, the extracellular matrix-treated Transwell inserts were placed in a 12-well plate. The flux of sodium fluorescein (MW, 376 Da) and fluorescein isothiocyanate (FITC)-labeled albumin (MW, 67 kDa) across endothelial monolayers were determined as previously reported^{22,23}. The basolateral compartments of 12-wells contained 1.5 mL Ringer solution (118 mmol/L NaCl, 4.8 mmol/L KCl, 2.5 mmol/L CaCl₂, 1.2 mmol/L MgSO₄, 5.5 mmol/L glucose, 20 mmol/L HEPES, pH 7.4). In the apical chamber, culture medium was replaced by 500 μL Ringer's solution containing 10 $\mu\text{g/mL}$ sodium fluorescein and 165 $\mu\text{g/mL}$ FITC-BSA. The inserts were transferred to a new well containing Ringer's solution at 20 min, 40 min and 60 min. FITC and fluorescein levels were measured by microplate reader (Molecular Devices Corp, Sunnyvale, USA) with emission at 488 nm, excitation at 535 nm and emission at 525 nm, excitation

at 440 nm. Flux across cell-free and cell-containing inserts were both measured. Transport was expressed as μL of donor (luminal) compartment volume from which the tracer is completely cleared. The transendothelial permeability coefficient (Pe) was calculated as previously described^{23,24}. FITC-BSA concentrations in the basolateral chamber, $[\text{BSA}]_{\text{basolateral}}$, were determined from a calibration curve of the fluorescence signal, which was linearly dependent upon BSA concentration.

Permeability, P , was calculated according to a linear approximation of Fick's Law:

$$PS = \frac{J}{[\text{BSA}]_{0,\text{apical}} - [\text{BSA}]_{0,\text{basolateral}}} \quad (2)$$

where S is the surface area of the transwell insert and $[\text{BSA}]_{0,\text{apical}}$ and $[\text{BSA}]_{0,\text{basolateral}}$ are the initial concentration of FITC-BSA in the top and bottom chamber, respectively. The flux, J , was calculated as the product of the volume of the basolateral chamber times the rate of increase in basolateral FITC-BSA concentration determined from the linear region of $[\text{BSA}]_{\text{basolateral}}$ versus time plot. Permeability was measured for each experimental group of endothelial monolayers, as well as for membrane supports in the absence of monolayers. Diffusional Pe were calculated by correction for the permeability of the membrane support in series with the monolayer:

$$\frac{1}{PeS} = \frac{1}{P_{\text{total}}S} - \frac{1}{P_{\text{blank}}S} \quad (3)$$

PeS divided by the surface area (1 cm^2 for Transwell-12) generated the endothelial permeability coefficient (Pe , 10^{-3} cm/min).

2.8. Analysis of levels of γ -glutamyl transpeptidase (γ -GT) and alkaline phosphatase (ALP)

The levels of γ -GT and ALP were measured in the cells using the commercially available assay kits. The kits were purchased from Jiancheng Chemical Factory (Nanjing, China). hBMECs were collected by scrapping and low-speed centrifugation (1000 rpm, 10 min) and measured according to manufacturer's protocols.

2.9. Statistical analysis

Data were expressed as mean \pm SEM and analyzed by one-way analysis of variance (ANOVA) with LSD *post-hoc* correction used for multiple comparisons in our experiment using Graph Pad Prism version 4.0 (GraphPad Inc., La Jolla, USA). Values of $P < 0.05$ are considered statistically significant and F values are given.

3. Results

3.1. Quercetin increased cell viability against $fA\beta_{1-40}$ -induced toxicity in hBMECs

In this study, the protective effects of quercetin on hBMECs against $fA\beta_{1-40}$ -induced toxicity were investigated in three cytotoxicity assays. Firstly, we examined the effect of quercetin on cell viability of hBMECs subjected to $fA\beta_{1-40}$ in the MTS assay. Cell viability was found to be significantly decreased in the presence of 20 $\mu\text{mol/L}$ $fA\beta_{1-40}$ (Fig. 2A, $P < 0.001$). Quercetin at 0.3 $\mu\text{mol/L}$, 3.0 $\mu\text{mol/L}$ and 30.0 $\mu\text{mol/L}$ dose-dependently increased cell viability (Fig. 2A, $P < 0.05$, $P < 0.001$, $P < 0.001$). Quercetin did

not show obvious effects on cell viability of hBMECs without fA β ₁₋₄₀ injury at any of the evaluated concentrations.

The cytoprotective effects of quercetin were also demonstrated by detecting LDH leakage caused by fA β ₁₋₄₀. Treatment with 20 μ mol/L fA β ₁₋₄₀ led to significant enzyme leakage from hBMECs (Fig. 2B, $P < 0.001$). However, when hBMECs were treated with quercetin at 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L in the presence of fA β ₁₋₄₀, the intensity of fluorescence based on the release of LDH in hBMECs decreased significantly (Fig. 2B, $P < 0.001$). No obvious effects of quercetin on the release of LDH from cells without fA β ₁₋₄₀ at 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L was found.

Similar effects were seen in the Hoechst 33342 staining assay. Treatment with fA β ₁₋₄₀ induced nuclear changes, indicating heterogeneous intensities and chromatin condensation in the nuclei of the fA β ₁₋₄₀-injured hBMECs (Fig. 2C). These cytotoxic effects were alleviated both in nuclear morphological condensation and fluorescence intensity through treatment with quercetin at 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L in a dose-dependent manner (Fig. 2D, $P < 0.05$, $P < 0.001$, $P < 0.001$). Quercetin at the same concentrations did not damage the nuclei of control cells.

Based on the above results, quercetin at 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L is found to significantly increase the cell viability, decrease LDH leakage, and ameliorate the nuclear injury induced by fA β ₁₋₄₀. Moreover, quercetin at the same concentrations did not show any cytotoxicity. Summing up the above, we selected concentrations ranging from 0.3 μ mol/L to 30.0 μ mol/L for further investigation.

3.2. Quercetin affected fA β ₁₋₄₀-induced oxidative stress in hBMECs

Oxidative stress plays an important role in A β -induced cytotoxicity. Therefore, we examined the effect of quercetin on fA β ₁₋₄₀-induced oxidative imbalance in hBMECs. The intracellular ROS level of fA β ₁₋₄₀-treated cells showed a remarkable increase (Fig. 3A, $P < 0.001$), while quercetin scavenged ROS generation to protect cells at the concentrations of 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L (Fig. 3A, $P < 0.01$, $P < 0.001$, $P < 0.001$) in the presence of fA β ₁₋₄₀. The results also revealed that fA β ₁₋₄₀ treatment induced the redox imbalance through decreasing SOD activity (Fig. 3B, $P < 0.001$). However, when the cells were treated with quercetin at 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L, SOD activity was remarkably increased in a dose-dependent manner (Fig. 3B, $P < 0.05$, $P < 0.01$, $P < 0.01$).

3.3. Quercetin improved barrier function of hBMECs against fA β ₁₋₄₀-induced toxicity

The BBB is a specialized barrier that maintains the integrity of brain by restricting permeability across the brain endothelium. In this study, the barrier integrity of BMECs and paracellular permeability were determined by the measurement of TEER and the flux of fluorescein sodium and FITC-albumin. As shown in Fig. 4A, the values of TEER in fA β ₁₋₄₀-treated cells were significantly reduced (Fig. 4A, $P < 0.001$). Treatment with




Figure 2 Quercetin increased cell viability against fA β ₁₋₄₀-induced toxicity in hBMECs. (A) Quercetin increased cell viability as evaluated by MTS assay. $n = 8$, $F(7, 56) = 9.541$, $P < 0.001$. (B) Quercetin decreased the levels of extracellular LDH released from hBMECs against fA β ₁₋₄₀-induced toxicity. $n = 6$, $F(7, 40) = 24.144$, $P < 0.001$. (C) The representative nuclear images stained with Hoechst 33342. (D) Quercetin inhibited the nuclear mean fluorescence intensity in hBMECs against fA β ₁₋₄₀-induced toxicity. $n = 4$, $F(7, 24) = 11.761$, $P < 0.001$. Data are expressed as mean \pm SEM, *** $P < 0.001$ versus control group, # $P < 0.05$, ## $P < 0.01$, ### $P < 0.001$ versus fA β ₁₋₄₀ group.




Figure 3 Quercetin affected fA β_{1-40} -induced oxidative stress in hBMECs. (A) Quercetin decreased the generation of ROS in hBMECs in the presence of fA β_{1-40} . $n=6$, $F(4, 25)=45.616$, $P<0.001$. (B) Quercetin improved the SOD activity in hBMECs against fA β_{1-40} -induced toxicity. $n=4$, $F(4, 15)=12.539$, $P<0.001$. Data are expressed as mean \pm SEM, *** $P<0.001$ versus control group, # $P<0.05$, ## $P<0.01$, ### $P<0.001$ versus fA β_{1-40} group.




Figure 4 Quercetin improved barrier function of hBMECs against fA β_{1-40} -induced toxicity. (A) Quercetin at concentrations of 3.0 $\mu\text{mol/L}$ and 30.0 $\mu\text{mol/L}$ decreased TEER. $n=4$, $F(4, 15)=28.676$, $P<0.001$. (B) Quercetin at concentration of 30.0 $\mu\text{mol/L}$ decreased transendothelial permeability for sodium fluorescein. $n=4$, $F(4, 15)=14.654$, $P<0.001$. (C) Quercetin at concentrations of 3.0 $\mu\text{mol/L}$ and 30.0 $\mu\text{mol/L}$ decreased FITC-albumin. $n=4$, $F(4, 15)=13.999$, $P<0.001$. Data are expressed as mean \pm SEM, ** $P<0.01$, *** $P<0.001$ versus control group, # $P<0.05$, ## $P<0.01$, ### $P<0.001$ versus fA β_{1-40} group.

quercetin significantly attenuated the reduction of TEER after being exposed to fA β_{1-40} at 3.0 $\mu\text{mol/L}$ and 30.0 $\mu\text{mol/L}$ (Fig. 4A, $P<0.01$, $P<0.001$).

Permeability and paracellular diffusion across confluent hBMEC monolayers was then evaluated by measuring the permeability coefficient of the permeability markers, fluorescein sodium and FITC-albumin. In the fA β_{1-40} -treated groups, the Pe values were both increased compared with the control group (Fig. 4B and C, $P<0.001$, $P<0.01$). After co-treatment with fA β_{1-40} and quercetin for 24 h, the Pe values of fluorescein sodium and FITC-albumin were both increased in the presence of 30.0 $\mu\text{mol/L}$ quercetin ($P<0.05$); meanwhile, the Pe value of FITC-albumin was increased at 3.0 $\mu\text{mol/L}$ quercetin as well ($P<0.05$).

3.4. Quercetin modulated the levels of γ -GT and ALP in fA β_{1-40} -treated hBMECs

γ -GT and ALP are known to be abundant in cerebral microvessels, and the levels of these two enzymes are involved in barrier formation. On the basis of the above results, we determined the effect of the modulation of the levels of γ -GT and ALP in fA β_{1-40} -treated hBMECs. Levels of γ -GT and ALP were significantly decreased in hBMECs after being exposed to fA β_{1-40} (Fig. 5, $P<0.01$, $P<0.001$), whereas quercetin reversed both changes at 3.0 $\mu\text{mol/L}$ and 30.0 $\mu\text{mol/L}$ (Fig. 5, $P<0.01$, $P<0.01$, $P<0.001$, $P<0.001$). Moreover, quercetin at 0.3 $\mu\text{mol/L}$ significantly increased ALP in presence of fA β_{1-40} , but not γ -GT.




Figure 5 Quercetin modulated levels of γ -GT and ALP in fA β ₁₋₄₀-treated hBMECs. Data are expressed as mean \pm SEM, $n=4$. γ -GT: $F(4, 15)=10.067$, $P<0.001$; ALP: $F(4, 15)=28.919$, $P<0.001$. ** $P<0.01$, *** $P<0.001$ versus control group, ## $P<0.01$, ### $P<0.001$ versus fA β ₁₋₄₀.

4. Discussion

The present study clarifies the beneficial effects of quercetin on AD-associated microvascular endothelial pathology. The present findings indicate that quercetin can protect hBMECs from fA β ₁₋₄₀-induced toxicity. In these effects, quercetin increases cell viability, reduces the amount of LDH release, relieves nuclear condensation, decreases oxidative stress, and strengthens the barrier integrity and functions through improving TEER, ameliorating transendothelial permeability and modulating barrier enzyme levels. These results suggest that quercetin protected endothelial cells against fA β ₁₋₄₀-induced endothelial dysfunction.

Recent advances in neuroscience research have contributed to the emerging concept that the structural and functional integrity of the central nervous system (CNS) depends on the coordinated activity of the neurovascular unit. BMECs, known to be essential for brain homeostasis, provide a dynamic interface between circulating blood and the brain parenchyma¹. Injuries to the endothelial cells of blood vessels triggered by inflammation and other insults (like A β) can result in endothelial death, which contributes to the pathogenesis of diverse neurological disorders²⁵. Accordingly, results from experimental animal models and clinical studies suggest that degenerative disorders are exacerbated, if not initiated, by brain microvasculature injury. Therefore, effective therapeutic strategies to protect neuronal functions, to preserve the integrity of the BBB, and to ensure brain repair should also be targeted toward the brain microvasculature.

Herein, we firstly evidenced that fA β ₁₋₄₀ treatment causes significant cell injury in the hBMEC experimental system²⁶. The toxic effect of fA β ₁₋₄₀ at 20 μ mol/L was achieved in hBMECs, involving decreasing cell viability, increasing LDH release, inducing nuclear injury, overproducing intracellular ROS, reducing SOD activity, and breaking the barrier function.

Appropriate administrative conditions for quercetin were determined using both control and fA β ₁₋₄₀-injured hBMECs. Quercetin at the determined optimal concentrations of 0.3 μ mol/L, 3.0 μ mol/L and 30.0 μ mol/L was found to significantly increase the viability of cells injured by fA β ₁₋₄₀. In line with the increased viability, a decrease in LDH release and relief of nuclear injury also indicated the protective effects of quercetin in this process. In addition, treatment of control cells with quercetin showed no statistically significant differences, indicating that quercetin has no toxicity under basal conditions.

Excessive generation of ROS within endothelial cells in response to A β leads to oxidative stress and cellular injury²⁷⁻²⁹. *In vitro* studies have revealed that oxidative stress results in dysfunction of the endothelial cell, destroying the integrity of the vascular barrier and leading to increased endothelial permeability, mitochondrial dysfunction, chronic inflammatory processes and amyloid deposition in blood vessels, which are involved in the imbalance of endothelial transductions during the pathogenesis of Alzheimer's deficits³⁰⁻³². Neuroprotective interventions for AD should be effective in reducing the severity of oxidative injury and maintaining the integrity of the BBB.

In this study an obvious imbalance was detected after treatment with fA β ₁₋₄₂ alone, involving a marked increase in ROS generation and a reduction of SOD activity. Treatment with quercetin attenuated fA β ₁₋₄₂-induced ROS generation and protected SOD activity in hBMECs. Quercetin is a phenolic compound isolated from plants, and has been thought to be effective in scavenging free radicals³³. Previous studies indicated that quercetin can act as an antioxidant to protect the skin from oxidative stress induced by ultraviolet rays^{34,35}. Here, quercetin produced a significant antioxidant effect in hBMECs treated with fA β ₁₋₄₀. We hypothesize that quercetin might exert cytoprotective effects by suppressing oxidative stress induced by A β ¹⁸. Accumulating evidence suggests that treatment with hydrogen peroxide disturbs the permeability barrier of epithelial cells through disruption of tight junction proteins^{36,37}. As hydrogen peroxide is a member of ROS, we suggest that quercetin protects hBMECs, at least in part, by reducing the overproduction of ROS and improving anti-oxidant efficacy.

The restrictive nature of BMECs for forming the BBB is due to tight junctions among adjacent endothelial cells. The BBB allows for the regulation of ion flux and paracellular diffusion through the development of high transendothelial electrical resistance and tight barrier properties³⁸⁻⁴⁰. The microvascular barrier function was determined to be suitable for subsequent experiments, as shown by the changes of TEER value, permeability property, and characteristic enzymatic activity.

TEER value is an important indicator of the barrier tightness of interendothelial tight junctions. In this study, the reduction of the TEER value could be markedly attenuated by treatment with quercetin at higher concentrations, indicating that the preservation of monolayer integrity may contribute to its protective effects against fA β ₁₋₄₀-induced damage. In addition, quercetin treatment at the highest concentrations attenuated apical-to-basolateral diffusion of both fluorescein sodium and FITC-albumin. Although increased permeability can allow deleterious molecules into the brain and aggravate the endothelial injury, it also may help therapeutic agents cross the BBB. A relevant effect of quercetin on the BBB function has been reported⁴¹, and the mechanisms attributed to the A β ₂₅₋₃₅-induced toxicity in mice have been suggested to result from suppression of MAPK transduction. In combination with our studies, it provides evidence for the explanation that quercetin can cross the BBB and has molecular effects in the brain.

γ -GT and ALP are the important membranous components of the BBB as well, known to be abundant on the apical surface of microvascular endothelial cells⁴¹. The activities of these enzymes can be altered in situations such as oxidative stress⁴², inflammation⁴³ and hypoxia⁴⁴. The levels of γ -GT and ALP changed consistently with TEER value and barrier permeability when subjected to fA β ₁₋₄₀, and they could be preserved with the quercetin treatment in a dose-dependent manner at 3.0 μ mol/L

and 30.0 $\mu\text{mol/L}$. These effects are consistent with the preservation of monolayer integrity based on TEER detection and transendothelial permeability. Thus, we extrapolate that quercetin is capable of protecting the barrier integrity and function of microvascular endothelial monolayers treated with $\text{fA}\beta_{1-40}$.

5. Conclusions

In summary, we have presented novel evidence to show that quercetin effectively prevented $\text{fA}\beta_{1-40}$ -induced hBMEC damage, which was characterized by reduced cell viability, increased LDH leakage and aggravated nuclear condensation. Furthermore, we found that quercetin showed sufficient activity in regulating the redox imbalance and strengthening barrier integrity through the preservation of permeability and characteristic enzymatic activities in the $\text{fA}\beta_{1-40}$ -treated hBMECs. To our knowledge, this is the first direct evidence for an effect of quercetin on brain microvascular endothelial cells. Our results provide a rational basis for the therapeutic application of this compound in the prevention/treatment of AD.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (Nos. 81373388, 81473374 and 81102830).

References

- Kalaria RN. The blood-brain barrier and cerebral microcirculation in Alzheimer disease. *Cerebrovasc Brain Metab Rev* 1992;4:226–60.
- Ghiso J, Fossati S, Rostagno A. Amyloidosis associated with cerebral amyloid angiopathy: cell signaling pathways elicited in cerebral endothelial cells. *J Alzheimers Dis* 2014;42 Suppl. 3: S167–76.
- Parham C, Auckland L, Rachwal J, Clarke D, Bix G. Perlecan domain V inhibits amyloid- β induced brain endothelial cell toxicity and restores angiogenic function. *J Alzheimers Dis* 2014;38:415–23.
- Sagare AP, Bell RD, Zlokovic BV. Neurovascular defects and faulty amyloid- β vascular clearance in Alzheimer's disease. *J Alzheimers Dis* 2013;33 Suppl. 1:S87–100.
- Liu R, Meng FR, Zhang L, Liu AL, Qin HL, Lan X, et al. Luteolin isolated from the medicinal plant *Elsholtzia rugulosa* (Labiatae) prevents copper-mediated toxicity in β -amyloid precursor protein Swedish mutation overexpressing SH-SY5Y cells. *Molecules* 2011;16:2084–96.
- Grammas P, Yamada M, Zlokovic B. The cerebrovasculature: a key player in the pathogenesis of Alzheimer's disease. *J Alzheimers Dis* 2002;4:217–23.
- Grinberg LT, Thal DR. Vascular pathology in the aged human brain. *Acta neuropathol* 2010;119:277–90.
- Deane R, Zlokovic BV. Role of the blood-brain barrier in the pathogenesis of Alzheimer's disease. *Curr Alzheimer Res* 2007;4:191–7.
- Deane R, Du Yan S, Subramanian RK, LaRue B, Jovanovic S, Hogg E, et al. RAGE mediates amyloid- β peptide transport across the blood-brain barrier and accumulation in brain. *Nat Med* 2003;9:907–13.
- Liu R, Wu CX, Zhou D, Yang F, Tian S, Zhang L, et al. Pinocembrin protects against β -amyloid-induced toxicity in neurons through inhibiting receptor for advanced glycation end products (RAGE)-independent signaling pathways and regulating mitochondrion-mediated apoptosis. *BMC Med* 2012;10:105.
- Yan SD, Chen X, Fu J, Chen M, Zhu H, Roher A, et al. RAGE and amyloid- β peptide neurotoxicity in Alzheimer's disease. *Nature* 1996;382:685–91.
- Morris AW, Carare RO, Schreiber S, Hawkes CA. The cerebrovascular basement membrane: role in the clearance of β -amyloid and cerebral amyloid angiopathy. *Front Aging Neurosci* 2014;6:251.
- Fossati S, Ghiso J, Rostagno A. Insights into caspase-mediated apoptotic pathways induced by amyloid- β in cerebral microvascular endothelial cells. *Neurodegener Dis* 2012;10:324–8.
- Lee JK, Jin HK, Park MH, Kim BR, Lee PH, Nakauchi H, et al. Acid sphingomyelinase modulates the autophagic process by controlling lysosomal biogenesis in Alzheimer's disease. *J Exp Med* 2014;211: 1551–70.
- More SS, Vince R. Potential of a γ -glutamyl-transpeptidase-stable glutathione analogue against amyloid- β toxicity. *ACS Chem Neurosci* 2012;3:204–10.
- Ansari MA, Abdul HM, Joshi G, Opii WO, Butterfield DA. Protective effect of quercetin in primary neurons against $\text{A}\beta(1-42)$: relevance to Alzheimer's disease. *J Nutr Biochem* 2009;20:269–75.
- Choi RC, Zhu JT, Leung KW, Chu GK, Xie HQ, Chen VP, et al. A flavonol glycoside, isolated from roots of *Panax notoginseng*, reduces amyloid-beta-induced neurotoxicity in cultured neurons: signaling transduction and drug development for Alzheimer's disease. *J Alzheimers Dis* 2010;19:795–811.
- Jiménez-Aliaga K, Bermejo-Bescós P, Benedí J, Martín-Aragón S. Quercetin and rutin exhibit anti-amyloidogenic and fibril-disaggregating effects *in vitro* and potent antioxidant activity in APPswe cells. *Life Sci* 2011;89:939–45.
- Liu R, Zhang TT, Zhou D, Bai XY, Zhou WL, Huang C, et al. Quercetin protects against the $\text{A}\beta(25-35)$ -induced amnesic injury: involvement of inactivation of rage-mediated pathway and conservation of the NVU. *Neuropharmacology* 2013;67:419–31.
- Huebbe P, Wagner AE, Boesch-Saadatmandi C, Sellmer F, Wolfram S, Rimbach G. Effect of dietary quercetin on brain quercetin levels and the expression of antioxidant and Alzheimer's disease relevant genes in mice. *Pharmacol Res* 2010;61:242–6.
- Tongjaroenbuangam W, Ruksee N, Chantiratikul P, Pakdeenarong N, Kongbuntad W, Govitrapong P. Neuroprotective effects of quercetin, rutin and okra (*Abelmoschus esculentus* Linn.) in dexamethasone-treated mice. *Neurochem Int* 2011;59:677–85.
- Veszelka S, Pásztoi M, Farkas AE, Krizbai I, Ngo TK, Niwa M, et al. Pentosan polysulfate protects brain endothelial cells against bacterial lipopolysaccharide-induced damages. *Neurochem Int* 2007;50:219–28.
- Zhao L, Hou L, Sun HJ, Yan X, Sun XF, Li JG, et al. Apigenin isolated from the medicinal plant *Elsholtzia rugulosa* prevents β -amyloid 25–35-induced toxicity in rat cerebral microvascular endothelial cells. *Molecules* 2011;16:4005–19.
- Vargo MA, Voss OH, Poustka F, Cardounel AJ, Grotewold E, Doseff. Apigenin-induced-apoptosis is mediated by the activation of PKC δ and caspases in leukemia cells. *Biochem Pharmacol* 2006;72:681–92.
- Deli MA, Veszelka S, Csizsár B, Tóth A, Kittel A, Csete M, et al. Protection of the blood-brain barrier by pentosan against amyloid- β -induced toxicity. *J Alzheimers Dis* 2010;22:777–94.
- Liu R, Li JZ, Song JK, Sun JL, Li YJ, Zhou SB, et al. Pinocembrin protects human brain microvascular endothelial cells against fibrillar amyloid- $\beta(1-40)$ injury by suppressing the MAPK/NF- κB inflammatory pathways. *BioMed Res Int* 2014;2014:470393.
- Deng J, Qi XL, Guan ZZ, Yan XM, Huang Y, Wang YL. Pretreatment of SH-SY5Y cells with dicaffeoylquinic acids attenuates the reduced expression of nicotinic receptors, elevated level of oxidative stress and enhanced apoptosis caused by β -amyloid peptide. *J Pharm Pharmacol* 2013;65:1736–44.
- Ashabi G, Ahmadiani A, Abdi A, Abraki SB, Khodagholi F. Time course study of $\text{A}\beta$ formation and neurite outgrowth disruption in differentiated human neuroblastoma cells exposed to H_2O_2 : protective role of autophagy. *Toxicol In Vitro* 2013;27:1780–8.
- Carrano A, Hoozemans JJM, van der Vies SM, Rozemuller AJM, van Horsen J, de Vries HE. Amyloid beta induces oxidative stress-mediated blood-brain barrier changes in capillary amyloid angiopathy. *Antioxid Redox Signal* 2011;15:1167–78.
- Strazielle N, Ghersi-Egea JF, Ghiso J, Dehouck MP, Frangione B, Patlak C, et al. *In vitro* evidence that β -amyloid peptide 1–40 diffuses across the blood-brain barrier and affects its permeability. *J Neuro-pathol Exp Neurol* 2000;59:29–38.

31. Bell RD, Zlokovic BV. Neurovascular mechanisms and blood-brain barrier disorder in Alzheimer's disease. *Acta Neuropathol* 2009;**118**:103–13.
32. Thal DR, Griffin WS, de Vos RA, Ghebremedhin E. Cerebral amyloid angiopathy and its relationship to Alzheimer's disease. *Acta Neuropathol* 2008;**115**:599–609.
33. Song Y, Liu J, Zhang F, Zhang J, Shi T, Zeng Z. Antioxidant effect of quercetin against acute spinal cord injury in rats and its correlation with the p38MAPK/iNOS signaling pathway. *Life Sci* 2013;**92**:1215–21.
34. Liu D, Hu H, Lin Z, Chen D, Zhu Y, Hou S, et al. Quercetin deformable liposome: preparation and efficacy against ultraviolet B induced skin damages *in vitro* and *in vivo*. *J Photochem Photobiol B* 2013;**127**:8–17.
35. Yin Y, Li W, Son YO, Sun L, Lu J, Kim D, et al. Quercitrin protects skin from UVB-induced oxidative damage. *Toxicol Appl Pharmacol* 2013;**269**:89–99.
36. Suganya N, Bhakkiyalakshmi E, Suriyanarayanan S, Paulmurugan R, Ramkumar KM. Quercetin ameliorates tunicamycin-induced endoplasmic reticulum stress in endothelial cells. *Cell Prolif* 2014;**47**:231–40.
37. Nekohashi M, Ogawa M, Ogihara T, Nakazawa K, Kato H, Misaka T, et al. Luteolin and quercetin affect the cholesterol absorption mediated by epithelial cholesterol transporter niemann-pick c1-like 1 in caco-2 cells and rats. *PLoS One* 2014;**9**:e97901.
38. Seok SM, Kim JM, Park TY, Baik EJ, Lee SH. Fructose-1,6-bisphosphate ameliorates lipopolysaccharide-induced dysfunction of blood-brain barrier. *Arch Pharm Res* 2013;**36**:1149–59.
39. Fischer S, Renz D, Schaper W, Karliczek GF. *In vitro* effects of fentanyl, methohexital, and thiopental on brain endothelial permeability. *Anesthesiology* 1995;**82**:451–8.
40. Gloor SM, Wachtel M, Bolliger MF, Ishihara H, Landmann R, Frei K. Molecular and cellular permeability control at the blood-brain barrier. *Brain Res Brain Res Rev* 2001;**36**:258–64.
41. Lawrenson R, Williams T, Farmer R. Clinical information for research: the use of general practice databases. *J Public Health Med* 1999;**21**:299–304.
42. Kugelman A, Choy HA, Liu R, Shi MM, Gozal E, Forman HJ. gamma-Glutamyl transpeptidase is increased by oxidative stress in rat alveolar L2 epithelial cells. *Am J Respir Cell Mol Biol* 1994;**11**:586–92.
43. Das M, Singh SV, Mukhtar H, Awasthi YC. Differential inhibition of rat and human glutathione S-transferase isoenzymes by plant phenols. *Biochem Biophys Res Commun* 1986;**141**:1170–6.
44. Yu S, Iwatsuki H, Ichinohe N, Mori F, Shoumura K. 'In vivo perfusion Turnbull's reaction' for Fe (II) histochemistry in non-anoxic/non-ischemic and anoxic/ischemic cat brains. *Neurosci Lett* 2001;**308**:79–82.