



Arabidopsis LEAFY COTYLEDON1 controls cell fate determination during post-embryonic development

Mingkun Huang^{1,2}, Yilong Hu^{1,2}, Xu Liu¹, Yuge Li¹ and Xingliang Hou^{1*}

¹ Key Laboratory of South China Agricultural Plant Molecular Analysis and Genetic Improvement, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China, ² University of the Chinese Academy of Sciences, Beijing, China

Arabidopsis LEAFY COTYLEDON1 (LEC1) transcription factor is a master regulator that shapes plant embryo development and post-embryonic seedling establishment. Loss-of-function of *LEC1* alters the cotyledon identity, causing the formation of ectopic trichomes, which does not occur in wild-type seedlings, implying that LEC1 might regulate embryonic cell fate determination during post-embryonic development. To test this hypothesis, we compared the expression of trichome development-related genes between the wild-type and the *lec1* mutant. We observed that transcripts of GLABROUS1 (*GL1*), *GL2*, and *GL3*, genes encoding the positive regulators in trichome development, were significantly upregulated, while the TRICHOMELESS1 (*TCL2*), ENHANCER OF TRY AND CPC1 (*ETC1*), and *ETC2* genes, encoding the negative regulators in trichome development, were downregulated in the *lec1* mutant. Furthermore, overexpression of *LEC1* activated the expressions of *TCL2*, CAPPICE (*CPC*), and *ETC1*, resulting in production of cotyledonary leaves with no or fewer trichomes during vegetative development. In addition, we demonstrated that LEC1 interacts with *TCL2* in yeast and *in vitro*. A genetic experiment showed that loss-of-function of *GL2* rescued the ectopic trichome formation in the *lec1* mutant. These findings strongly support that LEC1 regulates trichome development, providing direct evidence for the role of LEC1 in cell fate determination during post-embryonic development.

OPEN ACCESS

Edited by:

Anja Geitmann,
Université de Montréal, Canada

Reviewed by:

Sergey Morozov,
Moscow State University, Russia
Biswapriya Biswas Misra,
University of Florida, USA

*Correspondence:

Xingliang Hou
houxl@scib.ac.cn

Specialty section:

This article was submitted to
Plant Cell Biology,
a section of the journal
Frontiers in Plant Science

Received: 22 July 2015

Accepted: 20 October 2015

Published: 03 November 2015

Citation:

Huang M, Hu Y, Liu X, Li Y and Hou X
(2015) *Arabidopsis* LEAFY
COTYLEDON1 controls cell fate
determination during post-embryonic
development. *Front. Plant Sci.* 6:955.
doi: 10.3389/fpls.2015.00955

Keywords: *Arabidopsis*, LEAFY COTYLEDON1, trichome, cell fate determination, post-embryonic development

INTRODUCTION

In higher plants, embryogenesis generally terminates with a dormancy period for future sporophyte growth. Once conditions are favorable, seeds germinate, and undergo post-embryonic development, during which cells acquire specific fates. Extensive studies have shown that the specific cell fate determination during post-embryonic development is precisely controlled by multiple transcription factors (Peris et al., 2010; Perianez-Rodriguez et al., 2014). The plant Nuclear Factor Y (NF-Y) transcription factor LEAFY COTYLEDON1 (LEC1), a master regulator controlling embryogenesis (Meinke, 1992; West et al., 1994; Harada, 2001; Lee et al., 2003), was recently revealed to play a potential role in regulating post-embryonic development (Junker and Baumlein, 2012). *LEC1* is expressed in both developing embryo and post-embryonic seedlings, but

not in the vegetative true leaves (Kwong et al., 2003; Warpeha et al., 2007; Le et al., 2010). Loss-of-function of *LEC1* causes a pleiotropic phenotype, including embryo desiccation intolerance and defects in the accumulation of seed storage compounds in developing seeds, as well as cotyledon with trichomes, premature development of vascular tissue and mesophyll cells, and short hypocotyls in early seedlings (Meinke, 1992; Meinke et al., 1994; West et al., 1994; Brocard-Gifford et al., 2003). By contrast, ectopic expression *LEC1* allows seedlings to remain in the embryonic state with an unexpanded cotyledon after germination and is sufficient to induce conversion of true leaves into embryonic structures that lack trichomes (Lotan et al., 1998; Junker et al., 2012). These findings support the multifunctional roles of *LEC1* in embryonic and post-embryonic development. Trichomes originate from the epidermal cell layer whose nuclei have undergone multiple rounds of endoreduplication, which then enlarge and expand away from the surface (Hulskamp et al., 1994; Pattanaik et al., 2014). The regulation of trichome formation has been well documented in many studies. The R2R3 MYB transcription factor *GLABROUS1* (*GL1*) was the first identified positive regulator involved in trichome development (Marks and Feldmann, 1989; Oppenheimer et al., 1991). *GL1* interacts with the bHLH transcription factor *GLABRA3* (*GL3*), its close homolog *ENHANCER OF GLABRA3* (*EGL3*), and the WD40-repeat factor *TRANSPARENT TESTA GLABRA1* (*TTG1*), resulting in the formation of the MYB-bHLH-WD-repeat trichome-positive transcription complex (Zhao et al., 2008). This protein complex subsequently induces trichome formation by activating the expression of the downstream gene *GLABRA2* (*GL2*), encoding a homeodomain-leucine zipper (HD-Zip) transcription factor (Rerie et al., 1994; Morohashi et al., 2007). Contrastingly, a small single-repeat R3 MYB protein family, including *TRICHOMELESS1* (*TCL1*), *TCL2*, *TRIPTYCHON* (*TRY*), *CAPPICE* (*CPC*), *ENHANCER OF TRY AND CPC1* (*ETC1*), and *ETC2*, plays redundant roles in the negative regulation of trichome development. These proteins compete with *GL1* for binding to *GL3/EGL3*, resulting in the formation of an inactive protein complex that fails to activate *GL2* expression and thus represses trichome formation (Wang and Chen, 2014; Zhou et al., 2014).

Trichome formation is considered a specific characteristic that distinguishes true leaves from cotyledons in *Arabidopsis* (Chandler, 2008). A hypothesis for the ectopic trichome formation on the cotyledons of the *lec1* mutant is that the loss-of-function of *LEC1* might result in the disturbance of embryonic cell fate determination, allowing the precocious post-germination events in which cotyledons are partially converted into true leaves (West et al., 1994; Chandler, 2008). However, the mechanism by which *LEC1* functions in cell fate determination during post-embryonic development remains elusive.

Given that *LEC1* mutation causes remarkable ectopic trichome growth on cotyledons, the trichome-related genes might serve as good candidates to study the function of *LEC1* in the post-embryonic development phase. Here, we demonstrated that *LEC1* represses trichome cell differentiation during post-embryonic development by regulating trichome-related genes. In addition, *LEC1* was shown to interact with *TCL2*, a repressor

of trichome development, which indicated that *LEC1* might couple with other transcription factors to co-regulate trichome formation. A genetic experiment showed that loss-of-function of *GL2* rescued the ectopic trichome development phenotype of the *lec1* mutant. These results strongly supported the view that *LEC1* functions in cell fate determination during the post-embryonic phase, providing new insights into *LEC1*'s role beyond the embryogenesis.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

All *Arabidopsis* plants used in this study are in the Col genetic background. The mutant *gl2* was described previously (Wang et al., 2010). The mutant *lec1-4* (Salk_095699) was obtained from the *Arabidopsis* Biological Resource Center (<http://www.arabidopsis.org>). The estradiol-inducible *pER10:LEC1-MYC* transgenic line was obtained by kanamycin selection after transformation (Mu et al., 2008). The double mutant was generated by genetic crossing. After surface sterilization, seeds were sown on the half-strength Murashige and Skoog medium containing 0.8% agar and incubated at 4°C in the dark for 3 days. The seedlings were then transferred to the growth chamber at 22°C under long day conditions (16 h light/8 h dark). For estradiol treatment, 10 μM estradiol was added to the half-strength Murashige and Skoog medium plate and dimethyl sulfoxide, the solvent for estradiol, served as the mock treatment. After treatment, these seedlings were harvested for further analyses.

Microscopy

The formation of trichomes on the cotyledons and leaves was observed and photographed using a LEICA M165C stereoscope (Leica, Wetzlar, Germany).

Statistical Analysis

Five-day-old seedlings were observed to calculate frequency of the appearance of ectopic trichomes on cotyledons. At least 15 Col and 50 *lec1-4* seedlings were used in each experiment. Data represent mean ± SD of three independent repeats. To calculate the frequency of different trichome types on *lec1-4* cotyledons, at least 50 trichomes were analyzed in each experiment. Data represent the mean ± SD of seven independent repeats.

Gene Expression Analysis

Total RNA was extracted from 5-day-old *Arabidopsis* seedlings using the Plant RNA Kit (OMEGA, Atlanta, GA, USA), following the manufacturer's instructions. One microgram of total RNA was used for the reverse transcription reaction. The first-strand cDNA was synthesized using the M-MLV reverse transcriptase (Promega, Madison, WI, USA), according to the manufacturer's instructions. Quantitative real-time PCR was performed using the LightCycler480 system (Roche, Basel, Switzerland) in a total volume of 10 μl, with 0.25 μl of each primer (10 μM), 1 μl of cDNA product and 5 μl of SYBR Premix ExTaq (Takara, Tokyo, Japan). The PCR program included an initial denaturation step at

94°C for 1 min, followed by 40 cycles of 10 s at 94°C and 1 min at 60°C. Each sample was quantified at least in triplicate and normalized using the *TUB2* gene as the internal control. Primers used in this study are listed in Supplementary Table S1.

Yeast Two-Hybrid Assay

The full-length coding sequences of *LEC1* and *TCL2* were amplified and cloned into vectors pGBKT7 and pGADT7 (Clontech, Palo Alto, CA, USA), respectively. Primers are listed in Supplementary Table S1. Yeast two-hybrid assays were performed following the manufacturer's instructions of the Yeastmaker Yeast Transformation System 2 (Clontech). In brief, a single colony of yeast *AH109* was incubated at 30°C overnight ($OD_{600} > 1.0$). The cells were harvested by centrifugation and then resuspended in 25 mL ddH₂O. After re-harvesting the cells by centrifugation, a 1.5-ml sterile 1× Tris-EDTA/LiAc solution was added to prepare yeast-competent cells. For the yeast two-hybrid assay, the bait (0.5 µg) and/or prey (0.5 µg) plasmids with 0.1 mg of carrier DNA were co-transformed into the yeast-competent cells using Polyethylene glycol/LiAc solution. After incubation at 30°C for 30 min, 70 µL dimethyl sulfoxide was added and incubation continued at 42°C for 15 min. The cells were centrifuged and washed using Tris-EDTA solution. After transformation, the yeast cells were grown on SD/-Trp/-Leu/-His/-Ade medium for the interaction test.

Pull-Down Assay

To produce glutathione-S-transferase (GST)-TCL1, GST-TCL2, GST-CPC, GST-ETC1, His-LEC1, and His-GFP recombinant proteins, the full-length coding sequences of the tested genes were cloned into vector pGEX-4T-1 (Pharmacia, Piscataway, NJ, USA) or pQE30 (QIAGEN, Dusseldorf, Germany), respectively. Primers are listed in Supplementary Table S1. These constructs were transformed into the *Escherichia coli* Rosetta strain DE3 (Novagen, Billerica, MA, USA). One hundred milliliters of ($OD_{600} \approx 0.5$) Rosetta strains harboring various vectors were incubated at 16°C for 16 h with 0.1 mM isopropyl β-D-1-thiogalactopyranoside. The soluble GST fusion proteins were purified using glutathione sepharose beads (Amersham Biosciences, Piscataway, NJ, USA), while His fusion proteins were purified using Ni-nitrilotriacetic acid agarose beads (QIAGEN), according to the manufacturer's instructions, respectively. For pull-down assays, 2 µg of His-LEC1 or His-GFP were incubated with immobilized GST or GST fusion proteins in the binding buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, and 1 mM EDTA) at 4°C for 4 h. After washing with binding buffer three times, proteins retained on the beads were subsequently resolved by sodium dodecyl sulfate polyacrylamide electrophoresis and detected with an anti-His antibody (GIBCO, Beijing, China).

RESULTS

Ectopic Trichome Formation on the Cotyledons of the *lec1* Mutant

Extensive studies have shown that *LEC1* serves as a key regulator in embryo development, which is consistent with its main



expression in the developing seed (Kwong et al., 2003). However, recent observations suggested a potential role of *LEC1* in post-embryonic cell differentiation, including hypocotyl elongation and the formation of vascular tissue, mesophyll cells and trichomes (Warpeha et al., 2007; Junker and Baumlein, 2012; Junker et al., 2012; Supplementary Figure S1A). Serving as a general model system in the study of cell fate determination in *Arabidopsis* (Yang and Ye, 2013), we focused on trichome formation to reveal the function of *LEC1* in post-embryonic development. A desiccation-tolerant leaky allele *lec1-4* whose mature seeds germinate normally (unpublished data) was used in this study. Consistent with previous reports, the wild-type cotyledons were glabrous, while ~85% of *lec1* cotyledons produced a varied number of ectopic trichomes (Figure 1A; Supplementary Table S2). Moreover, unlike the mostly three-branched trichomes developed on true leaves, the trichomes on *lec1* cotyledons were mainly unbranched (~44%) or two-branched (~55%), and rarely three-branched (less than 1%; Figure 1B; Supplementary Table S3). These observations implied incomplete conversion of the *lec1* cotyledons into true leaves. In addition, there was no significant difference observed in trichome numbers or the morphology of rosette leaves between the wild-type and *lec1*, which is consistent with the observation that no *LEC1* transcripts accumulated in the vegetative tissues (Supplementary Figures S1A,B).

Misregulation of Trichome-Related Genes in the *lec1* Mutant

Trichome formation on leaves has been well studied during the past two decades and a number of trichome-related genes were identified, which could be divided into two groups: trichome-positive and -negative genes (Yang and Ye, 2013). To examine whether *LEC1*'s involvement in the inhibition of trichome formation on cotyledons occurs by mediating the regulation of trichome-related genes, five trichome-positive genes, and six trichome-negative genes were selected for quantitative real-time PCR analysis in wild-type and *lec1* seedlings. Interestingly, among the trichome-positive genes, the expressions of *GL1*, *GL2*, and *GL3* were significantly upregulated in the *lec1* mutant compared with the wild-type, while those of *EGL3* and *TTG1* showed no change (Figure 2A). Moreover, the expressions of the trichome-negative genes *TCL2*, *ETC1*, and *ETC2* were significantly downregulated in the *lec1* mutant compared with the wild-type, while those of *TRY* and *CPC* showed little change. The *TCL1* transcript was hardly detected in either the wild-type or *lec1* seedlings (Figure 2B). These results indicated that the ectopic trichome formation might be caused by misregulation of the trichome-related genes in the *lec1* mutant.

Overexpression of *LEC1* Activates Several Trichome-Negative Transcription Factors

Given that loss-of-function of *LEC1* resulted in misregulation of trichome-related genes, we wondered whether *LEC1* overexpression could activate or repress these genes. *LEC1* overexpression, driven by the 35S promoter, caused complex




phenotypes and embryonic lethality (Lotan et al., 1998). Thus, instead, we employed an inducible transgenic plant harboring *pER10:LEC1-MYC*, in which *LEC1* expression could be significantly induced by estradiol (Mu et al., 2008; Supplementary Figure S2). Unexpectedly, the trichome-positive

genes *GL1*, *GL2*, *GL3*, *EGL3*, and *TTG1* were not regulated by induced *LEC1* (data not shown). Nevertheless, overexpression of *LEC1* activated the expressions of *TCL2*, *CPC*, and *ETC1*, but had no effect on those of *TCL1*, *TRY*, and *ETC2* (Figure 3A), which revealed the role of *LEC1* in the regulation of trichome-negative genes. A previous study reported that *TCL2*, *CPC*, and *ETC1* overexpression lines all presented a glabrous leaves phenotype (Wang and Chen, 2014); therefore, we asked whether ectopic *LEC1* expression affects trichome formation on true leaves. The investigation demonstrated that cotyledons of *pER10:LEC1-MYC* under either mock or estradiol treatment displayed a similar glabrous phenotype to the wild-type (data not shown). Strikingly, under long-term induction by estradiol (14 days), the *pER10:LEC1-MYC* seedlings developed fewer trichomes and cotyledon-like rosette leaves (Figure 3B). Together with a previous report that the single-repeat R3 MYB quintuple mutant *try cpc etc1 etc3 tcl1* produced ectopic trichomes on cotyledons similar to the *lec1* mutant (Wang et al., 2008), our observations supported the hypothesis that *LEC1* suppresses trichome formation via activation of *TCL2*, *CPC*, and *ETC1* expression.

LEC1 Physically Interacts with *TCL2* *In Vitro* and in Yeast


LEAFY COTYLEDON1 belongs to the NF-Y family, whose members interact with other transcription factors to co-regulate target gene expression (Yamamoto et al., 2009). To investigate whether *LEC1* couples with cofactors to mediate trichome formation, we performed a yeast two-hybrid screening assay. Interestingly, the trichome transcriptional repressor *TCL2* was observed to interact with *LEC1* in yeast (Figure 4A). We then performed a pull-down assay to confirm the interaction between *LEC1* and *TCL2* using purified GST- and His-tagged proteins (Supplementary Figure S3). As expected, GST-*TCL2* interacted with His-*LEC1*, but not with the His-GFP protein, while GST



alone did not precipitate either of the His-tagged proteins (Figure 4B). TCL2 and its homologs TCL1, TRY, CPC, ETC1, and ETC2 exert a master role in repressing trichome formation during the whole life of plants; therefore, we further tested the binding between LEC1 and these homologous proteins. The pull-down results showed that LEC1 also interacted with TCL1, CPC, and ETC1 *in vitro* (Supplementary Figure S4); however, their bindings were not observed in the yeast two-hybrid assay, probably because of weak interactions in yeast (data not shown). These results indicated that, besides regulating the expression of the single-repeat R3 MYB genes, LEC1 might mediate trichome formation via direct protein interactions with these repressors, a result that requires further investigation.

GL2 Genetically Acts Downstream of LEC1


GLABRA2 functions as the key transcription activator in trichome development, whose expression is activated by the trichome-positive MYB-bHLH-WD and repressed by the trichome-negative single-repeat R3 MYB-bHLH-WD transcription complexes (Zhao et al., 2008). Loss-of-function of *GL2* completely abolished trichome formation on the first pair of true leaves in early seedlings (Figure 5). The elevated *GL2* expression in the *lec1* (Figure 2A) implied an epistatic effect



of *GL2* on *LEC1* in trichome formation on cotyledons. To test this hypothesis, we created a *gl2 lec1* double mutant and found that similar to *gl2*, the *gl2 lec1* double mutant displayed glabrous true leaves (Figure 5). In contrast, the *gl2* mutation repressed the ectopic trichome formation on cotyledons of *lec1* seedlings remarkably (Figure 5), confirming the primary role of *GL2* in trichome formation, regardless of the developmental stage. These results strongly supported the hypothesis that *GL2* is required for *LEC1*-mediated repression of trichome formation on cotyledons.

DISCUSSION

Post-embryonic development is a vital transition state from embryonic status to vegetative growth, during which seedlings sense various environmental cues, such as light and temperature, to enable young sporophyte survival. At this stage, cell fate determination occurs to promote post-embryonic organogenesis that arises from the primary meristems, during which the visible features, such as cotyledons extension, hypocotyl elongation, and the emergence of true leaves with trichomes, can be observed in seedlings. Studies have shown that multiple endogenous factors, including phytohormones and transcription regulators, are involved in post-embryonic seedling development (Perianez-Rodriguez et al., 2014). LEC1, the master regulator in embryogenesis, was recently reported to be involved in light-mediated post-embryonic hypocotyl elongation (Junker and



Baumlein, 2012). In this study, we provide molecular evidence that *LEC1* and its genetic downstream partner *GL2* are required for the repression of trichome formation on embryo-derived cotyledons, the characteristic that distinguishes them from true leaves. Mutation of *LEC1* resulted in the misregulation of several trichome-related genes and thus triggered trichome formation on cotyledons. By contrast, ectopic *LEC1* activates the expression of the single-repeat *R3 MYB* genes and induces embryonic leaves with few or no trichomes. Furthermore, the results indicated that *LEC1* might regulate trichome formation by interacting with other transcription factors (e.g., *TCL2* and its homologs). Taken together, our findings support the hypothesis that *LEC1* plays an essential role in cell fate determination during the post-embryonic development.

The fact that *LEC1* is expressed in both developing seeds and post-embryonic seedlings, combined with the pleiotropic phenotype of the *lec1* mutant, supports the multifunctional roles of *LEC1* in plant development (Junker and Baumlein, 2012; Supplementary Figure S1A); however, less is known about the downstream proteins of the *LEC1*-mediated regulatory process. A recent report indicated that *LEC1* might regulate directly a number of genes involved in light, phytohormone signaling, and embryogenesis, including several bHLH and MYB transcription factors (Junker et al., 2012). However, using a chromatin immunoprecipitation assay, we observed that *LEC1* does not target the selected trichome-related genes directly (data not shown), indicating that these genes might not act immediately downstream of *LEC1*. Instead, we observed that *LEC1* interacts with the trichome development-related transcription repressors *TCL1*, *TCL2*, *CPC*, and *ETC1* *in vitro*. *TCL1* was reported to be highly expressed in developing seeds and its protein targets the *GL1* promoter, while *TCL2* is highly expressed in cotyledons, rather than in developing

seeds, implying distinct roles of these two homologous genes in trichome regulation (Wang et al., 2007; Gan et al., 2011). The interactions between *LEC1* and these repressors provide a possible mechanism by which *LEC1* regulates embryonic cell fate determination spatio-temporally, via coupling with different transcription factors. Nevertheless, whether *LEC1* and *TCL2* together determine the post-embryonic trichome initiation requires further investigation.

Phytohormones serve as crucial endogenous integrators to mediate plant development by controlling precisely exact cell division and differentiation (Lumba et al., 2010; Vanstraelen and Benkova, 2012). Gibberellin (GA) has been reported to act as a positive endogenous factor to promote trichome formation, because the GA biosynthesis defective mutant *ga1* displays fewer trichomes on its leaves (Perazza et al., 1998; Qi et al., 2014). *LEC2* (Stone et al., 2001) and *FUSCA3* (*FUS3*, Keith et al., 1994), two B3 domain-containing transcription factors, also defined as the *LEC* genes, promote *GL1* expression via direct activation of the GA biosynthesis gene, *ATGA3ox2*. When crossed to *ga1*, the ectopic trichome formation on *lec2* and *fus3* cotyledons was completely suppressed (Curaba et al., 2004). However, the ectopic trichomes were not abolished in *lec1 ga1* double mutant, which suggested that *LEC1*-mediated trichome formation is independent of GA (Curaba et al., 2004). Given that *LEC2* and *FUS3* act as potential downstream candidates of *LEC1* (Junker et al., 2010), GA's effect on the function of *LEC* factors might be under the control of a complicated regulatory network. Abscisic acid is also involved in *LEC1*-mediated trichome formation. A previous study reported that the conversion of leaves into cotyledon-like structures is enhanced in the *LEC1* overexpression line when coupled with abscisic acid treatment (Junker et al., 2012). In addition, a number of abscisic acid-responsive genes were shown to be the direct targets of *LEC1* (Junker et al., 2012). Nevertheless, the role of phytohormones in *LEC1*-mediated cell fate determination remains elusive and requires further investigation.

Accurate control of *LEC1* expression is critical for plants to regulate cell fate determination during development. Epigenetic regulation plays key roles in the exclusion of *LEC1* in the vegetative growth stage. *PICKLE*, a CHD3 chromatin-remodeling factor, represses *LEC1* transcriptional activation in vegetative tissue by modifying histone methylation. In the *pickle* mutant, the seedling root produced an embryonic structure, termed the "PICKLE root", caused by the ectopic expression of *LEC1* (Ogas et al., 1997, 1999; Li et al., 2005). In addition, the polycomb group proteins *CLF* and *SWN* were demonstrated to couple with *PICKLE* to repress the conversion from vegetative tissue to an embryonic structure (Chanvivattana et al., 2004; Aichinger et al., 2009). Histone deacetylases (e.g., *HDA6* and *HDA19*) also contribute to the repression of embryonic properties by regulating several embryogenesis regulators, including *LEC1* (Tanaka et al., 2008). In addition to these epigenetic factors, some transcription factors, such as *VP1/ABI3-LIKE* and *MYB115/118*, were also reported as negative regulators of *LEC1* (Suzuki et al., 2007; Wang et al., 2009; Jia et al., 2013). Embryonic development originates from the zygote after fertilization in higher plants (Lau et al., 2012). High

expression during embryogenesis indicates that LEC1 is released from the suppressing environment of the vegetative tissue, thus allowing it to control embryonic and post-embryonic cell fate determination.

CONCLUSION

In this study, we revealed that LEC1 interacts with other transcription factors and regulates trichome-related gene expression to repress the trichome formation on cotyledons. These findings provide evidence that supports the hypothesis that LEC1, probably by integrating phytohormone signals and epigenetic regulation, mediates cell fate determination during the post-embryonic development phase, providing new insights into the role of LEC1 beyond embryogenesis.

ACCESSION NUMBERS

Sequence data from this article can be found in the *Arabidopsis* Genome Initiative database under the following accession numbers: *LEC1* (AT1G21970), *GL1* (AT3G27920), *GL2* (AT1G79840), *GL3* (AT5G41315), *EGL3* (AT1G63650), *TCL1* (AT2G30432), *TCL2* (AT2G30424), *TRY* (AT5G53200), *CPC* (AT2G46410), *ETC1* (AT1G01380), *ETC2* (AT2G30420), and *TUB2* (AT5G62690).

REFERENCES

- Aichinger, E., Villar, C. B. R., Farrona, S., Reyes, J. C., Hennig, L., and Kohler, C. (2009). CHD3 proteins and polycomb group proteins antagonistically determine cell identity in *Arabidopsis*. *PLoS Genet.* 5:605. doi: 10.1371/journal.pgen.1000605
- Brocard-Gifford, I. M., Lynch, T. J., and Finkelstein, R. R. (2003). Regulatory networks in seeds integrating developmental, abscisic acid, sugar, and light signaling. *Plant Physiol.* 131, 78–92. doi: 10.1104/pp.011916
- Chandler, J. W. (2008). Cotyledon organogenesis. *J. Exp. Bot.* 59, 2917–2931. doi: 10.1093/jxb/Ern167
- Chanvivattana, Y., Bishopp, A., Schubert, D., Stock, C., Moon, Y. H., Sung, Z. R., et al. (2004). Interaction of polycomb-group proteins controlling flowering in *Arabidopsis*. *Development* 131, 5263–5276. doi: 10.1242/Dev.01400
- Curaba, J., Moritz, T., Blervaque, R., Parcy, F., Raz, V., Herzog, M., et al. (2004). AtGA3ox2, a key gene responsible for bioactive gibberellin biosynthesis, is regulated during embryogenesis by LEAFY COTYLEDON2 and FUSCA3 in *Arabidopsis*. *Plant Physiol.* 136, 3660–3669. doi: 10.1104/pp.104.047266
- Gan, L. J., Xia, K., Chen, J. G., and Wang, S. C. (2011). Functional characterization of TRICHOMELESS2, a new single-repeat R3 MYB transcription factor in the regulation of trichome patterning in *Arabidopsis*. *BMC Plant Biol.* 11:176. doi: 10.1186/1471-2229-11-176
- Harada, J. J. (2001). Role of *Arabidopsis* LEAFY COTYLEDON genes in seed development. *J. Plant Physiol.* 158, 405–409. doi: 10.1078/0176-1617-00351
- Hulskamp, M., Misera, S., and Jurgens, G. (1994). Genetic dissection of trichome cell-development in *Arabidopsis*. *Cell* 76, 555–566. doi: 10.1016/0092-8674(94)90118-X
- Jia, H. Y., Mccarty, D. R., and Suzuki, M. (2013). Distinct roles of LAF1 network genes in promoting the embryonic seedling fate in the absence of VAL repression. *Plant Physiol.* 163, 1293–1305. doi: 10.1104/pp.113.220988
- Junker, A., and Baumlein, H. (2012). Multifunctionality of the LEC1 transcription factor during plant development. *Plant Signal. Behav.* 7, 1718–1720. doi: 10.4161/psb.22365

AUTHOR CONTRIBUTIONS

MH and XH designed the research. MH, YH, XL, and YL performed experiments. MH, YH, and XH analyzed data. MH and XH wrote the paper.

FUNDING

This work was supported by grants from the National Science Foundation of China (No. 31370342, 31301055, and 31300239) and “Hundred Talents” program of the Chinese Academy of Sciences.

ACKNOWLEDGMENT

We thank Shucui Wang for providing *gl2* mutant seeds, the *Arabidopsis* Biological Resource Centre for *lec1-4* seeds, and Jianru Zuo for *pER10:LEC1-MYC* plasmid.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2015.00955>

- Junker, A., Hartmann, A., Schreiber, F., and Baumlein, H. (2010). An engineer's view on regulation of seed development. *Trends Plant Sci.* 15, 303–307. doi: 10.1016/j.tplants.2010.03.005
- Junker, A., Monke, G., Rutten, T., Keilwagen, J., Seifert, M., Thi, T. M., et al. (2012). Elongation-related functions of LEAFY COTYLEDON1 during the development of *Arabidopsis thaliana*. *Plant J.* 71, 427–442. doi: 10.1111/j.1365-313X.2012.04999.x
- Keith, K., Kraml, M., Dengler, N. G., and Mccourt, P. (1994). *fusca3* – a heterochronic mutation affecting late embryo development in *Arabidopsis*. *Plant Cell* 6, 589–600. doi: 10.1105/tpc.6.5.589
- Kwong, R. W., Bui, A. Q., Lee, H., Kwong, L. W., Fischer, R. L., Goldberg, R. B., et al. (2003). LEAFY COTYLEDON1-LIKE defines a class of regulators essential for embryo development. *Plant Cell* 15, 5–18. doi: 10.1105/tpc.006973
- Lau, S., Slane, D., Herud, O., Kong, J. X., and Jurgens, G. (2012). Early embryogenesis in flowering plants: setting up the basic body pattern. *Annu. Rev. Plant Biol.* 63, 483–506. doi: 10.1146/annurev-arplant-042811-105507
- Le, B. H., Cheng, C., Bui, A. Q., Wagmaster, J. A., Henry, K. F., Pelletier, J., et al. (2010). Global analysis of gene activity during *Arabidopsis* seed development and identification of seed-specific transcription factors. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8063–8070. doi: 10.1073/pnas.1003530107
- Lee, H. S., Fischer, R. L., Goldberg, R. B., and Harada, J. J. (2003). *Arabidopsis* LEAFY COTYLEDON1 represents a functionally specialized subunit of the CCAAT binding transcription factor. *Proc. Natl. Acad. Sci. U.S.A.* 100, 2152–2156. doi: 10.1073/pnas.0437909100
- Li, H. C., Chuang, K., Henderson, J. T., Rider, S. D., Bai, Y., Zhang, H., et al. (2005). PICKLE acts during germination to repress expression of embryonic traits. *Plant J.* 44, 1010–1022. doi: 10.1111/j.1365-313X.2005.02602.x
- Lotan, T., Ohto, M., Yee, K. M., West, M. A., Lo, R., Kwong, R. W., et al. (1998). *Arabidopsis* LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* 93, 1195–1205. doi: 10.1016/S0092-8674(00)81463-4
- Lumba, S., Cutler, S., and Mccourt, P. (2010). Plant nuclear hormone receptors: a role for small molecules in protein-protein interactions. *Annu. Rev. Cell Dev. Biol.* 26, 445–469. doi: 10.1146/annurev-cellbio-100109-103956

- Marks, M. D., and Feldmann, K. A. (1989). Trichome development in *Arabidopsis-thaliana*. 1. T-DNA tagging of the *Glabrous1* gene. *Plant Cell* 1, 1043–1050. doi: 10.1105/tpc.1.11.1043
- Meinke, D. W. (1992). A homeoctic mutant of *Arabidopsis thaliana* with leafy cotyledons. *Science* 258, 1647–1650. doi: 10.1126/science.258.5088.1647
- Meinke, D. W., Franzmann, L. H., Nickle, T. C., and Yeung, E. C. (1994). Leafy cotyledon mutants of *Arabidopsis*. *Plant Cell* 6, 1049–1064. doi: 10.1105/tpc.6.8.1049
- Morohashi, K., Zhao, M. Z., Yang, M. L., Read, B., Lloyd, A., Lamb, R., et al. (2007). Participation of the *Arabidopsis* bHLH factor GL3 in trichome initiation regulatory events. *Plant Physiol.* 145, 736–746. doi: 10.1104/pp.107.104521
- Mu, J., Tan, H., Zheng, Q., Fu, F., Liang, Y., Zhang, J., et al. (2008). LEAFY COTYLEDON1 is a key regulator of fatty acid biosynthesis in *Arabidopsis*. *Plant Physiol.* 148, 1042–1054. doi: 10.1104/pp.108.126342
- Ogas, J., Cheng, J. C., Sung, Z. R., and Somerville, C. (1997). Cellular differentiation regulated by gibberellin in the *Arabidopsis thaliana* pickle mutant. *Science* 277, 91–94. doi: 10.1126/science.277.5322.91
- Ogas, J., Kaufmann, S., Henderson, J., and Somerville, C. (1999). PICKLE is a CHD3 chromatin-remodeling factor that regulates the transition from embryonic to vegetative development in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A* 96, 13839–13844. doi: 10.1073/pnas.96.24.13839
- Oppenheimer, D. G., Herman, P. L., Sivakumaran, S., Esch, J., and Marks, M. D. (1991). A myb gene required for leaf trichome differentiation in *Arabidopsis* is expressed in stipules. *Cell* 67, 483–493. doi: 10.1016/0092-8674(91)90523-2
- Pattanaik, S., Patra, B., Singh, S. K., and Yuan, L. (2014). An overview of the gene regulatory network controlling trichome development in the model plant, *Arabidopsis*. *Front. Plant Sci.* 5:259. doi: 10.3389/Fpls.2014.00259
- Perazza, D., Vachon, G., and Herzog, M. (1998). Gibberellins promote trichome formation by up-regulating GLABROUS1 in *Arabidopsis*. *Plant Physiol.* 117, 375–383. doi: 10.1104/PP.117.2.375
- Perianez-Rodriguez, J., Manzano, C., and Moreno-Risueno, M. A. (2014). Post-embryonic organogenesis and plant regeneration from tissues: two sides of the same coin? *Front. Plant Sci.* 5:219. doi: 10.3389/Fpls.2014.00219
- Peris, C. I. L., Rademacher, E. H., and Weijers, D. (2010). Green beginnings - pattern formation in the early plant embryo. *Plant Dev.* 91, 1–27. doi: 10.1016/S0070-2153(10)91001-6
- Qi, T. C., Huang, H., Wu, D. W., Yan, J. B., Qi, Y. J., Song, S. S., et al. (2014). *Arabidopsis* DELLA and JAZ proteins bind the WD-Repeat/bHLH/MYB complex to modulate gibberellin and jasmonate signaling synergy. *Plant Cell* 26, 1118–1133. doi: 10.1105/tpc.113.121731
- Rerie, W. G., Feldmann, K. A., and Marks, M. D. (1994). The *Glabra2* gene encodes a homeo domain protein required for normal trichome development in *Arabidopsis*. *Genes Dev.* 8, 1388–1399. doi: 10.1101/gad.8.12.1388
- Stone, S. L., Kwong, L. W., Yee, K. M., Pelletier, J., Lepiniec, L., Fischer, R. L., et al. (2001). LEAFY COTYLEDON2 encodes a B3 domain transcription factor that induces embryo development. *Proc. Natl. Acad. Sci. U.S.A.* 98, 11806–11811. doi: 10.1073/pnas.201413498
- Suzuki, M., Wang, H. H. Y., and McCarty, D. R. (2007). Repression of the LEAFY COTYLEDON 1/B3 regulatory network in plant embryo development by VP1/ABSCISIC ACID INSENSITIVE 3-LIKE B3 genes. *Plant Physiol.* 143, 902–911. doi: 10.1104/pp.106.092320
- Tanaka, M., Kikuchi, A., and Kamada, H. (2008). The *Arabidopsis* histone deacetylases HDA6 and HDA19 contribute to the repression of embryonic properties after germination. *Plant Physiol.* 146, 149–161. doi: 10.1104/pp.107.111674
- Vanstraelen, M., and Benkova, E. (2012). Hormonal interactions in the regulation of plant development. *Annu. Rev. Cell Dev. Biol.* 28, 463–487. doi: 10.1146/annurev-cellbio-101011-155741
- Wang, S. C., Barron, C., Schiefelbein, J., and Chen, J. G. (2010). Distinct relationships between *GLABRA2* and single-repeat R3 MYB transcription factors in the regulation of trichome and root hair patterning in *Arabidopsis*. *New Phytol.* 185, 387–400. doi: 10.1111/j.1469-8137.2009.03067.x
- Wang, S. C., and Chen, J. G. (2014). Regulation of cell fate determination by single-repeat R3 MYB transcription factors in *Arabidopsis*. *Front. Plant Sci.* 5:133. doi: 10.3389/Fpls.2014.00133
- Wang, S. C., Hubbard, L., Chang, Y., Guo, J. J., Schiefelbein, J., and Chen, J. G. (2008). Comprehensive analysis of single-repeat R3 MYB proteins in epidermal cell patterning and their transcriptional regulation in *Arabidopsis*. *BMC Plant Biol.* 8:81. doi: 10.1186/1471-2229-8-81
- Wang, S. C., Kwak, S. H., Zeng, Q. N., Ellis, B. E., Chen, X. Y., Schiefelbein, J., et al. (2007). TRICHOMELESS1 regulates trichome patterning by suppressing *GLABRA1* in *Arabidopsis*. *Development* 134, 3873–3882. doi: 10.1242/Dev.009597
- Wang, X. C., Niu, Q. W., Teng, C., Li, C., Mu, J. Y., Chua, N. H., et al. (2009). Overexpression of *PGA37/MYB118* and *MYB115* promotes vegetative-to-embryonic transition in *Arabidopsis*. *Cell Res.* 19, 224–235. doi: 10.1038/Cr.2008.276
- Warpeha, K. M., Upadhyay, S., Yeh, J., Adamiak, J., Hawkins, S. I., Lapik, Y. R., et al. (2007). The GCR1, GPA1, PRN1, NF-Y signal chain mediates both blue light and abscisic acid responses in *Arabidopsis*. *Plant Physiol.* 143, 1590–1600. doi: 10.1104/pp.106.089904
- West, M., Yee, K. M., Danao, J., Zimmerman, J. L., Fischer, R. L., Goldberg, R. B., et al. (1994). LEAFY COTYLEDON1 is an essential regulator of late embryogenesis and cotyledon identity in *Arabidopsis*. *Plant Cell* 6, 1731–1745. doi: 10.1105/tpc.6.12.1731
- Yamamoto, A., Kagaya, Y., Toyoshima, R., Kagaya, M., Takeda, S., and Hattori, T. (2009). *Arabidopsis* NF-YB subunits LEC1 and LEC1-LIKE activate transcription by interacting with seed-specific ABRE-binding factors. *Plant J.* 58, 843–856. doi: 10.1111/j.1365-313X.2009.03817.x
- Yang, C. X., and Ye, Z. B. (2013). Trichomes as models for studying plant cell differentiation. *Cell. Mol. Life Sci.* 70, 1937–1948. doi: 10.1007/s00018-012-1147-6
- Zhao, M., Morohashi, K., Hatlestad, G., Grotewold, E., and Lloyd, A. (2008). The TGT1-bHLH-MYB complex controls trichome cell fate and patterning through direct targeting of regulatory loci. *Development* 135, 1991–1999. doi: 10.1242/Dev.016873
- Zhou, L. M., Zheng, K. J., Wang, X. Y., Tian, H. N., Wang, X. L., and Wang, S. C. (2014). Control of trichome formation in *Arabidopsis* by poplar single-repeat R3 MYB transcription factors. *Front. Plant Sci.* 5:262. doi: 10.3389/Fpls.2014.00262

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Huang, Hu, Liu, Li and Hou. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.