

Comparison of clinicopathologic features and survival between eastern and western population with esophageal squamous cell carcinoma

Jie Zhang^{1,2*}, Yizhou Jiang^{1*}, Chunxiao Wu^{3*}, Shuang Cai¹, Rui Wang⁴, Ying Zhen³, Sufeng Chen^{1,2}, Kuaile Zhao⁵, Yangle Huang¹, James Luketich⁶, Haiquan Chen^{2,4}

¹Department of Oncology, Fudan University Shanghai Medical College, Shanghai 200032, China; ²Department of Thoracic Surgery, Fudan University Shanghai Cancer Center, Shanghai 200032, China; ³Department of Cancer Control & Prevention, Shanghai Municipal Center for Disease Control & Prevention, Shanghai 200336, China; ⁴Department of Thoracic Surgery, Shanghai Chest Hospital, Shanghai Jiaotong University, Shanghai 200032, China; ⁵Department of Radiotherapy, Fudan University Shanghai Cancer Center, Shanghai 200032, China; ⁶Department of Cardiothoracic Surgery, University of Pittsburgh Medical Center, Pittsburgh 15213, USA

Contributions: (I) Conception and design: J Zhang, Y Jiang, C Wu, R Wang, H Chen; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: J Zhang, Y Jiang; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*These authors contributed equally to this work.

Correspondence to: Haiquan Chen, MD. Department of Thoracic Surgery, Fudan University Shanghai Cancer Center, 399 Lingling Street, Shanghai 200032, China. Email: hqchen1@yahoo.com; Jie Zhang, MD. Department of Thoracic Surgery, Fudan University Shanghai Cancer Center, 399 Lingling Street, Shanghai 200032, China. Email: zhangjie2289@hotmail.com.

Background: Esophageal squamous cell carcinoma (ESCC) is the major histologic subtype of esophageal cancer, characterized by a high mortality rate and geographic differences in incidences. It is unknown whether there is difference between “eastern” ESCC and “western” ESCC. This study is attempted to demonstrate the hypothesis by comparing ESCC between Chinese residents and Caucasians living in the US.

Methods: The data sources of this study are from United States SEER limited-use database and Shanghai Cancer Registries by Shanghai Municipal Center for Disease Control (SMCDC). Consecutive, non-selected patients with pathologically diagnosed ESCC, between January 1, 2002 and December 31, 2006, were included in this analysis. 1-year, 3-year and 5-year survival estimates were computed and compared between two populations. A Cox proportional hazards model was used to determine factors affecting survival differences.

Results: A total of 1,718 Chinese, 1,624 Caucasians ESCC patients with individual American Joint Commission on Cancer (AJCC) staging information were included in this study. The Caucasian group had a significantly higher proportion of female patients than Chinese (38.24% *vs.* 18.68% $P < 0.01$). ESCC was diagnosed in Chinese patients at an earlier age and stage than Caucasians. Generally, Chinese patients had similar overall survival rate with Caucasian by both univariate and multivariate analysis. Overall survival was significantly worse only in male Caucasians compared to Chinese patients (median survival time, 12.4 *vs.* 14.5 months, $P < 0.01$, respectively).

Conclusions: ESCC from eastern and western countries might have some different features. These differences need to be taken into account for the management of ESCC patients in different ethnic groups.

Keywords: Esophageal squamous cell carcinoma (ESCC); race; prognostic factors; survival

Submitted Jul 09, 2015. Accepted for publication Sep 21, 2015.

doi: 10.3978/j.issn.2072-1439.2015.10.39

View this article at: <http://dx.doi.org/10.3978/j.issn.2072-1439.2015.10.39>

Introduction

Approximately 460,000 people are diagnosed with esophageal cancer worldwide annually, with ~380,000 people die of the disease. This makes it the eighth most common cancer and the sixth most common cause of cancer mortality (1). Esophageal squamous cell carcinoma (ESCC) is the major histologic subtype and is characterized by a high mortality rate and geographic differences in incidence (1-3). To date, there has been no comparative study for ESCC between eastern and western countries.

ESCC occurs more commonly in persons of lower socioeconomic status so that the availability of medical treatment is problematic (4). If there are differences in incidence and outcome, it could be as a result of differences in ethnic groups, living conditions, socioeconomic strata, treatment options and choices, and differences in the health care systems (5-9). A recent large genome-wide association study (GWAS) of ESCC has identified two susceptibility genes, *PLCE1* and *C20orf54*, which were highly associated with ESCC in the Chinese population. *PLCE1* regulates cell growth, differentiation, apoptosis and angiogenesis. *C20orf54* is responsible for transporting riboflavin, and deficiency of riboflavin has been documented as a risk factor for ESCC. *PLCE1* and *C20orf54* have important biological implications for ESCC and might account for the pathogenesis of ESCC (10). However, to date, no genetic susceptibility to ESCC in Caucasians has been identified.

The goals of our study were as follows: First, to use registry data to provide a detailed picture of ESCC cases, describing gender, age, primary site, stage, grade, and prognosis. Second is to compare ESCC survival for Chinese resident patients and Caucasians living in the US.

Materials and methods

Data source

Data for this analysis came from two sources. Data for Caucasians were from the Surveillance, Epidemiology and End Results (SEER) limited-use database. SEER collects cancer incidence and survival data from population-based cancer registries covering approximately 26% of the U.S. population. The routine data collection includes detailed information on demographics, diagnosis, tumor characteristics and treatment information (11). Data for Chinese were from Shanghai Cancer Registry (SCR) compiled by Shanghai Municipal Center for Disease Control (SMCDC) (12). A total of 175 hospitals, qualified

for cancer treatment, in the greater Shanghai area are responsible for reporting all newly diagnosed cancer cases, using a standardized report card. A patient consent form was obtained, following an approved, institutional-reviewed protocol (13). Death certificates for all cancer patients are obtained monthly from the Vital Statistics Section of the SMCDC.

Subjects

From the SEER database, we identified 1,624 Caucasians (tumor site codes 15.0 to 15.5, 15.8, 15.9 and ICD-O-3 morphology code 8070/3), who were diagnosed between January 1st, 2002 and December 31st, 2006, with individual American Joint Commission on Cancer (AJCC) staging information. Similarly, we identified 1,718 Chinese ESCC patients with staging information according to the same criteria. Thus, two groups of patients constitute the subjects of this analysis: Chinese group (1,718 patients) and Caucasian group (1,624 patients). The end follow-up data was December 31, 2009. For the Chinese group, the mean and median follow-up times were 755 and 435 days, respectively (range, 21-1,410 days). For the Caucasian group, the mean and median follow-up times were 725 and 456 days, respectively (range, 29-1,401 days). Like most of population based study, disease-free survival is not available in this study.

Definition of variables

The variables in this analysis included race/ethnicity, gender, age group, tumor stage and primary site. Tumor stage at diagnosis was defined according to the AJCC staging classification, 6th edition. Primary site of the tumor was defined by ICD-O-3 codes and was classified as cervical esophagus (C15.0), upper third of esophagus (C15.3), middle third of esophagus (C15.1, C15.4), lower third of esophagus (C15.2, C15.5), overlapping lesions (C15.8) of esophagus, NOS (C15.9) (2).

Statistical analysis

Differences in the distribution of gender, age group, tumor stage and primary site between two groups were evaluated using the Chi-square test (14).

One-year, 3-year and 5-year overall survival rate and median survival time were calculated using the life-table method. ESCC-specific survival curves were calculated

Table 1 Tumor characteristics of ESCC patients

Variable	Chinese resident (N=1,718)		Caucasians (N=1,624)		P
	Freq	%	Freq	%	
Gender					<0.01
Male	1,397	81.32	1,003	61.76	
Female	321	18.68	621	38.24	
Age group					<0.01
<60	592	34.46	353	21.74	
60~69	470	27.36	450	27.71	
70+	656	38.18	821	50.55	
AJCC tumor stage					<0.01
1, 2	868	50.52	728	44.83	
3, 4	850	49.48	896	55.17	
Primary site					<0.01
Cervical esophagus	3	0.17	101	6.22	
Upper esophagus	63	3.67	206	12.69	
Middle esophagus	585	34.05	595	36.64	
Lower esophagus	282	16.41	497	30.60	
Overlapping lesion of esophagus	158	9.20	91	5.60	
Esophagus, NOS	627	36.55	134	8.25	

ESCC, esophageal squamous cell carcinoma; AJCC, American Joint Commission on Cancer.

according to the Kaplan-Meier method and compared by the log-rank test (12).

To explain possible survival differences and identify factors affecting survival, the Cox proportional hazards model was used. For each population group, multivariate associations of gender, age group, tumor stage and primary site with survival were determined. The assumption of proportional hazards for Cox models was checked by plotting the log of the negative log of the survival density functions *vs.* the log of survival time. The plotted lines were roughly parallel over time and no violations of the proportional hazards assumption were found (8).

All analyses were performed with the SPSS version 17.0 (SPSS Inc., Chicago, IL, USA). All P values were two-tailed and $P < 0.05$ was considered statistically significant.

Results

Tumor characteristics of ESCC patients from the two populations

The distribution of cases by gender, age group, primary site and stage, is shown for each population in *Table 1*. The

Caucasian group had a significantly higher proportion of female patients than Chinese (38.24% *vs.* 18.68%, $P < 0.01$). Chinese patients were younger, with only 31.18% of patient above 70 years old while Caucasians with 50.55% above 70 years ($P < 0.01$). A total of 49.48% patients were diagnosed in the advanced stages (stages III and IV) in the Chinese resident population, while the majority of Caucasians (55.17%) presented with advanced stage disease ($P < 0.01$).

Survival analysis by age, gender and tumor stage

One-year, 3-year and 5-year overall survival rate and median survival time are shown in *Table 2*. Median survival time was within a narrow range (14.2-15.3 months) in the two populations. Generally, Chinese patients had similar overall survival rate with Caucasians, (median survival time, 15.3 *vs.* 14.2 months, $P = 0.40$). Overall survival was significantly lower in male (but not female) patients from Caucasians compared to Chinese patients (median survival time, 12.8 *vs.* 14.5 months, $P < 0.01$, respectively, *Table 3*). No statistically significant difference in survival was seen among patients of each age group or tumor stage from the two populations (*Tables 2-4*).

Table 2 Survival analysis of ESCC patients, Chinese resident and Caucasians by age

Variable	Chinese resident				Caucasians				P
	1-year	3-year	5-year	Median survival time (m)	1-year	3-year	5-year	Median survival time (m)	
Survival rate (%)	55.2	28.4	13.0	15.3	42.7	19.0	13.8	14.2	0.13
Age									
<60	56.1	28.4	13.3	15.4	49.1	22.0	18.6	13.2	0.18
60~69	60.3	33.5	15.2	17.3	49.0	23.4	16.6	16.0	0.13
70+	50.2	24.8	10.0	13.1	36.4	15.1	10.0	12.4	0.08

ESCC, esophageal squamous cell carcinoma.

Table 3 Survival analysis of ESCC patients, Chinese resident and Caucasians by gender

Variable	Chinese resident				Caucasians				P
	1-year	3-year	5-year	Median survival time (m)	1-year	3-year	5-year	Median survival time (m)	
Survival rate (%)	55.2	28.4	13.0	15.3	42.7	19.0	13.8	14.2	0.13
Gender									
Male	53.2	26.7	17.1	14.5	40.4	17.0	12.0	12.8	<0.01
Female	62.5	36.6	6.3	18.5	46.5	22.2	16.5	16.9	0.14

ESCC, esophageal squamous cell carcinoma.

Table 4 Survival analysis of ESCC patients, Chinese resident and Caucasians by stage

Variable	Chinese resident				Caucasians				P
	1-year	3-year	5-year	Median survival time (m)	1-year	3-year	5-year	Median survival time (m)	
Survival rate (%)	55.2	28.4	13.0	15.3	42.7	19.0	13.8	14.2	0.13
Stage									
1	60.8	34.3	19.4	16.3	59.7	33.9	19.6	20.8	0.09
2	71.4	41.5	17.0	24.4	60.4	31.6	19.4	22.6	0.14
3	49.3	21.2	9.5	12.8	46.3	22.1	10.2	10.8	0.12
4	30.0	9.7	3.5	7.9	24.4	6.6	4.5	8.0	0.08

ESCC, esophageal squamous cell carcinoma.

Multivariate analysis in ESCC patients

Among age, sex, stage and, tumor site, age and stage were found to be independent prognostic factors in each population after adjustment using the Cox proportional hazards model ($P < 0.01$).

A Cox proportional hazards model built jointly for all ESCC patients suggested no significant advantage in survival of Chinese compared to Caucasian patients [relative risk (RR) = 1.10, $P = 0.21$]. Multivariate analysis disclosed age and stage were independent prognostic factors in all ESCC patients ($P < 0.01$, respectively, *Table 5*).

Multivariate analysis in all male ESCC patients

Since univariate analysis showed overall survival was

significantly lower in male patients from Caucasians compared to Chinese patients, multivariate analysis was further performed to confirm this conclusion. Multivariate analysis among all male ESCC patients suggested age and stage were independent prognostic factors ($P < 0.01$). Compared to male Chinese, Caucasians had a worse prognosis (RR = 1.54, $P < 0.01$, *Table 6*).

Discussion

ESCC remains the predominant histological subtype of esophageal cancer world-widely (2,3). However, previous studies had provided scant and discordant data about the differences in tumor characteristics and prognoses between different ethnical groups (15). Several interesting results

Table 5 Multivariate analysis among all ESCC patients

Variable	All ESCC patients (N=3,342)		
	RR	95% CI	P
Race/ethnicity			
Chinese resident	1.00	–	–
Caucasians	1.10	0.65-1.42	0.21
Gender			
Male	1.00	–	–
Female	0.98	0.79-1.16	0.29
Age group			
<60	1.00	–	–
60~69	1.28	1.13-1.32	<0.01
70+	2.49	2.22-2.86	<0.01
AJCC tumor stage			
1	1.00	–	–
2	1.18	0.65-1.62	0.34
3	6.21	5.47-6.89	<0.01
4	13.50	11.32-15.80	<0.01
Primary site			
Middle third of esophagus	1.00	–	–
Cervical esophagus	0.98	0.70-1.12	0.29
Upper third of esophagus	1.09	0.60-1.49	0.38
Lower third of esophagus	1.16	0.74-1.44	0.16
Overlapping lesion of esophagus	0.95	0.78-1.10	0.14
Esophagus, NOS	1.03	0.89-1.15	0.10

ESCC, esophageal squamous cell carcinoma; AJCC, American Joint Commission on Cancer; RR, relative risk; CI, confidence interval.

are disclosed in the current study: Chinese patients were diagnosed with ESCC at an earlier age and stage than Caucasians (Tables 2,6). Another interesting finding was that Chinese patients had similar overall survival rate with Caucasian by both univariate and multivariate analysis. The median survival time of male Chinese patients was significantly longer than that of Caucasians (Tables 3,6).

Compared to Caucasians, Chinese patients develop the disease at an earlier age and earlier stage for the entire population (Table 2) and for males-only (Table 6). This may be attributed to different causes of the disease between the populations. A familial aggregation tendency in high-risk areas suggests that genes may play a role in ESCC pathogenesis (16,17). As mentioned in Introduction, several genes have been considered to contribute to ESCC in Chinese, such as *PLCE1* and *C20orf54* (10), and *ADH1B*

Table 6 Multivariate analysis among all male ESCC patients

Variable	All male ESCC patients (N=2,400)		
	RR	95% CI	P
Race/ethnicity			
Male Chinese residents	1.00	–	–
Male Caucasians	1.54	1.26-1.88	<0.01
Age group			
<60	1.00	–	–
60~69	1.26	1.18-1.35	<0.01
70+	2.78	2.40-2.96	<0.01
AJCC tumor stage			
1	1.00	–	–
2	1.19	0.49-1.91	0.38
3	5.83	5.22-6.37	<0.01
4	12.15	9.82-14.26	<0.01
Primary site			
Middle third of esophagus	1.00	–	–
Cervical esophagus	0.94	0.64-1.21	0.33
Upper third of esophagus	1.13	0.73-1.61	0.45
Lower third of esophagus	1.09	0.89-1.25	0.19
Overlapping lesion of esophagus	0.89	0.68-1.14	0.25
Esophagus, NOS	0.95	0.85-1.06	0.13

ESCC, esophageal squamous cell carcinoma; AJCC, American Joint Commission on Cancer; RR, relative risk; CI, confidence interval.

and *ALDH2* in Japanese (18). This genetic susceptibility among Asian populations may partly explain the early onset in Chinese patients. There has been research concerning environmental factors as well. Some studies have demonstrated that smoking, alcohol consumption, and high body mass index (BMI) are more important risk factors of ESCC in western countries than in China (3,19,20). Thus, for the American patients, it would take longer to develop ESCC, reflecting the cumulative effects of these factors. On the other hand, for Chinese, nutritional imbalance, intake of pickled vegetables, nitrosamine-rich and mycotoxin-contaminated foods seem more important (21,22). Such factors may take less time to develop neoplasia.

We also noted that Chinese male patients lived longer than Caucasians (Tables 3,6). Younger patients usually have fewer comorbid diseases, less complications from treatment, may have more of an economic ability to afford treatment, and more willingness to invest in health. Thus, younger

patients could have better outcomes if their diseases are not too aggressive (23,24). Besides, a considerable percentage of the western patients get ESCC after radiotherapy on mediastinum for their lymphoma. These tumors resulting from radiation grow with more invasive and malignant biological characteristics. This may be another explanation of the relatively poorer prognoses of Caucasians.

We also noticed the difference in distribution between males and females. In comparison with Chinese, Caucasians had a significantly higher proportion of females (*Table 1*). This difference may partly be attributed to higher smoking and drinking prevalence among Caucasian women.

The strength of this study is that the analysis of these characteristics and comparing Chinese resident patients and Caucasians living in the US, has not previously been done. However, our study was limited to available variables and comparable databases, so that not all possible factors affecting survival were considered. In addition, socio-economic status, family history of ESCC, and lifestyle factors, such as alcohol consumption and smoking, were also not studied. Further prospective studies are also needed to define the genetic susceptibility for ESCC, especially in western countries.

Acknowledgements

Funding: We acknowledge supported by key technology support project of the field of medicine and agriculture science from Shanghai Municipal Science and Technology Commission (15411951600); Shanghai science and technology commission foundation key project; Longhua Medical Project (LYTD-25); Ministry of Education Returned Scientific Research Foundation; Trans-Century Training Programme Foundation for the Talents by the State Education Commission (to J Zhang).

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Parkin DM, Bray F, Ferlay J, et al. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;55:74-108.
2. Daly JM, Fry WA, Little AG, et al. Esophageal cancer: Results of an American College of Surgeons patient care evaluation study. *J Am Coll Surg* 2000;190:562-72; discussion 572-3.
3. Hongo M, Nagasaki Y, Shoji T. Epidemiology of esophageal cancer: Orient to Occident. Effects of chronology, geography and ethnicity. *J Gastroenterol Hepatol* 2009;24:729-35.
4. van Loon AJ, Brug J, Goldbohm RA, et al. Differences in Cancer Incidence and Mortality among Socioeconomic Groups. *Scand J Soc Med* 1995;23:110-20.
5. Shavers VL, Brown ML. Racial and ethnic disparities in the receipt of cancer treatment. *J Natl Cancer Inst* 2002;94:334-57.
6. Ayanian JZ, Zaslavsky AM, Fuchs CS, et al. Use of adjuvant chemotherapy and radiation therapy for colorectal cancer in a population-based cohort. *J Clin Oncol* 2003;21:1293-300.
7. Cronin-Fenton DP, Sharp L, Carsin AE, et al. Patterns of care and effects on mortality for cancers of the oesophagus and gastric cardia: A population-based study. *Eur J Cancer* 2007;43:565-75.
8. von Bueren AO, Shalaby T, Oehler-Jänne C, et al. RNA interference-mediated c-MYC inhibition prevents cell growth and decreases sensitivity to radio- and chemotherapy in childhood medulloblastoma cells. *BMC Cancer* 2009;9:10.
9. O'Malley CD, Le GM, Glaser SL, et al. Socioeconomic status and breast carcinoma survival in four racial/ethnic groups-A population-based study. *Cancer* 2003;97:1303-11.
10. Wang LD, Zhou FY, Li XM, et al. Genome-wide association study of esophageal squamous cell carcinoma in Chinese subjects identifies susceptibility loci at PLCE1 and C20orf54. *Nat Genet* 2010;42:759-U46.
11. Das A, Singh V, Fleischer DE, et al. A comparison of endoscopic treatment and surgery in early esophageal cancer: An analysis of surveillance epidemiology and end results data. *Am J Gastroenterol* 2008;103:1340-5.
12. Zhang J, Chen SF, Zhen Y, et al. Multicenter Analysis of Lung Cancer Patients Younger Than 45 Years in Shanghai. *Cancer* 2010;116:3656-62.
13. Bao PP, Zheng Y, Wang CF, et al. Time Trends and Characteristics of Childhood Cancer Among Children Age 0-14 in Shanghai. *Pediatr Blood Cancer* 2009;53:13-6.
14. Greenstein AJ, Litle VR, Swanson SJ, et al. Effect of the number of lymph nodes sampled on postoperative survival of lymph node-negative esophageal cancer. *Cancer* 2008;112:1239-46.
15. Esophageal cancer: epidemiology, pathogenesis and prevention. *Nat Clin Pract Gastroenterol Hepatol* 2008;5:517-26.

16. Guohong Z, Min S, Duenmei W, et al. Genetic heterogeneity of oesophageal cancer in high-incidence areas of southern and northern China. *PLoS One* 2010;5:e9668.
17. Zhang W, Bailey-Wilson JE, Li W, et al. Segregation analysis of esophageal cancer in a moderately high-incidence area of northern China. *Am J Hum Genet* 2000;67:110-9.
18. Cui R, Kamatani Y, Takahashi A, et al. Functional Variants in ADH1B and ALDH2 Coupled With Alcohol and Smoking Synergistically Enhance Esophageal Cancer Risk. *Gastroenterology* 2009;137:1768-75.
19. Tran GD, Sun XD, Abnet CC, et al. Prospective study of risk factors for esophageal and gastric cancers in the Linxian General Population Trial cohort in China. *Int J Cancer* 2005;113:456-63.
20. Chow WH, Blot WJ, Vaughn TL, et al. Body mass index and risk of adenocarcinomas of the esophagus and gastric cardia. *J Natl Cancer Inst* 1998;90:150-5.
21. Siassi F, Ghadirian P. Riboflavin deficiency and esophageal cancer: A case control-household study in the Caspian Littoral of Iran. *Cancer Detect Prev* 2005;29:464-9.
22. Wang J, Wang S, Su J, et al. Food contamination of fumonisin B1 in high-risk area of esophageal and liver cancer. *Toxicol Sci* 2003;72:188.
23. Steyerberg EW, Neville B, Weeks JC, et al. Referral patterns, treatment choices, and outcomes in locoregional esophageal cancer: A population-based analysis of elderly patients. *J Clin Oncol* 2007;25:2389-96.
24. Greenstein AJ, Litle VR, Swanson SJ, et al. Prognostic significance of the number of lymph node metastases in esophageal cancer. *J Am Coll Surg* 2008;206:239-46.

Cite this article as: Zhang J, Jiang Y, Wu C, Cai S, Wang R, Zhen Y, Chen S, Zhao K, Huang Y, Luketich J, Chen H. Comparison of clinicopathologic features and survival between eastern and western population with esophageal squamous cell carcinoma. *J Thorac Dis* 2015;7(10):1780-1786. doi: 10.3978/j.issn.2072-1439.2015.10.39