

Hepatorenal syndrome: Update on diagnosis and treatment

Olga Baraldi, Chiara Valentini, Gabriele Donati, Giorgia Comai, Vania Cuna, Irene Capelli,
Maria Laura Angelini, Maria Iliara Moretti, Andrea Angeletti, Fabio Piscaglia, Gaetano La Manna

Olga Baraldi, Chiara Valentini, Gabriele Donati, Giorgia Comai, Vania Cuna, Irene Capelli, Maria Laura Angelini, Maria Iliara Moretti, Andrea Angeletti, Gaetano La Manna, Department of Experimental, Diagnostic, Specialty Medicine, Nephrology, Dialysis, and Renal Transplant Unit, S. Orsola University Hospital, 40138 Bologna, Italy

Fabio Piscaglia, Division of Internal Medicine, Department of Digestive Diseases and Internal Medicine, Sant'Orsola-Malpighi Hospital, University of Bologna, 40138 Bologna, Italy

Author contributions: Baraldi O and La Manna G designed the aim of the article, wrote the manuscript and made critical revisions; Valentini C, Donati G, Comai G, Cuna V, Capelli I, Angelini ML, Moretti MI, Angeletti A, Piscaglia F contributed equally to this work, generated the figures and wrote the manuscripts.

Conflict-of-interest statement: The authors do not have any disclosures to report.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Gaetano La Manna, MD, PHD, Professor, Department of Experimental, Diagnostic, Specialty Medicine - Dialysis, Nephrology and Transplantation Unit, S. Orsola University Hospital, Via G. Massarenti 9, 40138 Bologna, Italy. gaetano.lamanna@unibo.it
Telephone: +39-51-2143255
Fax: +39-51-344439

Received: April 14, 2015
Peer-review started: April 15, 2015
First decision: May 13, 2015
Revised: August 13, 2015
Accepted: September 16, 2015
Article in press: September 18, 2015
Published online: November 6, 2015

Abstract

Acute kidney injury (AKI) is a common complication in patients with end-stage liver disease and advanced cirrhosis regardless of the underlying cause. Hepatorenal syndrome (HRS), a functional form of kidney failure, is one of the many possible causes of AKI. HRS is potentially reversible but involves highly complex pathogenetic mechanisms and equally complex clinical and therapeutic management. Once HRS has developed, it has a very poor prognosis. This review focuses on the diagnostic approach to HRS and discusses the therapeutic protocols currently adopted in clinical practice.

Key words: Hepatorenal syndrome; Cirrhosis; Acute kidney injury; Diagnosis; Treatment; Terlipressin; Liver support system

© The Author(s) 2015. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Hepatorenal syndrome is a functional and potentially reversible form of kidney failure. The pathophysiological bases of this disease are complex and not fully understood. The aim of this review is to focus the current diagnostic approach and the updated therapeutic protocols adopted in clinical practice.

Baraldi O, Valentini C, Donati G, Comai G, Cuna V, Capelli I, Angelini ML, Moretti MI, Angeletti A, Piscaglia F, La Manna G. Hepatorenal syndrome: Update on diagnosis and treatment. *World J Nephrol* 2015; 4(5): 511-520 Available from: URL: <http://www.wjgnet.com/2220-6124/full/v4/i5/511.htm> DOI: <http://dx.doi.org/10.5527/wjn.v4.i5.511>

PATHOGENESIS

Hepatorenal syndrome (HRS) can be considered the final stage of a pathophysiological condition charac-




Figure 1 Hepatorenal syndrome: Pathogenesis. In cirrhotic patients portal hypertension can lead to markedly dilated splanchnic arterial vessels. The bacterial translocation of intestinal germs, the gradual decrease in systemic vascular resistances, the hepatic vascular neoformation are potential risk factors. The fall in mean arterial pressure is compensated by increase in cardiac output and by activation of renin-angiotensin-aldosterone system (RAAS) and sympathetic nervous system (SNS) to improve systemic vascular resistance. The response mechanisms to the decreased effective circulating volume caused by Enhanced vascular capacitance (so-called “arterial underfilling”) include the non-osmotic release of vasopressin accounting for renal tubular sodium resorption and water retention leading to the onset of ascites, edema and hypervolemic hyponatremia. These compensatory mechanisms ultimately have repercussions on kidney function causing reduced glomerular filtration rate (GFR) and further water retention thereby worsening the water overload.

terized by decreased renal blood flow resulting from deteriorating liver function in patients with cirrhosis and ascites^[1-5].

Hemodynamic changes associated with endothelial shear stress occur before the onset of ascites and are sustained by an increase in pro-angiogenic factors like the vascular endothelial growth factor and platelet-derived growth factor and vasodilators (carbon monoxide, endocannabinoids and nitric oxide) able to promote the formation of hepatic, splanchnic and porto-systemic collateral vessels^[6-11] (Figure 1).

The ensuing hemodynamic instability may give rise to many clinical events that further interfere with the compensatory mechanisms. These include the onset of spontaneous bacterial peritonitis, gastrointestinal bleeding and post-paracentesis circulatory dysfunction^[12].

The renal impairment is worsened by a progressive cardiac dysfunction known as cirrhotic cardiomyopathy. The latter is characterized by diastolic impairment with septal ventricular hypertrophy, blunted ventricular response to stress, systolic and diastolic dysfunction, and electrophysiological abnormalities (prolongation of QT interval)^[7]. Systolic dysfunction is due to impairment of both β -adrenergic receptor and increasing in endogenous cannabinoids and cardiosuppressants such as nitric oxide and inflammatory cytokines and myocyte apoptosis. Furthermore it is possible that several intracellular signaling pathways are involved.

On the other hand the activation of renin-angiotensin system and salt retention play a role in diastolic

disfunction. Recent studies have stated myocardial dysfunction in cirrhosis as a contributing, or even a precipitant factor, of HRS^[13,14].

EPIDEMIOLOGY

According to Fede *et al.*^[15], approximately 20% of cirrhotic patients with diuretic-resistant ascites potentially develop HRS, while a prospective study by Ginès *et al.*^[4] on 229 patients with cirrhosis found an 18% incidence of HRS at one year, rising to 39% at five years after initial diagnosis.

HRS may also arise in patients with acute liver failure as shown in Akriviadis *et al.*^[16]: They considered 101 patients with alcoholic hepatitis of whom 28 developed HRS after a four-week follow-up. Planas *et al.*^[17], in a study enrolling 263 cirrhotic patients with a follow-up of 41 ± 3 mo after the onset of ascites, found prevalence rates of 2.6% and 5% for HRS types I and II respectively, with a cumulative probability of 11.4% at five years. The prevalence of HRS increases with liver disease progression, Wong *et al.*^[18] reporting a rate of 48% in patients on the waiting list for liver transplant.

Despite discrepancies in literature data, the prevalence of HRS has dropped in recent years, probably as a result of a better understanding of its pathophysiology and improved clinical management^[19]. Nonetheless the long-term survival of HRS patients remains poor and the only effective treatment for this condition is liver transplantation.

Table 1 Diagnostic criteria for hepatorenal syndrome

Cirrhosis with ascites
Serum Creatinine > 1.5 mg/dL
Absence of shock
No improvement of serum creatinine (decrease to a level of 1.5 mg/dL or less) after at least 2 d of diuretic withdraw and volume expansion with albumin (The recommended dose of albumin is 1 g/kg of body weight per day up to a maximum of 100 g/d)
No current or recent exposure to nephrotoxic drugs
Absence of parenchymal disease as indicated by proteinuria > 500 mg/d, microscopic hematuria (50 red blood cells per high power field) and abnormal renal ultrasonography

HRS: Hepatorenal syndrome.

DIAGNOSIS

The diagnostic criteria for HRS were initially defined by the International Ascites Club (IAC) in 1994^[20-22]. Since then, advances in our understanding of HRS pathogenesis and the introduction of new therapies led to repeated revisions of the criteria. The latest version of 2007 excludes the use of creatinine clearance (due to its poor correlation with kidney function in patients with cirrhosis), and has eliminated minor criteria (sodium excretion fraction, urinary output) deemed less sensitive and specific. Concomitant bacterial infection does not rule out a diagnosis of HRS but it is crucial to identify the absence of septic shock^[1] (Table 1).

Two forms of HRS, types I and II, have been described. They differ in severity and rate of progression and can be considered two separate clinico-pathological entities^[23] (Table 2).

Type I HRS is characterized by acute onset and rapidly progressing kidney failure with a doubling of serum creatinine to > 2.5 mg/dL (corresponding to a 50% reduction in the creatinine clearance rate) in less than 2 wk, usually associated with multiorgan damage. The prognosis is poor with only 10% of patients surviving longer than 90 d^[4].

This type of HRS can develop spontaneously but more often tends to follow a precipitating event, mostly spontaneous bacterial peritonitis or other infections like pneumonia, urinary tract infections or cellulitis^[24]. Other potential risk factors include viral, alcoholic, toxic or ischemic hepatitis (*e.g.*, TIPS), gastrointestinal bleeding and surgical procedures (Table 3).

Type II HRS represents the final kidney response to hemodynamic impairments in cirrhosis. This type presents as a less severe and more gradual decline in renal function associated with refractory ascites. The increase in creatinine is gradual with mean values of 1.5-2.0 mg/dL. Type II HRS predisposes patients to the development of type I HRS after a precipitating event. The average survival rate is six to eight months after onset.

The differential diagnosis between the two types of HRS is based on the rate of progression and extent of renal impairment, whereas the pathophysiological

Table 2 Characteristics of type I and type II hepatorenal syndrome

HRS I	Doubling of serum creatinine in < 2 wk	A precipitating event is present in the most of case	No history of diuretic resistant ascites	10% survival in 90 d without treatment
HRS II	Renal impairment gradually progressive	No precipitating events	Always ascites diuretic resistance	Median survival 6 mo

HRS: Hepatorenal syndrome.

differences have not yet been fully clarified.

A spontaneous recovery is rare in both cases unless there is a significant improvement in liver function.

The differential diagnosis between HRS, other causes of kidney disease and septic shock remain extremely difficult. Despite the widespread circulation of the IAC criteria, a serum creatinine cut-off of 1.5 mg/dL appears limited as it does not take into account its physiological fluctuations. In addition, creatinine values \leq 1.5 mg/dL may overestimate the true reduction in GFR^[25].

The AKI network (AKIN) has proposed a new definition of AKI for the diagnosis of HRS designed to implement the traditional IAC criteria for prompt recognition of kidney damage. AKI is defined as the abrupt loss of kidney function resulting in a 0.3 mg/dL increase in serum creatinine in 48 h or a 50% increase over the basal value. The aim is to apply the AKI criteria to decompensated cirrhotic patients for an early identification of kidney failure and thereby implementing prompt aggressive treatment^[26].

Two recent prospective studies assessed the applicability of the AKI criteria in patients with cirrhosis. The study by Fagundes *et al.*^[27] on 375 patients and another by Piano *et al.*^[28] on 233 cirrhotic patients both divided the populations into two groups based on kidney function. The first group comprised patients with a serum creatinine increase \geq 0.3 mg/dL but below the threshold of 1.5 mg/dL, whereas the second enrolled patients with creatinine > 1.5 mg/dL. In both cases renal decline and mortality rates were significantly higher in the group with serum creatinine > 1.5 mg/dL, with a lower probability of kidney disease regression. These results suggest that AKI with serum creatinine values < 1.5 mg/dL is a relatively benign and potentially reversible condition, whereas the progression of renal deterioration to a significant decrease in GFR (values > 1.5 mg/dL) carries a poor prognosis^[27,28].

Nonetheless, a recent editorial by Arroyo *et al.*^[29] pointed to a lack of evidence demonstrating the real advantage of the IAC guidelines with respect to AKI criteria. The stratification of cirrhotic patients according to single organ damage (kidney, liver or brain) appears to simplify the complex changes occurring in patients with decompensated liver failure.

Mindikoglu *et al.*^[2] proposed a new classification

Table 3 Risk factors for the onset of hepatorenal syndrome

Spontaneous bacterial peritonitis
Large volume paracentesis (> 5 L) with inadequate albumin substitution
NSAID and other nephrotoxic drugs, iv contrast
Bleeding from esophageal varices
Post TIPS syndrome
Diuretic treatment

Spontaneous bacterial peritonitis are leading trigger of HRS. One-third of patients with SBP develop HRS in the absence of septic shock. Diuretic treatment has been suggested as a potential trigger of HRS, but there are no clear supportive data for this. HRS: Hepatorenal syndrome; NSAID: Non-steroidal anti-inflammatory drug; TIPS: Transjugular intrahepatic portosystemic shunt.

associating GFR measurement and renal blood flow to stratify renal dysfunction, introducing the new concept of "pre-HRS", *i.e.*, patients with reduced renal blood flow but still normal or slightly reduced GFR. However, further studies are required to establish the clinical utility of this concept^[30].

In all patients with acute renal failure and even more in patients with cirrhosis, serum creatinine may not reflect the reduction of kidney function with a significant difference between male and female. Because of that it was proposed using cystatin C as alternative marker of renal function.

Seo *et al.*^[31] and Sharawey *et al.*^[32] showed that serum cystatin C level is a good marker for predicting HRS and survival in patients with cirrhotic ascites.

In the last 2 years the IAC organised a consensus development meeting in order to analyse the new definition of AKI in patients with cirrhosis and HRS: All the experts agreed on the removal of a fixed cut-off value of serum creatinine from the diagnostic criteria of HRS and they didn't suggest to evaluate Cystatin C determination^[33] (Table 1).

As there are currently no specific tests to identify HRS, diagnosis rests on the exclusion of other causes of kidney failure. It is important to establish the etiology of kidney injury in order to institute the appropriate treatment.

The onset of AKI in patients with cirrhosis enters into the differential diagnosis with other forms of kidney injury: Pre-renal (45%), organic, including acute tubular necrosis and glomerulonephritis (32%), and less frequently obstructive nephropathy (< 1%)^[34,35] (Table 4).

The parameters traditionally used to distinguish AKI from chronic kidney disease (CKD) (urinary sodium concentration, serum and urine osmolarity) are not applicable in patients with cirrhosis and ascites. Likewise, serum urea values are usually reduced in cirrhotic patients due to the impaired hepatic synthesis.

Belcher *et al.*^[36] proposed the use of urinary biomarkers of AKI to improve the diagnostic process: urinary levels of neutrophil gelatinase-associated lipocalin (NGAL), interleukin 18 (IL-18), kidney injury

Table 4 Differential diagnosis of renal failure in cirrhosis

Pre-renal	History of fluid loss, gastrointestinal bleeding, treatment with diuretics or non-steroidal anti-inflammatory drugs
Organic	Medical history, laboratory tests (cryoglobulinemia, complementemia, <i>etc.</i>)
Obstructive	Ultrasound imaging
Chronic kidney disease	Anemia, proteinuria, secondary hyperparathyroidism, ultrasound evidence of renal cortical thinning

molecule-1 and liver fatty acid-binding protein are elevated in liver disease patients with kidney injury due to acute tubular necrosis.

Two recent trials studied patients admitted to hospital for cirrhosis-induced complications. They both demonstrated that raised urinary levels of NGAL may serve to distinguish functional kidney damage from acute tubular necrosis or necrosis arising in HRS^[37,38].

Barreto *et al.*^[39] confirmed that urinary NGAL predicts clinical outcome, namely persistent kidney injury and mortality at three months in hospitalized patients with cirrhosis and bacterial infections. Although further clinical trials are required, NGAL appears to predict short-term mortality in cirrhotic patients.

Renal biopsy is not used for diagnostic purposes but can be entertained when a decline in renal function is associated with active urinary sediment or clinical status not corresponding to IAC criteria or unresponsive to therapy.

TREATMENT

Despite improvements in the clinical management of HRS patients in the past twenty years, currently available treatments enhance patients' short-term survival but offer little benefit in the longer term.

The current therapeutic armamentarium includes drugs with specific vasoconstrictive effects on the splanchnic circulation in addition to renal and liver replacement therapies which can be artificial or natural (liver transplantation). Liver transplant remains the only truly effective treatment but is limited by the high mortality rate in HRS patients and the shortage of available organs.

A recent literature review by Fabrizi *et al.*^[40] noted that pre-transplant kidney function is the most important predictor for patient survival after liver transplant. Pharmacological treatment and medical care serve as a "bridge" to transplant to improve the patient's prognosis.

Prevention and general patient management

The cirrhotic patient with ascites must be closely monitored to prevent and treat precipitating factors^[41-45] (Table 5).

If multiorgan damage is present, some patients, especially those with type I HRS, may require a high level of care, and admission to an intensive care facility. In addition, a patient-tailored diet and physical rehabi-

Table 5 Prevention of hepatorenal syndrome and general patient management strategies

Avoid drugs that reduce renal perfusion or nephrotoxic substances
Minimize exposure to organ-iodated contrast agents
Intravenous albumin is recommended for volemic filling after large volume paracentesis (8 g of albumin for each liter of ascites removed)
Diuretic therapy should be suspended
Pentoxifylline as drug's anti-TNF α activity
Antibiotic prophylaxis to prevent infections reducing intestinal bacterial translocation (norfloxacin 400 mg/d)
Intravenous albumin administered in association with ceftriaxone in SPB
Adrenal insufficiency should be identified and treated
Drug dosages must be adjusted according to renal function

ligation program should be planned and each patient assessed for eligibility for liver transplantation to avoid aggressive treatment.

The aim of treatment must be to stabilize patients until liver transplantation and optimize their clinical condition for a successful transplant^[6].

Medical management

Medical management is targeted at the pathogenetic mechanisms underlying HRS. The ideal treatment is designed to improve liver function by exerting splanchnic vasoconstriction and renal vasodilation to reduce portal hypertension and raise systemic arterial pressure^[34]. The specific drug approach is based on the use of vasoconstrictor agents (terlipressin, norepinephrine, midodrine) to correct circulatory changes.

As reported in a review by Davenport *et al*^[41], intravenous administration of terlipressin and albumin is currently the treatment of choice for patients with type I and type II HRS, resulting in an overall reduction in short-term mortality rates.

The vasopressin synthetic analogue terlipressin is a V1 agonist of the receptors expressed on vascular smooth muscle cells in the splanchnic circulation. It is enzymatically transformed from the inactive to biologically active form (lysine-vasopressin) with a longer half-life than other vasopressin analogues, *e.g.*, ornipressin. Terlipressin's long half-life accounts for its initial administration as an intravenous bolus, now replaced by continuous infusion^[46,47]. The vasoconstrictive effect of terlipressin corrects the circulatory dysfunction typical of end-stage liver disease, indirectly rebalancing intrarenal vasoconstriction and lowering levels of renin, noradrenaline and ultimately serum creatinine. As a result, the kidney regains control of its self-regulatory system. In addition, terlipressin has a major impact on the portal circulation reducing portal venous flow and porto-systemic pressure with a concomitant increase in hepatic arterial blood flow and an improvement in hepatocellular oxygenation.

Terlipressin can be administered as an intravenous bolus starting from a dose of 0.5 mg every 4-6 h or as a continuous infusion (2 mg/d). The dosage can be doubled after three days of treatment if there is no improvement in serum creatinine (*i.e.*, a reduction of at

least 25%)^[12]. The total daily dose should not exceed 2 mg IV bolus every 4-6 h or 12 mg/d in continuous infusion^[40]. Continuous infusion is associated with a better clinical response and fewer side-effects^[48].

Terlipressin should be associated with albumin (at a dose of 1 g/kg per day on the first day, without exceeding 100 g/d, followed by 20-40 g/d). Albumin serves to expand the circulating plasma volume by raising the oncotic pressure. In addition, it has metabolic, immune and vasoconstrictor effects by binding to endotoxins, nitric oxide, bilirubin and fatty acids^[49,50]. The terlipressin-albumin association improves renal function by 40%-60%^[48], increasing the number of patients eligible for liver transplant thereby enhancing their outcome^[51-53]. When serum creatinine values reach < 1.5 mg/dL, treatment is deemed complete^[48]. The average recovery time is seven days up to a maximum of two weeks after which terlipressin should be suspended if there is no improvement in kidney function^[54]. Even when there is a complete response, HRS recurrence is common (50% of cases) and treatment should be resumed.

Terlipressin has an acceptable side-effects profile. Side effects include abdominal pain with cramps and diarrhea until intestinal ischemia; cardiac tachyarrhythmias and chest pain can be observed, in generale ECG monitoring is recommended. Vasoconstriction induced by terlipressin may cause also cyanosis, livedo reticularis, necrosis of the skin and extremities^[53]. Terlipressin could also associated with hyponatremia but without impairment of patients' survival^[55].

If patient shows side effects the dosage should be reduced or administration discontinued. Continuous infusion is safer and less burdened by side effects^[52].

The incidence of ischemic events ranges from 5% to 30% even though many studies exclude patients at risk of cardiovascular ischemia. Fabrizi *et al*^[56]'s literature meta-analysis of 243 patients compared the effects of terlipressin vs placebo on kidney function and survival in HRS patients. Their data confirm the regression of HRS in a significant number of treated patients but no effect on survival rates.

The association albumin and terlipressin showed an improvement of survival rates for positive effects of albumin on cardiac function, on the reduction of nitric oxid and on improving the responsiveness of arterial wall to vasoconstrictors. Other studies in patients treated with terlipressin and differents colloids didn't showed the same positive response^[52,53].

Terlipressin is not available in the United States and Canada so therapeutic protocols with other vasoconstrictor agents need to be considered in those countries.

The alpha-adrenergic receptor agonist norepinephrine has proved effective in the treatment of HRS. Continuous norepinephrine infusion (at a dose of 0.5-3 mg/h) must be associated with albumin administered as an IV bolus at least twice daily (1 g/kg up to a maximum of 100 g/d). The aim is to raise mean

arterial pressure by 10 mmHg and urinary output > 200 mL every four hours. The maximum period of treatment must not exceed 2 wk^[57,58].

A pilot study by Ghosh *et al.*^[59] compared terlipressin vs noradrenaline in 46 patients with type II HRS. Neither treatment proved superior to the other and the outcome was broadly the same in terms of HRS regression. Noradrenaline can be deemed as effective as terlipressin but its lower costs makes it an interesting option for the treatment of HRS.

Another alpha-adrenergic agent, midodrine, can be considered a good alternative to terlipressin and is the drug most commonly used in the United States. Midodrine is a prodrug metabolized by the liver into its active metabolite (desglymidodrine) and then excreted in the urine. When administered in association with octreotide (a somatostatin analogue and splanchnic vasodilator) it has a positive effect on renal function in HRS patients with 50% likelihood of disease reversal^[49,60-62].

Midodrine can be administered orally (initial dose 7.5 mg every 8 h up to a maximum of 12.5 mg three times daily) or octreotide can be given by continuous infusion (50 mcg/h) or subcutaneously (from 100 to 200 mcg 12.5 mg three times daily). Albumin must be associated at the usual dose^[6]. Midodrine dosage has a major effect on its effectiveness: Patients treated at the maximum dose have shown a complete response to therapy, whereas octreotide administered alone has no effect on kidney function^[62].

Transjugular intrahepatic portosystemic shunt

The creation of a portosystemic shunt to treat refractory ascites can improve renal function in cirrhotic patients as it increases venous return of splanchnic blood to the right heart thereby raising the effective arterial blood volume and reducing hepatic sinusoidal pressure. Although literature reports on the use of transjugular intrahepatic portosystemic shunt (TIPS) in HRS patients are scant, Brensing *et al.*^[63] analyzed the trend of creatinine clearance in patients treated with TIPS, finding a twofold increase in clearance values from 9 to 27 mL/min two weeks after the procedure. Despite its side-effects (namely the high incidence of hepatic encephalopathy), TIPS can be used in the short term to gain potential benefits in patients awaiting liver transplant^[64,65].

Renal replacement therapy

The indications for renal replacement therapy (RRT) in patients with HRS are the same as those for AKI patients without cirrhosis. HRS patients, particularly those with type I, may need to undergo dialysis because of metabolic acidosis or hyperkalemia due to water or sodium retention or less frequently uremic intoxication.

RRT is among the so-called bridging therapies designed to support patients awaiting liver transplant, but there is no evidence that dialysis improves the long-term survival of patients not eligible for trans-

plantation^[66].

By definition, patients with cirrhosis are at higher risk of bleeding and hemodynamic complications (hypotension, arrhythmias) hampering the decision to initiate and manage dialysis treatment. Cirrhotic patients on RRT have a 2%-8% higher mortality rate than other patients^[67].

Continuous renal replacement therapy (CRRT) is usually preferred to intermittent dialysis due to its greater hemodynamic stability ensuring fewer fluctuations in intracranial pressure. However, prospective studies show that the choice of RRT has no significant effect on survival rates in patients awaiting liver transplantation^[68-71]. Anticoagulation of the extracorporeal circuit is needed to maintain the filter patency without increasing the risk of hemorrhage. Regional citrate anticoagulation emerged as possible alternative but no specific protocols are currently recommended for patients with liver diseases^[72].

Peritoneal dialysis is an option to resolve ascites and correct other complications of cirrhosis without exposing patients to the complications of hemodialysis^[73,74].

The precise timing and dose of RRT have yet to be established but some studies demonstrate that the early initiation and maintenance of a constantly negative fluid balance have a positive effect on survival rates^[75].

Extracorporeal artificial liver support therapy

More complex therapies known as liver support measures may be required to replace the liver's detoxifying system. RRT removes water-soluble toxins whereas most of the molecules accumulated in the course of liver failure are linked to albumin and hence are not removed by conventional hemodialysis.

Liver support systems are designed to enhance and optimize these results, increasing the removal of water-soluble toxins and those linked to albumin.

To date these treatments have served as bridging therapies for patients awaiting liver transplantation.

Molecular adsorbent recirculating system

Molecular adsorbent recirculating system (MARS) combines the conventional CRRT monitor or a standard hemodialysis machine with an albumin dialysate circuit. The system is based on the removal of albumin-bound toxins (bile acids and nitric oxide) and water-soluble cytokines (IL-6 and TNF- α) to stabilize liver function and improve organ damage.

The MARS system consists of an albumin-impermeable membrane separating the patient's blood from the albumin dialysate solution. The free albumin in the dialysate attracts and binds the liver toxins in the patient's blood. The albumin dialysate, in its turn, is regenerated by a low flux dialysis filter and two adsorber cartridges, one filled with activated charcoal, the other with an anion exchanger resin. The regenerated albumin solution is then ready for new uptake of toxins from the blood, entering the circuit through a high permeability filter to undergo standard dialysis to remove water-

soluble toxins.

Some studies have reported better survival rates in patients treated with MARS compared to conventional CRRT, but overall survival remains very poor (37% at 7 d and 25% at 30 d). The main factor affecting survival is the patient's clinical status before treatment^[75,76].

In 2000, a trial by Mitzner *et al.*^[75] assessed survival rates in 13 patients with type I HRS. The eight patients treated with MARS had significantly better survival rates at 30 d than patients receiving standard medical therapy. By contrast, the randomized RELIEF study failed to show any significant differences in terms of survival in 189 patients treated with MARS vs standard medical therapy even though some benefit was noted in the management of encephalopathy in patients with type I HRS who underwent MARS^[77].

After a one-year follow-up, Donati *et al.*^[78] reported that among 64 patients treated with MARS, the best survival rates were found in the 11 patients who subsequently underwent liver transplant. The same authors observed an improvement in both systolic and diastolic blood pressure in 5 patients with type 2 HRS treated with MARS and standard medical therapy.

Fractionated plasma separation and adsorption (Prometheus)

The Prometheus system consists of a primary circuit (plasma filter and dialyzer) and a secondary circuit (adsorbent filters to remove bilirubin) for the combined removal of toxin albumin-bound and water-soluble molecules using a fractionated plasma separation and adsorption (FPSA) system. Unlike MARS, the plasma is separated from the blood through a high cut-off point polysulfone membrane (250 kDa, albumin permeable) and purified from the albumin-bound toxins by direct adsorption on resin-containing cartridges. The purified plasma is then returned to the blood circuit through a high efficiency dialyzer to remove water-soluble toxins.

The HELIOS study on 179 patients with liver failure treated with standard medical therapy vs extracorporeal treatment showed no significant advantage in terms of overall survival except in the subgroup of patients with type I HRS treated with FPSA who had a significant survival benefit^[79].

Liver transplantation

Liver transplantation remains the treatment of choice in HRS patients despite its mortality rate which is particularly high in patients with type I HRS whose survival is so poor that many die while awaiting transplant.

Recovery of renal function is not universal: Marik *et al.*^[80], in a study on 28 patients, noted a complete recovery of kidney function in only 58% of transplanted patients, a partial recovery in 15% and no recovery in 25% (observation time 110 d). Renal sodium excretion, serum creatinine and neurohormonal levels may normalize within a month whereas renovascular resistance may take more than a year to return to

normal after transplantation^[81,82].

Organ allocation is mainly based on the MELD score, a system devised to stratify disease severity on the basis of laboratory parameters (serum creatinine, bilirubin and INR) to assign organs according to the so-called sickest first policy^[83].

Considering all liver transplant recipients, those with HRS are more exposed to post-transplant complications, at greater risk of developing CKD and have a shorter overall survival^[84,85]. Those patients who fail to recover renal function and need to continue hemodialysis have an even worse survival rate (70% mortality at one year)^[86].

RRT prior to liver transplant is an important predictive factor. Patients undergoing hemodialysis for more than eight weeks have a markedly reduced probability of renal recovery and a combined liver-kidney transplant is recommended in these cases^[87,88].

Vasopressor treatment of HRS before liver transplant does not seem to affect subsequent patient outcome^[89]. Nonetheless, Angeli *et al.*^[83] reported that liver transplantation may be delayed in patients treated with vasopressors following a response to treatment and hence an improvement in clinical and hemodynamic status. This paradoxical situation must be avoided and the clinical criteria adopted to establish the priority of patients on the waiting list for transplant (first and foremost the MELD score) must always refer to the patient's initial condition and not to the status reached after treatment.

There are no specific recommendations as to post-transplant immunosuppressive therapy, but it may be advisable to delay the start of cyclosporine or tacrolimus to 48-72 h after transplantation to enhance renal recovery as suggested by Guevara and Arroyo^[90].

CONCLUSION

HRS is a life-threatening complication arising in patients with liver cirrhosis and triggered by a series of complex hemodynamic and neurohormonal changes linked to the liver disease. The condition carries a very poor prognosis and high morbidity and mortality rates.

Recent years have seen a reduction in HRS prevalence and an improvement in patient outcome probably reflecting a better understanding of HRS pathophysiology and advances in therapeutic strategies.

Treatment consists of medical management (mainly based on vasopressor administration), surgery (TIPS) or instrumental therapies (*e.g.*, renal replacement and liver support systems). Although the therapeutic armamentarium at our disposal will control the syndrome and obtain temporary remission, there is no guarantee of disease resolution.

The only effective treatment offering patients the hope of complete recovery is liver transplantation or combined kidney-liver transplant in selected cases. The decision to embark on transplantation must be carefully assessed in HRS patients considering all the potential

factors likely to influence transplant surgery and its outcome.

REFERENCES

- Salerno F, Gerbes A, Ginès P, Wong F, Arroyo V. Diagnosis, prevention and treatment of hepatorenal syndrome in cirrhosis. *Gut* 2007; **56**: 1310-1318 [PMID: 17389705]
- Mindikoglu AL, Weir MR. Current concepts in the diagnosis and classification of renal dysfunction in cirrhosis. *Am J Nephrol* 2013; **38**: 345-354 [PMID: 24107793 DOI: 10.1159/000355540]
- Liangpunsakul S, Agarwal R. Renal failure in cirrhosis: is it time to change the diagnosis and classification? *Am J Nephrol* 2013; **38**: 342-344 [PMID: 24107717 DOI: 10.1159/000355570]
- Ginès P, Guevara M, Arroyo V, Rodés J. Hepatorenal syndrome. *Lancet* 2003; **362**: 1819-1827 [PMID: 14654322 DOI: 10.1016/S0140-6736(03)14903-3]
- Ginès P, Schrier RW. Renal failure in cirrhosis. *N Engl J Med* 2009; **361**: 1279-1290 [PMID: 19776409 DOI: 10.1056/NEJMra0809139]
- Pillebout E. Hepatorenal syndrome. *Nephrol Ther* 2014; **10**: 61-68 [PMID: 24388293 DOI: 10.1016/j.nephro.2013.11.005]
- Wadei HM. Hepatorenal syndrome: a critical update. *Semin Respir Crit Care Med* 2012; **33**: 55-69 [PMID: 22447261 DOI: 10.1055/s-0032-1301735]
- Wiest R, Garcia-Tsao G. Bacterial translocation (BT) in cirrhosis. *Hepatology* 2005; **41**: 422-433 [PMID: 15723320 DOI: 10.1002/hep.20632]
- Francés R, González-Navajas JM, Zapater P, Muñoz C, Caño R, Pascual S, Santana F, Márquez D, Pérez-Mateo M, Such J. Translocation of bacterial DNA from Gram-positive microorganisms is associated with a species-specific inflammatory response in serum and ascitic fluid of patients with cirrhosis. *Clin Exp Immunol* 2007; **150**: 230-237 [PMID: 17822441 DOI: 10.1111/j.1365-2249.2007.03494.x]
- Bajaj JS, Hylemon PB, Ridlon JM, Heuman DM, Daita K, White MB, Monteith P, Noble NA, Sikaroodi M, Gillevet PM. Colonic mucosal microbiome differs from stool microbiome in cirrhosis and hepatic encephalopathy and is linked to cognition and inflammation. *Am J Physiol Gastrointest Liver Physiol* 2012; **303**: G675-G685 [PMID: 22821944 DOI: 10.1152/ajpgi.00152.2012]
- Oliver JA, Verna EC. Afferent mechanisms of sodium retention in cirrhosis and hepatorenal syndrome. *Kidney Int* 2010; **77**: 669-680 [PMID: 20147888 DOI: 10.1038/ki.2010.4]
- Fagundes C, Ginès P. Hepatorenal syndrome: a severe, but treatable, cause of kidney failure in cirrhosis. *Am J Kidney Dis* 2012; **59**: 874-885 [PMID: 22480795 DOI: 10.1053/j.ajkd.2011.12.032]
- Chayanupatkul M, Liangpunsakul S. Cirrhotic cardiomyopathy: review of pathophysiology and treatment. *Hepatology* 2014; **8**: 308-315 [PMID: 25221635 DOI: 10.1007/s12072-014-9531-1]
- Mocarzel L, Lanzieri P, Nascimento J, Peixoto C, Ribeiro M, Mesquita E. Hepatorenal syndrome with cirrhotic cardiomyopathy: case report and literature review. *Case Reports Hepatol* 2015; **2015**: 573513 [PMID: 25874140 DOI: 10.1155/2015/573513]
- Fede G, D'Amico G, Arvaniti V, Tsochatzis E, Germani G, Georgiadis D, Morabito A, Burroughs AK. Renal failure and cirrhosis: a systematic review of mortality and prognosis. *J Hepatol* 2012; **56**: 810-818 [PMID: 22173162 DOI: 10.1016/j.jhep.2011.10.016]
- Akriviadis E, Botla R, Briggs W, Han S, Reynolds T, Shakil O. Pentoxifylline improves short-term survival in severe acute alcoholic hepatitis: a double-blind, placebo-controlled trial. *Gastroenterology* 2000; **119**: 1637-1648 [PMID: 11113085 DOI: 10.1053/gast.2000.20189]
- Planas R, Montoliu S, Ballesté B, Rivera M, Miquel M, Masnou H, Galeras JA, Giménez MD, Santos J, Cirera I, Morillas RM, Coll S, Solà R. Natural history of patients hospitalized for management of cirrhotic ascites. *Clin Gastroenterol Hepatol* 2006; **4**: 1385-1394 [PMID: 17081806 DOI: 10.1016/j.cgh.2006.08.007]
- Wong LP, Blackley MP, Andreoni KA, Chin H, Falk RJ, Klemmer PJ. Survival of liver transplant candidates with acute renal failure receiving renal replacement therapy. *Kidney Int* 2005; **68**: 362-370 [PMID: 15954928 DOI: 10.1111/j.1523-1755.2005.00408.x]
- European Association for the Study of the Liver. EASL clinical practice guidelines on the management of ascites, spontaneous bacterial peritonitis, and hepatorenal syndrome in cirrhosis. *J Hepatol* 2010; **53**: 397-417 [PMID: 20633946 DOI: 10.1016/j.jhep.2010.05.004]
- Hkecer R, Sherlock S. Electrolyte and circulatory changes in terminal liver failure. *Lancet* 1956; **271**: 1121-1125 [PMID: 13377688 DOI: 10.1016/S0140-6736(56)90149-0]
- Koppel MH, Coburn JW, Mims MM, Goldstein H, Boyle JD, Rubini ME. Transplantation of cadaveric kidneys from patients with hepatorenal syndrome. Evidence for the functional nature of renal failure in advanced liver disease. *N Engl J Med* 1969; **280**: 1367-1371 [PMID: 4890476 DOI: 10.1056/NEJM196906192802501]
- Arroyo V, Ginès P, Gerbes AL, Dudley FJ, Gentilini P, Laffi G, Reynolds TB, Ring-Larsen H, Schölmerich J. Definition and diagnostic criteria of refractory ascites and hepatorenal syndrome in cirrhosis. International Ascites Club. *Hepatology* 1996; **23**: 164-176 [PMID: 8550036 DOI: 10.1002/hep.510230122]
- Arroyo V, Colmenero J. Ascites and hepatorenal syndrome in cirrhosis: pathophysiological basis of therapy and current management. *J Hepatol* 2003; **38** Suppl 1: S69-S89 [PMID: 12591187 DOI: 10.1016/S0168-8278(03)00007-2]
- Fasolato S, Angeli P, Dallagnese L, Maresio G, Zola E, Mazza E, Salinas F, Donà S, Fagioli S, Sticca A, Zanus G, Cillo U, Frasson I, Destro C, Gatta A. Renal failure and bacterial infections in patients with cirrhosis: epidemiology and clinical features. *Hepatology* 2007; **45**: 223-229 [PMID: 17187409 DOI: 10.1002/hep.21443]
- Moore K. Acute kidney injury in cirrhosis: a changing spectrum. *Hepatology* 2013; **57**: 435-437 [PMID: 22886711 DOI: 10.1002/hep.26003]
- Nadim MK, Kellum JA, Davenport A, Wong F, Davis C, Pannu N, Tolwani A, Bellomo R, Genyk YS. Hepatorenal syndrome: the 8th International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care* 2012; **16**: R23 [PMID: 22322077 DOI: 10.1186/cc11188]
- Fagundes C, Barreto R, Guevara M, Garcia E, Solà E, Rodríguez E, Graupera I, Ariza X, Pereira G, Alfaro I, Cárdenas A, Fernández J, Poch E, Ginès P. A modified acute kidney injury classification for diagnosis and risk stratification of impairment of kidney function in cirrhosis. *J Hepatol* 2013; **59**: 474-481 [PMID: 23669284 DOI: 10.1016/j.jhep.2013.04.036]
- Piano S, Rosi S, Maresio G, Fasolato S, Cavallin M, Romano A, Morando F, Gola E, Frigo AC, Gatta A, Angeli P. Evaluation of the Acute Kidney Injury Network criteria in hospitalized patients with cirrhosis and ascites. *J Hepatol* 2013; **59**: 482-489 [PMID: 23665185 DOI: 10.1016/j.jhep.2013.03.039]
- Arroyo V. Acute kidney injury (AKI) in cirrhosis: should we change current definition and diagnostic criteria of renal failure in cirrhosis? *J Hepatol* 2013; **59**: 415-417 [PMID: 23727236 DOI: 10.1016/j.jhep.2013.05.035]
- Wong F, Nadim MK, Kellum JA, Salerno F, Bellomo R, Gerbes A, Angeli P, Moreau R, Davenport A, Jalan R, Ronco C, Genyk Y, Arroyo V. Working Party proposal for a revised classification system of renal dysfunction in patients with cirrhosis. *Gut* 2011; **60**: 702-709 [PMID: 21325171 DOI: 10.1136/gut.2010.23613]
- Seo YS, Jung ES, An H, Kim JH, Jung YK, Kim JH, Yim HJ, Yeon JE, Byun KS, Kim CD, Ryu HS, Um SH. Serum cystatin C level is a good prognostic marker in patients with cirrhotic ascites and normal serum creatinine levels. *Liver Int* 2009; **29**: 1521-1527 [PMID: 19725889 DOI: 10.1111/j.1478-3231]
- Sharaway MA, Shawky EM, Ali LH, Mohammed AA, Hassan HA, Fouad YM. Cystatin C: a predictor of hepatorenal syndrome in patients with liver cirrhosis. *Hepatology* 2011; **5**: 927-933 [PMID: 21484118 DOI: 10.1007/s12072-011-9266-y]
- Angeli P, Ginès P, Wong F, Bernardi M, Boyer TD, Gerbes A, Moreau R, Jalan R, Sarin SK, Piano S, Moore K, Lee SS, Durand F, Salerno F, Caraceni P, Kim WR, Arroyo V, Garcia-Tsao G. Diagnosis and management of acute kidney injury in patients with cirrhosis: revised consensus recommendations of the International Club of Ascites. *J Hepatol* 2015; **62**: 968-974 [PMID: 25638527]

- DOI: 10.1016/j.jhep.2014]
- 34 **Møller S**, Krag A, Bendtsen F. Kidney injury in cirrhosis: pathophysiological and therapeutic aspects of hepatorenal syndromes. *Liver Int* 2014; **34**: 1153-1163 [PMID: 24673771 DOI: 10.1111/liv.12549]
 - 35 **García-Tsao G**, Parikh CR, Viola A. Acute kidney injury in cirrhosis. *Hepatology* 2008; **48**: 2064-2077 [PMID: 19003880 DOI: 10.1002/hep.22605]
 - 36 **Belcher JM**, Sanyal AJ, Peixoto AJ, Perazella MA, Lim J, Thiessen-Philbrook H, Ansari N, Coca SG, Garcia-Tsao G, Parikh CR. Kidney biomarkers and differential diagnosis of patients with cirrhosis and acute kidney injury. *Hepatology* 2014; **60**: 622-632 [PMID: 24375576 DOI: 10.1002/hep.26980]
 - 37 **Verna EC**, Brown RS, Farrand E, Pichardo EM, Forster CS, Soladell Valle DA, Adkins SH, Sise ME, Oliver JA, Radhakrishnan J, Barasch JM, Nickolas TL. Urinary neutrophil gelatinase-associated lipocalin predicts mortality and identifies acute kidney injury in cirrhosis. *Dig Dis Sci* 2012; **57**: 2362-2370 [PMID: 22562534 DOI: 10.1007/s10620-012-2180-x]
 - 38 **Fagundes C**, Pépin MN, Guevara M, Barreto R, Casals G, Solà E, Pereira G, Rodríguez E, García E, Prado V, Poch E, Jiménez W, Fernández J, Arroyo V, Ginès P. Urinary neutrophil gelatinase-associated lipocalin as biomarker in the differential diagnosis of impairment of kidney function in cirrhosis. *J Hepatol* 2012; **57**: 267-273 [PMID: 22521351 DOI: 10.1016/j.jhep.2012.03.015]
 - 39 **Barreto R**, Elia C, Solà E, Moreira R, Ariza X, Rodríguez E, Graupera I, Alfaro I, Morales-Ruiz M, Poch E, Guevara M, Fernández J, Jiménez W, Arroyo V, Ginès P. Urinary neutrophil gelatinase-associated lipocalin predicts kidney outcome and death in patients with cirrhosis and bacterial infections. *J Hepatol* 2014; **61**: 35-42 [PMID: 24613364 DOI: 10.1016/j.jhep.2014.02.023]
 - 40 **Fabrizi F**, Aghemo A, Messa P. Hepatorenal syndrome and novel advances in its management. *Kidney Blood Press Res* 2013; **37**: 588-601 [PMID: 24356549 DOI: 10.1159/000355739]
 - 41 **Davenport A**, Ahmad J, Al-Khafaji A, Kellum JA, Genyk YS, Nadim MK. Medical management of hepatorenal syndrome. *Nephrol Dial Transplant* 2012; **27**: 34-41 [PMID: 22287700 DOI: 10.1093/ndt/gfr736]
 - 42 **Ginès A**, Fernández-Esparrach G, Monescillo A, Vila C, Domènech E, Abecasis R, Angeli P, Ruiz-Del-Arbol L, Planas R, Solà R, Ginès P, Terg R, Inglada L, Vaqué P, Salerno F, Vargas V, Clemente G, Quer JC, Jiménez W, Arroyo V, Rodés J. Randomized trial comparing albumin, dextran 70, and polygeline in cirrhotic patients with ascites treated by paracentesis. *Gastroenterology* 1996; **111**: 1002-1010 [PMID: 8831595 DOI: 10.1016/S0016-5085(96)70068-9]
 - 43 **García-Compeán D**, Blanc P, Larrey D, Daures JP, Hirtz J, Mendoza E, Maldonado H, Michel H. Treatment of cirrhotic tense ascites with Dextran-40 versus albumin associated with large volume paracentesis: a randomized controlled trial. *Ann Hepatol* 2002; **1**: 29-35 [PMID: 15114293]
 - 44 **Fernández J**, Navasa M, Planas R, Montoliu S, Monfort D, Soriano G, Vila C, Pardo A, Quintero E, Vargas V, Such J, Ginès P, Arroyo V. Primary prophylaxis of spontaneous bacterial peritonitis delays hepatorenal syndrome and improves survival in cirrhosis. *Gastroenterology* 2007; **133**: 818-824 [PMID: 17854593 DOI: 10.1053/j.gastro.2007.06.065]
 - 45 **Sort P**, Navasa M, Arroyo V, Aldeguer X, Planas R, Ruiz-del-Arbol L, Castells L, Vargas V, Soriano G, Guevara M, Ginès P, Rodés J. Effect of intravenous albumin on renal impairment and mortality in patients with cirrhosis and spontaneous bacterial peritonitis. *N Engl J Med* 1999; **341**: 403-409 [PMID: 10432325 DOI: 10.1056/NEJM199908053410603]
 - 46 **Fernández J**, Escorsell A, Zabalza M, Felipe V, Navasa M, Mas A, Lacy AM, Ginès P, Arroyo V. Adrenal insufficiency in patients with cirrhosis and septic shock: Effect of treatment with hydrocortisone on survival. *Hepatology* 2006; **44**: 1288-1295 [PMID: 17058239 DOI: 10.1002/hep.21352]
 - 47 **Barbano B**, Sardo L, Gigante A, Gasperini ML, Liberatori M, Giraldo GD, Lacanna A, Amoroso A, Cianci R. Pathophysiology, diagnosis and clinical management of hepatorenal syndrome: from classic to new drugs. *Curr Vasc Pharmacol* 2014; **12**: 125-135 [PMID: 24678726 DOI: 10.2174/157016111201140327163930]
 - 48 **Alessandria C**, Ottobrelli A, Debernardi-Venon W, Todros L, Cerenzia MT, Martini S, Balzola F, Morgando A, Rizzetto M, Marzano A. Noradrenalin vs terlipressin in patients with hepatorenal syndrome: a prospective, randomized, unblinded, pilot study. *J Hepatol* 2007; **47**: 499-505 [PMID: 17560680 DOI: 10.1016/j.jhep.2007.04.010]
 - 49 **Angeli P**, Volpin R, Gerunda G, Craighero R, Roner P, Merenda R, Amodio P, Sticca A, Caregato L, Maffei-Faccioli A, Gatta A. Reversal of type 1 hepatorenal syndrome with the administration of midodrine and octreotide. *Hepatology* 1999; **29**: 1690-1697 [PMID: 10347109 DOI: 10.1002/hep.510290629]
 - 50 **Martín-Llahí M**, Pépin MN, Guevara M, Díaz F, Torre A, Monescillo A, Soriano G, Terra C, Fábrega E, Arroyo V, Rodés J, Ginès P. Terlipressin and albumin vs albumin in patients with cirrhosis and hepatorenal syndrome: a randomized study. *Gastroenterology* 2008; **134**: 1352-1359 [PMID: 18471512 DOI: 10.1053/j.gastro.2008.02.024]
 - 51 **Sanyal AJ**, Boyer T, Garcia-Tsao G, Regenstein F, Rossaro L, Appenrodt B, Blei A, Gülberg V, Sigal S, Teuber P. A randomized, prospective, double-blind, placebo-controlled trial of terlipressin for type 1 hepatorenal syndrome. *Gastroenterology* 2008; **134**: 1360-1368 [PMID: 18471513 DOI: 10.1053/j.gastro.2008.02.014]
 - 52 **Angeli P**. Review article: prognosis of hepatorenal syndrome--has it changed with current practice? *Aliment Pharmacol Ther* 2004; **20** Suppl 3: 44-6; discussion 47-8 [PMID: 15335400 DOI: 10.1111/j.1365-2036.2004.02113.x]
 - 53 **Gluud LL**, Christensen K, Christensen E, Krag A. Systematic review of randomized trials on vasoconstrictor drugs for hepatorenal syndrome. *Hepatology* 2010; **51**: 576-584 [PMID: 19885875 DOI: 10.1002/hep.23286]
 - 54 **Fabrizi F**, Dixit V, Martin P. Meta-analysis: terlipressin therapy for the hepatorenal syndrome. *Aliment Pharmacol Ther* 2006; **24**: 935-944 [PMID: 16948805 DOI: 10.1111/j.1365-2036.2006.03086.x]
 - 55 **Sugumaran A**, Lougher E, Czajkowski M, Yeoman A. Reduction of serum sodium (Na) in patients treated with terlipressin for variceal bleeding (VB) and hepatorenal syndrome (HRS). *Gut* 2014; **63**: A87-A88 [DOI: 10.1136/gutjnl-2014-307263.184]
 - 56 **Fabrizi F**, Dixit V, Messa P, Martin P. Terlipressin for hepatorenal syndrome: A meta-analysis of randomized trials. *Int J Artif Organs* 2009; **32**: 133-140 [PMID: 19440988]
 - 57 **Duvoux C**, Zanditenas D, Hézode C, Chauvat A, Monin JL, Roudot-Thoraval F, Mallat A, Dhumeaux D. Effects of noradrenalin and albumin in patients with type I hepatorenal syndrome: a pilot study. *Hepatology* 2002; **36**: 374-380 [PMID: 12143045 DOI: 10.1053/jhep.2002.34343]
 - 58 **Wadei HM**, Gonwa TA. Hepatorenal syndrome in the intensive care unit. *J Intensive Care Med* 2013; **28**: 79-92 [PMID: 21859679 DOI: 10.1177/0885066611408692]
 - 59 **Ghosh S**, Choudhary NS, Sharma AK, Singh B, Kumar P, Agarwal R, Sharma N, Bhalla A, Chawla YK, Singh V. Noradrenaline vs terlipressin in the treatment of type 2 hepatorenal syndrome: a randomized pilot study. *Liver Int* 2013; **33**: 1187-1193 [PMID: 23601499 DOI: 10.1111/liv.12179]
 - 60 **Prabhu MV**, Sukanya B, Santosh Pai BH, Reddy S. The hepatorenal syndrome - a review. *G Ital Nefrol* 2014; **31**: [PMID: 25030015]
 - 61 **Esraïlian E**, Pantangco ER, Kyulo NL, Hu KQ, Runyon BA. Octreotide/Midodrine therapy significantly improves renal function and 30-day survival in patients with type 1 hepatorenal syndrome. *Dig Dis Sci* 2007; **52**: 742-748 [PMID: 17235705 DOI: 10.1007/s10620-006-9312-0]
 - 62 **Wong F**, Pantea L, Sniderman K. Midodrine, octreotide, albumin, and TIPS in selected patients with cirrhosis and type 1 hepatorenal syndrome. *Hepatology* 2004; **40**: 55-64 [PMID: 15239086 DOI: 10.1002/hep.20262]
 - 63 **Breising KA**, Textor J, Perz J, Schiedermaier P, Raab P, Strunk H, Klehr HU, Kramer HJ, Spengler U, Schild H, Sauerbruch T. Long term outcome after transjugular intrahepatic portosystemic shunt in non-transplant cirrhotics with hepatorenal syndrome: a phase II study. *Gut* 2000; **47**: 288-295 [PMID: 10896924 DOI: 10.1136/gut.47.2.288]

- 64 **Guevara M**, Ginès P, Bandi JC, Gilabert R, Sort P, Jiménez W, Garcia-Pagan JC, Bosch J, Arroyo V, Rodés J. Transjugular intrahepatic portosystemic shunt in hepatorenal syndrome: effects on renal function and vasoactive systems. *Hepatology* 1998; **28**: 416-422 [PMID: 9696006 DOI: 10.1002/hep.510280219]
- 65 **Pipili C**, Cholongitas E. Renal dysfunction in patients with cirrhosis: Where do we stand? *World J Gastrointest Pharmacol Ther* 2014; **5**: 156-168 [PMID: 25133044 DOI: 10.4292/wjgpt.v5.i3.156]
- 66 **Keller F**, Heinze H, Jochimsen F, Passfall J, Schuppan D, Büttner P. Risk factors and outcome of 107 patients with decompensated liver disease and acute renal failure (including 26 patients with hepatorenal syndrome): the role of hemodialysis. *Ren Fail* 1995; **17**: 135-146 [PMID: 7644764 DOI: 10.3109/08860229509026250]
- 67 **Wilkinson SP**, Weston MJ, Parsons V, Williams R. Dialysis in the treatment of renal failure in patients with liver disease. *Clin Nephrol* 1977; **8**: 287-292 [PMID: 884909]
- 68 **Davenport A**. Renal replacement therapy in the patient with acute brain injury. *Am J Kidney Dis* 2001; **37**: 457-466 [PMID: 11228168 DOI: 10.1053/ajkd.2001.22068]
- 69 **Witzke O**, Baumann M, Patschan D, Patschan S, Mitchell A, Treichel U, Gerken G, Philipp T, Kribben A. Which patients benefit from hemodialysis therapy in hepatorenal syndrome? *J Gastroenterol Hepatol* 2004; **19**: 1369-1373 [PMID: 15610310 DOI: 10.1111/j.1440-1746.2004.03471.x]
- 70 **Gonwa TA**, Mai ML, Melton LB, Hays SR, Goldstein RM, Levy MF, Klntmalm GB. Renal replacement therapy and orthotopic liver transplantation: the role of continuous veno-venous hemodialysis. *Transplantation* 2001; **71**: 1424-1428 [PMID: 11391230 DOI: 10.1097/00007890-200105270-00012]
- 71 **Mackelaite L**, Alsaukas ZC, Ranganna K. Renal failure in patients with cirrhosis. *Med Clin North Am* 2009; **93**: 855-869, viii [PMID: 19577118 DOI: 10.1016/j.mcna.2009.03.003]
- 72 **Patel S**, Wendon J. Regional citrate anticoagulation in patients with liver failure--time for a rethink? *Crit Care* 2012; **16**: 153 [PMID: 22985662 DOI: 10.1186/cc11492]
- 73 **Pipili C**, Polydorou A, Pantelias K, Korfiatis P, Nikolakopoulos F, Grapsa E. Improvement of hepatic encephalopathy by application of peritoneal dialysis in a patient with non-end-stage renal disease. *Perit Dial Int* 2013; **33**: 213-216 [PMID: 23478376 DOI: 10.3747/pdi.2011.00271]
- 74 **Liu KD**, Himmelfarb J, Paganini E, Ikizler TA, Soroko SH, Mehta RL, Chertow GM. Timing of initiation of dialysis in critically ill patients with acute kidney injury. *Clin J Am Soc Nephrol* 2006; **1**: 915-919 [PMID: 17699307 DOI: 10.2215/CJN.01430406]
- 75 **Mitzner SR**, Stange J, Klammt S, Risler T, Erley CM, Bader BD, Berger ED, Lauchart W, Peszynski P, Freytag J, Hickstein H, Loock J, Löhr JM, Liebe S, Emmrich J, Korten G, Schmidt R. Improvement of hepatorenal syndrome with extracorporeal albumin dialysis MARS: results of a prospective, randomized, controlled clinical trial. *Liver Transpl* 2000; **6**: 277-286 [PMID: 10827226 DOI: 10.1053/lv.2000.6355]
- 76 **Mitzner SR**, Stange J, Klammt S, Peszynski P, Schmidt R, Nöldge-Schomburg G. Extracorporeal detoxification using the molecular adsorbent recirculating system for critically ill patients with liver failure. *J Am Soc Nephrol* 2001; **12** Suppl 17: S75-S82 [PMID: 11251037]
- 77 **Bañares R**, Nevens F, Larsen FS, Jalan R, Albillos A, Dollinger M, Saliba F, Sauerbruch T, Klammt S, Ockenga J, Pares A, Wendon J, Brünner T, Kramer L, Mathurin P, de la Mata M, Gasbarrini A, Müllhaupt B, Wilmer A, Laleman W, Eefsen M, Sen S, Zipprich A, Tenorio T, Pavesi M, Schmidt HH, Mitzner S, Williams R, Arroyo V. Extracorporeal albumin dialysis with the molecular adsorbent recirculating system in acute-on-chronic liver failure: the RELIEF trial. *Hepatology* 2013; **57**: 1153-1162 [PMID: 23213075 DOI: 10.1002/hep.26185]
- 78 **Donati G**, La Manna G, Cianciolo G, Grandinetti V, Carretta E, Cappuccilli M, Panicali L, Iorio M, Piscaglia F, Bolondi L, Coli L, Stefoni S. Extracorporeal detoxification for hepatic failure using molecular adsorbent recirculating system: depurative efficiency and clinical results in a long-term follow-up. *Artif Organs* 2014; **38**: 125-134 [PMID: 23834711 DOI: 10.1111/aor.12106]
- 79 **Kribben A**, Gerken G, Haag S, Herget-Rosenthal S, Treichel U, Betz C, Sarrazin C, Hoste E, Van Vlierberghe H, Escorsell A, Hafer C, Schreiner O, Galle PR, Mancini E, Caraceni P, Karvellas CJ, Salmhofer H, Knotek M, Ginès P, Kozik-Jaromin J, Rifai K. Effects of fractionated plasma separation and adsorption on survival in patients with acute-on-chronic liver failure. *Gastroenterology* 2012; **142**: 782-789.e3 [PMID: 22248661 DOI: 10.1053/j.gastro.2011.12.056]
- 80 **Marik PE**, Gayowski T, Starzl TE. The hepatoadrenal syndrome: a common yet unrecognized clinical condition. *Crit Care Med* 2005; **33**: 1254-1259 [PMID: 15942340 DOI: 10.1097/01.CCM.0000164541.12106.57]
- 81 **Navasa M**, Feu F, Garcia-Pagán JC, Jiménez W, Llach J, Rimola A, Bosch J, Rodés J. Hemodynamic and humoral changes after liver transplantation in patients with cirrhosis. *Hepatology* 1993; **17**: 355-360 [PMID: 8444409 DOI: 10.1002/hep.1840170302]
- 82 **Piscaglia F**, Zironi G, Gaiani S, Mazziotti A, Cavallari A, Gramantieri L, Valgimigli M, Bolondi L. Systemic and splanchnic hemodynamic changes after liver transplantation for cirrhosis: a long-term prospective study. *Hepatology* 1999; **30**: 58-64 [PMID: 10385639 DOI: 10.1002/hep.510300112]
- 83 **Angeli P**, Gines P. Hepatorenal syndrome, MELD score and liver transplantation: an evolving issue with relevant implications for clinical practice. *J Hepatol* 2012; **57**: 1135-1140 [PMID: 22749942 DOI: 10.1016/j.jhep.2012.06.024]
- 84 **Gonwa TA**, Mai ML, Melton LB, Hays SR, Goldstein RM, Levy MF, Klntmalm GB. End-stage renal disease (ESRD) after orthotopic liver transplantation (OLT) using calcineurin-based immunotherapy: risk of development and treatment. *Transplantation* 2001; **72**: 1934-1939 [PMID: 11773892 DOI: 10.1097/00007890-200112270-00012]
- 85 **Lee JP**, Heo NJ, Joo KW, Yi NJ, Suh KS, Moon KC, Kim SG, Kim YS. Risk factors for consequent kidney impairment and differential impact of liver transplantation on renal function. *Nephrol Dial Transplant* 2010; **25**: 2772-2785 [PMID: 20207711 DOI: 10.1093/ndt/gfq093]
- 86 **Ruiz R**, Barri YM, Jennings LW, Chinnakotla S, Goldstein RM, Levy MF, McKenna GJ, Randall HB, Sanchez EQ, Klntmalm GB. Hepatorenal syndrome: a proposal for kidney after liver transplantation (KALT). *Liver Transpl* 2007; **13**: 838-843 [PMID: 17539003 DOI: 10.1002/lt.21149]
- 87 **Marik PE**, Wood K, Starzl TE. The course of type 1 hepato-renal syndrome post liver transplantation. *Nephrol Dial Transplant* 2006; **21**: 478-482 [PMID: 16249201 DOI: 10.1093/ndt/gfi212]
- 88 **Gerbes AL**, Gülberg V, Wagershauser T, Holl J, Reiser M. Renal effects of transjugular intrahepatic portosystemic shunt in cirrhosis: comparison of patients with ascites, with refractory ascites, or without ascites. *Hepatology* 1998; **28**: 683-688 [PMID: 9731559 DOI: 10.1002/hep.510280313]
- 89 **Restuccia T**, Ortega R, Guevara M, Ginès P, Alessandria C, Ozdogan O, Navasa M, Rimola A, Garcia-Valdecasas JC, Arroyo V, Rodés J. Effects of treatment of hepatorenal syndrome before transplantation on posttransplantation outcome. A case-control study. *J Hepatol* 2004; **40**: 140-146 [PMID: 14672625 DOI: 10.1016/j.jhep.2003.09.019]
- 90 **Guevara M**, Arroyo V. Hepatorenal syndrome. *Expert Opin Pharmacother* 2011; **12**: 1405-1417 [PMID: 21480763 DOI: 10.1517/14656566]

P-Reviewer: Kelesidis T, Nechifor G, Trimarchi H, Trumper L, Yorioka N **S-Editor:** Tian YL **L-Editor:** A **E-Editor:** Wu HL





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

