

VOLUME 286 (2011) PAGES 17144–17155

DOI 10.1074/jbc.A110.210526

## **Neutralizing interleukin 1 $\beta$ (IL-1 $\beta$ ) induces $\beta$ -cell survival by maintaining PDX1 protein nuclear localization.**

Amin Ardestani, Nadine S. Sauter, Federico Paroni, Gitanjali Dharmadhikari, Jae-Hyoung Cho, Roberto Lupi, Piero Marchetti, José Oberholzer, Julie Kerr Conte, and Kathrin Maedler

This article has been retracted by the publisher. Fig. 1E contained images that had been reused from the following references (Glas, R., Sauter, N. S., Schulthess, F. T., Shu, L., Oberholzer, J., and Maedler, K. (2009) Purinergic P2X7 receptors regulate secretion of interleukin-1 receptor antagonist and  $\beta$  cell function and survival. *Diabetologia* **52**, 1579–1588 and Maedler, K., Schumann, D. M., Sauter, N., Ellingsgaard, H., Bosco, D., Baertschiger, R., Iwakura, Y., Oberholzer, J., Wollheim, C. B., Gauthier, B. R., and Donath, M. Y. (2006) Low concentration of interleukin-1 $\beta$  induces FLICE-inhibitory protein-mediated  $\beta$ -cell proliferation in human pancreatic islets. *Diabetes* **55**, 2713–2722), and these reused images represented different experimental conditions.

Authors are urged to introduce these corrections into any reprints they distribute. Secondary (abstract) services are urged to carry notice of these corrections as prominently as they carried the original abstracts.