

VOLUME 289 (2014) PAGES 23963–23976

DOI 10.1074/jbc.A114.571489

## NMR-based structural analysis of the complete rough-type lipopolysaccharide isolated from *Capnocytophaga canimorsus*.

Ulrich Zähringer, Simon Ittig, Buko Lindner, Hermann Moll, Ursula Schombel, Nicolas Gisch, and Guy R. Cornelis

PAGE 23969:

Some of the  $^{31}\text{P}$  NMR signals in Table 3 were erroneously assigned. The corrected Table 3 is shown here. These corrections do not affect the interpretation of the results or the conclusions of this work.

**TABLE 3**

$^{31}\text{P}$  NMR (283.7 MHz) data of compounds **1**, **2**, and **3**

$^{31}\text{P}$  signal resonances were assigned according to  $^1\text{H},^{31}\text{P}$ -HMQC and  $^1\text{H},^{31}\text{P}$ -HMQC-TOCSY spectroscopy.

| Residue                                | $^{31}\text{P}$ $\delta$ (ppm) |          |          |
|--|--------------------------------|----------|----------|
|  | <b>1</b>                       | <b>2</b> | <b>3</b> |
| GlcN→1- <i>P</i> -Etn                  | -                              | - 0.89   | - 1.25   |
| 6- <i>P</i> -Etn-Man <sup>I</sup> -(1→ | 0.83                           | 0.90     | 0.94     |
| 4- <i>P</i> -Etn-Kdo-(2→               | 0.09                           | 0.15     | 0.08     |

Authors are urged to introduce these corrections into any reprints they distribute. Secondary (abstract) services are urged to carry notice of these corrections as prominently as they carried the original abstracts.