

# High Prevalence and Genetic Heterogeneity of Rodent-Borne *Bartonella* Species on Heixiazi Island, China

Dong-Mei Li,<sup>a</sup> Yong Hou,<sup>b</sup> Xiu-Ping Song,<sup>a</sup> Ying-Qun Fu,<sup>b</sup> Gui-Chang Li,<sup>a</sup> Ming Li,<sup>b</sup> Marina E. Ereemeeva,<sup>c</sup> Hai-Xia Wu,<sup>a</sup> Bo Pang,<sup>a</sup> Yu-Juan Yue,<sup>a</sup> Ying Huang,<sup>a</sup> Liang Lu,<sup>a</sup> Jun Wang,<sup>a</sup> Qi-Yong Liu<sup>a</sup>

State Key Laboratory of Infectious Disease Prevention and Control, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Changping, Beijing, People's Republic of China<sup>a</sup>; Heilongjiang Entry-Exit Inspection and Quarantine Bureau, Harbin, Heilongjiang Province, People's Republic of China<sup>b</sup>; Jiann-Ping Hsu College of Public Health, Georgia Southern University, Statesboro, Georgia, USA<sup>c</sup>

**We performed genetic analysis of *Bartonella* isolates from rodent populations from Heixiazi Island in northeast China. Animals were captured at four sites representing grassland and brushwood habitats in 2011 and examined for the prevalence and genetic diversity of *Bartonella* species, their relationship to their hosts, and geographic distribution. A high prevalence (57.7%) and a high diversity (14 unique genotypes which belonged to 8 clades) of *Bartonella* spp. were detected from 71 rodents comprising 5 species and 4 genera from 3 rodent families. Forty-one *Bartonella* isolates were recovered and identified, including *B. taylorii*, *B. japonica*, *B. cooperplainsensis*, *B. grahamii*, *B. washoensis* subsp. *cynomysii*, *B. doshiae*, and two novel *Bartonella* species, by sequencing of four genes (*gltA*, the 16S rRNA gene, *ftsZ*, and *rpoB*). The isolates of *B. taylorii* and *B. grahamii* were the most prevalent and exhibited genetic difference from isolates identified elsewhere. Several isolates clustered with strains from Japan and far-eastern Russia; strains isolated from the same host typically were found within the same cluster. Species descriptions are provided for *Bartonella heixiaziensis* sp. nov. and *B. fuyuanensis* sp. nov.**

Significant numbers of zoonotic *Bartonella* species have been recognized in the past 25 years. Before 1991, *B. bacilliformis* was the only member of the *Bartonella* genus. At present, the genus *Bartonella* contains over 30 formally described species of *Bartonella* and many unnamed *Bartonella* spp. isolated from a variety of mammal species, most often from rodents, ruminants, and carnivores (1). Some *Bartonella* spp. are long-known etiological agents of human diseases (Carrion's disease, cat scratch disease, and trench fever) and some specific syndromes (verruca peruana, bacillary angiomatosis, endocarditis, pericarditis, and neuroretinitis) associated with these agents. Rodents are the largest source of *Bartonella* spp. There are more than 40 species of rodents known to be associated with 26 different species of *Bartonella*, seven of which have been implicated as occasional causes of human infections (1). People become infected with rodent-borne *Bartonella* incidentally, when they are exposed to habitats and ectoparasites of wild rodents harboring various *Bartonella* species.

Previous field and clinical studies in mainland China demonstrated circulation of a variety of different *Bartonella* spp. associated with peridomestic, feral, and exotic animals across the country (2–5) and low levels of human *Bartonella* infections, diagnosed primary as cat scratch disease due to *B. henselae*, in heavily populated areas (6). Heixiazi Island has a beautiful natural landscape and plentiful wildlife; it is expected to become a tourist attraction because of its special history and geographical position. This island was previously unpopulated and isolated from mainland northeast Asia before its recent opening to tourists. Thus, in the present study, to evaluate the potential risk for transmission of rodent-borne *Bartonella* on Heixiazi Island after opening it up to tourists, we conducted a baseline survey of the occurrence and prevalence of *Bartonella* infections in rodents on Heixiazi Island, China. The genetic diversity of *Bartonella* isolates and their relationship to hosts and geographic distributions were examined and

evaluated in the context of current knowledge about *Bartonella* species which are associated with rodents.

## MATERIALS AND METHODS

**Ethics statement.** This study was performed after review and approval by the Ethics Committee of the National Institute for Communicable Disease Control and Prevention of the Chinese CDC. All animals were treated according to the approved protocols and the guidelines for the Laboratory Animal Use and Care from the Chinese CDC and the Rules for the Implementation of Laboratory Animal Medicine (1998) from the Ministry of Health, China.

**Sampling sites and tissue collections.** Rodents were captured using the trap-at-night method in May 2011. Six hundred traps baited with peanuts were set up in four sites of Heixiazi Island consisting of grassland and brushwood, the predominant habitats on the island. The trapped animals were morphologically identified to species. Collection time, site, habitat, species, gender, weight, head-body length, and tail length were recorded. Liver and spleen were collected into 1.5-ml Nunc CryoTubes and stored in liquid nitrogen prior to processing and laboratory testing.

Received 20 June 2015 Accepted 3 September 2015

Accepted manuscript posted online 11 September 2015

**Citation** Li D-M, Hou Y, Song X-P, Fu Y-Q, Li G-C, Li M, Ereemeeva ME, Wu H-X, Pang B, Yue Y-J, Huang Y, Lu L, Wang J, Liu Q-Y. 2015. High prevalence and genetic heterogeneity of rodent-borne *Bartonella* species on Heixiazi Island, China. *Appl Environ Microbiol* 81:7981–7992. doi:10.1128/AEM.02041-15.

**Editor:** C. R. Lovell

Address correspondence to Qi-Yong Liu, liuqiyong@icdc.cn.

D.-M.L. and Y.H. contributed equally to this article.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AEM.02041-15>.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.

TABLE 1 Oligonucleotide primers used for PCR

Target gene	Primer	Primer direction	Primer sequence (5'–3')	Product length (bp)
<i>gltA</i>	BhCS.781p	Forward	GGG GAC CAG CTC ATG GTG G	379
	BhCS.1137n	Reverse	AAT GCA AAA AGA ACA GTA AAC A	
16S rRNA gene	fDI	Forward	AGA GTT TGA TCC TGG CTC AG	1,400–1,500
	rDI	Reverse	AAG GAG GTG ATC CAG CC	
<i>ftsZ</i>	Bfp1	Forward	ATT AAT CTG CAY CGG CCA GA	896
	Bfp2	Reverse	ACV GAD ACA CGA ATA ACA CC	
<i>rpoB</i>	1400F	Forward	CGC ATT GGC TTA CTT CGT ATG	866
	2300R	Reverse	GTA GAC TGA TTA AAC GCT G	

**Isolation and cultivation of *Bartonella* organisms, phenotypic characterization, and electron microscopy.** Approximately 20 mg of organ samples was disrupted using a Dounce–Potter homogenizer in 0.6 ml of tryptic soy broth (TSB) (BD), and suspensions were inoculated onto tryptic soy agar (TSA) plates (BD) supplemented with 5% defibrinated sheep blood. Plates were kept in a 5% CO<sub>2</sub> humidified atmosphere at 35°C for 30 days and were checked daily for bacterial growth starting at day 3 after inoculation. Small, round, gray-white colonies were morphologically identified as *Bartonella* and passaged onto fresh plates.

Material from a single colony was suspended in 100 µl of deionized water, heated for 10 min at 95°C, and cleared by centrifugation for 5 min at 6,000 × g at 4°C. The supernatant was used as a source of DNA template for PCR. The presence of *Bartonella* DNA was first determined by PCR detecting a portion of the citrate synthase gene (*gltA*) with the primers BhCS.781p and BhCS.1137n (Table 1) as described elsewhere (7). Bacterial isolates identified as *Bartonella* were cloned by limiting dilution, and established cultures were preserved in brain heart infusion (BHI) broth (BD) supplemented with 30% glycerol at –70°C. Isolates were passaged into flasks containing insect Schneider's medium and propagated according to previously established procedures to observe their growth characteristics (8, 9).

To prepare samples for electron microscopy, bacterial cells were suspended in deionized water, spread onto a water surface, absorbed onto Formvar stabilized with carbon support film, stained for 2 min in 1% uranyl acetate, and then air dried. Samples were examined with a transmission electron microscope (Hitachi, HT7700) at 80 kV with negative staining.

Biochemical testing was performed using commercially available reagents (Rapid ID 32 A; bioMérieux) according to the manufacturer's instructions. Fatty acid methyl esters were obtained by saponification, methylation, and extraction according to the MIS operation manual (Midi Inc., 2002) and then analyzed using Agilent Hp6890 as Chromatograph and ChemStation v A5.05.

**PCR amplification and sequencing of *Bartonella* genes.** DNA was extracted from bacterial isolates using the DNeasy tissue kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. For all identified isolates, four gene fragments were amplified and their sequences were obtained: *gltA*, the 16S rRNA gene (10), cell division protein gene *ftsZ* (11), and RNA polymerase beta-subunit gene *rpoB* (12). Nucleotide primers used in this study are listed in Table 1.

PCR amplification was performed using *Pfu* DNA polymerase (TransGen Biotech, Beijing, China) and 20 to 50 ng of genomic DNA in a final volume of 50 µl containing 0.2 µM each primer and 25 µl 2× TransTaq-T PCR SuperMix. The PCR amplification conditions were as follows: an initial step of 94°C for 2 min; 30 amplification cycles, each consisting of 94°C for 30 s and 53°C for 30 s; an elongation step of 72°C for 2 min; and a final incubation at 72°C for 5 min. Amplified products were analyzed by electrophoresis on 1% agarose gels supplemented with 0.005% of GoldView (SBS Genetech, Beijing, China) and visualized under UV light. PCR products of the expected length were

then purified and sequenced on both strands by Tsingke Biotechnology (Beijing, China).

**Phylogenetic analysis.** Individual gene sequences and concatenated sequences assembled for *gltA* (326 bp), 16S rRNA gene (1245 bp), *ftsZ* (767 bp), and *rpoB* (771 bp) fragments were aligned using ClustalW, and neighbor-joining analysis was performed using MEGA6 (13) with the Kimura 2-parameter model. A phylogeny was constructed based on the concatenated sequences using the maximum-likelihood method and HKY85 model implemented in PhyML software (14). Homologous sequences of *Brucella abortus* were used as an outgroup. Bootstrap analysis was carried out with 1,000 resamplings. Homologous sequences for reference *Bartonella* species were downloaded from the NCBI GenBank, and their accession numbers are listed in Table S1 in the supplemental material.

**Recombination analysis.** The multiple-alignment sequences (MAS) were scanned for intragenic and intergenic recombination using RDPv.3.44 (15), GENECONV, Bootscan, MaxChi, Chimaera, and SiScan as implemented in RDP with default settings. A phylogenetic tree was also inferred using ClonalFrame v1.1 (16) with the number of Markov chain Monte Carlo (MCMC) iterations set to 50,000, following a burn-in period of 50,000 iterations.

The Neighbor-Net implemented in the software SplitsTree 4.13 (17) with 1,000 bootstrap replicates was used to create the EqualAngle phylogenetic network for the concatenated sequences. The pairwise homoplasy index (PHI) in SplitsTree 4.13 was used to test the role of past recombination events in generating allelic variation.


**Population diversity and distribution analysis.** Nucleotide diversity indices ( $\pi$ ) and the polymorphic level (haplotype diversity [Hd]) were calculated using DNAsp v5.10 (18). The aligned segment of *gltA* was used to construct a phylogenetic network with the postprocessing of maximum parsimony (MP) calculation (19) using the median joining method (20) after the preprocessing of star contraction (21) implemented in the program Network 4.6.1.2. The parameters of characters were set at the default values (epsilon and weight values = 0 and 10). Homologous sequences for the *Bartonella* strains were downloaded from the NCBI GenBank, and their accession numbers are listed in Tables S1, S2, S3, and S4 in the supplemental material.

**Statistical analysis.** SPSS software (version 19.0) was used for statistical analysis. Chi-square and Fisher's exact tests were applied for assessing the differences of infection rates in the tissues tested. A *P* value of <0.05 was considered statistically significant.

**Nucleotide sequence accession numbers.** Sequences of the *Bartonella* gene fragments generated in this study were deposited in the NCBI GenBank database under accession numbers KJ175028 to KJ175068 for *gltA*, KJ361602 to KJ361642 for the 16S rRNA gene, KJ361684 to KJ361724 for *ftsZ*, and KJ361725 to KJ361765 for *rpoB*.

## RESULTS

**Study site description.** Heixiazhi Island (134°24'E, 48°22'N) (also called Fuyuan Delta or Bolshoi Ussuriysky Island), located at the



**FIG 1** Distribution of sampling sites. The map shows the locations of the geographic sites where the rodents were collected on Heixiazi Island, China, during 2011. Map created using ArcGis v10.2 (ESRI, Redland, CA).

Chinese-Russian border, is a natural sedimentary island at the confluence of Heilongjiang and Wusulijiang Rivers (called the Ussuri and Amur Rivers in Russia) (Fig. 1). Heixiazi Island serves as a natural border between China and Russia; it is 327 km<sup>2</sup> with an average altitude of about 40 m. China restored sovereign control over half of Heixiazi Island (171 km<sup>2</sup>) from Russia on 14 October 2008. The Chinese section of the island is part of Fuyuan County, Heilongjiang Province. The landscape consists mainly of wetlands and lush growth of elms, poplars, willows oak trees, sedges, and grasses.

**Animal trapping.** A total of 71 small mammals belonging to 6 species and 4 genera from 3 rodent families were captured in 4 sampling sites. Forty-three animals were trapped in woodland habitats and 28 animals in grassland habitats. The northern red-backed vole *Myodes rutilus* was the most prevalent animal species and accounted for 64.8% ( $n = 46$ ) of the rodents captured. Striped field mouse *Apodemus agrarius* ( $n = 12$ ), reed vole *Microtus fortis* ( $n = 5$ ), Maximowicz's vole *Microtus maximowiczii* ( $n = 1$ ), Korean field mouse *Apodemus peninsulae* ( $n = 6$ ), and Siberian chipmunk *Eutamias sibiricus* ( $n = 1$ ) were also collected and accounted for 16.9%, 7.0%, 1.4%, 8.5%, and 1.4%, respectively, of all trapped animals.

***Bartonella* prevalence and identification.** The prevalence of *Bartonella* in rodents was 57.7% (41/71) but there were no significant variations of infection rate ( $P > 0.05$ , Fisher's exact test) between the two species belonging to *Apodemus* spp. In total, 66 *Bartonella* isolates were established from tissue samples of 71 rodents, including 34 (48.6%) isolates from 70 liver tissues and 32 (47.8%) isolates from 67 spleen tissues. There were 25 paired isolates recovered from the spleens and livers, respectively, of 25 rodents. The other 16 isolates were obtained from livers (9 isolates) or spleens (7 isolates) of different rodents (Table 2; see Table S5 in the supplemental material). The prevalence of *Bartonella* infection in livers did not differ from the prevalence of *Bartonella* infection in spleens ( $P > 0.05$ , chi-square test). *Bartonella* growth could be observed as early as day 4 after inoculation of the tissue homogenate onto agar plates. Colony morphology and growth rate were consistent with the characteristics of *Bartonella* isolates described in the literature (22–24).

**Genetic identification and phylogenetic relationships of rodent *Bartonella* isolates from Heixiazi Island.** Based on sequence analysis of the 326-nucleotide (nt) *gltA* fragment, 41 newly established isolates of *Bartonella* belonged to six known *Bartonella* species and two potentially novel species. The isolates of *B. taylorii* (11 isolates), *B. japonica* (1 isolate), *B. coopersplainsensis* (3 isolates), *B. grahamii* (5 isolates), *B. washoensis* subsp. *cynomysii* (1 isolate), and *B. doshiae* (12 isolates) were obtained from voles, field mice, and chipmunks from Heixiazi Island. Seven isolates from voles (one from *M. maximowiczii* and 6 from *M. rutilus*) and one isolate from a mouse (*Apodemus agrarius*) had unique genotypes, and the representative strains were CR90HXZ and AA137HXZ. The isolates CR90HXZ and AA137HX, each shared less than 95% nucleotide sequence similarity in their *gltA* fragment with homologous genes of the nearest species of *Bartonella*, *B. vinsonii* subsp. *arupensis* and *B. grahamii*, respectively.


To further explore the relationship of the new *Bartonella* isolates from Chinese rodents with recognized species of *Bartonella*, we performed a phylogenetic analysis based on the alignment of the individual gene fragments of the *gltA*, 16S rRNA, *ftsZ*, and *rpoB* genes (see Fig. S1 in the supplemental material). Furthermore, 3,110-bp fragments of concatenated sequences of the same four conserved genes of 41 *Bartonella* isolates obtained in this study and homologous sequences of 26 *Bartonella* type strains retrieved from GenBank were analyzed. The same topology trees were produced for each individual gene fragment and for the concatenated sequence fragments using both the neighbor-joining (Fig. 2) and maximum-likelihood methods (see Fig. S2 in the supplemental material). Overall, the 41 isolates exhibited 14 distinct genotypes which were predicted using DnaSP v5.10.1; these 14 genotypes were clustered into 8 clades (I to VIII). The largest number of isolates was associated with clades formed by six from all recognized species of *Bartonella*, including *B. taylorii* (clade I), *B. japonica* (clade II), *B. coopersplainsensis* (clade III), *B. grahamii* (clade V), *B. washoensis* subsp. *cynomysii* (clade VI), and *B. doshiae* (clade VIII) (Fig. 2). Eleven Chinese isolates clustered with *B. taylorii* exhibited two distinct genotypes: one genotype comprised 8 isolates from *M. rutilus* (*Myodes* genotype) and a second genotype consisted of 3 isolates from 2 *A. peninsulae* animals and one *A. agrarius* animal (*Apodemus* genotype). Nucleotide sequences of another isolate from *A. peninsulae* (AP73HXZ) were identical to those of homologous gene fragments of *B. japonica* recovered from *A. argenteus* in Japan (AB242289). Furthermore, sequences

**TABLE 2** *Bartonella* infection rates in liver and spleen tissues of rodents collected on Heixiazi Island, China<sup>a</sup>

Rodent species	Liver		Spleen		Liver and spleen coisolation
	<i>n</i>	No. (%) positive	<i>n</i>	No. (%) positive	
<i>Microtus fortis</i>	5	1 (20.0)	5	1 (20.0)	1
<i>Microtus maximowiczii</i>	1	1 (100.0)	1	1 (100.0)	1
<i>Myodes rutilus</i>	46	25 (54.3)	43	22 (51.2)	19
<i>Apodemus agrarius</i>	11	4 (36.4)	11	5 (45.5)	2
<i>Eutamias sibiricus</i>	1	0	1	1 (100)	0
<i>Apodemus peninsulae</i>	6	3 (50)	6	2 (33.3)	2
Total	70	34 (48.6)	67	32 (47.8)	25

<sup>a</sup> There were no significant differences in the rates of *Bartonella* isolate recovery from liver and spleen ( $P > 0.05$ ).






of three isolates from *A. agrarius* (AA95HXZ, AA86HXZ, and AA123HXZ) were identical to each other and had the most similarity with the sequences of *B. cooperplainsensis* obtained from *Rattus leucopus* in Australia (EU111803). The *B. grahamii* clade (clade V) included 5 genetically distinct Chinese isolates recovered from *A. agrarius* (2 isolates), *M. rutilus* (2 isolates), and *M. fortis* (1 isolate). A single isolate, ES132HXZ from *E. sibiricus*, was grouped with *B. washoensis* subsp. *cynomysii* in cluster VI. Twelve isolates from *M. rutilus* had the same sequences and were most related to *B. doshiae*. Seven isolates from 6 *M. rutilus* and one from *M. maximowiczii* were identical to each other and were included in cluster VII, which represented a unique lineage within the trees evaluated. The nucleotide sequences of these isolates exhibited 4.6% nucleotide divergence from the homologous sequences of its closest neighbor, *B. washoensis* subsp. *cynomysii*. One isolate from *A. agrarius* formed cluster IV and had 4.5% genetic divergence from its nearest neighbor, *B. rattimassiliensis*. Based on these genetic differences, *Bartonella* isolates from *M. rutilus* and *M. maximowiczii* and an isolate from *A. agrarius* closest to *B. rattimassiliensis* represent novel species.

**Scanning of multiple sequence alignments for intragenic and intergenic mosaicism and effects of recombination on phylogenetic relationships.** Individual DNA alignments of *gltA*, the 16S rRNA gene, *ftsZ*, and *rpoB* were analyzed with RDPv.3.44 programs to detect putative interspecies recombinant sequences. Potential mosaic regions were considered likely recombinants only if they were detected by at least two programs. There were no significant interspecies recombination events detected in the individual DNA alignments. Accordingly, there was no interference from recombination for the phylogenetic trees constructed based on the four individual gene fragments.

To include more informative sites, the concatenated sequences of four genes were analyzed for occurrence of possible interspecies recombination events using the same method. Three significant recombinations were detected, which affected mostly the two different genotypes of *B. taylorii*. The recombination breakpoint position was located between nt 153 and 392 of the concatenated sequences of the *Myodes*-type isolates of *B. taylorii*, while the minor parental sequence originated from CR84HXZ of *B. grahamii*. These parts of the concatenated fragment were from the hypervariable 3' end of the *gltA* fragment (nt 153 to 326) and 16S rRNA gene fragment (nt 1 to 67). The topology of the phylogenetic tree was not changed upon exclusion of those nucleotides from the analysis. Furthermore, the split network also supported the same relationship among the isolates, and this was not reflected in the recombination breakpoint pattern. The fit, quite high at 98.2%, indicates that the data are very tree-like, further suggesting that the bootstrap split tree (Fig. 3) was congruent with the clustering in the phylogenetic binary tree (Fig. 2).

**FIG 2** Phylogenetic relationships of the *Bartonella* type strains and the *Bartonella* isolates from Heixiazhi Island based on the concatenation of fragments of *gltA*, the 16S rRNA gene, *ftsZ*, and *rpoB*. The phylogenetic tree was inferred using the neighbor-joining method based on the Kimura 2-parameter model in MEGA6 (13). The percentages of replicate trees (>70%) in which the associated taxa clustered together in the bootstrap test (1,000 replicates) are shown next to the branches. The analysis involved 3,110 concatenated nucleotides from 68 isolates. Roman numerals I to VIII on the right side of the tree correspond to different *Bartonella* genetic groups. Homologous sequences of *Brucella abortus* were used as an outgroup. All sequences were trimmed to the same size, and gaps were excluded from the analysis.



**FIG 3** Neighbor-Net tree of associations of *Bartonella* isolates obtained from different rodents inhabiting Heixiazi Island, China, and the *Bartonella* type strains based on concatenated conserved genes (*gltA*, 16S rRNA gene, *ftsZ*, and *rpoB*) representing a total of 3,110 nucleotides using SplitsTree4. The numbers in parentheses represent the number of isolates at each branch. The percentages of replicate trees in which the associated taxa clustered together in the bootstrap test (1,000 replicates) are shown next to the branches; only bootstrap values of >70% are indicated. Different colors refer to individual *Bartonella* genetic groups. Recombination is illustrated by the reticulation structures in the center among *Bartonella* strains. *Bartonella tamiae* was used as an outgroup. The fit parameter values were 98.2% in all cases, indicating that a substantial fraction of the phylogenetic signals could be depicted on the graph. The bar in the upper left corner of the network graph indicates the scale of branches.

For assessing the relative contributions of recombination and mutation in the delineation of various *Bartonella* genotypes from a common ancestor, we calculated the ratio of recombination to mutation value ( $r/m$  and  $\rho/\theta$  ratio) using ClonalFrame v1.1 (21). The inferred outputs were estimated at 0.83286 ( $r/m$ ) (95% confidence interval [CI]<sub>95</sub> = 0.507585 to 1.269109) and 0.036913 ( $\rho/\theta$ ) (CI<sub>95</sub> = 0.021735 to 0.058028), suggesting that nucleotide changes in these conserved genes occur more frequently by *de novo* mutation than by recombination.

**Genetic polymorphism of rodent-adapted *Bartonella* species.** Overall, the extent of the intraspecies variation within the different *Bartonella* clades was different and closely linked to their host associations based on the 4 conserved gene fragments. Twelve *B. doshaie* isolates from *M. rutilus* had an identical genotype. Similarly, three isolates of *B. cooperi* from *Apodemus* spp. exhibited the same genotype. In contrast, *B. taylorii*, *B. grahamii*, and *B. washoensis*-like isolates from Heixiazi Island exhibited noticeable intraspecies genetic diversity. Therefore, we examined the nucleotide polymorphisms of the 4 conserved genes in these isolates recovered from Heixiazi Island compared to the levels of inter- and intraspecies genetic variation among isolates of *B. taylorii*, *B. grahamii* and *B. washoensis* circulating in different parts of

the world, using sequences from GenBank. The level of nucleotide polymorphism differed between two groups, and the nucleotide diversities varied with the different alleles (Table 3). The 16S rRNA gene exhibited the lowest genetic diversity in each population of *Bartonella* isolates analyzed and had the lowest calculated genetic distance ( $D$ ) and nucleotide diversity ( $\pi$ ) values. The nucleotide diversity and genetic distance for the *gltA*, *ftsZ*, and *rpoB* gene fragments varied among the *B. taylorii*, *B. grahamii*, and *B. washoensis*-like isolates from Heixiazi Island. Analysis of the  $D$  and  $\pi$  parameters calculated for all *Bartonella* isolates from Heixiazi Island suggested that the highest polymorphism was within *gltA* and *rpoB*, similar to the conclusions derived when similar parameters were calculated for sequences of known *Bartonella* species. Our observations suggest that these two genes have more significant impact for defining species of *Bartonella* than the other two genes.

Isolates identified as *B. taylorii* belong to two genotypes which exhibit estimated genetic distances of 0.008: the *Apodemus*-associated genotype and the *Myodes*-associated genotype. The *Apodemus*-associated *B. taylorii* isolates were from two sympatric species of wood mice, *A. agrarius* and *A. peninsulae*. These isolates had identical nucleotide sequences for all four gene fragments se-

**TABLE 3** Nucleotide polymorphism and nucleotide diversity of rodent-associated isolates from Heixiazi Island (China) and type strains of *Bartonella* species<sup>a</sup>

<i>Bartonella</i> sp. (no. of isolates)	Gene	Size (bp)	<i>D</i>	<i>S</i>	<i>PI</i>	$\pi$ (mean $\pm$ SD)	Hd (mean $\pm$ SD)	<i>k</i>
<i>B. taylorii</i> (11)	<i>gltA</i>	326	0.033	0	23	0.03079 $\pm$ 0.01050	0.436 $\pm$ 0.133	10.036
	16S rRNA gene	1,247	0.002	0	5	0.00175 $\pm$ 0.00078	0.436 $\pm$ 0.133	2.182
	<i>ftsZ</i>	766	0.006	0	10	0.00570 $\pm$ 0.00217	0.436 $\pm$ 0.133	4.364
	<i>rpoB</i>	771	0.009	0	15	0.00847 $\pm$ 0.00303	0.436 $\pm$ 0.133	6.545
	Concatenate	3,110	0.008	0	53	0.00744 $\pm$ 0.00239	0.436 $\pm$ 0.133	23.127
<i>B. grahamii</i> (5)	<i>gltA</i>	326	0.007	1	6	0.01227 $\pm$ 0.00359	0.700 $\pm$ 0.218	4.000
	16S rRNA gene	1,245	0.000	0	1	0.00048 $\pm$ 0.00014	0.600 $\pm$ 0.175	0.600
	<i>ftsZ</i>	766	0.011	4	11	0.01069 $\pm$ 0.00260	1.000 $\pm$ 0.126	8.200
	<i>rpoB</i>	771	0.009	12	22	0.02335 $\pm$ 0.00480	1.000 $\pm$ 0.126	18.000
	Concatenate	3,108	0.010	17	40	0.00991 $\pm$ 0.00219	1.000 $\pm$ 0.126	30.800
<i>B. heixiaziensis</i> (7)	<i>gltA</i>	326	0.002	2	0	0.00175 $\pm$ 0.00120	0.286 $\pm$ 0.196	0.571
	16S rRNA gene	1,245	0.000	0	0	0.00000	0.000	0.000
	<i>ftsZ</i>	766	0.002	5	0	0.00186 $\pm$ 0.00128	0.286 $\pm$ 0.196	1.429
	<i>rpoB</i>	771	0.000	1	0	0.00037 $\pm$ 0.00026	0.286 $\pm$ 0.039	0.286
	Concatenate	3,107	0.001	8	0	0.00074 $\pm$ 0.00051	0.286 $\pm$ 0.196	2.286
All HXZ isolates (41)	<i>gltA</i>	326	0.110	15	78	0.09707 $\pm$ 0.00390	0.856 $\pm$ 0.033	31.645
	16S rRNA gene	1,245	0.006	6	26	0.00567 $\pm$ 0.00085	0.848 $\pm$ 0.032	7.040
	<i>ftsZ</i>	766	0.080	12	143	0.07373 $\pm$ 0.00327	0.860 $\pm$ 0.034	56.554
	<i>rpoB</i>	771	0.073	21	172	0.09479 $\pm$ 0.00339	0.860 $\pm$ 0.034	73.083
	Concatenate	3,106	0.057	54	419	0.05419 $\pm$ 0.00223	0.860 $\pm$ 0.034	168.322
All type strains (26)	<i>gltA</i>	326	0.145	49	104	0.12614 $\pm$ 0.00836	1.000 $\pm$ 0.011	41.123
	16S rRNA gene	1,236	0.015	52	43	0.01515 $\pm$ 0.00202	1.000 $\pm$ 0.011	18.723
	<i>ftsZ</i>	765	0.121	80	191	0.10947 $\pm$ 0.00837	1.000 $\pm$ 0.011	83.745
	<i>rpoB</i>	771	0.132	48	224	0.11947 $\pm$ 0.00560	1.000 $\pm$ 0.011	92.108
	Concatenate	3,097	0.081	230	560	0.07553 $\pm$ 0.00494	1.000 $\pm$ 0.011	233.609

<sup>a</sup> HXZ, Heixiazi Island; *D*, genetic distance; *S*, singleton variable sites; *PI*, parsimony-informative sites;  $\pi$ , nucleotide diversity; Hd, haplotype diversity; *k*, average number of nucleotide differences.

quenced, but these sequences were different from the sequences of the homologous genes of *M. rutilus*-associated *B. taylorii* isolates. No indel was found within the four gene fragments sequenced for all *B. taylorii* isolates from Heixiazi Island examined here; however, 53 polymorphic and parsimony informative sites were identified within 3,110 nt of concatenated *gltA*-16S rRNA gene-*ftsZ*-*rpoB* sequences of 11 *B. taylorii* isolates from the island. In particular, *gltA*, the 16S rRNA gene, *ftsZ* and *rpoB* contained 23, 5, 10 and 15 polymorphic sites, respectively, which accounted for 0.74%, 0.16%, 0.32%, and 0.48%, respectively, of the entire 3,110-nt concatenated fragment. The haplotype diversity (Hd),  $\pi$  value, and average number of nucleotide differences (*k*) are estimated to be 0.436, 0.00744, and 23.127, respectively. In comparison, analysis of the 258-nt sequence of *gltA* from the 108 entries listed as *B. taylorii* sequences in NCBI GenBank detected 63 polymorphic sites. Among them, there were 13 single nucleotide polymorphism (SNP) sites and 49 parsimony-informative sites; accordingly, the calculated  $\pi$  value per site was 0.03881, and the *k* value was 10.013. Furthermore, it was estimated that 53 haplotypes exist among previously identified isolates of *B. taylorii*; their predicted Hd value was 0.971.

Five Heixiazi Island isolates, including two isolates from *M. rutilus* (CR84HXZ and CR127HXZ), one isolate from *M. fortis* (MF74HXZ), and two isolates from *A. agrarius* (AA97HXZ and AA131HXZ) were clustered with *B. grahamii*, but all had distinct genotypes with an overall genetic distance of 0.011. Sequences of

four conserved genes compared for these isolates of *B. grahamii* had 57 polymorphic sites, no indels were present. There were 17 SNP and 40 parsimony-informative sites; they were found in *gltA*, the 16S rRNA gene, *ftsZ* and *rpoB* and accounted for 0.22% (7 nt), 0.03% (1 nt), 0.48% (15 nt), and 1.09% (34 nt), respectively, of the entire 3,109-nt concatenated fragment. The Hd,  $\pi$ , and *k* values were estimated to be 1.000, 0.00991, and 30.8, respectively. In comparison, analysis of the 307-nt sequence of *gltA* of the 64 entries identified as *B. grahamii* sequences found in GenBank detected 54 polymorphic sites. Among them, there were 27 SNP and were 27 parsimony-informative sites; the estimated  $\pi$  value per site was 0.02241, and the *k* value was 6.880. It was estimated that there are at least 36 haplotypes among previously identified isolates of *B. grahamii*; the predicted Hd value among known *B. grahamii* isolates was estimated to be 0.961.

Seven *Bartonella* isolates from Heixiazi Island, including six isolates from *M. rutilus* and one isolate from *M. maximowiczii*, exhibited two distinct genetic types, and both were clustered with *B. washoensis* as a nearest neighbor. Indel and parsimony-informative sites were not found in sequences generated for these isolates. The 16S rRNA gene sequences were conserved among all seven isolates. Eight SNP sites were found in the *gltA*, *ftsZ*, and *rpoB* sequences, which accounted for 0.06% (2 nt), 0.16% (5 nt), and 0.03% (1 nt) of the entire 3,107-nt concatenated fragment, respectively. The Hd,  $\pi$ , and *k* values among the two genotypes were 0.286, 0.00074, and 2.286, respectively. In comparison, anal-

ysis of the 306-nt sequence of *gltA* of the 82 of *B. washoensis* sequences found in GenBank detected 56 polymorphic sites. Among them, there were 17 SNP and 39 parsimony-informative sites; the estimated  $\pi$  value per site was 0.02984, and the  $k$  value was 9.132. Overall, there were 53 haplotypes among previously identified isolates of *B. washoensis*, and their predicted Hd is 0.9584.

**Relationship of *Bartonella*, hosts, and geographic distributions.** Since isolates of *B. taylorii*, *B. grahamii*, and *B. washoensis* are known for their ubiquitous distribution, the phylogenetic network of these *Bartonella* species was analyzed in the context of their geographic relationship and their host associations in order to define their influence on the *Bartonella* population genetic structure. Our analysis revealed that *Bartonella* isolates from Asia, Europe, and North America do not exhibit any obvious geographic clustering pattern; however, they could be easily grouped based on their host associations (Fig. 4).

The genetically different isolates of *B. taylorii* from voles and mice in Heixiazi Island belong to two different clades of *Bartonella*; one includes *B. taylorii* from Japan, and the other includes *B. taylorii* isolates from France, Greece, Japan, Lithuania, Poland, the Russian Far East, and Thailand (Fig. 4A1). This grouping of *B. taylorii* isolates matched the grouping based on their host associations (Fig. 4A2). Three *Apodemus* genotype isolates (AA139HXZ, AP119HXZ, and AP124HXZ) from Heixiazi Island were in the same cluster group as *B. taylorii* from Europe and the *Apodemus*- and *Myodes*-associated *B. taylorii* isolates from Japan. The eight *Myodes* genotype isolates (CR87HXZ, CR89HXZ, CR112HXZ, CR115HXZ, CR116HXZ, CR117HXZ, CR128HXZ, and CR142HXZ) grouped with the isolates from Asia and Europe, which have different animal species as hosts. Four isolates of *B. taylorii* from *Cenophthalmus lushuiensis*, one isolate from *Eothenomys miletus* trapped in Yunnan Province, China, and an isolate from Japanese *Microtus motibelli* clustered separately from the Heixiazi Island isolates (Fig. 4A1 and A2).

Five Heixiazi Island isolates of *B. grahamii* formed two clusters; one included 2 isolates (AA131HXZ and AA97HXZ) from *A. agrarius* and isolates from *Apodemus* spp. trapped in Japan and the Russian Far East (Fig. 4B1). The second cluster consisted of isolates originating from a variety of different hosts, including one from *M. fortis* (MF74HXZ) and two from *M. rutilus* (CR84HXZ and CR127HXZ) trapped on Heixiazi Island and four *Bartonella* strains isolated from a human, *A. agrarius*, and *M. ochrogaster* in Finland, the Russian Far East, and North America (Fig. 4B2).

Previously characterized isolates of *B. washoensis* associate with various species of squirrels in Asia and North America (25). The *B. washoensis* isolates from Hebei, Zhejiang Province, and Heixiazi Island in China belong to three groups, one of which also includes the Heixiazi Island isolates from *E. sibiricus* (Fig. 4C1). The other two clusters include 17 isolates of *B. washoensis* from *Spermophilus dauricus*, one isolate from *A. chevrieri*, and one isolate from a tick, *Haemaphysalis longicornis* (Fig. 4C2). *B. washoensis* isolates from North America exhibited a similar clustering pattern based on their host associations (Fig. 4C2).

**Host specificity of *Bartonella* isolates.** Overall, 28 *Bartonella* isolates of 4 different species were obtained from *M. rutilus* (Table 4). There were 12 isolates of *B. doshiae*, 8 of *B. taylorii*, 6 of *B. washoensis*, and 2 of *B. grahamii*. Eleven isolates of *B. taylorii* were established from *M. rutilus* (8 isolates; 72.7%), *A. peninsulae* (2 isolates, 18.2%), and *A. agrarius* (1 isolate, 9.1%). The vole was the predominant host of *B. taylorii*. Genetic variation among individ-

ual isolates of *B. taylorii* from Heixiazi Island separated the *Apodemus* genotype and the *Myodes rutilus* genotype. The majority of *B. washoensis*-like isolates originated from *M. rutilus* and *M. fortis* voles. *B. grahamii* isolates were recovered from *M. rutilus*, *A. agrarius*, and *M. fortis*. Except for *E. sibiricus*, five species of rodents trapped on the island harbored more than one species of *Bartonella*. Only one strain of *B. washoensis* was isolated from *E. sibiricus* from Heixiazi Island. A new genotype (VII) of *Bartonella* spp., which was isolated from 6 *M. rutilus* animals and 1 *M. maximowiczii* animal, appears to be predominantly associated with voles.

**Characterization of novel *Bartonella* isolates.** Light microscopy examination of the two isolates CR90HXZ and AA137HXZ revealed short, straight, or slightly curved Gram-negative bacilli. Electron microscopy with negative staining of isolate AA137HXZ detected small rods with unipolar surface pili; in contrast, pili were not observed for isolate CR90HXZ (Fig. 5).

Isolates CR90HXZ and AA137HXZ were catalase and oxidase negative and had no urease activity, similar to the case for other *Bartonella* isolates. Tests for  $\alpha$ -glucosidase,  $\alpha$ -L-fucosidase,  $\alpha$ -L-arabinofuranosidase, mannose, raffinose, and nitrate reduction and spot indole tests using both stains were negative. Amino acid amidase activity was observed with the following amino acids: arginine (Arg), phenylalanine (Phe), leucine (Leu), tyrosine (Tyr), alanine (Ala), glycine (Gly), histidine (His), and serine (Ser).

To further confirm that these novel isolates were typical of *Bartonella* species, their fatty acid profiles were determined. The common cellular fatty acid proportions of  $C_{18:1\omega7c}$ ,  $C_{18:0}$ ,  $C_{16:0}$ , and  $C_{18:1\omega7c}$  were, respectively, 62.39%, 18.7%, 10.82%, and 1.01% for isolate CR90HXZ and 56.29%, 12.66%, 22.79%, and 2.16% for isolate AA137HXZ.  $C_{18:1\omega9c}$  and  $C_{20:1\omega7c}$  accounted for, respectively, 1.71% and 2.06% in isolate CR90HXZ;  $C_{17:0}$  accounted for 4.02% of total fatty acids in isolate AA137HXZ, while  $C_{18:1\omega9c}$  and  $C_{20:1\omega7c}$  were absent.

## DISCUSSION

On Heixiazi Island, six species of rodents harbored eight *Bartonella* spp., including six known *Bartonella* species and two potentially novel *Bartonella* species. In particular, *Apodemus* rodents harbored the largest variety of *Bartonella* spp. (*B. cooperplainsensis*, *B. grahamii*, *B. taylorii*, *B. japonica*, and a novel species close to *B. rattimassiliensis*), *M. rutilus* harbored *B. taylorii*, *B. doshiae*, *B. grahamii*, and a novel species closely related to *B. washoensis*, *M. maximowiczii* harbored *B. grahamii* and the same novel species, and one *E. sibiricus* animal and one *M. fortis* animal harbored *B. washoensis* and *B. grahamii*, respectively. Since rodent species captured in our study widely inhabit the Sanjiang Plain (26), it is very likely that circulation of the same *Bartonella* spp. may be expected in areas surrounding the river and associated mainland territories.

Livers and spleens of infected animals had similar burdens of *Bartonella* as determined by the isolation results. These and previous observations suggest that these organs may act as depositories of harbored *Bartonella*, although they may be damaged as a result of a long-term *Bartonella* persistence in these tissues (27). These interactions underlie the primary etiology of *Bartonella*-caused hepatosplenomegaly and tissue inflammation in humans and some animals, including dogs (28–30). Nevertheless, the spleen is regarded as an organ for retaining and filtering infected erythrocytes rather than to be a sanctuary for persisting *Bartonella*,








TABLE 4 Rodent species from which the *Bartonella* isolates were established and their habitats on Heixiazi Island, China

Host	Nearest known <i>Bartonella</i> sp.	Similarity for 3,109 bp (%)	No. of isolates			Constituent ratio (%)
			Grassland	Woodland	Total	
<i>Apodemus agrarius</i>	<i>B. cooperplainsensis</i>	99.6	2	1	3	42.9
	<i>B. grahamii</i>	98.4	2	0	2	28.6
	<i>B. rattimassiliensis</i>	95.5	1	0	1	14.3
	<i>B. taylorii</i>	97.6	1	0	1	14.3
<i>Apodemus peninsulae</i>	<i>B. taylorii</i>	97.6	0	2	2	66.7
	<i>B. japonica</i>	99.9	1	0	1	33.3
<i>Myodes rutilus</i>	<i>B. taylorii</i>	97.8	4	4	8	28.6
	<i>B. doshiae</i>	99.5	1	11	12	42.9
	<i>B. grahamii</i>	99.7	2	0	2	7.1
	<i>B. washoensis</i> subsp. <i>cynomysii</i>	95.4	5	1	6	21.4
<i>Eutamias sibiricus</i>	<i>B. washoensis</i> subsp. <i>cynomysii</i>	97.1	1	0	1	100.0
<i>Microtus fortis</i>	<i>B. grahamii</i>	99.5	0	1	1	50.0
<i>Microtus maximowiczii</i>	<i>B. washoensis</i> subsp. <i>cynomysii</i>	95.4	1	0	1	50.0

as suggested by observations of chronic *B. birtlesii* infection in an *in vivo* mouse model (31); however, the precise mechanisms of *Bartonella* persistence in animals are not well understood.

Previous studies suggested that genes of rodent-borne *Bartonella* exhibit more frequent recombination events than those of the species typically associated with humans and cats (32, 33). Since any recombination event affecting housekeeping genes will compromise the accuracy of the phylogenetic trees (34), we examined both fragments of the individual genes and multigene concatenated sequences for occurrence of recombination events. No recombination events were found upon analyzing sequences of the four individual gene fragments, suggesting that these genes are suitable for the phylogenetic analysis of binary trees. Detection of recombination events in the multigene concatenated sequences is suggestive of an artifact due to concatenation or an indication of recombination in the regions adjacent to the primer annealing sites and external to the PCR fragments (15). Since the topology of the trees remained the same upon exclusion of these putative recombined portions of the fragments, it is unlikely that the phylogenetic relationship within this group of *Bartonella* species is affected by their inclusion in the concatenated fragments. Furthermore, the split-tree analysis also confirmed that the topology of the binary tree tends to be a tree, not a network. Although the multigene concatenation did not influence the results of phylogenetic tree construction in the present study, considering the illusion of recombination, owing to artificial connection of the multiple locus genes, could cause an inaccurate phylogenetic estimation of taxonomic relationships.

The extent of the genetic divergence established for different *Bartonella* populations matched the diversity of their associated hosts present on Heixiazi Island. Genetic diversity estimated for *B. grahamii* on Heixiazi Island is greater than the genetic diversity

established for *B. taylorii* from the same area (1 and 0.436); however, our findings are in contrast to the results of comparative analysis of *B. grahamii* and *B. taylorii* isolates of European and American origin (35). Furthermore, the information available for *gltA* fragment suggests that the genetic diversity of *B. taylorii* isolates ( $H_d = 0.971$ ,  $\pi = 0.03881$ ) is slightly greater than that of *B. grahamii* isolates ( $H_d = 0.961$ ,  $\pi = 0.02241$ ). Clear genetic diversity of *gltA* was demonstrated for *B. grahamii* upon comparison of 36 strains from Asia ( $H_d = 0.943$ ,  $\pi = 0.02154$ ) and 27 strains from Europe and North America ( $H_d = 0.889$ ,  $\pi = 0.01427$ ). The nucleotide diversity of *gltA* is consistent with the result of Berglund et al., who analyzed 26 strains from 11 species of wild rodents in seven countries (32, 36). Therefore, our analysis corroborates previous speculation that the European and American lineages of *Bartonella* strains evolved as an expansion of the genetically diverse Asian clade. Further in-depth spatial and temporal sampling with evenly distributed statistical sampling will be required to demonstrate the ecological significance and value of these observations. Multigene locus analysis targeting genes with different levels of genetic conservation at the population, species, and genus levels or whole-genome level comparisons would be ideal for such an approach, but this will require substantial computational effort and multicenter cooperation to assemble data for a representative collection of isolates.

Previous investigations of *Bartonella* infections of small mammals and their ectoparasites were conducted in southeast and southwest China, including Zhejiang (37), Fujian (38), and Yunnan (22, 23, 39, 40) Provinces. In the present study, we determined the prevalence of *Bartonella* in wild rodents under the undisturbed natural ecological conditions on an unpopulated sedimentary island at the Chinese-Russian border in northeast China. Analysis of the geographical distribution pattern of *Bartonella* isolates from

FIG 4 Median-joining networks of *gltA* genotypes established based on maximum-parsimony calculation for *Bartonella* isolates from Heixiazi Island and all other isolates of homologous species whose nucleotide sequences are available in the NCBI GenBank database. Genotypes are colored according to their geographical origins and host species. The smallest colored circle corresponds to a single isolate. Isolates from Heixiazi Island are indicated by a secondary dashed-line circle. A1, network of *B. taylorii* according to its geographic distribution; A2, network of *B. taylorii* according to its host species associations; B1, network of *B. grahamii* according to its geographic distribution; B2, network of *B. grahamii* according to its host species associations; C1, network of *B. washoensis* according to its geographic distribution; C2, network of *B. washoensis* according to its host species associations.




FIG 5 Electron micrographs of the two novel *Bartonella* strains, CR90HXZ without pili (A) and AA137HXZ showing the presence of polar pili (B). Bars, 0.5  $\mu\text{m}$ .

Heixiazi Island compared to that of *Bartonella* from other geographic regions determined that the occurrence of various genotypes of at least three species of *Bartonella* (*B. grahamii*, *B. taylorii*, and *B. washoensis*) was associated with the geographic locations of their animal hosts. The impacts of host availability at each particular geographic location were different among different species of *Bartonella*. Furthermore, distribution patterns of different genotypes in a given population were indeed related to the host species and appeared to have a more critical role than the geographical factors. This is confirmed by an observation that the isolates of *B. taylorii* and *B. grahamii* on Heixiazi Island were not present in the same phylogenetic clusters but instead were clustered together with the strains detected in or isolated from the same animal hosts in various geographic areas. Another example was the isolates of *B. grahamii* from North America which did not belong to the same cluster because of their different host origins. Obviously, the internal distribution pattern of genotypes originated from same hosts was impacted by geographic locations. For instance, it was easy to observe that all presently known isolates of *B. grahamii* from *Apodemus* spp. are split into two clades; one clade includes isolates from Asia, and another clade includes isolates from Europe and North America. Based on these observations, we speculate that hosts and environmental factors may have different ef-

fects on genetic variations observed among isolates of *Bartonella* spp. at the strain and species levels. Consequently, to identify the effects of geographical factors on *Bartonella* isolates in a given population, analysis of intra- and interspecies variations should be performed for isolates of the same genus or species of rodent hosts collected in different geographic locations.

Since *Bartonella* spp. are facultative intracellular pathogens, the reservoir animal hosts can be viewed as an ecological niche which captures the range of environmental conditions enabling isolated persistence of these species (41). In theory, the variation of intracellular pathogens is low, and it likely to be the first impacted by availability of hosts inhabiting specific niches within certain geographic parameters; in contrast, it may be determined by the particular effects of the infecting parasite on its vertebrate host. Therefore, for *Bartonella* sp. diversity, host diversity appears to play a more important role than geographic factors. This hypothesis will need further evaluation and will require statistically justified and supported sampling to ensure representative collection of the isolates.

To fulfill the rules of the International Code of Nomenclature of Bacteria (42), we provide the following descriptions of the novel *Bartonella* species characterized in this study.

**Description of *Bartonella heixiaziensis* sp. nov.** *Bartonella heixiaziensis* (Hei.xia.zi.en.sis. N.L. gen. n. from Heixiazi Island [Bolshoi Ussuriysky Island], a sedimentary island at the confluence of the Ussuri and Amur Rivers in northeast China's Heilongjiang Province where the voles [*Myodes rutilus* and *Microtus maximowiczii*] from which the first strains were isolated were trapped).

Following inoculation on blood TSA, small, off-white, translucent, smooth, round colonies about 1 to 2 mm in diameter appeared at day 5 of incubation at 35°C in a 5% CO<sub>2</sub> humidified atmosphere. Best growth was observed on blood-enriched solid medium or Schneider's insect medium (liquid) with 10% fetal bovine serum in a moist atmosphere containing 5% CO<sub>2</sub>. Electron microscopic examination revealed small Gram-negative bacilli without flagella or pili and approximately 1 to 2  $\mu\text{m}$  long and 0.5 to 0.8  $\mu\text{m}$  wide (Fig. 5A). Bacteria were oxidase and catalase negative, and exhibited no urease activity. Tests for  $\alpha$ -glucosidase,  $\alpha$ -L-fucosidase,  $\alpha$ -L-arabino-furanosidase, mannose, raffinose, nitrate reduction, and spot indole were negative. Amino acid-arylamidase activity was observed with the following amino acids: arginine (Arg), phenylalanine (Phe), leucine (Leu), tyrosine (Tyr), alanine (Ala), glycine (Gly), histidine (His) and, serine (Ser). This new species is distinguished from other *Bartonella* species by 16S rRNA gene (KJ361623), *gltA* (KJ175047), *ftsZ* (KJ361705), and *rpoB* (KJ361744) nucleotide sequences. The type strain CR90HXZ was isolated from the blood of a red-backed vole (*Myodes rutilus*) collected in May 2011; it was deposited in the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures as DSM 100695 and in the China General Microbiological Culture Collection Center as CGMCC1.15048. Its closest known relative is *B. washoensis* as determined by molecular comparison of the four gene fragments (43).

**Description of *Bartonella fuyuanensis* sp. nov.** *Bartonella fuyuanensis* (fu.yuan.en.sis. N.L. gen. n. from Fuyuan Delta, which is the ancient name for Heixiazi Island in northeast China's Heilongjiang Province, where the animals were trapped).

Following inoculation on blood TSA, small, off-white, translucent, smooth, round colonies about 1 to 2 mm in diameter ap-

peared at day 5 of incubation at 35°C in a 5% CO<sub>2</sub> humidified atmosphere. Best growth was observed on blood-enriched solid medium or Schneider's insect medium (liquid) with 10% fetal bovine serum in a moist atmosphere containing 5% CO<sub>2</sub>. Electron microscopic examination revealed small Gram-negative bacilli, approximately 1 to 2 μm long by 0.5 to 0.8 μm wide with polar pili (Fig. 2B). Bacteria were oxidase and catalase negative and exhibited no urease activity. Tests for α-glucosidase, α-L-fucosidase, α-L-arabinofuranosidase, mannose, raffinose, nitrate reduction, and spot indole were negative. Amino acid arylamidase activity was observed with the following amino acids: arginine (Arg), phenylalanine (Phe), leucine (Leu), tyrosine (Tyr), alanine (Ala), glycine (Gly), histidine (His), and serine (Ser). This species is distinguished from other *Bartonella* species by 16S rRNA gene (KJ361607), *gltA* (KJ175033), *ftsZ* (KJ361689), and *rpoB* (KJ361730) nucleotide sequences. The type strain AA137HXZ was isolated from the blood of a wild striped field mouse (*Apodemus agrarius*) collected in May 2011; it is deposited in the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures as DSM 100694 and in the China General Microbiological Culture Collection Center as CGMCC1.15047. Its closest known relative is *B. rattimassiliensis* as determined by molecular comparison of the four gene fragments (43).

#### ACKNOWLEDGMENTS

This work was supported by National Critical Project for Science and Technology on Infectious Disease grant 2012ZX10004219 and Special Funds for Public Welfare Projects grant 201310072, People's Republic of China.

We declare that no competing interests exist.

#### REFERENCES

1. Buffet JP, Kosoy M, Vayssier-Taussat M. 2013. Natural history of *Bartonella*-infecting rodents in light of new knowledge on genomics, diversity and evolution. *Future Microbiol* 8:1117–1128. <http://dx.doi.org/10.2217/fmb.13.77>.
2. Huang R, Liu Q, Li G, Li D, Song X, Birtles RJ, Zhao F. 2011. *Bartonella quintana* infections in captive monkeys, China. *Emerg Infect Dis* 17:1707–1709. <http://dx.doi.org/10.3201/eid1709.110133>.
3. Li H, Bai JY, Wang LY, Zeng L, Shi YS, Qiu ZL, Ye HH, Zhang XF, Lu QB, Kosoy M, Liu W, Cao WC. 2013. Genetic diversity of *Bartonella quintana* in macaques suggests zoonotic origin of trench fever. *Mol Ecol* 22:2118–2127. <http://dx.doi.org/10.1111/mec.12261>.
4. Sun J, Fu G, Lin J, Song X, Lu L, Liu Q. 2010. Seroprevalence of *Bartonella* in eastern China and analysis of risk factors. *BMC Infect Dis* 10:121. <http://dx.doi.org/10.1186/1471-2334-10-121>.
5. Yuan C, Zhu C, Wu Y, Pan X, Hua X. 2011. Bacteriological and molecular identification of *Bartonella* species in cats from different regions of China. *PLoS Negl Trop Dis* 5:e1301. <http://dx.doi.org/10.1371/journal.pntd.0001301>.
6. Liu Q, Eremeeva ME, Li D. 2012. *Bartonella* and *Bartonella* infections in China: from the clinic to the laboratory. *Comp Immunol Microbiol Infect Dis* 35:93–102. <http://dx.doi.org/10.1016/j.cimid.2012.01.002>.
7. Norman AF, Regnery R, Jameson P, Greene C, Krause DC. 1995. Differentiation of *Bartonella*-like isolates at the species level by PCR-restriction fragment length polymorphism in the citrate synthase gene. *J Clin Microbiol* 33:1797–1803.
8. Li DM, Miao ZG, Song XP, Wang J, Liu QY. 2012. Optimization of liquid growth conditions and determination of growth curves for *Bartonella* species. *Microbiol China* 39:1695–1702.
9. Riess T, Dietrich F, Schmidt KV, Kaiser PO, Schwarz H, Schäfer A, Kempf VA. 2008. Analysis of a novel insect cell culture medium-based growth medium for *Bartonella* species. *Appl Environ Microbiol* 74:5224–5227. <http://dx.doi.org/10.1128/AEM.00621-08>.
10. Daly JS, Worthington MG, Brenner DJ, Moss CW, Hollis DG, Weyant RS, Steigerwalt AG, Weaver RE, Daneshvar MI, O'Connor SP. 1993. *Rochalimaea elizabethae* sp. nov. isolated from a patient with endocarditis. *J Clin Microbiol* 31:872–881.
11. Zeaiter Z, Liang Z, Raoult D. 2002. Genetic classification and differentiation of *Bartonella* species based on comparison of partial *ftsZ* gene sequences. *J Clin Microbiol* 40:3641–3647. <http://dx.doi.org/10.1128/JCM.40.10.3641-3647.2002>.
12. Renesto P, Gouvernet J, Drancourt M, Roux V, Raoult D. 2001. Use of *rpoB* gene analysis for detection and identification of *Bartonella* species. *J Clin Microbiol* 39:430–437. <http://dx.doi.org/10.1128/JCM.39.2.430-437.2001>.
13. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 30:2725–2729. <http://dx.doi.org/10.1093/molbev/mst197>.
14. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307–321. <http://dx.doi.org/10.1093/sysbio/syq010>.
15. Martin DP, Lemey P, Lott M, Moulton V, Posada D, Lefevre P. 2010. RDP3: a flexible and fast computer program for analyzing recombination. *Bioinformatics* 26:2462–2463. <http://dx.doi.org/10.1093/bioinformatics/btq467>.
16. Didlot X, Falush D. 2007. Inference of bacterial microevolution using multilocus sequence data. *Genetics* 175:1251–1266.
17. Huson DH, Bryant D. 2006. Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol* 23:254–267.
18. Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25:1451–1452. <http://dx.doi.org/10.1093/bioinformatics/btp187>.
19. Polzin T, Daneshmand SV. 2001. On Steiner trees and minimum spanning trees in hypergraphs. *Oper Res Lett* 31:12–20.
20. Bandelt HJ, Forster P, Rohl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol* 16:37–48. <http://dx.doi.org/10.1093/oxfordjournals.molbev.a026036>.
21. Forster P, Röhl A, Lünemann P, Brinkmann C, Zerjal T, Tyler-Smith C, Brinkmann B. 2000. A short tandem repeat-based phylogeny for the human Y chromosome. *Am J Hum Genet* 67:182–196. <http://dx.doi.org/10.1086/302953>.
22. Li DM, Yu DZ, Liu QY, Gong ZD. 2004. Study on the prevalence of *Bartonella* species in rodent hosts from different environmental areas in Yunnan. *Zhonghua Liu Xing Bing Xue Za Zhi* 25:934–937.
23. Bai HM, Yang FL, Yang H, Zhang Q. 2005. Study on *Bartonella* species in rodents in western Yunnan, China. *Zhonghua Liu Xing Bing Xue Za Zhi* 26:868–870.
24. Song XP, Liu QY, Lu L, Zhao W, Li G-C, Li D-M, Sun JM. 2010. Isolation and sequence analysis of *Bartonella* in small mammals in Hainan province. *Chin J Vect Biol Control* 21:131–133.
25. Inoue K, Kabeya H, Hagiya K, Kosoy MY, Une Y, Yoshikawa Y, Maruyama S. 2011. Multi-locus sequence analysis reveals host specific association between *Bartonella washoensis* and squirrels. *Vet Microbiol* 148:60–65. <http://dx.doi.org/10.1016/j.vetmic.2010.08.007>.
26. Zhang XL, Zhao ZJ, Li M, Guo XM, Yang J, Qu Y-M, Li Y-J, Wen ZQ, Hu K-X, Sun XH. 2011. Survey of rodent diversity in different habitats in the central border area of China and Russia. *Chin J Hyg Insect Equip* 17:34–36.
27. Chiaraviglio L, Duong S, Brown DA, Birtles RJ, Kirby JE. 2010. An immunocompromised murine model of chronic *Bartonella* infection. *Am J Pathol* 176:2753–2763. <http://dx.doi.org/10.2353/ajpath.2010.090862>.
28. Liston TE, Je K. 1996. Granulomatous hepatitis and necrotizing splenitis due to *Bartonella henselae* in a patient with cancer: case report and review of hepatosplenic manifestations of *Bartonella* infection. *Clin Infect Dis* 22:951–957. <http://dx.doi.org/10.1093/clinids/22.6.951>.
29. Klotz SA. 2011. Cat-scratch disease. *Am Fam Physician* 83:152–155.
30. Hutchins RG, Breitschwerdt EB, Cullen JM, Bissett SA, Gookin JL. 2012. Limited yield of diagnoses of intrahepatic infectious causes of canine granulomatous hepatitis from archival liver tissue. *J Vet Diagn Invest* 24:888–894. <http://dx.doi.org/10.1177/1040638712453583>.
31. Deng HK, Le Rhun D, Lecuelle B, Le Naour E, Vayssier-Taussat M. 2012. Role of the spleen in *Bartonella* spp. infection. *FEMS Immunol Med Microbiol* 64:143–145. <http://dx.doi.org/10.1111/j.1574-695X.2011.00908.x>.
32. Berglund EC, Ellegaard K, Granberg F, Xie Z, Maruyama S, Kosoy MY, Birtles RJ, Sg A. 2010. Rapid diversification by recombination in *Bartonella grahamii* from wild rodents in Asia contrasts with low levels of



- genomic divergence in Northern Europe and America. *Mol Ecol* 19:2241–2255. <http://dx.doi.org/10.1111/j.1365-294X.2010.04646.x>.
33. Paziewska A, Siński E. 2011. Recombination within and between species of the alpha Proteobacterium *Bartonella* infecting rodents. *Microb Ecol* 61:134–145. <http://dx.doi.org/10.1007/s00248-010-9735-1>.
  34. Posada D, Ka C. 2002. The effect of recombination on the accuracy of phylogeny estimation. *J Mol Evol* 54:396–402. <http://dx.doi.org/10.1007/s00239-001-0034-9>.
  35. Buffet JP, Pisanu B, Brisse S, Roussel S, Félix B, Halos L, Chapuis JL, Vayssier-Taussat M. 2013. Deciphering *Bartonella* diversity, recombination, and host specificity in a rodent community. *PLoS One* 8:e68956. <http://dx.doi.org/10.1371/journal.pone.0068956>.
  36. Birtles RJ, Harrison TG, Molyneux DH. 1994. *Grahamella* in small woodland mammals in the U.K.: isolation, prevalence and host specificity. *Ann Trop Med Parasitol* 88:317–327.
  37. Liu Q, Sun J, Lu L, Fu G, Ding G, Song X, Meng F, Wu H, Yang T, Ren Z. 2010. Detection of *Bartonella* species in small mammals from Zhejiang Province, China. *J Wildl Dis* 46:179–185. <http://dx.doi.org/10.7589/0090-3558-46.1.179>.
  38. Ye X, Li GW, Yao ML, Luo W, Su LQ. 2009. Study on the prevalence and genotypes of *Bartonella* species in rodent hosts from Fujian coastal region. *Zhong Hua Liu Xing Bing Xue Za Zhi* 30:989–992.
  39. Ying B, Kosoy MY, Maupin GO, Tsuchiya KR, Gage KL. 2002. Genetic and ecologic characteristics of *Bartonella* communities in rodents in southern China. *Am J Trop Med Hyg* 66:622–627.
  40. Li DM, Liu QY, Yu DZ, Zhang JZ, Gong ZD, Song XP. 2007. Phylogenetic analysis of *Bartonella* detected in rodent fleas in Yunnan, China. *J Wildl Dis* 43:609–617. <http://dx.doi.org/10.7589/0090-3558-43.4.609>.
  41. Casadevall A. 2008. Evolution of intracellular pathogens. *Annu Rev Microbiol* 62:19–33. <http://dx.doi.org/10.1146/annurev.micro.61.080706.093305>.
  42. Lapage SP, Sneath PHA, Lessel EF, Skerman VBD, Seeliger HPR, Clark WA (ed). 1992. International code of nomenclature of bacteria: bacteriological code, 1990 revision. ASM Press, Washington, DC.
  43. La Scola B, Zeaiter Z, Khamis A, Raoult D. 2003. Gene-sequence-based criteria for species definition in bacteriology: the *Bartonella* paradigm. *Trends Microbiol* 11:318–321. [http://dx.doi.org/10.1016/S0966-842X\(03\)00143-4](http://dx.doi.org/10.1016/S0966-842X(03)00143-4).