

Fate of MgSiO₃ melts at core–mantle boundary conditions

Sylvain Petitgirard^{a,1}, Wim J. Malfait^b, Ryosuke Sinmyo^a, Ilya Kuznetsov^{a,c}, Louis Hennet^d, Dennis Harries^e, Thomas Dane^c, Manfred Burghammer^{c,f}, and Dave C. Rubie^a

^aBayerisches Geoinstitut, University of Bayreuth, Bayreuth D-95490, Germany; ^bLaboratory for Building Energy Materials and Components, Swiss Federal Laboratories for Materials Science and Technology, 8600 Dübendorf, Switzerland; ^cEuropean Synchrotron Radiation Facility, BP 220, Grenoble F-38043, France; ^dConditions Extrêmes et Matériaux: Haute Température et Irradiation, CNRS UPR 3079, Université d'Orléans, 45071 Orléans Cedex 2, France; ^eAnalytical Mineralogy of Micro- and Nanostructures, Institute of Geoscience, Friedrich Schiller University Jena, Jena 07745, Germany; and ^fDepartment of Analytical Chemistry, Ghent University, B-9000 Ghent, Belgium

Edited by Russell J. Hemley, Carnegie Institution of Washington, Washington, DC, and approved October 7, 2015 (received for review June 25, 2015)

One key for understanding the stratification in the deep mantle lies in the determination of the density and structure of matter at high pressures, as well as the density contrast between solid and liquid silicate phases. Indeed, the density contrast is the main control on the entrainment or settlement of matter and is of fundamental importance for understanding the past and present dynamic behavior of the deepest part of the Earth's mantle. Here, we adapted the X-ray absorption method to the small dimensions of the diamond anvil cell, enabling density measurements of amorphous materials to unprecedented conditions of pressure. Our density data for MgSiO₃ glass up to 127 GPa are considerably higher than those previously derived from Brillouin spectroscopy but validate recent ab initio molecular dynamics simulations. A fourth-order Birch–Murnaghan equation of state reproduces our experimental data over the entire pressure regime of the mantle. At the core–mantle boundary (CMB) pressure, the density of MgSiO₃ glass is 5.48 ± 0.18 g/cm³, which is only 1.6% lower than that of MgSiO₃ bridgmanite at 5.57 g/cm³, i.e., they are the same within the uncertainty. Taking into account the partitioning of iron into the melt, we conclude that melts are denser than the surrounding solid phases in the lowermost mantle and that melts will be trapped above the CMB.

silicate glass density | X-ray absorption | basal magma ocean

Variations in seismic velocities observed in the deep mantle could be attributed to melting phenomena (1), accumulation of dense matter (2), core–mantle interactions (3), or even a deep hidden geochemical reservoir that dates from the early differentiation of the Earth (4). Indeed, melting phenomena play a critical role in the formation and evolution of terrestrial planets. In the early Earth's history, large-scale melting events and magma oceans facilitated the segregation of the iron-rich core and the partitioning of elements between the different layers of the Earth (5). Catastrophic events like the moon-forming giant impact may have melted the entire Earth and modified the thermal and chemical state of the planet's interior (6). It has long been thought that the mantle crystallized from the bottom to the top, with crystals being denser than melts (7). However, the formation of a dense basal magma ocean (BMO) at the bottom of the mantle concomitant to the final accretion of the Earth was proposed as a consequence of partial melting (2). The formation of a BMO requires the accumulation of dense iron-rich silicate melts at the core–mantle boundary (CMB). In this scenario, the mantle starts to crystallize at an intermediate depth, and a raft of crystals continues to grow toward both the bottom and top of the mantle. In the modern mantle, seismic observations highlight ultralow velocity zones (ULVZs) near the CMB (8) that may be caused by melting of deep mantle material (9), potentially feeding sources of hot spots. Alternatively, the ULVZs may be dense remnants from an initial BMO (10, 11). Both scenarios for the formation of the Earth's mantle and its geodynamical evolution critically depend on the density and fate of silicate melts at high pressure. Indeed, the main parameter that controls the entrainment and/or settling of materials in the deep

Earth is the density contrast between crystals and melts at relevant pressures (P) and temperatures (T). Thus, knowledge of the densities of melts and their equation of state (EoS) is of prime importance to understand the past and present structures of the lowermost part of the mantle. However, high-pressure density data are lacking due to experimental challenges. Although many experiments on both melts and glasses densities have been carried out in large volume apparatus to pressures of 20 GPa (12, 13), the few studies conducted in the diamond anvil cell (DAC) were limited to 50–60 GPa (14, 15). Studying glasses is a potential alternative for understanding melts because many of their physical properties are similar to those of melt: from the atomistic scale, with the evolution of the silicon coordination number from fourfold to sixfold from 0 to 40 GPa (15), to the macroscopic scale with similarities in compressibility (16, 17). Using the X-ray absorption technique adapted to the DAC, we more than double the pressure range for density measurements on amorphous materials. We extend the dataset for MgSiO₃ glass to CMB pressures and propose an EoS for MgSiO₃ melts that spans the entire mantle.

The MgSiO₃ glass starting materials were prepared using an aerodynamic levitation setup (*SI Appendix, Fig. S1*) and measured in situ at high pressure at the ID13 beamline of the European Synchrotron Radiation Facility (ESRF; Grenoble, France) (*SI Appendix, Fig. S3 A and B*). Samples were pressurized in DACs and sealed in beryllium (Be) gaskets giving radial access for the X-ray beam. For pressures below 20 GPa, discs of MgSiO₃ glass were cut using a focus ion beam (FIB) (*SI Appendix, Fig. S2*), loaded in a Be gasket, and immersed in a methanol–ethanol mixture. For pressures above 20 GPa, the sample chamber was filled with MgSiO₃ glass powder only. Pressure

Significance

A new technique has been developed to measure in situ the density of amorphous material composed of light elements under extreme conditions of pressure using the X-ray absorption method. At core–mantle boundary (CMB) pressure, the densities of MgSiO₃ glass and melts are similar to the one of the crystalline bridgmanite, within uncertainty. Due to the affinity of iron oxide for silicate liquids, melting in the MgSiO₃–FeSiO₃ system will produce dense melts that could accumulate above the CMB, leading to the formation of a dense basal magma ocean in the early Earth's mantle.

Author contributions: S.P. and W.J.M. designed research; S.P., W.J.M., R.S., I.K., T.D., and M.B. performed research; S.P., L.H., and D.H. contributed new reagents/analytic tools; S.P. and W.J.M. analyzed data; and S.P., W.J.M., R.S., and D.C.R. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. Email: sylvain.petitgirard@uni-bayreuth.de.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1512386112/-DCSupplemental.




Fig. 3. (A) MgSiO_3 melt density along different isotherms. From the fourth-order BM EoS at 300 K (blue curve), we extrapolate our results to high temperature using the thermal expansion of MgSiO_3 molten. Also reported are the results for MgSiO_3 melts at 4,000 K calculated by Stixrude and Karki (21) (dashed black curve), as well as the EoS for MgSiO_3 bridgmanite (29) at 300 and 4,000 K (29). (B) The density cross-over pressure in the MgSiO_3 - FeSiO_3 system. The double arrow represents the range of experimental data for the lower mantle KD.

computed by de Koker et al. (22) for αK_T , which is independent of temperature but varies with volume and can be fitted using a power law curve (SI Appendix, Fig. S10). Our experimental isotherms agree with the isotherms from MD simulations along different isotherms (19, 21, 23) (Fig. 3A). At 4,000 K and 135 GPa, the density contrast between molten and crystalline MgSiO_3 (bridgmanite) is only 1.6% (Fig. 3A). The small density contrast highlights that density cross-overs are probable, and melts can be as dense or even denser than the surrounding solids in the deep mantle, regardless of the presence of a poly-amorphous phase transition near 133 GPa (11), which may lead to a density inversion at even higher pressure as highlighted by shock compression experiments (24) (SI Appendix, Fig. S11). The buoyancy of melts in the deep Earth is strongly influenced by the addition of iron as the main heavy element. Although the partitioning of iron (K_D) between solid and melts remains controversial at lower mantle pressures (9, 25), it is agreed that iron is an incompatible element. Therefore, iron is enriched in the melt phase with a K_D value as low as ~ 0.4 – 0.3 at 25 GPa (26, 27) and most likely even lower at higher pressures. Given that even in the absence of iron, MgSiO_3 melt is already nearly as dense as its crystalline counterpart at CMB conditions (Fig. 3A), melts enriched in iron will be more dense than their equilibrium minerals, regardless of the exact nature of the iron partitioning. With our MgSiO_3 density data at 4,000 K and FeSiO_3 density data at 4,000 K from Ramo and Stixrude (28), we calculated the density cross-over pressure in the MgSiO_3 - FeSiO_3 system as a function of iron partitioning. For the available iron partitioning data, a

density cross-over will occur in the lower mantle between solid bridgmanite (29) and liquid with a similar composition at 4,000 K (Fig. 3B). However, the thickness of the layer where melts are denser than the solids varies from 1,600 km to only a few tens/hundreds of kilometers depending on the iron partitioning behavior between melts and bridgmanite. For super-Earths, which may have stable postperovskite at pressures above 135 GPa and 4,000 K, the evaluation of density cross-over would require density data of postperovskite end members and iron partitioning behavior for this phase.

In conclusion, our experimental density data at CMB pressures confirm the viability of a BMO and the possibility of melts being trapped in the lowermost mantle. Thus, partial melting in the deepest part of the Earth will favor the enrichment and storage of incompatible radiogenic elements at the CMB (27). These trapped melts, or their Fe-rich crystallization products, may explain the large low shear velocity provinces (LLSVPs) area and ULVZs above the CMB and constitute an ideal candidate for a hidden geochemical reservoir.

ACKNOWLEDGMENTS. We thank H. Shultz for the polishing of the starting material prior to the focus ion beam cuttings; D. J. Frost, G. Steinle-Neumann, L. Dubrovinsky, and S. Jacobsen for fruitful discussion and comments; two anonymous reviewers for comments; the European Synchrotron Radiation Facility for provision of beamtime under Proposal E5-163 and beamlines ID27 and ID09a for both Raman systems, portable and off-line respectively. S.P. and R.S. are financed by the Humboldt Foundation. S.P. and D.C.R. are supported by the European Research Council Advanced Grant “ACCRETE” (Contract 290568).

- Fiquet G, et al. (2010) Melting of peridotite to 140 gigapascals. *Science* 329(5998):1516–1518.
- Labrosse S, Hernlund J, Coltice N (2007) A crystallizing dense magma ocean at the base of the Earth’s mantle. *Nature* 450(7171):866–869.
- Otsuka K, Karato S (2012) Deep penetration of molten iron into the mantle caused by a morphological instability. *Nature* 492(7428):243–246.
- Boyet M, Carlson R (2005) ^{142}Nd evidence for early (>4.53 Ga) global differentiation of the silicate Earth. *Science* 309(5734):341–348.
- Rubie DC, et al. (2011) Heterogeneous accretion, composition and core-mantle differentiation of the Earth. *Earth Planet Sci Lett* 301(1-2):31–42.
- Jacobson SA, et al. (2014) Highly siderophile elements in Earth’s mantle as a clock for the Moon-forming impact. *Nature* 508(7494):84–87.
- Solomatov VS, Stevenson DJ (1993) Suspension in convective layers and style of differentiation of a terrestrial magma ocean. *J Geophys Res* 98(E3):5375–5390.
- Williams Q, Garnero EJ (1996) Seismic evidence for partial melt at the base of Earth’s mantle. *Science* 273(5281):1528–1530.
- Andraut D, et al. (2012) Solid-liquid iron partitioning in Earth’s deep mantle. *Nature* 487(7407):354–357.
- Lay T, Garnero EJ, Williams Q (2004) Partial melting in a thermo-chemical boundary layer at the base of the mantle. *Phys Earth Planet Inter* 146(3-4):441–467.
- Murakami M, Bass JD (2011) Evidence of denser MgSiO_3 glass above 133 gigapascal (GPa) and implications for remnants of ultradense silicate melt from a deep magma ocean. *Proc Natl Acad Sci USA* 108(42):17286–17289.
- Malfait WJ, et al. (2014) Supervolcano eruptions driven by melt buoyancy in large silicic magma chambers. *Nat Geosci* 7(2):122–125.
- Sakamaki T, Suzuki A, Ohtani E (2006) Stability of hydrous melt at the base of the Earth’s upper mantle. *Nature* 439(7073):192–194.
- Sato T, Funamori N (2008) Sixfold-coordinated amorphous polymorph of SiO_2 under high pressure. *Phys Rev Lett* 101(25):255502.
- Sanloup C, et al. (2013) Structural change in molten basalt at deep mantle conditions. *Nature* 503(7474):104–107.
- Wang Y, et al. (2014) Atomistic insight into viscosity and density of silicate melts under pressure. *Nat Commun* 5:3241.
- Sakamaki T, et al. (2014) Contrasting sound velocity and intermediate-range structural order between polymerized and depolymerized silicate glasses under pressure. *Earth Planet Sci Lett* 391:288–295.
- Sanchez-Valle C, Bass JD (2010) Elasticity and pressure-induced structural changes in vitreous MgSiO_3 -enstatite to lower mantle pressures. *Earth Planet Sci Lett* 295(3-4):523–530.

19. Ghosh DB, Karki B, Stixrude L (2014) First-principles molecular dynamics simulations of MgSiO_3 glass: Structure, density, and elasticity at high pressure. *Am Mineral* 99(5-6): 1304–1314.
20. Rivers ML, Carmichael ISE (1987) Ultrasonic studies of silicate melts. *J Geophys Res* 92(B9):9247–9270.
21. Stixrude L, Karki B (2005) Structure and freezing of MgSiO_3 liquid in Earth's lower mantle. *Science* 310(5746):297–299.
22. de Koker N, Stixrude L (2009) Self-consistent thermodynamic description of silicate liquids, with application to shock melting of MgO periclase and MgSiO_3 perovskite. *Geophys J Int* 178(1):162–179.
23. Stixrude L, de Koker N, Sun N, Mookherjee M, Karki B (2009) Thermodynamics of silicate liquids in the deep Earth. *Earth Planet Sci Lett* 278(3-4):226–232.
24. Mosenfelder JL, Asimow PD, Frost DJ, Rubie DC, Ahrens TJ (2009) The MgSiO_3 system at high pressure: Thermodynamic properties of perovskite, postperovskite, and melt from global inversion of shock and static compression data. *J Geophys Res* 114(B01203):1–16.
25. Nomura R, et al. (2011) Spin crossover and iron-rich silicate melt in the Earth's deep mantle. *Nature* 473(7346):199–202.
26. Tronnes RG, Frost DJ (2002) Peridotite melting and mineral-melt partitioning of major and minor elements at 22–24.5 GPa. *Earth Planet Sci Lett* 197(1-2):117–131.
27. Hirose K, Shimizu N, van Westrenen W, Fei YW (2004) Trace element partitioning in Earth's lower mantle and implications for geochemical consequences of partial melting at the core-mantle boundary. *Phys Earth Planet Inter* 146(1-2): 249–260.
28. Ramo DM, Stixrude L (2014) Spin crossover in Fe_2SiO_4 liquid at high pressure. *Geophys Res Lett* 41(13):4512–4518.
29. Xu W, Lithgow-Bertelloni C, Stixrude L, Ritsema J (2008) The effect of bulk composition and temperature on mantle seismic structure. *Earth Planet Sci Lett* 275(1-2):70–79.