

SCIENTIFIC REPORTS



OPEN

Impact of a wastewater treatment plant on microbial community composition and function in a hyporheic zone of a eutrophic river

Received: 02 June 2015

Accepted: 28 October 2015

Published: 26 November 2015

Siavash Atashgahi^{1,2,3}, Rozelin Aydin^{1,*}, Mauricio R. Dimitrov^{1,*}, Detmer Sijkema^{1,*}, Kelly Hamonts^{2,*}, Leo Lahti^{1,4}, Farai Maphosa¹, Thomas Kruse¹, Edoardo Saccenti⁵, Dirk Springael³, Winnie Dejonghe² & Hauke Smidt¹

The impact of the installation of a technologically advanced wastewater treatment plant (WWTP) on the benthic microbial community of a vinyl chloride (VC) impacted eutrophic river was examined two years before, and three and four years after installation of the WWTP. Reduced dissolved organic carbon and increased dissolved oxygen concentrations in surface water and reduced total organic carbon and total nitrogen content in the sediment were recorded in the post-WWTP samples. Pyrosequencing of bacterial 16S rRNA gene fragments in sediment cores showed reduced relative abundance of heterotrophs and fermenters such as *Chloroflexi* and *Firmicutes* in more oxic and nutrient poor post-WWTP sediments. Similarly, quantitative PCR analysis showed 1–3 orders of magnitude reduction in phylogenetic and functional genes of sulphate reducers, denitrifiers, ammonium oxidizers, methanogens and VC-respiring *Dehalococcoides mccartyi*. In contrast, members of *Proteobacteria* adapted to nutrient-poor conditions were enriched in post-WWTP samples. This transition in the trophic state of the hyporheic sediments reduced but did not abolish the VC respiration potential in the post-WWTP sediments as an important hyporheic sediment function. Our results highlight effective nutrient load reduction and parallel microbial ecological state restoration of a human-stressed urban river as a result of installation of a WWTP.

Rivers in urban areas are often heavily impacted by untreated wastewater discharge containing excessive organic inputs that can induce anoxia, eutrophication of freshwater ecosystems and severe reduction in water quality¹. Half of the world's population lived in urban areas in 2008². This number is predicted to increase to 66% by 2050 mainly due to urban growth in less developed countries³, expected to lead to a similar increase in the production of nutrient rich urban waste water. Wastewater treatment plants (WWTPs) are one of the most common measures of modern environmental biotechnology to treat municipal wastewater. Wastewater treatment removes a large fraction of nutrients, before the resulting effluent is discharged into receiving water bodies⁴. WWTP-induced reduction in anthropogenic nutrient input into fresh water ecosystems is in line with the European Water Framework Directive (WFD) that

¹Laboratory of Microbiology, Wageningen University, Dreijenplein 10, 6703 HB Wageningen, The Netherlands.

²Flemish Institute for Technological Research (VITO), Separation and Conversion Technology, Boeretang 200, 2400 Mol, Belgium. ³KU Leuven, Division Soil and Water Management, Kasteelpark Arenberg 20, 3001 Heverlee, Belgium. ⁴Department of Veterinary Biosciences, University of Helsinki, PO Box 66, FI-00014, Finland.

⁵Laboratory of Systems and Synthetic Biology, Wageningen University, Dreijenplein 10, 6703 HB, Wageningen, The Netherlands. [†]Present address: Department of Bioengineering, Adana Science and Technology University, 01180 Adana, Turkey. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to S.A. (email: siavash.atashgahi@wur.nl)

aims at protection of aquatic ecosystems and optimisation of water quality to achieve good ecological and chemical status of these ecosystems⁵. Accordingly, 20–80% of WWTPs in European countries have implemented tertiary treatment that uses technological advances for chemical and biological removal of recalcitrant organic pollutants as well as inorganic nutrients such as nitrogen and phosphorus⁶.

Parallel to reducing nutrient load into downstream receiving ecosystems, complementary measures are mandatory to monitor their ecological state. How will reducing the nutrient concentrations of treated effluent affect the microbial structure and function in the receiving environment? How will it impact the trophic state of the ecosystem and processes dependent on otherwise high nutrient load? Allochthonous organic carbon for example fuels river biogeochemical activities⁷, which mainly take place in the riverbed sediment, either at or just below the surface, including the zone where shallow groundwater mixes with surface water known as the hyporheic zone⁸. The high organic matter content in hyporheic zones allows multiple critical reactions in biogeochemical cycling and hence natural attenuation of inorganic and organic pollutants present in upwelling contaminated groundwater plumes⁹. For instance, in recent years, the hyporheic zone has received much attention because of its capacity to retain or degrade groundwater contaminants like chlorinated aliphatic hydrocarbons (CAHs)^{10–13}.

In spite of intensive studies on microbial community composition and function in hyporheic zones^{14–25}, far less attention has been paid to the impact of WWTPs effluent on the microbial ecology of riverbed sediments^{26–28}. Decreased microbial diversity^{26,27} and altered biogeochemical cycling of nitrogen²⁸ were reported in downstream effluent impacted sediments compared to samples taken upstream of the WWTP. However, the outcome at other locations may differ from these observations, as the impact of a WWTP on the downstream sediment microbiota depends on different factors such as type of the WWTP and hence the composition of the effluent, the population size of the urban area, climate and geography of the region, the size and flow rate of the river, and the buffering and self-purification capacity of the stream sediments to stresses.

In the present study, we investigated microbial community composition and function across sediment depth in a CAH impacted hyporheic zone of a short stretch of the Belgian Zenne River, as a response to an upstream WWTP installation. The Zenne river was notorious for being one of Belgium's most polluted rivers receiving untreated municipal effluents from the Brussels Capital Region leading to severe eutrophication of the river. In previous studies performed at that river stretch between 2004 and 2007, during which the Zenne surface water was eutrophic, the highly reducing and organic rich sediments were shown to be conducive to natural attenuation of CAHs present in the discharging groundwater mainly by organohalide respiration (OHR)^{12,13,29}. Since March 2007, a technologically advanced WWTP was installed approximately one km upstream of the test site to treat the municipal wastewater produced by 1.4 million inhabitants prior to its discharge into the river (www.aquaris.be). The biological treatment line of the plant consists of a primary settling stage and a subsequent modern tertiary treatment technology for elimination of nitrogen, phosphorus and carbon-containing pollution by an activated sludge process. The implementation of the WWTP resulted in important chemical changes in the surface water composition such as decrease in organic carbon content and increase in dissolved oxygen concentration. For a conclusive assessment of the impact of the installation of a WWTP on the receiving river ecosystem, we studied the sediment microbial community as indicator for system recovery and impact on particular ecosystem function i.e. vinyl chloride (VC) respiration. In contrast to former studies based on differences in upstream and downstream sediment microbial community where no information was available about the pre-WWTP conditions^{26–28}, we sampled our intensively studied test site two years before (2005) and three and four years after (2010 and 2011, respectively) construction of the WWTP.

Results

Chemical characterization of the study site. No obvious trends were observed in surface water temperature, concentrations of anions and cations (except Fe) in the surface water during the experimental period (Table 1): none of the q -values for the Spearman ρ correlation was found significant at the 0.25 level after Benjamini-Hochberg correction for multiple testing (Supplementary Table S3). However, dissolved organic carbon (DOC) ($\rho = -0.89$, q -value = 0.09), oxygen ($\rho = +0.71$, q -value = 0.09), pH ($\rho = -0.66$, q -value = 0.09), Fe ($\rho = -0.7$, q -value = 0.14) and conductivity ($\rho = -0.52$, q -value = 0.24) showed significant differences. In particular we observed that the dissolved oxygen content in the surface water increased to above 3 mg/L in 2007 and on average above 4 mg/L after 2008 (q -value = 0.2, Kruskal-Wallis test). DOC concentrations in the surface water decreased from an average of 51 ± 27 mg/L in the period Dec 2005-May 2006 to an average of 7 ± 1 mg/L in May-Sept 2011 (q -value = 0.24). The DOC and sulphate values in interstitial water (ISW) collected by means of permanent pore water samplers were not significantly different (Table 2). The concentrations of nitrate and nitrite were always below detection limit. The total organic carbon (TOC) significantly decreased with depth (Fig. 1a, Supplementary Table S4,5) in most cores except for core-A in 2010 ($\rho = -0.83$, q -value = 0.29) and core-B in 2011 ($\rho = 0.29$, q -value = 0.87). The total nitrogen (TN) content on the other hand decreased with depth only for core-B in 2005 ($\rho = -0.67$, q -value = 0.19) and core-A in 2010 ($\rho = -0.89$, q -value = 0.19). A significant decrease over time for both TOC and TN values was observed only in the top 20 cm sediment horizon (Kruskal-Wallis q -values of 0.01 and 0.01, respectively) (Supplementary Table S6).

Date	Concentration (mg/L)									DOC (mg/L)	O ₂ (mg/L)	Temp (°C)	pH	Conductivity (µS/cm)
	PO ₄	SO ₄	NO ₂	NO ₃	Cl	Fe	Na	Mg	K					
Dec 2005	— ^a	—	—	—	—	—	—	—	—	32	1.2	—	—	—
May 2006	2.32	214	BD ^b	0.68	258	0.51	90	15	18	70	—	17.8	7.88	1126
Dec 2006	—	—	—	—	—	—	—	—	—	—	1.9	14.5	7.62	1576
Mar 2007	0.67	103	BD	7.52	95	—	—	—	—	—	3.6	12.9	7.62	1542
May 2007	1.97	58	BD	BD	148	0.58	80.3	14.9	16.2	—	3.7	13.2	7.58	1368
Nov 2007	0.82	101	BD	6.70	79	0.09	44.7	13.7	9.6	—	—	7.5	8.03	791
Oct 2008	—	73	BD	BD	61	0.11	—	—	—	—	5.9	15.7	8.46	952
Feb 2010	—	65	—	13.9	92	0.08	63	11	10	15.85	5.5	11.5	7.32	911
May 2010	—	111	0.56	10.4	104	0.02	81	17	13	19.73	4.2	13.9	7.47	1122
Sep 2010	1.80	102	BD	6.14	53	0.01	71	14	15	13.12	4.4	18.2	7.42	962
Jan 2011	—	—	—	—	—	—	—	—	—	—	6.5	7.4	7.42	1070
May 2011	—	100	—	—	140	0.17	—	—	—	6.55	4.7	18	7.38	1211
Sep 2011	1.10	78	0.15	2.4	79	0.01	75	12	9.7	7.86	5.3	17.1	7.47	777

Table 1. Physico-chemical parameters of surface water. The dates of sediment core samplings are shown in boldface and the date of WWTP installation (Mar 2007) is shown in bold/italic. ^aNo data. ^bBelow detection limit.

Date	Sampling depth (cm)	Concentration (mg/L)				
		DOC	SO ₄	NO ₂	NO ₃	VC
Dec 2005	20	5	170	BD ^a	BD	0.02
	60	4	198	BD	BD	1.2
May 2010	20	3.7	145	BD	BD	0
	60	3.5	144	BD	BD	0
Sep 2011	20	6.3	140	BD	BD	0
	60	2.6	140	BD	BD	0

Table 2. Physico-chemical parameters of interstitial water sampled from the permanent pore water samplers in the riverbed. ^aBD: Below the detection limit.

Quantification of phylogenetic and functional gene markers in sediment cores. The highest bacterial 16S rRNA gene numbers were detected in surficial layers of both cores taken in 2005 with numbers above 10^7 copies per gram of sediment (Fig. 1b, Supplementary Table S4). Bacterial 16S rRNA gene numbers in both 2005 cores significantly declined with depth to below 10^5 per gram of sediment in deeper layers while no significant depth-related reduction was noted in post-WWTP cores (Supplementary Table S5). Bacterial 16S rRNA gene numbers showed a significant reduction over time in post-WWTP cores both in top and bottom sediment horizons (Kruskall-Wallis q-values of 0.01 and 0.07, respectively) (Supplementary Table S6). Similar to the bacterial trends, 16S rRNA gene copy numbers of *Dehalococcoides mccartyi* showed significant reduction with depth in the pre-WWTP core samples (Supplementary Table S5) while reduction over time was only significant in the top sediment horizon (Kruskall-Wallis q-value = 0.01). The abundance of functional gene biomarkers for sulphate reducers (*dsrB*), methanogens (*mcrA*), bacterial/archaeal ammonium oxidizers (bacterial/archaeal *amoA*) and denitrifiers (*nirK*, *nirS*, *nosZ*) significantly decreased over time in post-WWTP (Supplementary Fig. S2, Supplementary Table S6–8).

Occurrence of VC respiration activity in sediment microcosms. VC respiration was compared between samples taken before and after the installation of the WWTP in microcosms prepared from the top 20 cm sediments and groundwater from the SB1 monitoring well (Fig. 2). VC dechlorination was observed in microcosms containing sediment material sampled in 2005 and in 2010 (Fig. 2a) after seven days and, stoichiometric quantities of ethene were produced within 28 days (Fig. 2b). VC dechlorination in microcosms prepared with sediment material sampled in 2011 started after a lag phase of 35 days, and complete VC dechlorination to ethene was observed after 70 days. Addition of tree bark to sediments

from 2011 reduced the lag phase to 14 days, and all VC was stoichiometrically dechlorinated to ethene within 35 days.

Composition and distribution of the most abundant bacterial taxa. Prior to pyrosequencing, DNA extracts of a total of 80 sediment core slices at 5 cm intervals were analyzed by 16S rRNA gene-based PCR-DGGE bacterial community fingerprinting. Based on the obtained patterns (Supplementary Fig. S1), a total of 40 representative samples were selected for pyrosequencing of the PCR-amplified V1-V2 region of the bacterial 16S rRNA gene. A total of 309191 reads was obtained for all 40 samples. After quality filtering, denoising, and chimera removal, a total of 269411 reads was recovered ranging from 967 to 20335 reads per sample (Supplementary Table S9), with average and median of 6739 and 6068 reads per sample, respectively.

Proteobacteria, *Firmicutes*, *Chloroflexi*, *Bacteroidetes*, *Nitrospirae*, *Fusobacteria*, and *Actinobacteria* were the seven most abundant phyla in all samples (Fig. 3). The *Chloroflexi* phylum was the most abundant in the surface layers of the pre-WWTP samples and generally declined with depth (Fig. 3, right), which was statistically significant in core-B ($\rho = -0.77$, q-value = 0.18). In contrast, *Nitrospira* had higher relative abundance in deeper sediment layers ($\rho = 0.8$, q-value = 0.16). No significant depth-related stratification of bacterial community composition was noted in post-WWTP samples at the phylum level (data not shown). However, significant differences were observed between pre- and post-WWTP samples. The relative abundance of *Chloroflexi* and *Firmicutes* significantly decreased in post-WWTP samples while those of *Nitrospirae* and *Proteobacteria* increased significantly (Supplementary Table S10). At the order level, *Anaerolineales* (*Anaerolineae* class, Supplementary Fig. S3), *Lactobacillales* (*Bacilli* class), *Clostridiales* (*Clostridia* class), *Rhodocyclales* (*Betaproteobacteria* class) and *Desulfobacterales* (*Deltaproteobacteria* class) showed a significant reduction in relative abundance in post-WWTP samples (Supplementary Fig. S4, Supplementary Table S10). In contrast, *Burkholderiales* (*Betaproteobacteria* class) and *Xanthomonadales* (*Gammaproteobacteria* class) significantly increased in relative abundance over time in post-WWTP samples (Supplementary Table S10).

Bacterial diversity and richness indices. In the sediment cores sampled in 2005, the alpha diversity (Shannon) and richness (Chao1 and OTU count) indices did not show significant difference between two cores (Fig. 4, Supplementary Table S11). Similarly, the alpha diversity and richness of the post-WWTP samples of core-B of 2010 and core-B of 2011 did not show significant differences with the pre-WWTP samples. However, the alpha diversity and richness indices were the highest and lowest in the core-A of 2010 and core-A of 2011, respectively (Fig. 4, Supplementary Table S11).

Correlation between environmental parameters and microbial community composition. We performed multivariate analysis (RDA) to study the relationship of environmental parameters and bacterial community composition at the order level (Fig. 5). The 2005 samples were mainly correlated with the organic carbon and nitrogen load, i.e. DOC content of surface water and TOC and TN contents in the sediment. The 2011 samples showed the opposite pattern while being positively correlated with the O₂ concentration in surface water. The samples of 2010 were distinctly grouped in the centre and along the second canonical axis (i.e., the y axis in Fig. 5) with the lowest between-sample variation. In parallel, there was a clear trend on positioning of the bacterial orders in relation to environmental parameters. The bacterial orders *Xanthomonadales*, *Rhodobacterales*, *Burkholderiales*, *Bacillales*, *Rhizobiales*, *Sphingomonadales* and *Pseudomonadales* were positively correlated with the oxygen concentration in surface water while they were negatively correlated with DOC concentration in surface water and TOC and TN contents in the sediment. In contrast, *Desulfovibrionales*, *Anaerolineae*, *Lactobacillales*, *Desulfobacterales*, *Syntrophobacterales* and *Rhodocyclales* showed negative correlation to oxygen concentration in surface water while being positively correlated with DOC content of surface water and TOC content of the sediment (Fig. 5).

Discussion

The objective of this study was to assess the long-term impact of the implementation of a WWTP on the size, structure and function of microbial communities residing in the hyporheic zone of a stretch of the Belgian Zenne River impacted by a VC-contaminated groundwater plume. Previous studies on riverbed sediments impacted by WWTP effluent showed decreased microbial diversity^{26,27} and an altered nitrogen cycle²⁸ in the receiving rivers. However, the potential impact of the WWTP effluent on the downstream river can be influenced by a number of criteria, in particular the applied treatment technology that has been shifting towards advanced tertiary treatment in European countries in recent years.

Decades of municipal sewage input to the Zenne River at various locations in the vicinity of the test area caused high enrichment of TOC and TN in the pre-WWTP sediments. In turn, the establishment of the WWTP led to significant DOC decrease and O₂ increase in the Zenne surface water. This translated into effective reduction of TOC and TN in post-WWTP sediments in particular in the surficial sediment horizons (Fig. 1a, Supplementary Table S4–6). Concomitant with this transition from nutrient-rich to nutrient-poor conditions, we observed a remarkable reduction in abundance of phylogenetic and functional biomarkers. qPCR analysis showed the highest copy numbers of bacterial 16S




Figure 1. Sediment core characteristics. Total organic carbon and total nitrogen content (a) and 16S rRNA gene copy numbers of bacteria and *D. mccartyi* (as measured by qPCR) (b) in duplicate (filled (A) and open (B) symbols) sediment core taken in 2005 (top), 2010 (middle) and 2011 (bottom). Each qPCR value represents the average value obtained from triplicate reactions. Sediment-surface water interface was considered as the zero-depth. cmbsf: centimetres below surface.

rRNA genes and genetic biomarkers associated with sulphate reducers (*dsrB*), denitrifiers (*nirS*, *nirK*, *nosZ*) and methanogens (*mcrA*) in pre-WWTP sediments that in most cases significantly decreased in post-WWTP sediments (Supplementary Table S6). Depletion of labile organic carbon and nitrogen sources and establishment of more oxic conditions in the post-WWTP sediment could be the driver of this pattern as especially sulphate reducers and methanogens show highest activity under nutrient-rich and highly reducing conditions³⁰. The higher abundance of ammonium-oxidising bacteria and archaea (*amoA*) in pre-WWTP sediments could be due to ample presence of nitrogen resources (Fig. 1a).

Consistent with this trend, we observed reduced relative abundance of bacterial guilds that are known for their heterotrophic, syntrophic and fermentative lifestyle. On the other hand, the nutrient-poor conditions following the construction of the WWTP in our study likely favored members of the community that have higher substrate affinity and are more competitive under low nutrient conditions characterized by the presence of more recalcitrant complex organic substrates. For example, the pre-WWTP sediment and in particular the nutrient-rich shallow layers contained high relative abundance of *Chloroflexi* and in particular members of the order *Anaerolineales* that are known heterotrophs capable of metabolizing carbohydrates and peptides³¹. We noticed a strong correlation between *Anaerolineales* and DOC content in surface water and the sediment TOC and TN content (Fig. 5). The decreased relative abundance of *Chloroflexi* in the riverbed sediment downstream of a WWTP was reported previously where a lower organic carbon load was noted²⁶. Moreover, a recent community genomics study expanded the potential contribution of *Chloroflexi* in sediment carbon cycling to respiration of sugars, fermentation, and




Figure 2. Dechlorination in sediment microcosms. VC dechlorination (a) and accumulation of ethene (b) in microcosms prepared from top 20 cm riverbed sediment from 2005, 2010, 2011 and 2011 stimulated with tree bark (TB). Sediment bottles of 2005 were re-spiked with VC on day 28 after initial dechlorination of residual VC contamination in the original sample. The day of VC spike was considered as the day zero on the x-axis.

acetogenesis³². Besides *Anaerolineales*, the high loads of organic carbon coupled to high microbial activity (and hence subsequent oxygen depletion) in the pre-WWTP hyporheic sediments favored other bacterial community members that are active under anoxic and sub-oxic conditions. Accordingly, bacterial fermenters, such as members of the *Lactobacillales* and *Clostridiales*, sulphate-reducing *Desulfobacterales* and denitrifying members of *Rhodocyclales* showed strong positive correlations with organic carbon content of the surface water and sediment (Fig. 5). Besides, these microbial guilds had the highest relative abundances in pre-WWTP sediments (Supplementary Fig. S4) and significantly reduced in post-WWTP samples (Supplementary Table S10) that is consistent with the quantified biomarkers (for SRBs and denitrifiers, Supplementary Fig. S4).

The low-organic carbon concentrations after the installation of the WWTP also correlated with reduced abundance of *D. mccartyi*, which belongs to the *Chloroflexi* and is currently the only OHRB known to completely dechlorinate CAHs to ethene. Similarly, non-dechlorinating anaerobes (e.g. SRBs, methanogens, fermenters) that are commonly found to coexist with OHRB, and have been suggested to drive OHR by syntrophic interactions and exchange of essential metabolites^{32–34}, were reduced in relative abundance in post-WWTP samples^{33–35}. However, this did not impact VC dechlorination potential in the microcosms prepared from the sediments of 2010 as compared with the microcosms of 2005. Although diminished but consistent VC dechlorination was noted for the 2011 sediments, subsequent stimulation of VC dechlorination by tree bark addition showed that the OHRB and supporting non-dechlorinators were still present, albeit at lower initial abundance, and could be stimulated by re-establishment of organic-rich conditions.

In contrast, the reduced nutrient availability in post-WWTP sediments coincided with enrichment of certain proteobacterial orders. For instance, the proportion of *Burkholderiales* sequences showed a significant increase over time especially in the deeper sediment horizons (Supplementary Fig. S4) with a positive correlation to nutrient-poor and oxic conditions (Fig. 5). Members of the *Burkholderiales* have versatile catabolic traits enabling them to degrade recalcitrant and aromatic compounds and survive in environments with limited nutrient availability^{36,37}. Bacterial orders of *Xanthomonadales*, *Pseudomonadales*, and *Sphingomonadales* that are known to be active under oxic conditions and can survive nutrient-poor conditions were positively correlated with oxygen concentration in surface water, and showed a negative correlation to organic carbon content in surface water and sediment (Fig. 5). The wide metabolic flexibility and possession of high affinity transporters for substrates and nutrients makes *Pseudomonadales* capable of thriving in oligotrophic environments³⁸. Similarly *Xanthomonadales* members were proposed to survive in niches where nutrients are limited by decomposing recalcitrant carbon




Figure 3. Dominant bacterial phyla. Log-scale relative abundance of the dominant bacterial phyla at different depths in the duplicate sediment core samples taken in 2005 (left), 2010 (middle) 2011 (right). Sediment-surface water interface was considered as the zero-depth.

sources such as hemicellulose³⁹ and members of the *Sphingomonadales* were considered to be adapted to conditions of low availability of metabolizable substrates⁴⁰.

When we compared the bacteria alpha diversity (Shannon) and richness (Chao1, OTU count) between the pre- and post-WWTP sediment samples, one of the 2010 cores showed significantly higher values compared to the remaining cores, whereas one of the cores of 2011 had significantly lower richness and diversity. Conflicting results have been reported for the impact of WWTP on sediment bacterial diversity with some showing an increase in bacterial diversity downstream of a WWTP effluent input^{28,41} while others found decreased diversity^{26,27}. It should be noted that we cannot exclude the impact of other parameters such as season-dependent community dynamics in our study as the three sampling campaigns were performed in three different seasons (winter 2005, summer 2010, and fall 2011). Sediment microbial communities have been shown to be spatially heterogeneous and correlated with seasonal changes in physicochemical parameters such as temperature^{15,16,21,25}, DOC content^{15,16,25}, nitrate content^{15,23}, and phosphate concentration¹⁵. Moreover, the decreased concentrations of CAHs reaching the river could also impact the community members such as OHRB.

Our study showed that operation of a technologically advanced WWTP effectively promoted low-nutrient conditions in the longer term. This was translated into transition of the bacterial community in the riverbed hyporheic sediment from members favoured under nutrient-rich and reducing conditions to those adapted to nutrient-poor and oxic conditions, indicating that microbial communities are good sentinels and integrators of environmental change. The low-organic condition did not have a detrimental impact on VC respiration as one of the important hyporheic functions at CAH-contaminated sites fuelled by organic matter. Overall, the results of this study showed a positive impact of the technologically advanced WWTP on the sediment bacterial community, in line with sustainable water resource management aims.

Methods

Site description. The study site is a short transect of the Zenne River located in an industrial area near Brussels, Belgium. The test site was previously described in detail¹², and a simplified schematic presentation is given in Fig. 6 Briefly, at the study site, the Zenne River is relatively straight, 12–15 m wide, 0.5–2 m deep with a stream flow of 5–10 m³/s in dry weather conditions¹³. Based on data from October 2004 until January 2007, the nearby aquifer was shown to be contaminated with tetrachloroethene (PCE), trichloroethene (TCE) and 1,1,1-trichloroethane, originating from several sources¹². Moreover, it was noted that *cis*-dichloroethene and in particular VC as degradation products of reductive dechlorination of PCE and TCE, were reaching to the Zenne riverbed sediment^{12,29}. Since 2009 the concentration of CAHs that reach the river has decreased while the concentration of contaminants remained stable in the source area. It is unclear, however, whether this is due to increased OHR activity during plume flow from the source towards the river, or rather due to plume diversion.




Figure 4. Diversity indices. Alpha diversity (Shannon) and richness (Chao1 and OTU count) indices at different depths in the sediment core samples.

Sample collection. Sediment cores were collected from the top 80-cm layer of the riverbed sediment in December 2005, May 2010 and September 2011, at post 25 (P25, Fig. 6) using a 4-cm-diameter piston sediment sampler. At all three time-points, triplicate sediment cores were collected at a distance of approximately 50 cm from each other, randomly designated as cores A, B, or C. Two of the sediment cores (A and B) were preserved immediately on dry ice to conserve the spatial structure of the sediment layers. These cores were kept at -20°C until DNA extraction in November 2011 as described below. Core C was transferred directly to the lab in a cooler and used to set up microcosms to study OHR potential. Groundwater, surface water, sediment pore water sampling, as well as preservation were performed as reported previously¹³. Briefly, groundwater samples were collected from the monitoring well SB-1 (Fig. 6), and surface water samples were collected from the river as grab samples in glass bottles, without leaving a headspace. Electrical conductivity, dissolved oxygen and pH were measured regularly (Table 1) in the surface water using an electrometric multimeter (MultiLine F/SET3, WTW, Weilheim, Germany) and appropriate electrodes (TetraCon 325, Cellox 325 and Sen Tix 41 from WTW, Weilheim, Germany, respectively). Sediment interstitial water (ISW) was collected from Teflon pore water samplers that were permanently installed in the test area near the right riverbank at two different depths in the river sediment (20 and 60 cm) as reported previously¹³.




Figure 5. Redundancy analysis (RDA). Triplot of the relationship between microbial composition at order level and environmental parameters. Orders are indicated with filled triangles, environmental parameters with black arrows. The samples of 2005, 2010, and 2011 are shown as circle, star and diamonds, respectively, with core-a samples as filled and core-b samples as empty symbols. The eigenvalues of the first (x) and second (y) canonical axes are 0.262 and 0.194, respectively. The arrow length and direction corresponds to the variance that can be explained by the environmental variable. The direction of an arrow indicates an increasing magnitude of the environmental variable. The perpendicular distance between orders and environmental variable axes in the plot reflects their correlations. The smaller the distance, the stronger the correlation. SW: surface water; ISW: interstitial water.




Figure 6. Schematic representation of the test site. The locations of interstitial water samplers (ISW) and sediment core sampling at post 25 and of groundwater monitoring wells (large black dots) monitored in this study. Flow directions of the CAH plume and Zenne River are indicated by arrows.

Microcosm set up. Riverbed sediment microcosms were prepared directly after each sampling campaign with samples taken from the top (0–20 cm depth) sediment layer of the core C samples. The microcosms were prepared in an anoxic glove box (Don Whitley Scientific Ltd, West Yorkshire, UK) containing an atmosphere of N₂. Approximately 37 g (wet) sediment was transferred to 160-ml glass serum bottles containing 70 ml groundwater from the monitoring well SB-1. All microcosms were prepared in duplicate. Two extra microcosms prepared from the core C sample of 2011 were supplemented with 1 g tree bark as a source of solid organic carbon and electron donor as described elsewhere⁴². All microcosms were spiked with VC giving a final aqueous concentration of 2 mg/L. The microcosms containing sediment from 2005 were first allowed to degrade residual VC that was present in the groundwater, before VC was added.

DNA extraction, PCR-DGGE and qPCR. All A and B sediment cores were divided into 1-cm slices until a depth of 60–70 cm. At 5 cm intervals, a total of 80 sediment samples of approximately 2 g were used for DNA extraction as described previously⁴³. DGGE fingerprinting of the bacterial 16S rRNA gene pool in these 80 samples was performed as described previously⁴⁴, and 40 samples were chosen for further molecular analysis. The 16S rRNA gene of bacteria and *D. mccartyi* and selected protein-encoding genes commonly used as biomarkers for specific functional guilds, i.e. sulphate reducers (*dsrB*), denitrifiers (*nirK*, *nirS*, *nosZ*), methanogens (*mcrA*), and bacterial/archaeal ammonium oxidizers (bacterial/archaeal *amoA*) were quantified for the 40 selected samples. The qPCR measurements were performed in triplicate in 25- μ L reactions in an iQ5 iCycler (Bio-Rad, Veenendaal, the Netherlands) using the iQ SYBR Green Supermix kit (Bio-Rad). See Supplementary Table S1 for the lists of the primers and thermal cycling conditions used for qPCRs in this study. Standard curves were obtained using serial dilutions of a known amount of plasmid DNA containing a fragment of the respective genes. Genomic DNA from *Methanosarcina mazei* (ATCC BAA-159; GenBank accession number NC_003901) was used as standard for *mcrA* quantification.

Bacterial 16S rRNA gene amplicon pyrosequencing. Bacterial 16S rRNA genes were first amplified from the DNA extracts using the general 27F/1492R primer set⁴⁵ (Supplementary Table S2). In a second PCR barcoded amplicons covering the V1-V2 region of the bacterial 16S rRNA gene were generated using the 27F-DegS primer as the forward primer⁴⁶ and an equimolar mix of the two reverse primers, 338R I and II⁴⁷. The PCR mix (100 μ L final volume) contained 20 μ L of 5 \times HF buffer (Finnzymes, Vantaa, Finland), 2 μ L PCR Grade Nucleotide Mix (Roche Diagnostic GmbH, Mannheim, Germany), 1 μ L of Phusion Hot start II High-Fidelity DNA polymerase (2 U/ μ L) (Finnzymes), 500 nM of the reverse primer mix and the forward primer (Biolegio BV, Nijmegen, The Netherlands), 2 μ L (i.e. 40 ng) template, and 65 μ L nuclease free water. PCR was performed using the following conditions: 98 °C for 30 s, followed by 30 cycles consisting of denaturation at 98 °C for 10 s, annealing at 56 °C for 20 s, elongation at 72 °C for 20 s, and a final extension at 72 °C for 10 min. The PCR product was quality-checked on a 1% agarose gel, purified from gel using the High Pure PCR Cleanup Micro Kit (Roche Diagnostics) and its concentration was determined by a Qubit 2.0 Fluorometer (Life Technologies, Darmstadt, Germany). Amplicons were sequenced using an FLX genome sequencer in combination with titanium chemistry (GATC-Biotech, Konstanz, Germany).

Analysis of the pyrosequencing data. Pyrosequencing data were processed and analysed using QIIME v1.8.0⁴⁸. Raw sequencing data was first pre-processed according to default options in QIIME to remove low quality and ambiguous reads. Pre-processed data was denoised⁴⁹ to minimize sequencing errors in the dataset and chimeric sequences were removed using UCHIME⁵⁰. Operational taxonomic units (OTUs) were defined at a 97% identity level using uclust⁵¹ and the SILVA 111 database⁵² as a reference. The full pre-processed data was transformed into relative abundances before comparing taxon abundances between samples. Note that the relative abundance comparisons between the different taxa are not independent, and may be therefore affected by the compositionality effect⁵³. Bacterial diversity and richness indices were determined with R (3.1.2)⁵⁴ using the phyloseq package (1.9.15)⁵⁵ from the complete dataset without rarefaction⁵⁶.

Analytical methods. Concentrations of CAHs, ethene, and ethane were determined via head-space analysis on a Varian GC-FID (CP-3800) as described previously⁵⁷. Total organic carbon (TOC) and total nitrogen (TN) were measured using the oxidative digestion method (C/N analyzer EA1110). Concentrations of sulphate, nitrate, phosphate, and chloride were analyzed by ion chromatography using a Dionex DX-120 ion chromatograph equipped with a Dionex AS14A column (Dionex, Sunnyvale, CA). Dissolved organic carbon (DOC) was determined from samples as the difference between total dissolved carbon and dissolved inorganic carbon, measured with a Shimadzu TOC-5000 analyzer equipped with an ASI-5000 auto-sampler.

Statistical analyses. Univariate statistical analysis (Spearman correlation and Kruskal-Wallis test) were performed within the Matlab environment (Version 2015a, The Mathworks, Natick, MA), unless

otherwise stated. For all statistical analysis, Benjamini-Hochberg⁵⁸ correction for multiple testing was applied for false discovery rate (FDR)-controlling; a q-value < 0.25 was deemed significant^{59,60}. The phylogenetic profiling data was analysed with R (3.1.2)⁵⁴ using the phyloseq package (1.9.15)⁵⁵. We included OTUs that were present in at least three samples and contained one or more reads. This resulted in a final set of 1303 detected OTUs (out of 21067 total OTUs). In order to relate the changes in microbial communities (at order level) with environmental variables, redundancy analysis (RDA) was used as implemented in the CANOCO 5 software (Biometris, Wageningen, The Netherlands). The environmental variables tested were DOC and oxygen in surface water, TOC and TN in the sediment cores, and DOC and VC concentration in ISW. Nitrite and nitrate in ISW were not considered as their levels were always below detection limit. The community structure was visualized via ordination triplots with scaling focused on intersample differences.

References

- Carey, R. O. & Migliaccio, K. W. Contribution of wastewater treatment plant effluents to nutrient dynamics in aquatic systems. *Environ Manage* **44**, 205–217 (2009).
- Grimm, N. B. *et al.* Global change and the ecology of cities. *Science* **319**, 756–760 (2008).
- United Nations. Department of Economic and Social Affairs, Population Division 2014. *World Urbanization Prospects: The 2014 Revision, Highlights (ST/ESA/SER.A/352)*. (2014).
- Gücker, B., Brauns, M. & Pusch, M. T. Effects of wastewater treatment plant discharge on ecosystem structure and function of lowland streams. *J North Am Benthol Soc* **25**, 313–329 (2006).
- European Commission. Establishing a framework for community action in the field of water policy. Directive 2000/60/EC of the European Parliament and of the Council. *Official Journal of the European Communities L* **327**, 1–72 (2000).
- European Environment Agency Assessment on Urban Waste Water Treatment Indicators, ref: CSI 024. Available at http://themes.eea.europa.eu/IMS/ISpecs/ISpecification20041007132045/IAssessment1196343193294/view_content (June 2009). (2009).
- Tank, J. L., Rosi-Marshall, E. J., Griffiths, N. A., Entekin, S. A. & Stephen, M. L. A review of allochthonous organic matter dynamics and metabolism in streams. *J North Am Benthol Soc* **29**, 118–146 (2010).
- Boulton, A. J., Humphreys, W. F. & Eberhard, S. M. Imperilled subsurface waters in Australia: Biodiversity, threatening processes and conservation. *Aquatic Ecosyst Health Manage* **6**, 41–54 (2003).
- Biksey, T. M. & Gross, E. D. The hyporheic zone: Linking groundwater and surface water—understanding the paradigm. *Remediation* **12**, 55–62 (2001).
- Abe, Y., Aravena, R., Zopf, J., Parker, B. & Hunkeler, D. Evaluating the fate of chlorinated ethenes in streambed sediments by combining stable isotope, geochemical and microbial methods. *J Contam Hydrol* **107**, 10–21 (2009).
- Ellis, P. A. & Rivett, M. O. Assessing the impact of VOC-contaminated groundwater on surface water at the city scale. *J Contam Hydrol* **91**, 107–127 (2007).
- Hamonts, K. *et al.* Factors determining the attenuation of chlorinated aliphatic hydrocarbons in eutrophic river sediment impacted by discharging polluted groundwater. *Environ Sci Technol* **43**, 5270–5275 (2009).
- Hamonts, K. *et al.* Temporal variations in natural attenuation of chlorinated aliphatic hydrocarbons in eutrophic river sediments impacted by a contaminated groundwater plume. *Water Res* **46**, 1873–1888 (2012).
- Bai, Y. *et al.* Bacterial communities in the sediments of Dianchi lake, a partitioned eutrophic waterbody in China. *PLoS ONE* **7**, e37796 (2012).
- Febria, C. M., Beddoes, P., Fulthorpe, R. R. & Williams, D. D. Bacterial community dynamics in the hyporheic zone of an intermittent stream. *ISME J* **6**, 1078–1088 (2012).
- Febria, C. M., Fulthorpe, R. R. & Williams, D. D. Characterizing seasonal changes in physicochemistry and bacterial community composition in hyporheic sediments. *Hydrobiologia* **647**, 113–126 (2010).
- Feris, K. P. *et al.* Seasonal dynamics of shallow-hyporheic-zone microbial community structure along a heavy-metal contamination gradient. *Appl Environ Microbiol* **70**, 2323–2331 (2004).
- Feris, K. P., Ramsey, P. W., Frazar, C., Rillig, M. C., Gannon, J. E. & Holben, W. E. Structure and seasonal dynamics of hyporheic zone microbial communities in free-stone rivers of the western United States. *Appl Environ Microbiol* **46**, 200–215 (2003).
- Findlay, S., Strayer, D., Goumbala, C. & Gould, K. Metabolism of streamwater dissolved organic carbon in the shallow hyporheic zone. *Limnol Oceanogr* **38**, 1493–1499 (1993).
- Findlay, S. E. G., Sinsabaugh, R. L., Sobczak, W. V. & Hoostal, M. Metabolic and structural response of hyporheic microbial communities to variations in supply of dissolved organic matter. *Limnol Oceanogr* **48**, 1608–1617 (2003).
- Hullar, M. A. J., Kaplan, L. A. & Stahl, D. A. Recurring seasonal dynamics of microbial communities in stream habitats. *Appl Environ Microbiol* **72**, 713–722 (2006).
- Iribar, A., Sánchez-Pérez, J. M., Lyautey, E. & Garabétian, F. Differentiated free-living and sediment-attached bacterial community structure inside and outside denitrification hotspots in the river-groundwater interface. *Hydrobiologia* **598**, 109–121 (2008).
- Lowell, J. L., Gordon, N., Engstrom, D., Stanford, J. A., Holben, W. E. & Gannon, J. E. Habitat heterogeneity and associated microbial community structure in a small-scale floodplain hyporheic flow path. *Microb Ecol* **58**, 611–620 (2009).
- Pérez-Jiménez, J. R. & Kerkhof, L. J. Phylogeography of sulfate-reducing bacteria among disturbed sediments, disclosed by analysis of the dissimilatory sulfite reductase genes (*dsrAB*). *Appl Environ Microbiol* **71**, 1004–1011 (2005).
- Sliva, L. & Williams, D. D. Exploration of riffle-scale interactions between abiotic variables and microbial assemblages in the hyporheic zone. *Can J Fish Aquat Sci* **62**, 276–290 (2005).
- Drury, B., Rosi-Marshall, E. & Kelly, J. J. Wastewater treatment effluent reduces the abundance and diversity of benthic bacterial communities in urban and suburban rivers. *Appl Environ Microbiol* **79**, 1897–1905 (2013).
- Lu, X. M. & Lu, P. Z. Characterization of bacterial communities in sediments receiving various wastewater effluents with high-throughput sequencing analysis. *Microb Ecol* **67**, 1–12 (2014).
- Wakelin, S. A., Colloff, M. J. & Kookana, R. S. Effect of wastewater treatment plant effluent on microbial function and community structure in the sediment of a freshwater stream with variable seasonal flow. *Appl Environ Microbiol* **74**, 2659–2668 (2008).
- Kuhn, T. K. *et al.* Assessment of the intrinsic bioremediation capacity of an eutrophic river sediment polluted by discharging chlorinated aliphatic hydrocarbons: A compound-specific isotope approach. *Environ Sci Technol* **43**, 5263–5269 (2009).
- Muyzer, G. & Stams, A. J. M. The ecology and biotechnology of sulphate-reducing bacteria. *Nat Rev Microbiol* **6**, 441–454 (2008).
- Yamada, T. & Sekiguchi, Y. Cultivation of uncultured *Chloroflexi* subphyla: Significance and ecophysiology of formerly uncultured *Chloroflexi* 'subphylum i' with natural and biotechnological relevance. *Microb Environ* **24**, 205–216 (2009).
- Hug, L. A. *et al.* Community genomic analyses constrain the distribution of metabolic traits across the *Chloroflexi* phylum and indicate roles in sediment carbon cycling. *Microbiome* **1**, 22 (2013).

33. Brisson, V. L., West, K. A., Lee, P. K. H., Tringe, S. G., Brodie, E. L. & Alvarez-Cohen, L. Metagenomic analysis of a stable trichloroethene-degrading microbial community. *ISME J* **6**, 1702–1714 (2012).
34. Duhamel, M., Mo, K. & Edwards, E. A. Characterization of a highly enriched *Dehalococcoides*-containing culture that grows on vinyl chloride and trichloroethene. *Appl Environ Microbiol* **70**, 5538–5545 (2004).
35. Hug, L. A., Beiko, R. G., Rowe, A. R., Richardson, R. E. & Edwards, E. A. Comparative metagenomics of three *Dehalococcoides*-containing enrichment cultures: the role of the non-dechlorinating community. *BMC Genomics* **13**, 327 (2012).
36. Li, R., Khafipour, E., Krause, D. O., Entz, M. H. & De Kievit, T. R. Fernando WGD. Pyrosequencing reveals the influence of organic and conventional farming systems on bacterial communities. *PLoS ONE* **7**, e51897 (2012).
37. Suárez-Moreno, Z. R., Caballero-Mellado, J., Coutinho, B. G., Mendonça-Previato, L., James, E. K. & Venturi, V. Common features of environmental and potentially beneficial plant-associated *Burkholderia*. *Microb Ecol* **63**, 249–266 (2012).
38. Martins Dos Santos, V. A. P., Heim, S., Moore, E. R. B., Strätz, M. & Timmis, K. N. Insights into the genomic basis of niche specificity of *Pseudomonas putida* KT2440. *Environ Microbiol* **6**, 1264–1286 (2004).
39. Déjean, G. *et al.* The xylan utilization system of the plant pathogen *Xanthomonas campestris* pv *campestris* controls epiphytic life and reveals common features with oligotrophic bacteria and animal gut symbionts. *New Phytologist* **198**, 899–915 (2013).
40. Pang, C. M. & Liu, W. T. Biological filtration limits carbon availability and affects downstream biofilm formation and community structure. *Appl Environ Microbiol* **72**, 5702–5712 (2006).
41. Marti, E. & Balcázar, J. L. Use of pyrosequencing to explore the benthic bacterial community structure in a river impacted by wastewater treatment plant discharges. *Res Microbiol* **165**, 468–471 (2014).
42. Atashgahi, S. *et al.* Evaluation of solid polymeric organic materials for use in bioreactive sediment capping to stimulate the degradation of chlorinated aliphatic hydrocarbons. *Appl Microbiol Biotechnol* **98**, 2255–2266 (2014).
43. Hendrickx, B. *et al.* Dynamics of an oligotrophic bacterial aquifer community during contact with a groundwater plume contaminated with benzene, toluene, ethylbenzene, and xylenes: An *in situ* mesocosm study. *Appl Environ Microbiol* **71**, 3815–3825 (2005).
44. Uyttebroek, M. *et al.* Distribution of the *Mycobacterium* community and polycyclic aromatic hydrocarbons (PAHs) among different size fractions of a long-term PAH-contaminated soil. *Environ Microbiol* **8**, 836–847 (2006).
45. Lane, D. 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial Systematics*. Stackebrandt, E., and Goodfellow, M. (eds). New York, USA, Wiley, pp. 115–148 (1991).
46. Van den Bogert, B., De Vos, W. M., Zoetendal, E. G. & Kleerebezem, M. Microarray analysis and barcoded pyrosequencing provide consistent microbial profiles depending on the source of human intestinal samples. *Appl Environ Microbiol* **77**, 2071–2080 (2011).
47. Daims, H., Brühl, A., Amann, R., Schleifer, K. H. & Wagner, M. The domain-specific probe EUB338 is insufficient for the detection of all bacteria: Development and evaluation of a more comprehensive probe set. *System Appl Microbiol* **22**, 434–444 (1999).
48. Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* **7**, 335–336 (2010).
49. Bragg, L., Stone, G., Imelfort, M., Hugenholtz, P. & Tyson, G. W. Fast, accurate error-correction of amplicon pyrosequences using Acacia. *Nat Methods* **9**, 425–426 (2012).
50. Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**, 2194–2200 (2011).
51. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461 (2010).
52. Pruesse, E. *et al.* SILVA: A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* **35**, 7188–7196 (2007).
53. Aitchison, J. The Statistical Analysis of Compositional Data. *J R Stat Soc Series B (Methodol)* **44**, 139–177 (1982).
54. R., Core Team. R: A language and environment for statistical computing. (ed[^](eds). *R Foundation for Statistical Computing* (2013).
55. McMurdie, P. J. & Holmes, S. Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* **8**, e61217 (2013).
56. McMurdie, P. J. & Holmes, S. Waste not, want not: why rarefying microbiome data is inadmissible. *PLoS Comput Biol* **10**, e1003531 (2014).
57. Atashgahi, S., Maphosa, F., Doğan, E., Smidt, H., Springael, D. & Dejonghe, W. Small-scale oxygen distribution determines the vinyl chloride biodegradation pathway in surficial sediments of riverbed hyporheic zones. *FEMS Microbiol Ecol* **84**, 133–142 (2013).
58. Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing under dependency. *Annal Stat* **29**, 1165–1188 (2001).
59. Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. *Gen Biol* **13**, R79 (2012).
60. Schnabel, R. B. *et al.* The relation of genetic and environmental factors to systemic inflammatory biomarker concentrations. *Circ Cardiovasc Gen* **2**, 229–237 (2009).

Acknowledgements

This study was supported by EU-projects SQUAREHAB (grant 226565) and SEDBARCAH (grant 511254), the EcoLinc project of the Netherlands Genomics Initiative and BE-BASIC (grant F08.004.01). M.R.D. was supported by the Strategic Research Fund of the WIMEK graduate school. L.L. was supported by the Academy of Finland (grant 256950). We thank Ricardo J. Eloy Alves and Christa Schleper for providing bacterial and archaeal *amoA* clones for qPCR.

Author Contributions

W.D., D.S. and H.S. planned the research; S.A., K.H., R.A., F.M. and T.K. did the experiments; S.A., D.S., M.R.D., L.L. and E.S. performed data analysis; and all authors wrote and reviewed the manuscript.

Additional Information

Accession codes: The complete dataset is available at the European Bioinformatics Institute (www.ebi.ac.uk) under accession number PRJEB8703.

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Atashgahi, S. *et al.* Impact of a wastewater treatment plant on microbial community composition and function in a hyporheic zone of a eutrophic river. *Sci. Rep.* **5**, 17284; doi: 10.1038/srep17284 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>