



HHS Public Access

Author manuscript

Birth Defects Res A Clin Mol Teratol. Author manuscript; available in PMC 2017 January 01.

Published in final edited form as:

Birth Defects Res A Clin Mol Teratol. 2016 January ; 106(1): 55–60. doi:10.1002/bdra.23389.

Maternal Occupational Exposure to Polycyclic Aromatic Hydrocarbons and Craniosynostosis among Offspring in the National Birth Defects Prevention Study

Jacqueline L. O'Brien¹, Peter H. Langlois², Christina C. Lawson³, Angela Scheuerle⁴, Carissa M. Rocheleau³, Martha A. Waters³, Elaine Symanski⁵, Paul A. Romitti⁶, A.J. Agopian⁵, and Philip J. Lupo⁷ **The National Birth Defects Prevention Study**

¹Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, Texas, USA

²Birth Defects Epidemiology and Surveillance Branch, Texas Department of State Health Services, Austin, Texas, USA

³National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Cincinnati, Ohio, USA

⁴Department of Pediatrics, Division of Genetics and Metabolism, University of Texas Southwestern Medical Center, Dallas, TX, USA

⁵Division of Epidemiology, Human Genetics and Environmental Sciences, University of Texas School of Public Health, Houston, Texas, USA

⁶Department of Epidemiology, College of Public Health, University of Iowa, Iowa City, USA

⁷Department of Pediatrics, Section of Hematology-Oncology, Baylor College of Medicine, Houston, Texas, USA

Abstract

Background—Evidence in animal models and humans suggests that exposure to polycyclic aromatic hydrocarbons (PAHs) may lead to birth defects. To our knowledge, this relationship has not been evaluated for craniosynostosis, a birth defect characterized by the premature closure of sutures in the skull. We conducted a case-control study to examine associations between maternal occupational exposure to PAHs and craniosynostosis.

Methods—We used data from craniosynostosis cases and control infants in the National Birth Defects Prevention Study (NBDPS) with estimated delivery dates from 1997–2002. Industrial hygienists reviewed occupational data from the computer-assisted telephone interview and assigned a yes/no rating of probable occupational PAH exposure for each job from one month before conception through delivery. We used logistic regression to assess the association between occupational exposure to PAHs and craniosynostosis.

Corresponding author: Philip J. Lupo, Address: One Baylor Plaza, Houston TX, 77030, BCM305, Phone: 713- 798-2960, Philip.Lupo@bcm.edu.

Conflicts of interest: none declared

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention or the National Institute for Occupational Safety and Health.

Results—The prevalence of exposure was 5.3% in case mothers (16/300) and 3.7% in control mothers (107/2,886). We observed a positive association between exposure to PAHs during the one month before conception through the third month of pregnancy and craniosynostosis [odds ratio (OR) = 1.75; 95% confidence interval (CI): 1.01, 3.05] after adjusting for maternal age and maternal education. The number of cases for each craniosynostosis subtype limited subtype analyses to sagittal craniosynostosis; the odds ratio remained similar (OR = 1.76, 95% CI: 0.82–3.75), but was not significant.

Conclusions—Our findings support a moderate association between maternal occupational exposure to PAHs and craniosynostosis. Additional work is needed to better characterize susceptibility and the role PAHs may play on specific craniosynostosis subtypes.

Keywords

craniosynostosis; polycyclic aromatic hydrocarbon (PAH); birth defects; case-control

INTRODUCTION

Polycyclic aromatic hydrocarbons (PAHs) are a group of toxic chemicals that are ubiquitous in certain occupational settings, including the oil and gas industry, coal-fired and other power plants, and restaurants (Sjaastad & Svendsen, 2009). These toxicants have been linked to adverse birth outcomes such as intrauterine growth retardation (Choi, Rauh, Garfinkel, Tu, & Perera, 2008), preterm birth (Singh et al., 2008), decreased infant growth (Choi et al., 2006; Langlois et al., 2014), as well as certain birth defects (e.g., gastroschisis (Lupo, Langlois, et al., 2012), oral clefts (Langlois et al., 2013), and neural tube defects (Ren et al., 2011)). However, no studies have examined the relationship between PAHs and craniosynostosis, a condition characterized by the premature closure of one or more of the sutures in the infant skull (Nagaraja, Anslow, & Winter, 2013). This relationship is of particular interest as previous assessments suggest craniosynostosis is associated with maternal smoking (Hackshaw, Rodeck, & Boniface, 2011), a major source of environmental PAHs.

Given the growing evidence of the teratogenic potential of PAHs, and the need to better understand modifiable risk factors for craniosynostosis—especially with well-designed, large-scale population-based studies—we investigated associations between maternal occupational exposure to PAHs and risk of craniosynostosis among offspring using data from the National Birth Defects Prevention Study (NBDPS).

MATERIALS AND METHODS

The study population included craniosynostosis case and control infants from the NBDPS, with estimated delivery dates from October 1, 1997 through December 31, 2002. Details of the NBDPS have been published previously (Yoon et al., 2001). Briefly, the NBDPS is a multi-center, population-based, case-control study of major birth defects. During the period 1997–2002, case infants with one or more major birth defects were identified through eight birth defect surveillance systems in the United States (Arkansas, California, Georgia, Iowa, Massachusetts, New Jersey, New York, and Texas). The cases identified included live

births, stillbirths, and induced pregnancy terminations. Clinical geneticists used standardized case definitions and confirmatory diagnostic procedures (Rasmussen et al., 2003) to classify case infants. Infants and fetuses with single gene disorders or chromosomal abnormalities were excluded, and each eligible case was classified as having an isolated defect (no additional major birth defect) or multiple defects (one or more additional unrelated major birth defects). The sutures involved were also identified (i.e., sagittal, coronal, metopic, lambdoid, multiple sutures, or unknown). Control infants (live born infants without major birth defects) were randomly selected from birth certificates or birth hospital records from the same geographic populations that gave rise to the cases.

Case and control mothers completed a 1-hour computer-assisted telephone interview (CATI) between six weeks and two years after the estimated date of delivery. The interview included sections on maternal health, lifestyle and behavioral characteristics, and occupational history. For each job, mothers were asked to describe: job title, typical job tasks, employer, what their employer made or did, any specific equipment or chemicals used, typical hours per week worked, and starting and stopping dates. Jobs were classified according to the Standard Occupational Classification (SOC) System (United States Department of Labor Bureau of Labor Statistics, 2000) and the North American Industry Classification System (United States Department of Labor Bureau of Labor Statistics, 1997). Two trained industrial hygienists, working independently and blinded to case-control status, reviewed job descriptions from the CATI and assigned a dichotomous (yes/no) rating of probable occupational PAH exposure for each job. Any discrepancies were resolved in a consensus meeting, which included a third hygienist. The approach used to estimate PAH exposure has been described previously (Rocheleau et al., 2011) and is described in detail in other NBDPS studies (Langlois et al., 2014; Langlois et al., 2013; Langlois et al., 2012; Lupo, Langlois, et al., 2012; Lupo, Symanski, et al., 2012).

Prior to the interview, all mothers provided verbal consent for study participation. The Centers for Disease Control and Prevention Institutional Review Board (IRB), along with the IRBs for each participating center, approved the NBDPS.

Our analysis was limited to case infants with a diagnosis of craniosynostosis and all control infants with exposure information. Specifically, case and control mothers were eligible if they worked in part-time or full-time jobs (paid or volunteer) for at least 1 month from the 3 months before conception through the end month of pregnancy. Of the 4,584 eligible mothers with estimated delivery dates between 1997–2002, 1,275 (28%) were excluded from the analysis because they did not work during this time period. For our primary analysis, a mother was classified as exposed if she had one or more jobs rated as exposed during the critical period defined as one month before conception through the third month of pregnancy, and was classified as unexposed if all of her jobs during this time period were rated as unexposed. Although the critical period for exposure is likely in early pregnancy, we conducted a secondary analysis where the exposure window was defined as one month before conception through delivery, because suture closure is not completed until after birth.

Information about maternal demographic and infant characteristics was obtained from the CATI, and included maternal age at delivery (<20, 20–24, 25–29, 30–34, 35 years); pre-

pregnancy body mass index (BMI, kg/m²) (<18.5, 18.5–24.9, 25–29.9, 30); parity (0, 1); maternal race or ethnicity (non-Hispanic white, Non-Hispanic black, Hispanic, Other); maternal education (<12, 12, >12 years); gestational diabetes (yes/no); alcohol use one month before pregnancy through the third month of pregnancy (yes/no); smoking history one month before pregnancy through the third month of pregnancy (nonsmoker, light moderate, heavy); secondhand smoke at home (yes/no); secondhand smoke at work (yes/no), and infant sex (male/female). We calculated frequency distributions of these maternal demographics for cases and controls, and the crude odds ratios and 95% CIs for the associations between each maternal factor and craniosynostosis.

To examine the association between exposure to PAHs in the workplace and craniosynostosis, we used unconditional logistic regression models to calculate crude and multivariable-adjusted odds ratios (ORs) and 95% confidence intervals (CIs). The variables considered for our final model were determined *a priori* based on a review of the craniosynostosis literature, and all variables were established or suspected risk factors for craniosynostosis (maternal age, maternal education, race/ethnicity, smoking history, exposure to secondhand smoke at home, and body mass index) (Boulet, Rasmussen, & Honein, 2008; Carmichael et al., 2008; Lee et al., 2012). Variables were incorporated as confounders in the final model if inclusion resulted in 10% change in the estimate of the effect between PAHs and craniosynostosis. Due to the numbers of cases for each craniosynostosis subtype, we were only able to estimate a separate multiple logistic regression model for sagittal craniosynostosis.

RESULTS

The study population included 316 craniosynostosis cases and 2,993 control infants, with delivery dates from October 1, 1997 through December 31, 2002. Maternal occupational exposure to PAHs was 5.3% among case mothers and 3.7% among control mothers during the month before conception through the third month of pregnancy. Mothers of case infants were older (mean age case mothers = 30.0 years; mean age control mothers = 28.2 years) and more likely to be exposed to secondhand smoke at home (Table 1). Case infants were more likely to be males.

In the unadjusted analysis (Table 2), maternal occupational exposure to PAHs during the one month before conception through the third month of pregnancy was positively, but not significantly, associated with craniosynostosis (OR = 1.44, 95% CI: 0.84, 2.47). However, in the final model, adjusted for maternal age and education, mothers of cases were 75% (95% CI: 1.01, 3.05) more likely to be exposed to occupational PAHs when compared to control mothers. In the fully adjusted model where exposure was defined as one month before conception through delivery, the effect was attenuated (OR = 1.55, 95% CI: 0.91, 2.70).

Our analyses of specific craniosynostosis subtypes were limited by sample size [sagittal: 163 cases (52%); metopic: 50 cases (16%); coronal: 57 cases (18%); lambdoid: 15 cases (5%); multiple types: 28 cases (9%); unspecified: 3 cases (1%)]. Therefore, we only analyzed the largest subtype (i.e., sagittal). Specifically, the OR from the multivariable-adjusted model

for sagittal craniosynostosis suggested increased risk, but it was not statistically significant (OR = 1.76, 95% CI: 0.82, 3.75).

DISCUSSION

Using data from the largest population-based case-control study of birth defects in the United States, we observed a modest association between maternal occupational exposure to PAHs during the early months of pregnancy and risk of craniosynostosis among offspring. Our findings are in keeping with a growing body of literature suggesting the teratogenic potential of PAHs (Ren et al., 2011).

While there have been no previous assessments of maternal occupational exposure to PAHs and craniosynostosis, these toxicants have been linked to birth defects in other epidemiologic research. For example, studies have reported associations between PAHs and gastroschisis (Lupo, Langlois, et al., 2012), small for gestational age (Langlois et al., 2014), and neural tube defects (Ren et al., 2011). There is also biological evidence that PAHs may be teratogenic. PAHs are lipophilic and readily absorbed by human tissue, including the placenta (Agency for Toxic Substances and Disease Registry, 1995). Once inhaled or ingested, they are metabolized into compounds capable of binding to DNA. These PAH-DNA adducts can trigger cellular mutations that are teratogenic (Wells, McCallum, Lam, Henderson, & Ondovcik, 2010). Furthermore, PAH-DNA adducts have been detected in fetal organs, umbilical cord blood, and amniotic fluid (Agency for Toxic Substances and Disease Registry, 1995). Finally, birth defects and other adverse fetal outcomes have been reported in animal models after maternal exposure to PAHs (Barbieri, Ognio, Rossi, Astigliano, & Rossi, 1986; Incardona, Collier, & Scholz, 2004; Lambert & Nebert, 1977).

Our findings must be considered in light of certain limitations. There is the potential for exposure misclassification when using expert assessments of job descriptions, although this method is generally considered superior to self-reported exposures that depend on participant awareness of PAH exposures and accurate recall. Furthermore, assessing occupational exposure to PAHs, even if captured without any exposure misclassification, is not a complete assessment of exposure, given how ubiquitous PAHs are in the environment, and we lacked additional information on other potential environmental sources. That said, occupational exposure to PAHs often exceeds the level of exposure in the environment (Brandt & Watson, 2003). We did not have information on the intensity and frequency of exposure, which prevented an analysis of between- and within- job exposure variability, as well as an exposure-response analysis.

Because this is an observational study, uncontrolled or residual confounding could explain the relation between exposure to PAHs and craniosynostosis. However, detailed information on the health and lifestyle of the participants permitted evaluation of many potential confounders. We did not observe any change to our conclusions when smoking (or exposure to secondhand smoke) was considered in our models. The sample size limited this and other stratified and subtype analyses. For example, we were only able to evaluate associations among sagittal craniosynostosis. Larger sample sizes are needed to examine the effects of PAHs on other craniosynostosis subtypes.

Strengths of our study include the use of data from the NBDPS, the largest population-based case-control study of birth defects, which has extensive occupational PAH exposure information from 1997–2002, as well as detailed information on potential confounders. Furthermore, the thorough case classification process in the NBDPS by clinical geneticists excluded cases due to single gene disorders or chromosomal abnormalities, resulting in a more homogeneous craniosynostosis case group.

CONCLUSIONS

To our knowledge, this is the first study to assess maternal occupational exposure to PAHs and craniosynostosis in offspring. Our analysis suggests that PAH exposure during early pregnancy is associated with a moderately increased risk of craniosynostosis in offspring. Because PAHs are ubiquitous in our environment, it is important to consider their effects on human health, and increase awareness of potential harm so that those at greatest risk may employ measures to minimize exposure, especially during pregnancy. Future investigations could be improved by incorporating additional measures of exposure and the inclusion of additional cases for a closer examination of PAHs and craniosynostosis subtype.

Acknowledgments

We thank the families involved in the NBDPS for their participation in the study. The authors would like to acknowledge the birth defects surveillance centers in the following states: Arkansas, California, Georgia, Iowa, Massachusetts, New Jersey, New York, and Texas).

Grant information: This project was supported by NIH R03DE021739 (P.J. Lupo). This work was also supported in part through a cooperative agreement (U01DD000494) between the Centers for Disease Control and Prevention and the Texas Department of State Health Services (DSHS) and by the National Institute for Occupational Safety and Health (contract 200-2000-08018). Funding also came from Title V Maternal and Child Health Block Grants Funds from the Office of Title V and Family Health, Texas DSHS.

Literature cited

- Agency for Toxic Substances and Disease Registry. Toxicological Profile for Polycyclic Aromatic Hydrocarbons (PAHs) (Update): US Department of Health and Human Services. Public Health Service, Agency for Toxic Substances and Disease Registry; 1995.
- Barbieri O, Ognio E, Rossi O, Astigiano S, Rossi L. Embryotoxicity of benzo(a)pyrene and some of its synthetic derivatives in Swiss mice. *Cancer Res.* 1986; 46(1):94–98. [PubMed: 3753553]
- Boulet SL, Rasmussen SA, Honein MA. A population-based study of craniosynostosis in metropolitan Atlanta, 1989–2003. *Am J Med Genet A.* 2008; 146A(8):984–991.10.1002/ajmg.a.32208 [PubMed: 18344207]
- Brandt HC, Watson WP. Monitoring human occupational and environmental exposures to polycyclic aromatic compounds. *Ann Occup Hyg.* 2003; 47(5):349–378. [PubMed: 12855487]
- Carmichael SL, Ma C, Rasmussen SA, Honein MA, Lammer EJ, Shaw GM. Craniosynostosis and maternal smoking. *Birth Defects Res A Clin Mol Teratol.* 2008; 82(2):78–85.10.1002/bdra.20426 [PubMed: 18050313]
- Choi H, Jedrychowski W, Spengler J, Camann DE, Whyatt RM, Rauh V, Perera FP. International studies of prenatal exposure to polycyclic aromatic hydrocarbons and fetal growth. *Environ Health Perspect.* 2006; 114(11):1744–1750. [PubMed: 17107862]
- Choi H, Rauh V, Garfinkel R, Tu Y, Perera FP. Prenatal exposure to airborne polycyclic aromatic hydrocarbons and risk of intrauterine growth restriction. *Environ Health Perspect.* 2008; 116(5): 658–665.10.1289/ehp.10958 [PubMed: 18470316]

- Hackshaw A, Rodeck C, Boniface S. Maternal smoking in pregnancy and birth defects: a systematic review based on 173 687 malformed cases and 11.7 million controls. *Hum Reprod Update*. 2011; 17(5):589–604.10.1093/humupd/dmr022 [PubMed: 21747128]
- Incardona JP, Collier TK, Scholz NL. Defects in cardiac function precede morphological abnormalities in fish embryos exposed to polycyclic aromatic hydrocarbons. *Toxicol Appl Pharmacol*. 2004; 196(2):191–205.10.1016/j.taap.2003.11.026 [PubMed: 15081266]
- Lambert GH, Nebert DW. Genetically mediated induction of drug-metabolizing enzymes associated with congenital defects in the mouse. *Teratology*. 1977; 16(2):147–153.10.1002/tera.1420160206 [PubMed: 412268]
- Langlois PH, Hoyt AT, Desrosiers TA, Lupo PJ, Lawson CC, Waters MA, National Birth Defects Prevention Study. Maternal occupational exposure to polycyclic aromatic hydrocarbons and small for gestational age offspring. *Occup Environ Med*. 2014; 71(8):529–535.10.1136/oemed-2013-101833 [PubMed: 24893704]
- Langlois PH, Hoyt AT, Lupo PJ, Lawson CC, Waters MA, Desrosiers TA, Lammer EJ. Maternal occupational exposure to polycyclic aromatic hydrocarbons and risk of oral cleft-affected pregnancies. *Cleft Palate Craniofac J*. 2013; 50(3):337–346.10.1597/12-104 [PubMed: 23136939]
- Langlois PH, Hoyt AT, Lupo PJ, Lawson CC, Waters MA, Desrosiers TA, Lammer EJ. Maternal occupational exposure to polycyclic aromatic hydrocarbons and risk of neural tube defect-affected pregnancies. *Birth Defects Res A Clin Mol Teratol*. 2012 In Press.
- Lee HQ, Hutson JM, Wray AC, Lo PA, Chong DK, Holmes AD, Greensmith AL. Changing epidemiology of nonsyndromic craniosynostosis and revisiting the risk factors. *J Craniofac Surg*. 2012; 23(5):1245–1251.10.1097/SCS.0b013e318252d893 [PubMed: 22976622]
- Lupo PJ, Langlois PH, Reefhuis J, Lawson CC, Symanski E, Desrosiers TA, Shaw GM. Maternal occupational exposure to polycyclic aromatic hydrocarbons: effects on gastroschisis among offspring in the National Birth Defects Prevention Study. *Environ Health Perspect*. 2012; 120(6):910–915.10.1289/ehp.1104305 [PubMed: 22330681]
- Lupo PJ, Symanski E, Langlois PH, Lawson CC, Malik S, Gilboa SM, Mitchell LE. Maternal occupational exposure to polycyclic aromatic hydrocarbons and congenital heart defects among offspring in the national birth defects prevention study. *Birth Defects Res A Clin Mol Teratol*. 2012; 94(11):875–881.10.1002/bdra.23071 [PubMed: 22945317]
- Nagaraja S, Anslow P, Winter B. Craniosynostosis. *Clin Radiol*. 2013; 68(3):284–292.10.1016/j.crad.2012.07.005 [PubMed: 22939693]
- Rasmussen SA, Olney RS, Holmes LB, Lin AE, Keppler-Noreuil KM, Moore CA. Guidelines for case classification for the National Birth Defects Prevention Study. *Birth Defects Res A Clin Mol Teratol*. 2003; 67(3):193–201.10.1002/bdra.10012 [PubMed: 12797461]
- Ren A, Qiu X, Jin L, Ma J, Li Z, Zhang L, Zhu T. Association of selected persistent organic pollutants in the placenta with the risk of neural tube defects. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(31):12770–12775.10.1073/pnas.1105209108 [PubMed: 21768370]
- Rocheleau CM, Lawson CC, Waters MA, Hein MJ, Stewart P, Correa A, Reefhuis J. Inter-Rater Reliability of Assessed Prenatal Maternal Occupational Exposures to Solvents, Polycyclic Aromatic Hydrocarbons, and Heavy Metals. *J Occup Environ Hyg*. 2011 In press.
- Singh VK, Singh J, Anand M, Kumar P, Patel DK, Krishna Reddy MM, Javed Siddiqui MK. Comparison of polycyclic aromatic hydrocarbon levels in placental tissues of Indian women with full- and preterm deliveries. *Int J Hyg Environ Health*. 2008; 211(5–6):639–647.10.1016/j.ijheh.2007.11.004 [PubMed: 18308633]
- Sjaastad AK, Svendsen K. Exposure to polycyclic aromatic hydrocarbons (PAHs), mutagenic aldehydes, and particulate matter in Norwegian la carte restaurants. *Ann Occup Hyg*. 2009; 53(7):723–729. doi: mep059 [pii] 10.1093/annhyg/mep059. [PubMed: 19759173]
- United States Department of Labor Bureau of Labor Statistics. North American Industry Classification System (NAICS). 1997. Retrieved 27 October, 2009, from <http://www.bls.gov/bls/naics.htm>
- United States Department of Labor Bureau of Labor Statistics. Standard Occupational Classification. 2000. Retrieved 27 October, 2009, from http://www.bls.gov/soc/soc_majo.htm

Wells PG, McCallum GP, Lam KC, Henderson JT, Ondovcik SL. Oxidative DNA damage and repair in teratogenesis and neurodevelopmental deficits. *Birth Defects Res C Embryo Today*. 2010; 90(2):103–109.10.1002/bdrc.20177 [PubMed: 20544694]

Yoon PW, Rasmussen SA, Lynberg MC, Moore CA, Anderka M, Carmichael SL, Edmonds LD. The National Birth Defects Prevention Study. *Public Health Rep*. 2001; 116(Suppl 1):32–40. [PubMed: 11889273]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Distribution of maternal factors among employed^a mothers of craniosynostosis case infants and control infants, National Birth Defects Prevention Study, 1997–2002 [*n* (%)]

Characteristic	Cases (n=316) n (%)	Controls (n=2,993) n (%)	OR (95% CI)
<i>Infant</i>			
Sex			
Male	223 (71)	1,495 (50)	2.40 (1.86, 3.09)
Female	93 (29)	1,496 (50)	1.00 (Ref)
<i>Maternal</i>			
Age (years)			
<20	14 (4)	240 (8)	1.00 (Ref)
20–24	56 (18)	645 (22)	1.49 (0.81, 2.72)
25–29	67 (21)	801 (27)	1.43 (0.79, 2.60)
30–34	104 (33)	858 (29)	2.08 (1.17, 3.70)
35	75 (24)	449 (15)	2.86 (1.58, 5.17)
Pregnancy BMI (kg/m ²)			
Underweight (<18.5)	16 (5)	153 (5)	1.02 (0.59, 1.75)
Normal weight (18.5–24.9)	172 (55)	1,676 (57)	1.00 (Ref)
Overweight (25–29.9)	73 (23)	664 (23)	1.07 (0.80, 1.43)
Obese (≥30)	53 (17)	439 (15)	1.18 (0.85, 1.63)
Parity			
0	139 (44)	1,331 (44)	1.00 (Ref)
1	177 (56)	1,661 (56)	1.02 (0.81, 1.29)
Race/ethnicity			
Non-Hispanic white	242 (77)	1,940 (65)	1.00 (Ref)
Non-Hispanic black	14 (4)	377 (13)	0.30 (0.17, 0.52)
Hispanic	45 (14)	528 (18)	0.68 (0.49, 0.95)
Other	15 (5)	141 (5)	0.85 (0.49, 1.48)
Education (years)			
<12	21 (7)	296 (10)	1.00 (Ref)
12	73 (23)	741 (25)	1.39 (0.84, 2.30)
>12	221 (70)	1,951 (65)	1.60 (1.00, 2.54)
Gestational diabetes			
No	288 (95)	2,804 (96)	1.00 (Ref)
Yes	16 (5)	118 (4)	1.32 (0.77, 2.26)
Smoking ^b			
Nonsmoker	243 (77)	2,378 (79)	1.00 (Ref)
Light (1–24 cigarettes/day)	50 (16)	423 (14)	1.15 (0.84, 1.60)
Moderate (15–24 cigarettes/day)	21 (7)	159 (5)	1.29 (0.80, 2.08)
Heavy (≥25 cigarettes/day)	2 (0.6)	33 (1)	0.59 (0.14, 2.29)
Secondhand smoke at home ^b			

Characteristic	Cases (n=316) n (%)	Controls (n=2,993) n (%)	OR (95% CI)
No	245 (78)	2,456 (82)	1.00 (Ref)
Yes	71 (22)	535 (18)	1.33 (1.01, 1.76)
Secondhand smoke at work ^b			
No	262 (83)	2,412 (81)	1.00 (Ref)
Yes	54 (17)	572 (19)	0.87 (0.64, 1.18)

BMI=body mass index; CI=confidence interval; OR=odds ratio

^aEmployed for at least one month from the three months before conception through the end of pregnancy.

^bOne month before conception through the third month of pregnancy.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Associations between maternal occupational exposure to polycyclic aromatic hydrocarbons and craniosynostosis in offspring, National Birth Defects Prevention Study, 1997–2002

Table 2

	Cases (n)	Controls (n)	Crude OR		Adjusted OR ^a	
			OR	95% CI	OR	95% CI
Exposure Status						
No PAH exposure	300	2,886	1.0	(Reference)	1.0	(Reference)
PAH exposure	16	107	1.44	(0.84,2.47)	1.75	(1.01, 3.05)

CI=confidence interval; OR=odds ratio; PAH=polycyclic aromatic hydrocarbon

^a Adjusted for maternal age and maternal education.