



Published in final edited form as:

Neuroepigenetics. 2015 October 1; 4: 12–27. doi:10.1016/j.nepig.2015.10.002.

Tet1 Oxidase Regulates Neuronal Gene Transcription, Active DNA Hydroxy-methylation, Object Location Memory, and Threat Recognition Memory

Dinesh Kumar^{1,4}, Milan Aggarwal¹, Garrett A. Kaas¹, John Lewis¹, Jing Wang¹, Daniel L. Ross², Chun Zhong³, Andrew Kennedy¹, Hongjun Song³, and J. David Sweatt^{1,*}

¹Department of Neurobiology and Evelyn F. McKnight Brain Institute, University of Alabama at Birmingham, Birmingham, AL 35294, USA

²Department of Pharmacology & Toxicology, University of Alabama at Birmingham, Birmingham, AL 35294, USA

³Institute for Cell Engineering, Department of Neurology, Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

Abstract

A dynamic equilibrium between DNA methylation and demethylation of neuronal activity-regulated genes is crucial for memory processes. However, the mechanisms underlying this equilibrium remain elusive. Tet1 oxidase has been shown to play a key role in the active DNA demethylation in the CNS. In this study, we used *Tet1* gene knockout (Tet1KO) mice to examine the involvement of Tet1 in memory consolidation and storage in the adult brain. We found that *Tet1* ablation leads to: altered expression of numerous neuronal activity-regulated genes, compensatory upregulation of active demethylation pathway genes, and upregulation of various epigenetic modifiers. Moreover, Tet1KO mice showed an enhancement in the consolidation and storage of threat recognition (cued and contextual fear conditioning) and object location memories. We conclude that Tet1 plays a critical role in regulating neuronal transcription and in maintaining the epigenetic state of the brain associated with memory consolidation and storage.

Keywords

Cytosine methylation; active demethylation; learning; memory; fear conditioning; threat conditioning; transcription factor; gene expression; chromatin; transcription; motor learning; epigenetic; HDAC; TET; DNA; hydroxymethylcytosine; metaplasticity; homeostatic plasticity; neuroepigenetics

*Corresponding Author: J. David Sweatt, Department of Neurobiology, University of Alabama at Birmingham, SHEL 1010, 1825 University Blvd, Birmingham, AL 35294, (205) 975-5196 (PH), dsweatt@uab.edu.

⁴Current Address: Computer Science and Artificial Intelligence Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139 and Broad Institute of MIT and Harvard, Cambridge, MA 02139, USA.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Recent findings have clearly implicated a role for DNA methylation in memory formation and storage (Miller and Sweatt 2007; Lubin et al. 2008; Feng et al. 2010a; Miller et al. 2010; Lesburgueres et al. 2011; Monsey et al. 2011; Sweatt 2013). Conventionally, DNA methylation has been considered to act as a transcriptional silencer (Jaenisch and Bird 2003; Bonasio et al. 2010; Feng et al. 2010b). However, recent studies point towards a more complex role of DNA methylation based on the cell type involved or the genomic context in which the methylation event occurs (Yu et al. 2013; Bahar Halpern et al. 2014; Jeltsch and Jurkowska 2014). Also, recent discoveries have identified the presence of DNA methylation at “unconventional” non-CpG (cytosine-guanine dinucleotide sequences) sites (Xie et al. 2012; Lister et al. 2013; Varley et al. 2013). Until recently, DNA methylation was considered to be essentially irreversible, however, new discoveries have shown that the methylation of memory-associated genes can be dynamic and reversible, strongly indicating the presence of an active DNA demethylation pathway in the adult brain (Miller and Sweatt 2007; Lubin et al. 2008; Ma et al. 2009; Guo et al. 2011). The concurrent discovery of the Ten-Eleven Translocation (Tet) family of proteins (Tahiliani et al. 2009) and the rediscovery of the 5hmC base in DNA derived from the CNS (Penn et al. 1972; Kriaucionis and Heintz 2009) suggested the presence of a TET-driven active DNA demethylation pathway in the brain.

Tet proteins (TET1, 2 and 3) were identified as Fe(II)-and 2-oxoglutarate-dependent dioxygenases that can oxidize 5mC to 5hmC (Iyer et al. 2009; Kriaucionis and Heintz 2009; Tahiliani et al. 2009; Globisch et al. 2010), and recent studies have given some insight into the role of Tet1 and Tet3 as drivers of active cytosine demethylation in the CNS (Guo et al. 2011; Kaas et al. 2013; Rudenko et al. 2013; Zhang et al. 2013; Li et al. 2014). Tet1KO mice have been shown to be developmentally normal (Dawlaty et al. 2011), and to manifest robust fear conditioning memory refractory to memory extinction (Rudenko et al. 2013). Recently, our lab showed that Tet1 expression exhibited a learning-associated down-regulation in vivo, and the overexpression of TET1 in adult dorsal hippocampus leads to a deficit in long-term fear conditioning memory (Kaas et al. 2013). These findings suggested that Tet1 might act as a negative memory regulator; therefore in the present studies we extended the findings of Kaas et. al. and tested the hypothesis that Tet1 deletion might improve memory acquisition, consolidation and storage in contextual and cued Pavlovian threat (fear) conditioning paradigms.

We, for the first time to our knowledge, report that Tet1KO mice have enhanced threat recognition (also known as fear conditioning) (Ledoux 2014) memory consolidation and storage. We also found that a virally mediated shRNA knockdown of *Tet1* in dorsal hippocampus led to an enhanced long-term memory for object location. Moreover, *Tet1* gene ablation led to alterations in various neuronal activity-regulated genes, including key genes from the cyclic AMP (cAMP) transcription-regulating pathway that have previously been shown to be critical in long term memory consolidation. We also observed that *Tet1*-deficient animals exhibit up-regulation of genes related to epigenomic modifications and DNA demethylation pathway including *Tet2* and *Tet3*.

Overall, we show in this study that Tet1 is crucial in maintaining the methylation status of the brain by controlling 5hmC production, and that it is also critical for the regulation of neuronal gene transcription. Most importantly, deletion of TET1 led to enhanced memory consolidation and storage. Therefore, Tet1 inhibition might serve as a useful pharmacologic target for cognitive enhancement.

Material and Methods

Animals and Genotyping

Tet1^{+/-} mice were purchased from The Jackson Laboratory (Strain Name: B6;129S4-Tet1tm1.1Jae/J, Stock Number: 017358) and were bred as heterozygotes at the University of Alabama at Birmingham. WT and KO male offspring of the heterozygote parents of age between 3–4 months were used for all the behavior and molecular experiments. For the Object Location Memory task, 10–12 week old C57BL/6 male mice from Harlan were injected rAAVs through stereotaxic surgeries. Animals were singly housed three days before the start of any behavior experiments. Animals were maintained under a 12h light/dark schedule with access to food and water *ad libitum*. All studies were performed in compliance with the University of Alabama Institutional Animal Care and Use Committee guidelines.

Cresyl Violet Staining

Tet1KO and WT males the age between 3–4 months were used for sectioning. Protocol for cresyl violet staining was adopted from (Almonte et al. 2013).

Quantitation of Global Modified Cytosines Using Mass Spectrometry

Extraction, hydrolysis and quantification of the cytosines and modified cytosines were done according to the method described previously in (Kaas et al. 2013). Tet1KO and WT (n=4 males/group) mice of age between 3–4 months were used for the tissue extraction. Statistical comparisons between two groups were performed using an unpaired t-test (two tailed). Statistical analysis between three or more groups was accomplished using one-way ANOVA with Bonferroni post hoc test.

Behavior Tasks

Baseline behavior assessments were performed as previously described (Chwang et al. 2007). Social approach paradigm was adopted from (Silverman et al. 2010; Ellegood et al. 2013). Regarding our nomenclature for Pavlovian associative conditioning to aversive foot-shock stimuli, in this paper we refer to that form of learning as “threat recognition training”. The protocols we used for threat recognition training are identical to those we have used previously and have referred to in prior publications as “fear conditioning” (see, e.g. Chwang et al., 2007). However, beginning with this manuscript we have updated our nomenclature to more accurately reflect what is likely occurring with foot-shock conditioning, that is threat response conditioning and not “fear” conditioning *per se*. The rationale behind this change in nomenclature was recently and clearly articulated by Joe LeDoux, and we fully ascribe to this new viewpoint (Ledoux 2014). For the threat recognition training (fear conditioning) paradigm, animals were first trained for a total of 3

minutes in a novel context. First 2 minutes were given as habituation period, after that an audio cue, 75dB, was played for 30 seconds, immediately followed by a foot shock (0.5mA, 1sec., for “light” and 0.8mA, 2sec., for the “strong” training). The animals were given an additional 30 seconds in the cage before removal. For the robust training paradigm, animals received 3 tone-shock pairings (75dB, 0.8mA, 2sec.), every 1 min after the first 2 minutes of habituation, and the animal was removed after a total of six and one-half minutes. For contextual testing, animals were placed back in the same context for 3 minutes after 1 hour of training for testing short-term memory, and after 24 hours, 15 days, and 30 days of training for assessing long-term memory storage. For cued testing, animals were placed in a modified context for 5 min, and the same audio cue was presented during the last 3 min. Percent freezing was scored manually with the evaluator blinded to genotype. Tet1KO and WT males (n=8 males/group) of the age between 3–4 months were used for all the behavior studies. Statistical comparisons between two groups were performed using an unpaired t-test (two tailed). Statistical analysis between three or more groups was accomplished using one-way ANOVA with Bonferroni post hoc test.

Object Location Memory (OLM) Task

The object location memory (OLM) task was adapted from (Stefanko et al. 2009; Haettig et al. 2011). Both, the OLM training and the testing sessions were recorded using TopScan (Clever Sys, Reston, VA) and the data were analyzed by the individual blinded to the genotype of each animal. The relative exploration time was recorded and expressed as a discrimination index (D.I. = $[t\text{-novel} - t\text{-familiar}] / [t\text{-novel} + t\text{-familiar}] \times 100$). C57BL/6 mice from Harlan, 10–12 weeks of age (n = 8 males/group) were used. Statistical comparisons between two groups were performed using an unpaired t-test (two tailed).

rAAVs and Stereotaxic Surgeries

High titers of recombinant AAV2/9 virus expressing either a HA-tagged human TET1 catalytic or a catalytically inactive TET1 mutant (H1671Y/D1673A) domain were generated as previously described (Guo et al. 2011). Viral injection into the dorsal hippocampus of 10–12 week old C57BL/6 was done using the following stereotaxic coordinates: –2 mm antero-posterior, ± 1.5 mm mediolateral, and –1.6 mm dorsoventral from bregma. A total of 1.5 μ l of viral solution per hemisphere was injected. Injections were performed using a 10ml Hamilton Gastight syringe controlled by a Pump 11 Elite Nanomite Programmable Syringe Pump (Harvard Apparatus). The injections proceeded at a speed of 150 nl min⁻¹ through a 32 gauge needle. The injection needle was left in place an additional 5 min to allow the fluid to diffuse. Behavioral experiments were performed 30 days following stereotaxic delivery of rAAVs. Electrophysiology experiments were conducted 14 days following AAV-eYFP, Tet1 or Tet1m viral injection.

Tissue Collection and Gene Expression

Hippocampal sub-dissections were done as described in (Lein et al. 2004). In the case of cortex, predominantly pre-frontal cortex (plus some adjacent tissue) was taken for all molecular analysis. All dissections were carried out under a dissecting scope and immediately frozen on dry ice and stored at –80°C until further processing. RNA was extracted using the RNeasy Mini Kit (Qiagen) following the manufacturer’s protocol. 150ng

of total RNA was converted to cDNA using the iScript cDNA synthesis Kit (Bio-Rad). Quantitative real time PCR was performed on an iQ5 real-time PCR detection system using iQ™ SYBR® Green Supermix and 300 M of primer. All qRT-PCR primers were designed using Primer Quest (Integrated DNA Technologies) to span exon-exon junctions or were acquired directly as pre-designed PrimeTime® qPCR Primer Assays (Integrated DNA Technologies). For all qRT-PCR reactions, glyceraldehyde 3-phosphate dehydrogenase (*Gapdh*) was used as an internal control. The gene expression analysis was done using the comparative Ct method adopted from (Livak and Schmittgen 2001; Pfaffl 2001). An R package, ComplexHeatmap (<https://github.com/jokergoo/ComplexHeatmap>) was used to make heatmaps from the qRT-PCR gene expression data. Statistical comparisons between two groups were performed using an unpaired t-test (two tailed).

Electrophysiology

Extracellular field potential recordings from hippocampal area CA1 Schaffer collateral synapses were obtained as described earlier (Feng et al. 2010a). Baseline synaptic transmission was plotted using stimulus intensities between (1–30mV), and evoked field EPSP slopes. Subsequent experimental stimuli were set to an intensity that evoked a fEPSP that had a slope of 50% or 25% of the maximum fEPSP slope. Various time intervals (10–300ms) between constant-stimulus paired pulses were used to measure Paired-Pulse Facilitation (PPF). For inducing long-term potentiation, three different stimuli were used: in the first two cases LTP was induced using one tetanus stimulus of 100Hz for either 1 sec or 0.1 sec, and in the third case LTP was induced using a one-time theta burst stimulation (one episode of theta burst stimulation, 10 bursts at 5 Hz, each with 4 stimuli at 100 Hz with 20 sec intervals). Data was recorded for 3 hours following stimulation. Field EPSPs were recorded every 20 sec. (traces were averaged for every 2 min interval). Animals (n=6 males/group, 7 slices each) of the age 3–4 months were used for making slices. Analysis was done using an unpaired t-test (two-tailed) and two-way ANOVA.

Statistical Analysis

All statistical analysis was done using GraphPad Prism version 4.00 for Windows (GraphPad Software, San Diego California USA).

Results

***Tet1* deletion does not affect overall adult brain morphology**

*Tet1*KO mice were originally generated by deletion of exon 4 and were shown to be grossly normal (Dawlaty et al. 2011; Rudenko et al. 2013). To check for any morphological defects in the brain, we utilized cresyl violet staining of brain sections of the WT and KO mice. *Tet1*KO mice showed no obvious morphological differences in comparison to WT (Fig. 1A–C). The loss of *Tet1* mRNA in *Tet1*KO animals was confirmed by quantitative real time PCR (qRT-PCR) ($p^{****} < 0.0001$, Fig. 1D).

5hmC is enriched in brain areas involved in active memory processing

The exact function of 5hmC is not yet known. However, relative 5hmC levels, but not 5mC levels, have been consistently shown to be highest in the brain, compared to all other tissues

(Kriaucionis and Heintz 2009; Globisch et al. 2010; Munzel et al. 2010). This implies that 5hmC may have important brain-specific functions. Given that memory processing (acquisition, consolidation, storage and retrieval) is a major function of the central nervous system, we determined if the 5hmC mark is differentially distributed in brain subareas involved in cognitive functions. We used a highly sensitive HPLC/MS technique to answer this question (Fig. 2A). We measured the percentage 5hmC, 5mC and 5C (unmodified cytosine) relative to total cytosine (5hmC + 5mC + 5C) in five different brain areas, and our measurements were consistent with earlier reported results (Globisch et al. 2010; Munzel et al. 2010). We found that 5hmC levels in area CA1 (0.70%) and cortex (0.74%) were significantly higher (p value <0.0001) than in the dentate gyrus (0.63%), area CA3 (0.55%), and cerebellum (0.40%), (Fig. 2B). Cerebellum was found to have the lowest amount of 5hmC (0.40%) among the brain regions tested (Fig. 2B). In contrast to 5hmC, we found a fairly uniform distribution of the 5mC mark, ranging from 7 to 8% (Fig. 2C), and the percentage of unmodified cytosines, around 92–93%, was also similar in the different brain areas (Fig. 2D). These results, consistent with other recent studies (Khare et al. 2012; Lister et al. 2013; Li et al. 2014), suggest the involvement of the 5hmC mark in learning and memory.

Tet1 deletion leads to a reduction in 5hmC levels

Tet proteins have been shown to convert 5mC to 5hmC (Tahiliani et al. 2009), therefore we wanted to investigate if Tet1 contributes to establishing baseline 5hmC levels in the brain and if 5mC and 5C (cytosine) levels are also affected by *Tet1* ablation. To determine this we sub-dissected WT and Tet1KO mice brain regions (CA1, DG, CA3, cortex and cerebellum) and then measured 5mC, 5hmC, and 5C levels using a quantitative HPLC/MS technique. We found a significant reduction ($p^{**}<0.05$, $p^{**}<0.005$, and $p^{****}<0.0001$) in 5hmC levels in Tet1KO mice in all brain areas investigated (Fig. 3A–E, left column). These data strongly support the idea that Tet1 acts catalytically on 5mC in the CNS in order to generate 5hmC (Kaas et al. 2013; Rudenko et al. 2013).

Since 5hmC can only be derived from oxidation of 5mC, it was possible that in the absence of Tet1-mediated demethylation in Tet1KO mice, there might be an increase in relative 5mC levels, given that we observed a decrease in 5hmC levels. However, we did not see a significant increase ($p > 0.05$) in 5mC levels except in the cortex ($p^*<0.05$) (Fig. 3A–E, center column). Interestingly in the cerebellum, both 5mC and 5hmC levels dropped significantly in Tet1KO mice (p^* and $p^{**}<0.05$) (Fig. 3E, left and center). We also did not observe a significant change ($p > 0.05$) in unmodified cytosine (5C) except in the case of cerebellum, where its levels in Tet1KO were slightly but significantly higher than the WT ($p^*<0.05$) (Fig. 3A–E, right column). These results are consistent with a recently published study (Li et al. 2014) indicating that Tet1 may not be the sole regulator of active 5mC demethylation in the brain.

Tet1KO mice show normal baseline behaviors, motor memory, and social interactions

We performed a series of baseline behavior tests on Tet1KO mice to determine if *Tet1* deletion affects basic exploratory, emotional and social behavior. In the open field paradigm, that measures general locomotor activity and anxiety, WT and Tet1KO did not show any

significant differences ($p > 0.05$) in the total distance travelled and mean velocity (Fig. 4A). Also, both genotypes spent similar amounts of time in the center and along the walls of the chamber (Fig. 4B), suggesting a lack of difference in anxiety levels ($p > 0.05$). Another paradigm, the elevated plus maze, was also used to assess anxiety and depression-like behaviors. Tet1KO and WT did not show any significant differences ($p > 0.05$) in time spent in the open or closed arms nor in the number of entries made into each arm ($p > 0.05$) (Fig. 4C). Overall these data indicate normal baseline locomotor and anxiety-related behavior in Tet1KO mice.

We also utilized a social approach paradigm described in (Silverman et al. 2010; Ellegood et al. 2013) using WT and Tet1KO animals to determine if Tet1KO mice exhibit any autistic-like behaviors. WT and Tet1KO showed no significant differences ($p > 0.05$) in their social interaction preferences; both preferred to spend more time with a novel mouse versus a novel object or empty chamber (Fig. 4D). Based on these results we conclude that Tet1KO mice have normal exploratory and social behavior, at least as assessed using these protocols.

In terms of molecular changes, *Tet1* ablation elicited a significant decrease not only in 5hmC but also in 5mC levels in the cerebellum. We wanted to check if this reduction in the levels of both these cytosine derivatives was associated with any effect on cerebellum-dependent motor memory in Tet1KO mice. Using the accelerating rotarod paradigm we found that Tet1KO and WT mice showed no significant differences in terms of the time-to-fall or velocity-at-fall in their rotarod performance on all three trial days (Fig. 4E). These observations suggest that Tet1KO mice have normal cerebellum-dependent motor memory.

Tet1KO mice exhibit enhanced threat recognition memory (contextual and cued fear conditioning)

Enrichment of 5hmC in brain areas (CA1 and Cortex) that are involved in memory processing propelled us to investigate Pavlovian threat recognition learning in Tet1-deficient animals. Therefore we assessed both cued and contextual fear conditioning in Tet1KO animals versus littermate WT controls.

To evaluate long-term memory consolidation and storage, both WT and Tet1KO cohorts were trained with two different shock protocols, using “light” (0.5mA for 1 sec.) or “strong” (0.8mA for 2 sec.) training in a novel context. Memory retention was tested 24 hours after training. Tet1KO mice exhibited significantly higher freezing ($p^* < 0.05$ and $p^{**} < 0.005$) than littermate WT mice (Fig. 5B–C, left column) with both “light” and “strong” training paradigms. However, we observed no significant difference ($p > 0.05$) in freezing levels at 24 hours after robust training using a more intense shock protocol (0.8 mA, 2sec., repeated 3 times) (Fig. 5D), demonstrating that WT animals and Tet1KO animals have similar maximal learning capacities.

It has been shown that two weeks after memory acquisition, memories undergo an extensive systems consolidation and are transferred to the cortex for long-term storage. These memories then become independent of the hippocampus and are referred to as “remote” memories (Dudai 2004). To study if *Tet1* ablation affects remote memory systems consolidation and long-term storage of remote memories, WT and Tet1KO mice were tested

in the training context 15 and 30 days after light or strong training. Interestingly, in both cases, after 15 days ($p^{***}<0.005$, $p^{**}<0.005$) and after 30 days ($p^{***}<0.0005$, $p^{**}<0.005$), Tet1KO animals still showed significantly higher freezing levels than WT (Fig. 5B–C, center and right columns). Together, these results indicate an enhanced long-term remote memory consolidation and storage in Tet1KO mice.

We also checked memory acquisition and short-term memory formation in Tet1KO animals. For this, we tested animals in the same context, one hour after the training. WT and Tet1KO did not show any significant difference ($p>0.05$) in the percentage freezing one hour after training, indicating that fear learning and short-term memory are normal in Tet1KO mice (Fig. 5E).

We also investigated if *Tet1* ablation had an effect on hippocampus-independent cue memory in which animals learn to associate an auditory cue with a foot shock. For this paradigm, we trained WT and KO mice using the same “light” and “strong” foot-shock protocols that were used for the contextual training, however using an auditory cue (75 dB) as the conditioned stimulus (CS). For testing, animals were exposed to a novel context for 5 minutes, 24 hours after training, during which the CS audio cue was played for the last three minutes (Fig. 6A). We observed significantly higher freezing ($p^* < 0.05$) in Tet1KO animals compared to WT mice using both “light” (Fig. 6B) and “strong” (Fig. 6C) protocols. We did not observe a significant difference ($p>0.05$) in freezing with the “robust” foot-shock training protocol used as an additional control (Fig. 6D). Overall, these results show an enhancement in both hippocampus-dependent and -independent threat memory in Tet1KO mice.

Virally-mediated knock-down of *Tet1* expression in the hippocampus enhances spatial memory for object location

The use of fear conditioning training in a global knockout mouse line as described above does not allow for the determination of whether or not the role of TET1 oxidase in threat memory is limited to the hippocampus or stems from a synergistic effect of TET1 activity spanning several brain regions. Moreover, the enhanced memory phenotype in the Tet1KO mice may be a manifestation of potential developmental changes. For these reasons, and to investigate whether observed phenotype is due to changes in neuronal function, an additional series of experiments were carried out, which combined the known hippocampus-selective, object location memory task together with the acute reduction of *Tet1* expression in the dorsal hippocampus. We selectively targeted dorsal hippocampus as this region has been shown to be actively involved in spatial memory processing. Knock down of *Tet1* mRNA levels was accomplished using AAVs engineered to express shRNAs specifically designed to target endogenous *Tet1* transcripts, as described in (Guo et al. 2011).

To validate the effectiveness of our approach, we first stereotaxically injected either AAV-*Tet1*-shRNA viral particles or a scrambled control (scr-shRNA), into the dorsal hippocampus. Following two weeks post-surgery, we observed robust expression throughout the dorsal CA1 and CA3 regions of the hippocampus (Fig 7A). In addition, RT-PCR analysis revealed a 46% reduction in *Tet1* mRNA levels in hippocampal tissue derived from AAV-*Tet1*-shRNA injected mice compared to scr-shRNA controls ($*p < 0.05$) (Fig. 7B). As

a biochemical control for *Tet1* knockdown we also measured the percentage of 5hmC and 5mC in microdissected hippocampal tissue from both groups, using mass spectrometry as described earlier. We did not see a significant decrease ($p > 0.05$) in the 5hmC levels in *Tet1*-shRNA infused mice compare to the control. However, consistent with a reduction in TET1 activity we observed a trend ($p = 0.06$) towards a decrease in global 5hmC levels (Fig. 7C) in *Tet1*-shRNA infused mice. No differences were found with regards to global 5mC levels (Fig. 7D).

Using the open field paradigm, we first tested whether a reduction in dorsal hippocampal *Tet1* expression in mice might affect their baseline behaviors or locomotion. We found no significant differences ($p > 0.05$) between the two groups in terms of total distance traveled or in time spent in the center or periphery of the open field. These results, consistent with our previous results with the Tet1KO mice, indicate that loss of *Tet1* expression does not affect basal exploratory and locomotor behavior in mice (Fig. 7E, F).

We next determined if *Tet1* knockdown in the dorsal hippocampus had any influence on object location memory (OLM). For this task, *Tet1*-shRNA and control scr-shRNA infused mice were first exposed to two novel objects for a total of 10 minutes (Fig. 7G). Mice were then tested for long-term memory of the original location of the objects 24 hours later (Fig. 7I). Importantly, during training, a comparison of the percentage of time spent exploring each of the two objects did not differ significantly between groups (Fig. 7H), indicating no bias towards either object or its location in the chamber. In contrast, during the testing phase, mice receiving injections of AAV-*Tet1*-shRNA showed a statistically significant enhancement in memory for the object in the familiar location, and spent significantly more time ($p^* < 0.05$) exploring the object in the novel location, compared with AAV-scr-shRNA controls (Fig. 7J). These results are consistent with the idea that Tet1 may serve as a critical memory suppressor and that inhibition of Tet1 leads to persistence of hippocampus-dependent long-term memories.

Tet1KO mice have normal basal synaptic transmission and LTP, and virally mediated TET1 overexpression in dorsal hippocampus does not affect LTP

We next determined if enhanced long-term threat recognition memory is accompanied by a facilitation of long-term potentiation (LTP) in Tet1KO mice. A recent study (Rudenko et al. 2013) showed normal long-term potentiation in Tet1KO mice using a relatively strong LTP induction protocol (two episodes of theta burst stimulation, 10 bursts at 5 Hz, each with 4 stimuli at 100 Hz with 10 seconds intertrain intervals). To complement these prior studies we used a relatively mild LTP induction stimulus (100Hz, 1sec., delivered once) to check for the possibility of enhanced long term potentiation in Tet1KO mice at near-threshold LTP induction stimuli. However, we did not observe any significant enhancement ($p > 0.05$) in LTP measured in area CA1 using this stimulus (Fig. 8A, bottom). Neither did we observe enhanced LTP using a sub-threshold stimulus intensity under our conditions (Fig. 8B, bottom). We also did not find any significant difference in paired pulse facilitation (PPF) ($p > 0.05$), suggesting that both WT and Tet1KO have normal presynaptic release probability (Fig. 8A–B, middle). Basal synaptic transmission was also found to be normal in Tet1KO mice (Fig. 8A–B, top).

To further investigate if Tet1 plays a role in regulating LTP, we used the virus-mediated overexpression approach described in (Guo et al. 2011; Kaas et al. 2013). Thus, we stereotaxically injected AAVs overexpressing a hemagglutinin (HA)-tagged catalytic domain of human TET1, or a catalytically inactive version (TET1m), into the dorsal hippocampus. At two weeks post-infection, hippocampal slices were prepared for LTP assessment. LTP was induced using one-time theta burst stimulation (one episode of theta burst stimulation, 10 bursts at 5 Hz, each with 4 stimuli at 100 Hz with 20 sec intervals). We found that neither the overexpression of the catalytically active (AAV-TET1) nor the catalytically inactive (AAV-TET1m) TET1 peptide led to any significant effect on LTP compared to control AAV-eYFP (Fig 8C, bottom) ($p > 0.05$). In addition, no significant differences in basal synaptic transmission ($p > 0.05$) (Fig 8C, top) and PPF (Fig 8C, middle) between the AAV-TET1, AAV-TET1m and AAV-eYFP infused mice.

Overall, these results indicate that despite Tet1KO mice displaying enhanced behavioral memory, hippocampal LTP remains normal in these animals – replicating the prior results of Rudenko et. al. (Rudenko et al. 2013) using two additional LTP induction protocols, and in addition testing an independent assessment of the effects of TET1 overexpression. Taken together, these various observations strongly suggest that the behavioral effects of Tet1 knockout or overexpression do not involve hippocampal LTP, but rather involve some other form of synaptic or cellular plasticity.

Tet1 deletion leads to altered transcription of memory-associated genes

De novo transcription and translation of activity-regulated genes is requisite for long-term memory storage. A variety of neuronal genes regulate neural plasticity in response to a learning experience, and altered transcription of these genes is crucial in neural development, learning, and memory (Leslie and Nedivi 2011; West and Greenberg 2011). We used qRT-PCR to investigate the expression of several activity-induced genes that have been implicated in various forms of synaptic plasticity and synapse development, in the different brain areas of Tet1KO mice under study: CA1, CA3, DG, cerebral cortex, and cerebellum.

We observed a significant decrease ($p^* < 0.05$, $p^{****} < 0.0001$) in the expression of *Arc* (activity-regulated cytoskeleton-associated protein) in all brain areas except cerebellum (Fig. 9). Interestingly, *Egr1* (early growth response protein 1) was significantly decreased in CA1 ($p^{***} < 0.0005$), but was significantly increased in cerebellum ($p^{**} < 0.005$), and did not show any significant change in expression in other brain areas (Fig. 9). A significant increase ($p^* < 0.05$, $p^{**} < 0.005$) in the expression of *Creb1* (cAMP responsive element binding protein 1) was observed in CA1, cortex and cerebellum (Fig. 9). We also probed *Bdnf* (brain-derived neurotrophic factor) transcripts by using exon IX primers (an exon present in all the expressed isoforms), and found a significant increase ($p^{***} < 0.0005$) in the expression of *Bdnf* in all the brain areas examined except cortex (Fig. 9). *c-Fos* (FBJ osteosarcoma oncogene) was significantly downregulated in area CA1, DG, and cortex. A significant increase in *calcineurin* expression ($p^* < 0.05$, $p^{**} < 0.005$) was seen in all the brain areas examined (Fig. 9). *reelin* (*reln.*) was significantly upregulated ($p^* < 0.05$, $p^{**} < 0.005$) only in DG and cerebellum (Fig. 9). Interestingly, *Homer1* (homer protein homolog 1) showed a

significant downregulation ($p^* < 0.05$) in area CA1 and cortex, and a significant upregulation ($p^* < 0.05$) in area CA3 (Fig. 9). *Cdk5* (cyclin-dependent kinase 5) was significantly upregulated ($p^* < 0.05$, $p^{**} < 0.005$) in all brain areas except DG (Fig. 9). *Nr4a2* (nuclear receptor related 1 protein) was also significantly upregulated ($p^* < 0.05$, $p^{**} < 0.005$, and $p^{****} < 0.0001$) in all brain areas except cerebellum (Fig. 9). *NPas4* (neuronal PAS domain protein 4) was significantly downregulated ($p^{**} < 0.005$, $p^{***} < 0.0005$, $p^{****} < 0.0001$) in all the brain areas examined except CA3 (Fig. 9). These results show that genetic deletion of *Tet1* leads to an extensive alteration in the expression of crucial neuronal-activity regulated genes, in a wide variety of memory-associated brain regions, and that *Tet1* may positively or negatively regulate the same gene depending upon the brain area.

Tet1 regulates the transcription of *Tet2*, *Tet3*, and other active demethylation pathway genes

Tet1 clearly has a role in maintaining 5hmC levels in the brain, as illustrated by our observation that loss of *Tet1* led to a significant reduction in the levels of 5hmC in memory-associated brain subregions (Fig. 3). However, the loss of *Tet1* and the reduction in the level of 5hmC do not translate to a significant increase in 5mC in *Tet1*KO mice, except in the cortex (Fig. 3). This suggests the possibility that *Tet2* and *Tet3* might be playing a compensatory role in the absence of *Tet1*, restoring normal baseline cytosine methylation levels, even in the face of loss of one driver of active demethylation. Therefore we checked the expression of *Tet2* and *Tet3* mRNAs in different brain areas of *Tet1*KO mice by qRT-PCR and found a significant ($p^* < 0.05$, $p^{**} < 0.005$, $p^{***} < 0.0005$, $p^{****} < 0.0001$) compensatory increase in the expression of both the transcripts (Fig. 10).

As described in the introduction, mounting evidence now points to the existence of an active demethylation pathway in cells (Fig. 10A), and the components of this pathway are conserved in the brain. The active demethylation pathway cycle consists of cytosine methylation by DNA methyltransferases (DNMTs), repetitive oxidation of the methyl group by Tet proteins (TET1, 2, and 3), and restoration of unmodified cytosines by DNA glycosylase mediated base excision repair (BER) or by nucleotide excision repair (NER) (Wu and Zhang 2014), (Fig. 10A). We decided to check if there is a compensatory alteration in the expression of the various genes involved in this pathway in various regions of the *Tet1*KO brain. We found a significant increase ($p^* < 0.05$, $p^{**} < 0.005$, $p^{***} < 0.0005$, $p^{****} < 0.0001$) in the expression of several genes involved in this pathway (Fig. 10): *Gadd45b* (growth arrest and DNA-damage-inducible beta), *Smug1* (single-strand selective monofunctional uracil DNA glycosylase), *Apobec1* (apolipoprotein B mRNA-editing enzyme 1), and *Tdg1* (thymine DNA glycosylase 1) were all upregulated in all the different brain areas investigated (Fig. 10). Interestingly, *Gadd45b* has also been characterized as a neuronal activity-induced immediate early gene (Ma et al. 2009). From these results, we conclude that *Tet1*KO mice have an overall compensatory upregulation of various active demethylation genes including *Tet2* and *Tet3*, suggesting a pathway in cells allowing the coordinated genetic regulation of the entire machinery of active DNA demethylation.

Tet1 deletion leads to up-regulation of other epigenetic modifiers

Different epigenetic marks (5mC, 5hmC, 5fC, and 5caC) have been shown to interact with common and specific epigenetic reader proteins that have gene regulation capability (Iurlaro et al. 2013; Spruijt et al. 2013). For this reason we analyzed the expression of different epigenetic modifiers after the ablation of *Tet1*, including DNMTs and methyl-cytosine binding proteins. Interestingly, we found that Tet1KO mice manifest significant increases ($p^* < 0.05$, $p^{**} < 0.005$, $p^{***} < 0.0005$ and $p^{****} < 0.0001$) in the expressions of *Dnmt1*, *3a*, *3b* (DNA methyltransferases); *Mecp2* (methyl CpG binding protein 2) and *Mbd3* and *4* (methyl-CpG-binding domain protein) transcripts in the five brain regions under study (the only exception being no change in *Mbd4* transcripts in CA1 and CA3) (Fig. 11). This observation also suggests the existence of coordinated transcriptional regulation of genes contributing to the pathways subserving epigenomic regulation in the CNS.

Discussion

A growing body of evidence suggests that DNA methylation in the adult brain is dynamically regulated and crucial for controlling memory formation and storage (Sweatt 2013). In this regard our studies further establish Tet1 oxidase as a driver of active demethylation in the adult CNS and a controller of memory consolidation and stabilization.

In the present studies we found the presence of 5hmC throughout the brain, however, we observed a significant enrichment of 5hmC levels in hippocampal area CA1 and cortex (Fig. 2B). Since CA1 and cortex are parts of the brain actively involved in memory processing, the specific enrichment of 5hmC in these areas is consistent with a role for 5hmC in cognitive function. Other recent studies using genome-wide sequencing have also pointed at the involvement of 5hmC in learning and memory (Khare et al. 2012; Lister et al. 2013; Li et al. 2014).

Tet1KO mice showed lower levels of 5hmC than WT in the different brain regions we assessed (Fig. 3A–E, left column), data that is directly supportive of the hypothesis that Tet1 catalyzes the formation of 5hmC (Tahiliani et al. 2009). However, deletion of TET1 did not lead to significant accumulation of 5mC except in cortex (Fig. 3A–E, center), suggesting possible compensation by TET2 and TET3 to restore normal levels of 5mC in DNA through ongoing demethylation. This interpretation is consistent with a recent study by Li et al. showing the involvement of TET3 in mediating rapid behavioral adaptation in the prefrontal cortex (Li et al. 2014).

One striking discovery in the present studies is the observation of memory enhancement in Tet1KO animals, particularly regarding threat recognition long-term memory and remote memory consolidation. Consistent with a previous study (Rudenko et al. 2013), we found normal threat memory acquisition in Tet1KO mice, and that short-term threat memory was also normal in Tet1KO mice (Fig. 5E). However, we found an enhancement in memory consolidation and long-term storage of hippocampus and cortex dependent memories in Tet1KO mice (Fig. 5B–C). We also found an enhancement in the hippocampus-independent, amygdala-dependent cue memory in Tet1KO mice (Fig. 6A–D). We further showed that a virus-mediated shRNA knock-down of *Tet1* in dorsal hippocampus led to an enhanced

hippocampus-dependent long-term spatial memory for object location (Fig. 7J). Recently, Zhang et al. reported that a deletion of Tet1 resulted in a hippocampus-dependent spatial memory impairment as assessed by the Morris water maze task (Zhang et al. 2013). These observations suggest a different role of Tet1 in different memory types. Although both Morris water maze and contextual fear conditioning are hippocampal-dependent tasks, Morris water maze may involve stronger and more aversive motivational factors than fear conditioning, and occurs over many more training trials of longer duration. These differences might account for differential susceptibilities to effects of Tet1 knockout in the water maze versus fear conditioning. Overall however, these new findings establish Tet1 as a regulator of associative conditioning and spatial memory

The cellular mechanisms through which Tet1 oxidase and active demethylation regulate memory formation and consolidation remain mysterious. However, the present results along with others recently published (Kaas et al. 2013; Rudenko et al. 2013) make it clear that hippocampal LTP is not a strong candidate to subserve this function. Thus, in the present studies *Tet1* gene deletion and TET1 catalytic domain overexpression in mice did not alter long-term potentiation (Fig. 8A–C). These findings suggest that TET1 is acting *via* some other form of synaptic or cellular plasticity. Two specific possibilities are: altered synaptic LTD (Rudenko et al. 2013) or altered neuron-wide synaptic homeostasis (Sweatt 2013; Meadows et al. 2015; Yu et al. 2015). Future studies will hopefully yield further insights into this important issue.

In our studies of transcriptional regulation in Tet1KO mice (Fig. 9), we investigated the expression of a wide variety of known memory-associated genes in Tet1KO mice. A number of these genes, (*Arc*, *Bdnf*, *calcineurin*, *reelin* and *Npas4*) have previously been shown to be either epigenetically regulated or shown to interact with cellular epigenetic signaling (Miller and Sweatt 2007; Lubin et al. 2008; Miller et al. 2010; Penner et al. 2011; Rudenko et al. 2013). Interestingly, in the present studies we found that the deletion of *Tet1* can lead to down-regulation or up-regulation of the same memory-associated gene depending upon the brain area; for example, *Egr1* is downregulated in CA1 and upregulated in cerebellum and *Homer1* is downregulated in CA1 and cortex whereas it is upregulated in CA3. We found a significant increase in *Creb1* in CA1 and in cortex; *Creb1* is an important component of the cAMP pathway and has been shown to be a positive regulator of long-term memory storage (Dash et al. 1990; Bartsch et al. 1998). We also found a significant increase in the expression of two important *Creb1* targets, *Bdnf* (in CA1) and *Nr4a2* (in both CA1 and Cortex). Both *Bdnf* and *Nr4a2* expression have been shown to increase following treatment with HDAC inhibitors that result in memory enhancement, and blocking *Nr4a* signaling affects the ability of HDAC inhibitors to enhance memory (Vecsey et al. 2007; Hawk et al. 2012). Finally, we observed a significant increase in *Cdk5* transcription both in CA1 and cortex, and recently *Cdk5* has been implicated in the cAMP pathway (Guan et al. 2011).

Our results investigating altered gene transcription in Tet1KO mice demonstrate broad secondary changes in the transcription of genes encoding epigenomic signaling enzymes including components of both the cytosine methylation and demethylation pathways. These findings suggest the model that active DNA demethylation pathways are coordinately regulated at the transcriptional level, perhaps as a cellular form of homeostatic plasticity at

the genomic level. Consistent with this idea, we found that in Tet1KO mice, not only *Tet2* and *Tet3* but also other known genes involved in the cytosine demethylation pathway (*Gadd45b*, *Smug1*, *Apobec1*, and *Tdg*) are significantly upregulated (Fig. 10). Interestingly, a recent study shows distinct roles of Tet1 and Tet2 in mouse embryonic stem cells, Tet1 mainly controlling 5hmC levels at gene promoters and transcription start sites, while Tet2 maintains 5hmC levels in gene bodies and at exon boundaries (Huang et al. 2014). In the future it will be interesting to determine if different *Tet* isoforms similarly perform different roles in epigenomic structure in the CNS (Li et al. 2014). However, the current results and those of (Kaas et al. 2013) strongly indicate the coordinated expression of genes encoding major components of the active DNA demethylation pathway.

Our results are in good agreement with recently published findings from the Tsai and Jaenisch laboratories characterizing the same line of *Tet1*-deficient mice. Specifically, our results independently replicate their findings concerning normal brain morphology, altered 5-hmC production, normal baseline behaviors, normal hippocampal synaptic transmission and LTP, and altered *Arc*, *c-Fos*, and *Npas4* gene transcription in *Tet1* $-/-$ mice. Thus our work complements these previous findings and also extended them by investigating additional brain subregions, social interaction behavior, and motor learning, thereby revealing an important new attribute of the *Tet1* $-/-$ mouse line, which is enhanced long-term and remote threat recognition memory (fear conditioning). Interestingly, while our paper was in revision a similar study came out from Nestler lab (Feng et al. 2015), however focusing on the role of Tet1 and 5hmC in cocaine action in the nucleus accumbens. They reported a down regulation of Tet1 in nucleus accumbens in response to chronic cocaine administration and also showed, using virally mediated manipulations, that Tet1 acts as a negatively regulator of cocaine reward memory.

Considering transcription changes vis-à-vis the memory enhancement we observed, there are several specific possibilities that may have contributed to the enhanced memory consolidation and storage in Tet1KO. These include: 1) upregulation of *Creb1* (a positive regulator of memory storage) and other activity-regulated genes involved in the cAMP pathway as described before; 2) decrease in *Npas4* expression in Tet1KO mice, which might have led to an increase in the number of excitatory circuits resulting in the enhancement in memory consolidation and storage, because *Npas4* is a transcription factor that has been implicated in inhibitory synapse development (Lin et al. 2008; Bloodgood et al. 2013); and 3) Increased expression of DNMTs in Tet1KO mice might have led to a secondary increase in memory capacity, because DNMT inhibition has been shown to diminish memory consolidation and storage. These possibilities are not mutually exclusive, and all the three changes might have together altered the epigenetic state of the Tet1KO brain such that memory consolidation and storage is favored.

Overall, our results implicate Tet1 as a suppressor of memory formation. Since the Tet1KO mice are developmentally normal and have no obvious changes in baseline behavior or motor activity yet exhibit memory enhancement, Tet1 inhibition may serve as a strong therapeutic target for memory restoration in neurodegenerative disorders such as Alzheimer's disease, in age-related cognitive decline, or in intellectual disability syndromes.

Acknowledgments

The authors thank Li-Huei Tsai and Joe LeDoux for helpful discussions, Alison Margolies for experimental assistance, and Cristin Gavin for help in preparing the figures and manuscript. We apologize to the many authors whose primary work was not directly cited, owing to limitations of space. This research was supported by funds from the NIMH (MH57014 and MH091122), the Ellison Medical Foundation, the Evelyn F. McKnight Brain Research Foundation, the Swedish Research Council (to DK), and the Swedish Society for Medical Research (to DK).

References

- Almonte AG, Qadri LH, Sultan FA, Watson JA, Mount DJ, Rumbaugh G, Sweatt JD. Protease-activated receptor-1 modulates hippocampal memory formation and synaptic plasticity. *J Neurochem.* 2013; 124:109–122. [PubMed: 23113835]
- Bahar Halpern K, Vana T, Walker MD. Paradoxical role of DNA methylation in activation of FoxA2 gene expression during endoderm development. *J Biol Chem.* 2014
- Bartsch D, Casadio A, Karl KA, Serodio P, Kandel ER. CREB1 encodes a nuclear activator, a repressor, and a cytoplasmic modulator that form a regulatory unit critical for long-term facilitation. *Cell.* 1998; 95:211–223. [PubMed: 9790528]
- Bloodgood BL, Sharma N, Browne HA, Trepman AZ, Greenberg ME. The activity-dependent transcription factor NPAS4 regulates domain-specific inhibition. *Nature.* 2013; 503:121–125. [PubMed: 24201284]
- Bonasio R, Tu S, Reinberg D. Molecular signals of epigenetic states. *Science.* 2010; 330:612–616. [PubMed: 21030644]
- Chwang WB, Arthur JS, Schumacher A, Sweatt JD. The nuclear kinase mitogen-and stress-activated protein kinase 1 regulates hippocampal chromatin remodeling in memory formation. *J Neurosci.* 2007; 27:12732–12742. [PubMed: 18003853]
- Dash PK, Hochner B, Kandel ER. Injection of the cAMP-responsive element into the nucleus of Aplysia sensory neurons blocks long-term facilitation. *Nature.* 1990; 345:718–721. [PubMed: 2141668]
- Dawlaty MM, Ganz K, Powell BE, Hu YC, Markoulaki S, Cheng AW, Gao Q, Kim J, Choi SW, Page DC, et al. Tet1 is dispensable for maintaining pluripotency and its loss is compatible with embryonic and postnatal development. *Cell Stem Cell.* 2011; 9:166–175. [PubMed: 21816367]
- Dudai Y. The neurobiology of consolidations, or, how stable is the engram? *Annu Rev Psychol.* 2004; 55:51–86. [PubMed: 14744210]
- Ellegood J, Babineau BA, Henkelman RM, Lerch JP, Crawley JN. Neuroanatomical analysis of the BTBR mouse model of autism using magnetic resonance imaging and diffusion tensor imaging. *Neuroimage.* 2013; 70:288–300. [PubMed: 23275046]
- Feng J, Shao N, Szulwach KE, Vialou V, Huynh J, Zhong C, Le T, Ferguson D, Cahill ME, Li Y, et al. Role of Tet1 and 5-hydroxymethylcytosine in cocaine action. *Nat Neurosci.* 2015; 18:536–544. [PubMed: 25774451]
- Feng J, Zhou Y, Campbell SL, Le T, Li E, Sweatt JD, Silva AJ, Fan G. Dnmt1 and Dnmt3a maintain DNA methylation and regulate synaptic function in adult forebrain neurons. *Nat Neurosci.* 2010a; 13:423–430. [PubMed: 20228804]
- Feng S, Jacobsen SE, Reik W. Epigenetic reprogramming in plant and animal development. *Science.* 2010b; 330:622–627. [PubMed: 21030646]
- Globisch D, Munzel M, Muller M, Michalakis S, Wagner M, Koch S, Bruckl T, Biel M, Carell T. Tissue distribution of 5-hydroxymethylcytosine and search for active demethylation intermediates. *PLoS One.* 2010; 5:e15367. [PubMed: 21203455]
- Guan JS, Su SC, Gao J, Joseph N, Xie Z, Zhou Y, Durak O, Zhang L, Zhu JJ, Clauser KR, et al. Cdk5 is required for memory function and hippocampal plasticity via the cAMP signaling pathway. *PLoS One.* 2011; 6:e25735. [PubMed: 21984943]
- Guo JU, Su Y, Zhong C, Ming GL, Song H. Hydroxylation of 5-methylcytosine by TET1 promotes active DNA demethylation in the adult brain. *Cell.* 2011; 145:423–434. [PubMed: 21496894]

- Haettig J, Stefanko DP, Multani ML, Figueroa DX, McQuown SC, Wood MA. HDAC inhibition modulates hippocampus-dependent long-term memory for object location in a CBP-dependent manner. *Learn Mem.* 2011; 18:71–79. [PubMed: 21224411]
- Hawk JD, Bookout AL, Poplawski SG, Bridi M, Rao AJ, Sulewski ME, Kroener BT, Manglesdorf DJ, Abel T. NR4A nuclear receptors support memory enhancement by histone deacetylase inhibitors. *J Clin Invest.* 2012; 122:3593–3602. [PubMed: 22996661]
- Huang Y, Chavez L, Chang X, Wang X, Pastor WA, Kang J, Zepeda-Martinez JA, Pape UJ, Jacobsen SE, Peters B, et al. Distinct roles of the methylcytosine oxidases Tet1 and Tet2 in mouse embryonic stem cells. *Proc Natl Acad Sci U S A.* 2014; 111:1361–1366. [PubMed: 24474761]
- Iurlaro M, Ficiz G, Oxley D, Raiber EA, Bachman M, Booth MJ, Andrews S, Balasubramanian S, Reik W. A screen for hydroxymethylcytosine and formylcytosine binding proteins suggests functions in transcription and chromatin regulation. *Genome Biol.* 2013; 14:R119. [PubMed: 24156278]
- Iyer LM, Tahiliani M, Rao A, Aravind L. Prediction of novel families of enzymes involved in oxidative and other complex modifications of bases in nucleic acids. *Cell Cycle.* 2009; 8:1698–1710. [PubMed: 19411852]
- Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat Genet.* 2003; 33(Suppl):245–254. [PubMed: 12610534]
- Jeltsch A, Jurkowska RZ. New concepts in DNA methylation. *Trends in biochemical sciences.* 2014; 39:310–318. [PubMed: 24947342]
- Kaas GA, Zhong C, Eason DE, Ross DL, Vachhani RV, Ming GL, King JR, Song H, Sweatt JD. TET1 controls CNS 5-methylcytosine hydroxylation, active DNA demethylation, gene transcription, and memory formation. *Neuron.* 2013; 79:1086–1093. [PubMed: 24050399]
- Khare T, Pai S, Koncevicus K, Pal M, Kriukiene E, Liutkeviciute Z, Irimia M, Jia P, Ptak C, Xia M, et al. 5-hmC in the brain is abundant in synaptic genes and shows differences at the exon-intron boundary. *Nature structural & molecular biology.* 2012; 19:1037–1043.
- Kriaucionis S, Heintz N. The nuclear DNA base 5-hydroxymethylcytosine is present in Purkinje neurons and the brain. *Science.* 2009; 324:929–930. [PubMed: 19372393]
- Ledoux JE. Coming to terms with fear. *Proc Natl Acad Sci U S A.* 2014; 111:2871–2878. [PubMed: 24501122]
- Lein ES, Zhao X, Gage FH. Defining a molecular atlas of the hippocampus using DNA microarrays and high-throughput in situ hybridization. *J Neurosci.* 2004; 24:3879–3889. [PubMed: 15084669]
- Lesburgueres E, Gobbo OL, Alaux-Cantin S, Hambucken A, Trifilieff P, Bontempi B. Early tagging of cortical networks is required for the formation of enduring associative memory. *Science.* 2011; 331:924–928. [PubMed: 21330548]
- Leslie JH, Nedivi E. Activity-regulated genes as mediators of neural circuit plasticity. *Prog Neurobiol.* 2011; 94:223–237. [PubMed: 21601615]
- Li X, Wei W, Zhao QY, Widagdo J, Baker-Andresen D, Flavell CR, D'Alessio A, Zhang Y, Bredy TW. Neocortical Tet3-mediated accumulation of 5-hydroxymethylcytosine promotes rapid behavioral adaptation. *Proc Natl Acad Sci U S A.* 2014; 111:7120–7125. [PubMed: 24757058]
- Lin Y, Bloodgood BL, Hauser JL, Lapan AD, Koon AC, Kim TK, Hu LS, Malik AN, Greenberg ME. Activity-dependent regulation of inhibitory synapse development by Npas4. *Nature.* 2008; 455:1198–1204. [PubMed: 18815592]
- Lister R, Mukamel EA, Nery JR, Urich M, Puddifoot CA, Johnson ND, Lucero J, Huang Y, Dwork AJ, Schultz MD, et al. Global epigenomic reconfiguration during mammalian brain development. *Science.* 2013; 341:1237905. [PubMed: 23828890]
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001; 25:402–408. [PubMed: 11846609]
- Lubin FD, Roth TL, Sweatt JD. Epigenetic regulation of BDNF gene transcription in the consolidation of fear memory. *J Neurosci.* 2008; 28:10576–10586. [PubMed: 18923034]
- Ma DK, Jang MH, Guo JU, Kitabatake Y, Chang ML, Pow-Anpongkul N, Flavell RA, Lu B, Ming GL, Song H. Neuronal activity-induced Gadd45b promotes epigenetic DNA demethylation and adult neurogenesis. *Science.* 2009; 323:1074–1077. [PubMed: 19119186]

- Meadows JP, Guzman-Karlsson MC, Phillips S, Holleman C, Posey JL, Day JJ, Hablitz JJ, Sweatt JD. DNA methylation regulates neuronal glutamatergic synaptic scaling. *Sci Signal*. 2015; 8:ra61. [PubMed: 26106219]
- Miller CA, Gavin CF, White JA, Parrish RR, Honasoge A, Yancey CR, Rivera IM, Rubio MD, Rumbaugh G, Sweatt JD. Cortical DNA methylation maintains remote memory. *Nat Neurosci*. 2010; 13:664–666. [PubMed: 20495557]
- Miller CA, Sweatt JD. Covalent modification of DNA regulates memory formation. *Neuron*. 2007; 53:857–869. [PubMed: 17359920]
- Monsey MS, Ota KT, Akingbade IF, Hong ES, Schafe GE. Epigenetic alterations are critical for fear memory consolidation and synaptic plasticity in the lateral amygdala. *PLoS One*. 2011; 6:e19958. [PubMed: 21625500]
- Munzel M, Globisch D, Bruckl T, Wagner M, Welzmler V, Michalakakis S, Muller M, Biel M, Carell T. Quantification of the sixth DNA base hydroxymethylcytosine in the brain. *Angew Chem Int Ed Engl*. 2010; 49:5375–5377. [PubMed: 20583021]
- Penn NW, Suwalski R, O'Riley C, Bojanowski K, Yura R. The presence of 5-hydroxymethylcytosine in animal deoxyribonucleic acid. *Biochem J*. 1972; 126:781–790. [PubMed: 4538516]
- Penner MR, Roth TL, Chawla MK, Hoang LT, Roth ED, Lubin FD, Sweatt JD, Worley PF, Barnes CA. Age-related changes in *Arc* transcription and DNA methylation within the hippocampus. *Neurobiol Aging*. 2011; 32:2198–2210. [PubMed: 20189687]
- Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*. 2001; 29:e45. [PubMed: 11328886]
- Rudenko A, Dawlaty MM, Seo J, Cheng AW, Meng J, Le T, Faull KF, Jaenisch R, Tsai LH. Tet1 is critical for neuronal activity-regulated gene expression and memory extinction. *Neuron*. 2013; 79:1109–1122. [PubMed: 24050401]
- Silverman JL, Yang M, Lord C, Crawley JN. Behavioural phenotyping assays for mouse models of autism. *Nat Rev Neurosci*. 2010; 11:490–502. [PubMed: 20559336]
- Spruijt CG, Gnerlich F, Smits AH, Pfaffeneder T, Jansen PW, Bauer C, Munzel M, Wagner M, Muller M, Khan F, et al. Dynamic readers for 5-(hydroxy)methylcytosine and its oxidized derivatives. *Cell*. 2013; 152:1146–1159. [PubMed: 23434322]
- Stefanko DP, Barrett RM, Ly AR, Reolon GK, Wood MA. Modulation of long-term memory for object recognition via HDAC inhibition. *Proc Natl Acad Sci U S A*. 2009; 106:9447–9452. [PubMed: 19470462]
- Sweatt JD. The emerging field of neuroepigenetics. *Neuron*. 2013; 80:624–632. [PubMed: 24183015]
- Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, Brudno Y, Agarwal S, Iyer LM, Liu DR, Aravind L, et al. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science*. 2009; 324:930–935. [PubMed: 19372391]
- Varley KE, Gertz J, Bowling KM, Parker SL, Reddy TE, Pauli-Behn F, Cross MK, Williams BA, Stamatoyannopoulos JA, Crawford GE, et al. Dynamic DNA methylation across diverse human cell lines and tissues. *Genome Res*. 2013; 23:555–567. [PubMed: 23325432]
- Vecsey CG, Hawk JD, Lattal KM, Stein JM, Fabian SA, Attner MA, Cabrera SM, McDonough CB, Brindle PK, Abel T, et al. Histone deacetylase inhibitors enhance memory and synaptic plasticity via CREB:CBP-dependent transcriptional activation. *J Neurosci*. 2007; 27:6128–6140. [PubMed: 17553985]
- West AE, Greenberg ME. Neuronal activity-regulated gene transcription in synapse development and cognitive function. *Cold Spring Harb Perspect Biol*. 2011; 3
- Wu H, Zhang Y. Reversing DNA Methylation: Mechanisms, Genomics, and Biological Functions. *Cell*. 2014; 156:45–68. [PubMed: 24439369]
- Xie W, Barr CL, Kim A, Yue F, Lee AY, Eubanks J, Dempster EL, Ren B. Base-resolution analyses of sequence and parent-of-origin dependent DNA methylation in the mouse genome. *Cell*. 2012; 148:816–831. [PubMed: 22341451]
- Yu DH, Ware C, Waterland RA, Zhang J, Chen MH, Gadkari M, Kunde-Ramamoorthy G, Nosavanh LM, Shen L. Developmentally programmed 3' CpG island methylation confers tissue- and cell-type-specific transcriptional activation. *Mol Cell Biol*. 2013; 33:1845–1858. [PubMed: 23459939]

- Yu H, Su Y, Shin J, Zhong C, Guo JU, Weng YL, Gao F, Geschwind DH, Coppola G, Ming GL, et al. Tet3 regulates synaptic transmission and homeostatic plasticity via DNA oxidation and repair. *Nat Neurosci.* 2015; 18:836–843. [PubMed: 25915473]
- Zhang RR, Cui QY, Murai K, Lim YC, Smith ZD, Jin S, Ye P, Rosa L, Lee YK, Wu HP, et al. Tet1 regulates adult hippocampal neurogenesis and cognition. *Cell Stem Cell.* 2013; 13:237–245. [PubMed: 23770080]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Tet1KO mice have normal brain morphology

(A), (B), and (C). Cresyl violet staining of the coronal sections (50 μ m) of Tet1+/+ and Tet1KO mice brain showing cerebrum (A), hippocampus (B), and cerebellum (C). (D) Absence of Tet1 expression in the different brain areas of Tet1KO mice was confirmed by quantitative real-time PCR (qRT-PCR) using WT and Tet1KO animals, bars represent the Tet1 mRNA levels relative to WT ($p^{****}<0.0001$, $n=6-8$ males/group), statistical comparisons were performed using an unpaired t-test (two tailed).




Figure 2. 5hmC levels are enriched in brain areas involved in active memory formation and storage, whereas 5mC levels are more or less uniformly distributed in different brain regions
 The quantification of the modified and unmodified cytosine bases was done by using highly sensitive LC-MS/MS-MRM techniques. (A) Standard curves for 5mC and 5hmC. The percentages of 5mC and 5hmC are plotted against the known ratios of methylated and hydroxymethylated DNA to the total amount of cytosine in the standard samples. (B) Percentage 5hmC relative to total cytosine in different brain areas; cortex (0.74%) and CA1 (0.70 %) have significantly higher 5hmC levels (p value < 0.0001) than DG (0.63%), CA3 (0.55%) and cerebellum (0.40%). (C) No significant difference ($p > 0.05$) was observed in the percentage 5mC relative to total cytosine in different brain regions. (D) Percentage unmodified cytosine (5C) of the total cytosine in different brain regions. ($n=4$ males, error bars \pm SEM), Statistical analysis was accomplished using one-way ANOVA with Bonferroni post hoc test.




Figure 3. Tet1 loss leads to a reduction in 5hmC levels

Representative graphs of percentage 5hmC, 5mC, and 5C levels in the areas CA1, DG, CA3, cortex and cerebellum, determined using LC-MS/MS-MRM. (A), (B), (C), (D), and (E) (left column), The levels of 5hmC in Tet1KO CA1, DG, CA3, cortex and cerebellum are significantly lower than the WT CA1, DG, CA3, cortex and cerebellum levels respectively ($p^* < 0.05$, $p^{**} < 0.005$, $p^{****} < 0.0001$). (A), (B), (C), (D), and (E) (center column), 5mC levels are significantly higher in Tet1KO cortex ($p^* < 0.05$), significantly lower in Tet1KO cerebellum ($p^* < 0.05$), and no significant differences were found between 5mC levels of the areas CA1, DG, and CA3 of the Tet1 KO and WT mice. (A), (B), (C), (D), and (E) (right column), 5C levels are slightly, but significantly higher in Tet1KO cerebellum ($p^{**} < 0.05$), and no significant differences were found between 5C levels of the areas CA1, DG, CA3, and cortex of the Tet1 KO versus WT mice. (n=4 males/group, error bars \pm SEM), statistical comparisons were performed using an unpaired t-test (two tailed).




Figure 4. Tet1 KO mice show normal baseline and social behavior

(A) Open field. No significant differences were found in the horizontal activity measured as distance travelled (left) and velocity (right) between WT and Tet1KO. (B) No significant differences were found between the WT and Tet1KO in terms of time spent in the center versus the time spent in periphery, taken as a measure of anxiety. (C) Elevated plus maze. No significant differences were found between WT and Tet1KO in the time spent in the open and closed arms and the number of entries made to each arm of the maze. (D) Both, WT and Tet1KO exhibited normal social interaction and preferred spending time with novel

mouse over novel object and central chamber. (E) No significant differences were found between the performance of the WT and Tet1KO in the accelerating rotarod test, left column shows the time to fall and right column represents the velocity at fall. (n=8 males/group, error bars \pm SEM),). Statistical comparisons between two groups were performed using an unpaired t-test (two tailed), statistical analysis between three or more groups was accomplished using one-way ANOVA with Bonferroni post hoc test.




Figure 5. Tet1KO has enhanced long term memory for contextual threat recognition training (fear conditioning)

(A) Schematic diagram of the contextual fear conditioning (B) Tet1 KO showed significantly higher freezing than WT in context testing after 24 hours ($p < 0.05$), 15 days ($p^{***} < 0.005$), and 30 days ($p^{***} < 0.0005$) of "light" training (0.5mA for 1 sec). (C), Tet1 KO showed significantly higher freezing than WT in context testing after 24 hours ($p^{**} < 0.005$), 15 days ($p^{**} < 0.005$), and 30 days ($p^{**} < 0.005$) of "strong" training (0.8mA for 2 sec). (D) and (E) No significant difference in freezing was observed between WT and Tet1KO in context testing after 24 hours of robust training (0.8 mA, 2sec., 3 times) and 1 hour of light training (0.5mA for 1 sec). (n=8 males/group, error bars \pm SEM), Statistical comparisons were performed using an unpaired t-test (two tailed).




Figure 6. Tet1KO has enhanced long term memory for cue threat recognition training (fear conditioning)

(A) Schematic diagram of Cue fear conditioning (B) and (C) Tet1KO showed significantly higher freezing ($p^* < 0.05$) in the cue testing after 24 hours of cue training using “light” (0.5 mA for 1 sec) and “strong” training (0.8 mA for 2 sec) protocol. (D) No significant difference ($p > 0.05$) in freezing was observed between WT and Tet1KO in cue testing after 24 hours of robust training (0.8 mA, 2 sec., 3 times). ($n = 8$ males/group, error bars \pm SEM). Statistical comparisons were performed using an unpaired t-test (two tailed).




Figure 7. Viral mediated reduction in hippocampal Tet1 mRNA enhances spatial memory (A) A representative image of AAV-mediated shRNA transgene expression; 14 days post injection. (B) Quantitative real time PCR analysis comparing the levels of Tet1 mRNA in mice 14 days following injection with either AAV-Tet1-shRNA or AAV-scrambled-shRNA control (n = 3–5/ group). (C) Global 5hmC and (D) global 5mC percentages relative to total cytosine content, as measured by HPLC/MS (n = 3 males/group). (E) Total distance traveled during 15 min in the open field. (F) The ratio of time spent in the center to time spent in the periphery of the open field, an indirect measure of anxiety. (G) Left, a schematic diagram of object location memory training. (H) Discrimination index for each group during training. (I) A schematic diagram of the object location memory test. (J) Discrimination index 24 hours after OLM training. For figures E–J, n = 8 for each group. Where applicable, NS = not statistically significant ($p \geq 0.05$), * $p < 0.05$, unpaired t test (two-tailed). All data are presented as \pm SEM.




Figure 8. Hippocampal long-term potentiation in Tet1KO mice and in mice with virally mediated Tet1 over-expression
 (A–B) (top), No significant differences were found in baseline synaptic transmission between WT and Tet1KO, as indicated in these plots representing input-output relationship of evoked fEPSP slope versus stimulus intensity. (C) (top), No significant differences were found in baseline synaptic transmission between AAV-Tet1, AAV-Tet1m and AAV-eYFP overexpressing mouse hippocampal brain slices, (AB) (middle), No significant differences were found in paired-pulse facilitation using different inter-event interval stimuli with normal and low intensity stimulation. (C) (middle), No significant differences were found in paired-pulse facilitation between AAV-Tet1, AAV-Tet1m and AAV-eYFP overexpressing mouse hippocampal brain slices, using normal amplitude stimulus intensity. (A–B) (bottom), No significant differences were found in long term potentiation between WT and Tet1KO, fEPSPs were recorded from area CA1 before and after tetanic stimuli, (100Hz, 1sec, 50% of the maximum slope) and (100Hz, 0.1sec, 25% of the maximum slope), of Schaffer collaterals. (C) (bottom), No significant differences were found in the long-term potentiation between AAV-Tet1, AAV-Tet1m and AAV-eYFP overexpressing mice hippocampal brain slices using theta-burst stimulation. (n=6 males/group, 7 slices each, error bars +/- SEM,). Statistical comparisons between two groups were performed using an unpaired t-test (two tailed); statistical analysis between three or more groups was accomplished using two-way ANOVA.




Figure 9. Tet1KO has altered expression of neuronal activity-regulated genes

The heatmap represents the qRT-PCR analysis of the mRNA transcripts. Color key represents relative gene expression (fold change) of the labeled gene in Tet1KO compared to the WT control. The “stars” represent the significance ($p^* < 0.05$, $p^{**} < 0.005$, $p^{***} < 0.0005$, $p^{****} < 0.0001$, $n=8$ males/group, error bars represented as \pm SEM). Statistical comparisons were performed using an unpaired t-test (two tailed).




Figure 10. Tet1KO has compensatory up-regulation of various active demethylation pathway genes

(A) Schematic presentation of the active DNA demethylation pathway in the adult CNS. (B) The heatmap represents the qRT-PCR analysis of the mRNA transcripts. Color key represents relative gene expression (fold change) of the labeled gene in Tet1KO compared to the WT control. The “stars” represent the significance ($p^* < 0.05$, $p^{**} < 0.005$, $p^{***} < 0.0005$, $p^{****} < 0.0001$, $n=8$ males/group, error bars represented as \pm SEM). Statistical comparisons were performed using an unpaired t-test (two tailed).




Figure 11. Genetic deletion of Tet1 leads to an increase in the expression of various epigenetic modifiers

The heatmap represents the qRT-PCR analysis of the mRNA transcripts. Color key represents relative gene expression (fold change) of the labeled gene in Tet1KO compared to the WT control. The “stars” represent the significance ($p^* < 0.05$, $p^{**} < 0.005$, $p^{***} < 0.0005$, $p^{****} < 0.0001$, $n=8$ males/group, error bars represented as \pm SEM). Statistical comparisons were performed using an unpaired t-test (two tailed).