



HHS Public Access

Author manuscript

Cancer Epidemiol Biomarkers Prev. Author manuscript; available in PMC 2016 December 01.

Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2015 December ; 24(12): 1898–1901. doi:
10.1158/1055-9965.EPI-15-0588.

Physical Activity and Risk of Male Breast Cancer

Hannah Arem¹, Louise A. Brinton¹, Steven C. Moore¹, Susan M. Gapstur², Laurel A. Habel³,
Kenneth Johnson⁴, Laurence N. Kolonel⁵, Valerie A. McCormack⁶, Karin B. Michels^{7,8,9},
Howard D. Sesso¹⁰, Giske Ursin¹¹, Stephen K. Van Den Eeden³, Elisabete
Weiderpass^{11,12,13,14}, Michael B. Cook^{1,*}, and Charles E. Matthews^{1,*}

¹National Cancer Institute, Division of Cancer Epidemiology and Genetics, Bethesda, MD
²American Cancer Society, Atlanta, GA ³Division of Research, Kaiser Permanente Northern
California, Oakland, CA ⁴Department of Epidemiology and Community Medicine, University of
Ottawa, Ottawa, Ontario, Canada ⁵Cancer Research Center of Hawaii, Honolulu, HI ⁶International
Agency for Research on Cancer, Lyon, France ⁷Obstetrics and Gynecology Epidemiology Center,
Department of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's
Hospital, Harvard Medical School, Boston, MA ⁸Department of Epidemiology, Harvard School of
Public Health, Boston, MA ⁹Channing Laboratory, Department of Medicine, Brigham and
Women's Hospital, Harvard Medical School, Boston, MA ¹⁰Divisions of Preventive Medicine and
Aging, Brigham and Women's Hospital, Boston, MA ¹¹Cancer Registry of Norway, Oslo, Norway
¹²Karolinska Institutet, Department of Medical Epidemiology and Biostatistics, Stockholm,
Sweden ¹³Department of Community Medicine, Faculty of Health Sciences, University of Tromsø,
The Arctic University of Norway, Tromsø, Norway ¹⁴Genetic Epidemiology Group, Folkhälsan
Research Center, Helsinki, Finland

Abstract

The association between leisure-time physical activity (LTPA) and male breast cancer risk (MBC) is unclear. In the MBC pooling project, with 449 cases and 13,855 matched controls, we used logistic regression with study stratification to generate adjusted odds ratios (OR) and 95% confidence intervals (CI) for LTPA tertiles and MBC risk. Compared with low LTPA, medium and high LTPA were not associated with MBC risk (OR=1.01, 95% CI 0.79–1.29; 0.90, 0.69–1.18, respectively). In joint-effects analyses, compared with the referent of high body mass index (BMI) (≥ 25 kg/m²)/low LTPA, neither medium nor high PA were associated with risk among high BMI men, but normal BMI men (<25 kg/m²) with low or medium LTPA were at a non-significant ~16% reduced risk and those with high LTPA were at a 27% reduced risk (OR=0.73, 95% CI 0.50–1.07). Physical activity alone may not confer protection against MBC risk.

Corresponding author: Hannah Arem, M.H.S., Ph.D., Aremhe2@mail.nih.gov, NCI Shady Grove, Division of Cancer Epidemiology and Genetics, 9609 Medical Center Drive Rm 6E314, MSC 9768, Bethesda, MD 20892-9768, phone: 240-276-7204 fax: 240-276-7837.

* Contributed equally to this manuscript

Conflicts of interest: none

Keywords

Male breast cancer; epidemiology; physical activity; pooled analysis

Introduction

Male breast cancer (MBC) is a rare disease, with a lifetime risk of 1 in 1,000. Due in part to the rarity of this cancer, relatively few studies have examined lifestyle-related etiologic factors. Leisure time physical activity (LTPA) has been consistently associated with a lower risk of postmenopausal female breast cancer(1, 2), but associations with MBC are unclear. Previous studies of physical activity and MBC risk suggest inverse—but not statistically significant—associations(3–6), yet were based on small case numbers and study characteristics or designs (proxy respondents(6), case-control(4, 5)) that may capture physical activity differently than a prospective design with self-reported data. In this analysis, with more cases and predominately prospectively collected data, we hypothesized that moderate to vigorous-intensity LTPA would be associated with a lower MBC risk.

Materials and Methods

Methods for the Male Breast Cancer Pooling Project (MBCPP) have been previously published(7). In short, the MBCPP identified all case-control or cohort studies with 10 cases using literature searches of PubMed, citations within published manuscripts, and advertisements at the National Cancer Institute Cohort Consortium meetings(7). Of the 21 studies identified for inclusion in the MBCPP, 10 had collected baseline information on LTPA (nine prospective nested case-control studies(8–16) and one retrospective case-control study(4)). The 2 cohort studies and 10 case-control studies that were not included in our analysis did not ask detailed information on LTPA and thus were not eligible for this analysis. Characteristics of included studies are presented in Supplementary Table S1. Cases were defined by the International Classification of Diseases, 10th Edition code C50 from cancer registries, medical record, or self-report. All studies contributed de-identified data following approved data sharing agreements, as well as National Cancer Institute and study center institutional review board clearances. Participants gave informed consent by nature of study participation.

LTPA was harmonized across studies into categories of low, medium and high. Studies with a single LTPA question on frequency of activity (AARP, JANUS, PHS, PLCO, Kaiser Permanente) ranged from four to six response levels. For example, in the PHS, the frequency question was “How often do you exercise vigorously enough to work up a sweat?”, with six categorical choices ranging from daily to rarely/never. In these studies we collapsed the categories to create a relatively even distribution for low, medium and high. For studies with line items for more than one type of activity (Canada, CPS II-NC, EPIC, HPFS, MEC), we assigned metabolic equivalent hours per week, using intensity values previously assigned by individual studies and by referencing the updated Ainsworth Compendium of Physical Activities (17). Example activities from HPFS included separate line items with 10 categorical responses ranging from none to 11+ h/wk for walking

outdoors, jogging, running, bicycling, lap swimming, tennis, squash, or calisthenics/rowing. We then divided the study-specific distributions into categories of low (<33rd percentile), medium (33–<66th percentile), and high (≥66th percentile) LTPA.

We used unconditional logistic regression with stratification by study to compute odds ratios (OR) and 95% confidence intervals (95% CI). We first created models adjusted for age, and then additionally adjusted for race, education, marital status, diabetes history, alcohol, smoking status, and body mass index (BMI). We created missing categories for race, education, marital status, diabetes, and alcohol, as these covariates were missing for 5% of study subjects, and for alcohol, for which 13% were missing. Sensitivity analyses excluding observations with missing data yielded unaltered results. To assess influence by individual studies, we also performed analyses excluding one study at a time.

Because BMI was previously associated with MBC risk in this population (7), and because joint effects of BMI and physical activity have been observed for female breast cancer risk, we created six categories to look at joint effects: high BMI (≥25 kg/m²) with 1) low (referent), 2) medium or 3) high activity levels, and normal BMI (<25 kg/m²) with 4) low, 5) medium or 6) high activity levels. We also stratified analyses by age, BMI, diabetes, smoking and family history of breast cancer and assessed interactions using the Wald test. All analyses were performed using SAS V9.3 (SAS Institute Inc, Cary, NC). P-values <0.05 were considered significant.

Results

This pooled analysis of 10 studies included 449 cases and 13,855 controls. More active men were more likely to be non-Hispanic white, college graduates, have a lower BMI, and were less likely to be current smokers or report diabetes (Table 1).

Compared with men reporting low LTPA, men reporting medium and high LTPA levels did not have a lower MBC risk after adjustment for other risk factors (OR=1.01, 95% CI 0.79–1.29; 0.90, 0.69–1.18, respectively) (Table 2). Results were similar after excluding one study at a time to test undue influence by an individual study (Supplementary Table S2).

In joint effects analyses of BMI and LTPA we found that among high BMI men (≥25 kg/m²) neither medium nor high LTPA were associated with risk (OR=1.01, 95% CI 0.76–1.34; 0.90, 0.66–1.24, respectively) compared with the referent group of low LTPA/high BMI; However, non-significant ~16% lower risks were observed among normal BMI men (<25 kg/m²) who reported either low or medium LTPA (OR=0.83, 95% CI 0.58–1.19; 0.84, 0.59–1.18, respectively), and a non-significant 27% lower risk was observed among men who had a normal BMI/ high LTPA (OR=0.73, 95% CI 0.50–1.07) (Table 2).

Stratified analyses did not suggest statistically significant interactions by median age, BMI, diabetes, or family history of breast cancer (all $p_{\text{interaction}}$ -values ≥0.05; Supplementary Table S3). There was a stronger inverse association between LTPA and MBC among current smokers (OR=0.25, 95% CI 0.07–0.85) than among never or former smokers, with the $p_{\text{interaction}}$ showing borderline significance ($p=0.05$).

Discussion

Previous studies of LTPA and female breast cancer risk have suggested stronger risk reductions of about 25% among postmenopausal normal weight women compared to no association for obese women(2), which is similar to the magnitude of association we found among active men with a normal BMI.

Published MBC studies have shown non-significant inverse associations with LTPA, but were based on limited numbers (range=81–178)(3–6) and faced challenges of collecting PA data from proxy respondents of deceased men(6), and retrospective (case-control designs), which may introduce recall bias(4, 5). Reasons for a possible interaction of LTPA with BMI remain unknown, but may reflect the cumulative effect of reduced estrogen exposure for lean active men, supported by other findings from the MBCPP of elevated risks being associated with both high BMI(7) and higher estradiol(18).

Strengths of this study include the relatively large sample size, prospective data collection in nine of the ten studies, and use of studies with sufficiently detailed information on LTPA. Limitations of this analysis include the inability to stratify by genetic MBC risk factors such as Klinefelter's syndrome or BRCA status, as this information was not available. We also had LTPA and BMI data only from baseline, which does not account for changes in these factors over time. We captured only leisure-time activity in this study, and were unable to examine occupational activity or sedentary time, each of which may affect risk. In summary, our findings suggest that physical activity alone may not be protective for MBC risk.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

Financial support: H. Arem, L.A. Brinton, S.C. Moore, M.B. Cook, C.E. Matthews are supported by the intramural research program at the National Institutes of Health, Bethesda, Maryland. The Physicians' Health Study was supported by grants CA 097193, CA 34944, CA 40360, HL 26490, and HL 34595 from the National Institutes of Health (H.D. Sesso). The Health Professionals' Follow-Up Study was supported by research grant CA167552 from the National Cancer Institute, National Institutes of Health (K.B. Michels). Support for MEC was provided by NIH/NCI under grant number R37CA54281 (L.N. Kolonel). The American Cancer Society funds the creation, maintenance, and updating of the Cancer Prevention Study II cohort (S.M. Gapstur).

References

1. Monninkhof EM, Elias SG, Vlems FA, van der Tweel I, Schuit AJ, Voskuil DW, et al. Physical activity and breast cancer: a systematic review. *Epidemiology*. 2007; 18:137–157. [PubMed: 17130685]
2. Friedenreich CM, Cust AE. Physical activity and breast cancer risk: impact of timing, type and dose of activity and population subgroup effects. *British journal of sports medicine*. 2008; 42:636–647. [PubMed: 18487249]
3. Brinton LA, Richesson DA, Gierach GL, Lacey JV, Park Y, Hollenbeck AR, et al. Prospective evaluation of risk factors for male breast cancer. *Journal of the National Cancer Institute*. 2008; 100:1477–1481. [PubMed: 18840816]
4. Johnson K, Pan S, Mao Y. Group CCRER. Risk factors for male breast cancer in Canada, 1994–1998. *European journal of cancer prevention*. 2002; 11:253–263. [PubMed: 12131659]

5. Ewertz M, Holmberg L, Tretli S, Pedersen BV, Kristensen A. Risk Factors for Male Breast Cancer? A Case-Control Study from Scandinavia. *Acta Oncologica*. 2001; 40:467–471. [PubMed: 11504305]
6. Hsing AW, McLaughlin JK, Cocco P, Chien HTC, Fraumeni JF Jr. Risk factors for male breast cancer (United States). *Cancer Causes & Control*. 1998; 9:269–275. [PubMed: 9684707]
7. Brinton LA, Cook MB, McCormack V, Johnson KC, Olsson H, Casagrande JT, et al. Anthropometric and hormonal risk factors for male breast cancer: male breast cancer pooling project results. *Journal of the National Cancer Institute*. 2014; 106:djt465. [PubMed: 24552677]
8. Schatzkin A, Subar AF, Thompson FE, Harlan LC, Tangrea J, Hollenbeck AR, et al. Design and Serendipity in Establishing a Large Cohort with Wide Dietary Intake Distributions The National Institutes of Health–American Association of Retired Persons Diet and Health Study. *American journal of epidemiology*. 2001; 154:1119–1125. [PubMed: 11744517]
9. Calle EE, Rodriguez C, Jacobs EJ, Almon ML, Chao A, McCullough ML, et al. The American cancer society cancer prevention study II nutrition cohort. *Cancer*. 2002; 94:2490–2501. [PubMed: 12015775]
10. Riboli E, Hunt K, Slimani N, Ferrari P, Norat T, Fahey M, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public health nutrition*. 2002; 5:1113–1124. [PubMed: 12639222]
11. Giovannucci E, Rimm EB, Liu Y, Leitzmann M, Wu K, Stampfer MJ, et al. Body mass index and risk of prostate cancer in US health professionals. *Journal of the National Cancer Institute*. 2003; 95:1240–1244. [PubMed: 12928350]
12. Cutler JL, Ramcharan S, Feldman R, Siegelaub A, Campbell B, Friedman GD, et al. 1. Methods and population. *Preventive medicine*. 1973; 2:197–206. [PubMed: 4723635]
13. Sesso HD, Gaziano JM, VanDenburgh M, Hennekens CH, Glynn RJ, Buring JE. Comparison of baseline characteristics and mortality experience of participants and nonparticipants in a randomized clinical trial: the Physicians' Health Study. *Controlled clinical trials*. 2002; 23:686–702. [PubMed: 12505246]
14. Prorok PC, Andriole GL, Bresalier RS, Buys SS, Chia D, Crawford ED, et al. Design of the prostate, lung, colorectal and ovarian (PLCO) cancer screening trial. *Controlled clinical trials*. 2000; 21:273S–309S. [PubMed: 11189684]
15. Jellum E, Andersen A, Lund-Larsen P, Theodorsen L, Orjasaeter H. The JANUS serum bank. *Science of the total environment*. 1993; 139:527–535. [PubMed: 8272856]
16. Kolonel LN, Henderson BE, Hankin JH, Nomura AM, Wilkens LR, Pike MC, et al. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. *American journal of epidemiology*. 2000; 151:346–357. [PubMed: 10695593]
17. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR Jr, Tudor-Locke C, et al. 2011 Compendium of Physical Activities: a second update of codes and MET values. *Medicine and science in sports and exercise*. 2011; 43:1575–1581. [PubMed: 21681120]
18. Brinton LA, Key TJ, Kolonel LN, Michels KB, Sesso HD, Ursin G, et al. Prediagnostic Sex Steroid Hormones in Relation to Male Breast Cancer Risk. *Journal of Clinical Oncology*. 2015 In Press.

Table 1

Baseline characteristics of the pooled study populations of male breast cancer cases (n=449) and controls (n=13,855)

Characteristic	Physical activity tertile ^a		
	Low	Medium	High
Cases/controls	157/3884	177/5757	115/4214
Age at entry (years), mean (SD) ^{b, c}	61.0 (7.9)	61.7 (7.6)	62.1 (7.7)
Race, n (%) white ^b	3499 (86.8)	5293 (89.9)	3882 (90.2)
Married, n (%)	3316 (85.3)	5064 (87.3)	3625 (87.5)
Body mass index (kg/m ²), mean (SD)	27.3 (4.4)	26.7 (3.8)	26.0 (3.3)
Smoking status, n (%)			
<i>Never</i>	1269 (32.3)	2006 (34.9)	1479 (35.2)
<i>Former</i>	1990 (50.7)	3131 (54.4)	2352 (56.0)
<i>Current</i>	668 (17.0)	615 (10.7)	372 (8.9)
Diabetes, n (%)			
<i>No</i>	3333 (91.0)	5280 (91.7)	3832 (93.0)
<i>Yes</i>	328 (8.9)	480 (8.3)	288 (7.0)
Alcohol (grams/day), mean (SD)	17.3 (40.6)	15.4 (33.7)	15.7 (30.3)
Educational level, n (%)			
<i>Less than high school</i>	683 (17.3)	457 (7.9)	389 (9.2)
<i>High school graduate</i>	625 (15.8)	803 (13.8)	650 (15.3)
<i>Some college/vocational school</i>	914 (23.1)	1485 (25.5)	985 (23.2)
<i>College graduate</i>	1728 (43.8)	3069 (52.8)	2229 (52.4)

^aPhysical activity tertiles were created from study specific distributions and then pooled.

^bCohort control subjects were incidence-density matched to cases on sex, race, study center, date of birth, date of entry, and exit date.

^cCase-control subjects were frequency matched on age, and were sampled from provincial health insurance plans in Canada (coverage of >95% of Canadians).

Table 2

Odds ratios and 95% confidence intervals for physical activity and risk of male breast cancer (n=449 cases/13855 controls)

Physical Activity level	Low	Medium	High	p-trend
N (case/control)	157/3884	177/5757	115/4214	
<i>Model 1^a</i>	1.00	1.04 (0.81–1.32)	0.92 (0.70–1.19)	0.515
<i>Model 2^b</i>	1.00	1.00 (0.79–1.28)	0.88 (0.68–1.15)	0.362
<i>Model 3^c</i>	1.00	1.01 (0.79–1.29)	0.90 (0.69–1.18)	0.443
<i>Joint effects^b</i>				
<i>High BMI (>25 kg/m²)</i>	1.00	1.01 (0.79–1.34)	0.90 (0.66–1.24)	
<i>Normal BMI (<25 kg/m²)</i>	0.83 (0.58–1.19)	0.84 (0.59–1.18)	0.73 (0.50–1.07)	

^aModel 1 included stratification by cohort and was adjusted for age.

^bModel 2 was additionally adjusted for race (white, black, other), educational level (high school or less, some college, completed college), marital status (married, not married), history of diabetes (yes, no), alcohol consumption (none, 1–2 drinks/day, >2 drinks/day), and smoking status (never, former, current).

^cModel 3 was additionally adjusted for body mass index (<18.5, 18.5–<25, 25–<30, 30+ kg/m²).