

CORRECTION

Correction: Severity of Old World Cutaneous Leishmaniasis Is Influenced by Previous Exposure to Sandfly Bites in Saudi Arabia

Karina Mondragon-Shem, Waleed S. Al-Salem, Louise Kelly-Hope, Maha Abdeladhim, Mohammed H. Al-Zahrani, Jesus G. Valenzuela, Alvaro Acosta-Serrano

There is an error in the third sentence of the “Detection of human anti-PpSP32 antibodies” sub-section of the Methods section. The concentration of PpSP32 should read 2 ug/ml = 0.2 ug/well, not 2 mg/ml = 0.1 mg/well.

Reference

1. Mondragon-Shem K, Al-Salem WS, Kelly-Hope L, Abdeladhim M, Al-Zahrani MH, Valenzuela JG, et al. (2015) Severity of Old World Cutaneous Leishmaniasis Is Influenced by Previous Exposure to Sandfly Bites in Saudi Arabia. PLoS Negl Trop Dis 9(2): e0003449. doi:[10.1371/journal.pntd.0003449](https://doi.org/10.1371/journal.pntd.0003449) PMID: [25646796](https://pubmed.ncbi.nlm.nih.gov/25646796/)



OPEN ACCESS

Citation: Mondragon-Shem K, Al-Salem WS, Kelly-Hope L, Abdeladhim M, Al-Zahrani MH, Valenzuela JG, et al. (2015) Correction: Severity of Old World Cutaneous Leishmaniasis Is Influenced by Previous Exposure to Sandfly Bites in Saudi Arabia. PLoS Negl Trop Dis 9(12): e0004294. doi:[10.1371/journal.pntd.0004294](https://doi.org/10.1371/journal.pntd.0004294)

Published: December 8, 2015

Copyright: © 2015 Mondragon-Shem et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.