

Elimination of routine chest radiographs following off-pump coronary artery bypass surgery: A randomized controlled trial study

Seyed Khalil Forouzannia, Ali Sarvi¹, Mohammadtaghi Sarebanhassanabadi², Reza Nafisi-Moghadam³

Tehran Heart Center, Tehran University of Medical Sciences, Tehran, ¹Department of Medicine, Islamic Azad University, Yazd Branch, ²Yazd Cardiovascular Research Center, ³Department of Radiology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Abstract

Background: Post cardiac surgery routine chest radiographs (CXRs), ordered without any clinical and laboratory indications, is a standard obligatory practice in many cardiothoracic centers. Routine CXRs incur cost, manpower, and radiation. The objective of this study is to assess early outcome in off-pump coronary artery bypass (OPCAB) patients with postoperative routine versus clinically indicated CXR protocols.

Materials and Methods: This study is a randomized clinical trial conducted on 231 OPCAB candidates in Afshar Cardiac Center, Yazd, Iran. Patients were categorized into two groups. All 118 patients in group A had routine postoperative CXRs. The 113 patients in group B were selectively exposed to CXR only on clinical indications. All patients were postoperatively followed up for 30 days. Data gathered from both groups were statistically analyzed.

Results: Routine postoperative CXRs obtained in 118 OPCAB group A candidates showed abnormal findings in 20 patients that did not require new intervention. One month follow-up of these patients showed no complications. In 113 OPCAB candidates of group B, 7 on-demand CXRs were obtained on clinical evaluation that required added intervention. In a 1-month follow-up of this group, five patients presented with symptomatic complaints. On re-examination, none needed readmission, intervention, or paraclinical evaluation. No complications were observed due to CXR elimination.

Conclusion: The study suggests that postoperative CXR selected on clinical grounds in place of routine CXR does not change early postoperative outcome of OPCAB procedure.

Key Words: Chest radiograph, off-pump coronary artery bypass, outcome

Address for correspondence:

Dr. Ali Sarvi, Islamic Azad University, Yazd Branch, Medical College, Yazd, Iran. E-mail: A.Sarvi.md@gmail.com

Received: 26.05.2014, Accepted: 09.09.2014

INTRODUCTION

Coronary artery bypass grafting (CABG), a landmark

procedure in cardiovascular history, is today a leading cardiac operation.^[1] The time-honored on-pump CABG has often been replaced by an equally effective, alternative, comparatively economical off-pump coronary artery bypass (OPCAB) procedure particularly performed in patients with co-morbid conditions including renal impairment and increased risk of cardiovascular accident.^[2] Pre- and postoperative orders for a frequently practiced procedure may need revision.

Postoperative routine chest radiograph (CXR) is a conventional protocol, practiced in most cardiothoracic

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.167966

Copyright: © 2015 Forouzannia. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: Forouzannia SK, Sarvi A, Sarebanhassanabadi M, Nafisi-Moghadam R. Elimination of routine chest radiographs following off-pump coronary artery bypass surgery: A randomized controlled trial study. *Adv Biomed Res* 2015;4:236.

centers, undertaken without any clinical or laboratory indications. Although postoperative CXR has been broadly evaluated in cardiac surgeries, in general, however, different cardiac procedures merit selective postoperative protocols as in OPCAB. Clinical cardiopulmonary findings are not always revealed on CXR.^[3] Its accuracy in visualizing and defining etiology of pulmonary opacity is moderate.^[4] Not all abnormal cardiopulmonary CXR findings after OPCAB are unexpected or require change of management.^[5,6] In addition, cost, manpower resources, radiation, and manipulation of the critically ill patients that may induce transient hemodynamic instability and endotracheal tube (ETT) and vascular catheter displacement make routine CXR debatable.^[6,7]

MATERIALS AND METHODS

This clinical trial was conducted at Afshar Cardiac Center, a university hospital in Yazd, Iran. The study was approved by the Hospital Research Ethics Committee. A total of 231 patients over a period of 6 months from March to September 2011 were subjected to the OPCAB procedure. Patients were randomized into two groups using simple randomization: Group A included 118 patients and group B 113 patients. Routine CXR was obtained for all 118 patients in group A. On-demand CXR was advised for selective patients in group B. All patients were carefully monitored postoperatively for cardiopulmonary complications. Hemodynamics, pulse oximetry, arterial blood gas analysis, frequent auscultation of lungs and control of chest tube drains, and position of ETT and vascular lines were among the critical care protocol.

Routine CXR was obtained without any clinical indication. On-demand CXR was obtained on clinical findings, either to confirm or clarify clinical diagnosis. Selection on clinical evaluation was based on findings of fever, dyspnea, abnormal pulmonary sounds, signs and symptoms of cardiac tamponade, lower extremity edema, abnormal chest tube bleed or air leak, and doubtful position of ETT and vascular lines. Final clinical evaluation for on-demand CXR rested on the cardiovascular surgeon and X-ray findings were reported by the radiologist. All patients were discharged 5-7 days after surgery. A 30-day follow-up included at least two visits. Patients were in constant contact with the cardiac clinic. Symptomatic patients were selectively re-examined to rule out complications.

A detailed questionnaire was completed for each patient. Clinical indication resulting in on-demand CXR, X-ray findings, and new intervention were

recorded. Data obtained from both groups were statistically analyzed by SPSS 11.5 (SPSS Inc, Chicago, IL) using Chi-square and Fisher's exact test. *P* value of 0.05 was considered statistically significant [Figure 1].

RESULTS

A total of 231 OPCAB candidates including 151 males (65.4%) and 80 females (34.6%) with ages ranging from 31 to 83 years (average 60.2) were categorized into two groups. Although efforts were made to balance the two groups, age and tobacco smoke exposure were greater and statistically significant in group B. These variables were not critical for the study. However, no significant differences were observed between preoperative variables of gender, diabetes mellitus, hypertension, hyperlipidemia, number of coronary vessels involved and, in particular, left main coronary artery (LMCA) lesion [Table 1].

All 118 patients of group A obtained CXR as a routine protocol. Twenty abnormal CXRs were observed including five atelectasis, five pulmonary opacities, nine mild pleural effusion, and one abnormal position of diaphragm. The remaining 98 group A patients had normal CXRs. None of the 20 patients with positive CXR findings required new intervention.




Figure 1: The flowchart of study groups

On-demand CXRs were obtained in seven group B patients on clinical grounds. These included three patients with abnormal pulmonary sounds, three patients with chest tube air leak, and one case with subcutaneous emphysema. Positive X-ray findings included pneumothorax, hydropneumothorax, massive pleural effusion, lobar atelectasis, and pulmonary congestion, clarifying or confirming clinical findings in each case. All seven on-demand CXRs prompted change in management. There was no statistically significant relationship between preoperative variables and indication for CXR in group B patients [Table 2].

A 30-day follow-up was undertaken for both groups. Group A had a normal follow-up. In group B, 5 patients presented with symptomatic complaints including three cases of dyspnea, one with fever and rigor, and one with chest pain. On re-examination, none needed readmission, intervention, or paraclinical evaluation. No complications were observed due to CXR elimination.

DISCUSSION

Routine postoperative CXR is a standard of care in many centers all over the world. However, the protocol has been often questioned by some authors. These include Roa *et al.*, Silverstein *et al.*, O'Brien *et al.*, and of recent, Ganapathy *et al.* and Tolsma *et al.*, who favor the elimination of routine CXR.^[8-14]

Most current literatures refer to routine CXR evaluation after cardiac surgeries in general. Protocols in elective cardiac operations differ from emergency post-traumatic cardiac operations wherein postoperative CXR would be mandatory. Accompanying trivial lung injuries in post-traumatic cases are often missed and may flare up in course of time, emphasizing postoperative screening with CXR.^[15]

Likewise, clinical settings may influence the rule of CXR. Pediatric examination, particularly in ICU, is more complex and less informative.^[16] CXR in these settings mostly reveal malposition of device. The effect of a minor intervention, performed on CXR findings, can be significant in the critically ill children.^[17] Also, each elective cardiac surgery would require a selective postoperative protocol. This is a study for routine and clinically indicated postoperative CXR in exclusively OPCAB candidates with regards to early postoperative outcome.

Findings on routine CXR may not always influence management. Several postoperative

Table 1: Preoperative variables in groups A and B

Variables	Group A (n=118)	Group B (n=113)	P value
	M:F=75:43	M:F=76:37	
DM			
Yes	37 (31.4)	38 (33.6)	0.712
No	81 (68.6)	75 (66.4)	
HTN			
Yes	56 (47.5)	52 (46)	0.826
No	62 (52.5)	61 (54)	
HLP			
Yes	42 (35.6)	40 (35.4)	0.975
No	76 (64.4)	73 (64.6)	
Smoking			
Yes	29 (24.6)	47 (41.6)	0.006
No	89 (75.4)	66 (58.4)	
Involved vessels			
0	0 (0)	1 (0.9)	0.409
1	9 (7.6)	4 (3.5)	
2	31 (26.3)	29 (25.7)	
3	78 (66.1)	79 (69.9)	
LMCA involvement			
Yes	3 (2.5)	3 (2.7)	1.000
No	115 (97.5)	110 (97.3)	

DM: Diabetes mellitus, HTN: Hypertension, HLP: Hyperlipidemia, LMCA: Left main coronary artery

Table 2: Relationship between preoperative variables and indication for CXR in group B

Variables	Indication for CXR		P value
	No (n=106)	Yes (n=7)	
	M:F=70:36	M:F=6:1	
Age, years			
31-54	26 (89.7)	3 (10.3)	0.476
55-64	28 (93.3)	2 (6.7)	
65-89	52 (96.3)	2 (3.7)	
Risk factors			
No	35 (94.6)	2 (5.4)	1.000
Yes	71 (93.4)	5 (6.6)	
Smoking			
No	64 (97)	2 (3)	0.125
Yes	42 (89.4)	5 (10.6)	
Involved vessels			
0	1 (100)	0 (0)	1.000
1	4 (100)	0 (0)	
2	27 (93.1)	2 (6.9)	
3	74 (93.7)	5 (6.3)	
LMCA involvement			
No	103 (93.6)	7 (6.4)	1.000
Yes	3 (100)	0 (0)	

Risk factors are diabetes mellitus, hyperlipidemia, and hypertension. LMCA: Left main coronary artery

radiological findings, including minor atelectasis and minimal-to-moderate pleural effusion, usually resolve spontaneously.^[5,6] In our study, all 20 abnormal routine CXR findings in group A did not prompt new intervention. On the other hand, clinically it indicated on-demand CXR changed management status.

In contrast to group A candidates, 106 patients in group B who were closely clinically followed did not require CXR, thus saving on cost, manpower, and avoiding unnecessary maneuver and cumulative radiation, both to attending staff and patients.^[6,7] No differences were observed in 30-day postoperative outcome between the both groups.

The use of CXR to confirm position of ETT and vascular lines remains controversial. O'Brien and colleagues demonstrated that clinical evaluation of ETT proved to be more sensitive than CXR.^[12] Palesty *et al.* and Gray *et al.* concluded that CXR was unnecessary for vascular line position,^[18,19] however, Gladwin *et al.* had contradictory thoughts, emphasizing the need for CXR.^[20]

Post cardiac surgery echocardiography in cardiac tamponade is a more reliable diagnostic tool than CXR. Since the pericardium is left open after surgery, the cardiac silhouette stands abnormal on CXR. A series of CXRs would be needed to detect changes in the abnormal cardiac shadow. In addition, cardiac dilatation cannot be ruled out from pericardial effusion. In contrast, even a 20-50 ml pericardial effusion is detected by echocardiography.^[21] Likewise, sonography more accurately detects diaphragmatic motility than CXR.^[22] Routine postoperative CXR shows limited reliability in these complications. The application of a more sensitive alternative method for evaluation in these situations is recommended.

CONCLUSION

Clinically indicated CXR in place of routine CXR after OPCAB procedure significantly reduced the number of CXRs without affecting the length of hospital stay, readmission rate, and mortality rate. Thus, it seems that postoperative on-demand CXR on clinical evaluation is an efficacious, cost-effective, and comparatively safe protocol than routine CXR.

In order to eliminate limitations of this study, greater number of patients sampling, evaluation at different cardiothoracic centers, and a postoperative follow-up beyond 30 days are recommended for future studies.

ACKNOWLEDGMENT

We would like to express our appreciation to the nursing staff of Afshar Cardiac Center for their invaluable assistance during the study. This study is an extract from a thesis.

REFERENCES

1. Shekar PS. On-pump and off-pump coronary artery bypass grafting. *Circulation* 2006;113:e51-2.

2. Nwaejike N, Mansha M, Bonde P, Campalani G. Myocardial revascularization by off-pump coronary bypass surgery (OPCABG): A ten year review. *Ulster Med J* 2008;77:106-9.
3. Wiener-Kronish JP. Postoperative pleural and pulmonary abnormalities in patients undergoing coronary artery bypass grafts. *Chest* 1992;102:1313-4.
4. Rubenfield GD, Caldwell E, Granton J, Hudson LD, Matthay MA. Interobserver variability in applying a radiographic definition for ARDS. *Chest* 1999;116:1347-53.
5. Labidi M, Baillet R, Dionne B, Lacasse Y, Maltais F, Boulet LP. Pleural effusions following cardiac surgery: Prevalence, risk factors, and clinical features. *Chest* 2009;136:1604-11.
6. Roa PS, Abid Q, Khan KJ, Meikle RJ, Natarajan KM, Morritt GN, *et al.* Evaluation of routine postoperative chest X-rays in the management of cardiac surgical patient. *Eur J Cardiothorac Surg* 1997;12:724-9.
7. Chahine-Malus N, Stewart T, Lapinsky SE, Marras T, Dancy D, Leung R, *et al.* Utility of routine chest radiographs in a medical-surgical intensive care unit: A quality assurance survey. *Crit Care* 2001;5:271-5.
8. Chong JL, Pillai R, Fisher A, Grebenik C, Sinclair M, Westaby S. Cardiac surgery: Moving away from intensive care. *Br Heart J* 1992;68:430-3.
9. Hornick PI, Harris P, Cousins C, Taylor KM, Keogh BE. Assessment of the value of the immediate postoperative chest radiograph after cardiac operation. *Ann Thorac Surg* 1995;59:1150-4.
10. Stevens JJ, Booth JV, Little J, Gopinath S, Ghosh S, Kneeshaw JD, *et al.* The value of immediate postoperative chest radiography in adult cardiac surgery. *Br J Anaesth* 1995;A110:56.
11. Silverstein DS, Livingston DH, Elcavage J, Kovar L, Kelly KM. The utility of routine chest radiography in the surgical intensive care unit. *J Trauma* 1993;35:643-6.
12. O'Brien W, Karski J, Cheng D, Sandler A. Does the immediate postoperative chest x-ray help in the clinical management of patients following cardiac surgery. *Can J Anaesth* 1995;42:A3-76.
13. Tolsma M, Rijpstra TA, Schultz MJ, Mulder PG, van der Meer NJ. Significant changes in the practice of chest radiography in Dutch intensive care units: A web-based survey. *Ann Intensive Care* 2014;4:10.
14. Ganapathy A, Adhikari NK, Spiegelman J, Scales DC. Routine chest x-rays in intensive care units: A systemic review and meta-analysis. *Crit Care* 2012;16:R68.
15. Ahangar AG, Dar AM, Mir IA, Bhat MA, Lone GN, Shah SH. Role of routine chest roentgenography in patients with open heart operation. *J K Science* 2003;5:104-6.
16. Hauser GJ, Pollack MM, Sivit CJ, Taylor GA, Balas DI, Guion CJ. Routine chest radiographs in pediatric intensive care: A prospective study. *Pediatrics* 1989;83:465-70.
17. Quasney MW, Goodman DM, Billow M, Chiu H, Easterling L, Frankel L, *et al.* Routine chest radiographs in pediatric intensive care units. *Pediatrics* 2001;107:241-8.
18. Palesty JA, Amshel CE, Dudrick SJ. Routine chest radiographs following central venous recatheterization over a wire are not justified. *Am J Surg* 1998;176:618-21.
19. Gray P, Sullivan G, Ostryzniuk P, McEwen TA, Rigby M, Roberts DE. Value of postprocedural chest radiographs in adult intensive care unit. *Crit Care Med* 1992;20:1513-8.
20. Gladwin MT, Slonim A, Landucci DL, Gutierrez DC, Cunnion RE. Canulation of the internal jugular vein: Is postprocedural chest radiography always necessary? *Crit Care Med* 1999;27:1819-23.
21. Hamid M, Khan MU, Bashour AC. Diagnostic value of chest x-ray and echocardiography for cardiac tamponade in post cardiac surgery patients. *J Pak Med Assoc* 2006;56:104-7.
22. Fedulla AJ, Lerner RM, Gibson J, Shayne DS. Sonographic measurement of diaphragmatic motion after coronary artery bypass surgery. *Chest* 1992;102:1683-6.

Source of Support: Nil, Conflict of Interest: None declared.