



Published in final edited form as:

J Sch Health. 2016 January ; 86(1): 61–71. doi:10.1111/josh.12353.

Health risk behaviors in a representative sample of bisexual and heterosexual female high school students in Massachusetts

Jaclyn M. White, MPH,

Research Analyst, The Fenway Institute, Fenway Health, Doctoral Student, Yale School of Public Health, Department of Chronic Disease Epidemiology, 1340 Boylston Street, 8th Floor, Boston, MA 02215, Tel: 508-340-7715, Fax: 617-267-0764

Katie B. Biello, PhD, MPH,

Research Scientist, The Fenway Institute, Fenway Health, Post-Doctoral Fellow, Harvard School of Public Health, Department of Epidemiology, 1340 Boylston Street, 8th Floor, Boston, MA 02215

Sari L. Reisner, ScD, MA,

Research Scientist, The Fenway Institute, Fenway Health, Post-Doctoral Fellow, Harvard School of Public Health, Department of Epidemiology, 1340 Boylston Street, 8th Floor, Boston, MA 02215

Amaya Perez-Brumer, MSc,

Doctoral Student, Columbia University, Mailman School of Public Health, Department of Sociomedical Sciences, 722 West 168th Street, New York, NY 10032, Tel: 719-330-7817

Katherine J. Heflin, and

Master's Student, Harvard School of Public Health, Department of Health Policy and Management, 4904 Glen Cove Pkwy, Bethesda, MD 20816, Tel: 785-383-4950

Matthew J. Mimiaga, ScD, MPH

Research Scientist, The Fenway Institute, Fenway Health, Associate Professor, Epidemiology, Harvard School of Public Health, Associate Professor, Psychiatry, Harvard Medical School/ Massachusetts General Hospital, 1 Bowdoin Square, 7th Floor, Boston, MA 02114

Jaclyn M. White: jwhite@fenwayhealth.org; Katie B. Biello: kbiello@fenwayhealth.org; Sari L. Reisner: SReisner@fenwayhealth.org; Amaya Perez-Brumer: agp2133@columbia.edu; Katherine J. Heflin: kjh030@mail.harvard.edu; Matthew J. Mimiaga: MMimiaga@fenwayhealth.org

Abstract

BACKGROUND—Differences in sexual health-related outcomes by sexual behavior *and* identity remain under-investigated among bisexual female adolescents.

METHODS—Data from girls (N = 875) who participated in the Massachusetts Youth Risk Behavior Surveillance survey were analyzed. Weighted logistic regression models were fit to examine sexual and psychosocial health by lifetime sexual behavior (behaviorally bisexual vs. behaviorally heterosexual) and sexual identity (bisexual vs. heterosexual) adjusting for grade and race/ethnicity.

RESULTS—Overall, 10.5% of girls reported lifetime bisexual behavior and 8.1% reported a bisexual identity. Behavior and identity were discordant for bisexual young women as 53.2% of behaviorally bisexual students had a bisexual identity and 46.8% had a heterosexual identity. Bisexual identity and behavior were associated with unprotected intercourse at last sexual encounter, early sexual debut, 4 or more lifetime partners, history of forced/unwanted sex, STI testing history, past-year depression, and past-month drug use (all $ps < .05$).

CONCLUSION—Bisexuality, whether defined by identity or behavior, is associated with adverse sexual and psychosocial health outcomes in adolescent girls. Studies that explore wellness across the lifespan, and are designed to recognize developmental differences burgeoning in adolescence, may provide insights into the differential sexual risk outcomes observed among bisexual girls.

Keywords

bisexual; women; adolescents; sexual risk

The sexual health of bisexual women has historically been understudied.¹ This is largely because bisexual and other sexual minority women (ie, women who have sex with women only, women who have sex with both women and men, and/or women that identify as lesbian or bisexual) were presumed to be at low-risk for HIV, sexually transmitted infections (STIs) and other sexual health-related indicators such as pregnancy.^{2, 3} However, as research in this area has expanded, studies have shown that bisexual women are at equal or greater risk for acquiring HIV and other STIs than heterosexual women.^{4–6} Whereas the mechanisms behind the excess risk observed among bisexual women remains understudied, numerous studies have documented a panoply of psychosocial stressors that bisexual and other sexual minority women experience (eg, depression, substance use, violence, abuse)^{7–14} and these factors have been shown to be associated with adverse sexual health outcomes in diverse lesbian, gay and bisexual (LGB) populations.^{15–19}

One challenge to understanding the specific pathways for sexual risk behavior among bisexual women is the failure of researchers to categorize lesbian and bisexual women as distinct. Indeed, much of the quantitative research to date has collapsed lesbian, bisexual and other women who have sex with women into a single analytic group (eg, lesbian/bisexual, women who have sex with women, sexual minority, LGB, etc.).^{19–21} Whereas amalgamating bisexual and lesbian women into a single category may be beneficial for conducting statistical analyses or developing more generalizable results, it is often problematic as these categories may obfuscate our understanding of the unique risk profiles of female subgroups.^{1, 22–25} Additionally, even when bisexual women have been examined independently, researchers have typically explored only a single aspect of sexual orientation (eg, identity or behavior),^{1, 26} which may also conceal behavioral distinctions in sexual risk relative to other groups. For example, studies that explore sexual risk by combining bisexual and lesbian women on the basis of behavior (ie, women who have sex with women) may falsely conclude that bisexual and lesbian women are at equal risk for HIV and other STIs,^{2, 3} despite the fact that behaviorally bisexual women – a subgroup known to engage in higher-risk receptive penile intercourse with male partners – also comprise the sample and may have risk profiles that are more similar to heterosexual women than lesbian women.⁶ However, examining sexual risk by identity alone (ie, not taking actual behavior into

account) may also lead to the misclassification of health disparities as research indicates that sexual orientation identity is often fluid,^{1, 25, 27} especially among women and adolescents who are in the process of identity exploration and formation.^{1, 28}

Adolescence is not only a key developmental period with regard to sexual identity, but also sexual behavior, as behavioral patterns emerging in adolescence often lead to behavioral patterns in adulthood that perpetuate HIV or STI risk.²¹ Indeed, understanding the sexual experiences of school-aged youth is important, particularly among sexual minority young women as research shows that compared to their heterosexual peers, young sexual minority women report more unprotected sex,^{21, 29} sex at younger ages,^{21, 29} greater use of drugs or alcohol in general,^{21, 30} and during sex,¹⁸ more sexual partners,^{29, 31} higher prevalence of forced or coerced sexual contact,^{21, 29, 30} higher prevalence of pregnancy,^{21, 31} and more STIs.³² Sexual minority youth are also at risk for bullying^{18, 33} and physical victimization,^{19, 34} which is associated with a number of adverse health outcomes including depression,³⁵ substance use,¹⁸ and sexual risk behavior.¹⁹ However, as with the empirical literature among adult sexual minority populations, sexual health research with adolescents has focused largely on young men who have sex with men or examined lesbian, bisexual and gay adolescents or lesbian and gay young women as a homogenous group.^{1, 19, 36, 37} Moreover, little research has examined young sexual minority women's use of HIV and STI testing services – a preventative strategy indicative of risk awareness, with implications for secondary prevention.^{38, 39} Given that sexual minority youth are at greater risk for a variety of negative health outcomes, relative to their heterosexual counterparts, and research has shown that sexual orientation identity may change over time,²⁸ it is important to simultaneously examine both behavior and identity when seeking new insights on the HIV and STI risk profiles of adolescent girls.

The current study addressed several of the aforementioned gaps in the literature by examining differences in sexual risk behaviors and psychosocial factors by both sexual orientation identity and behavior using a large representative sample of high school girls in Massachusetts. The following research questions were examined: (1) Are bisexual girls at greater risk for adverse sexual risk behavior (unprotected sex, 4 or more sexual partners, early sexual debut, pregnancy, substance use during sex, HIV or STI diagnosis) and psychosocial outcomes (depression, binge drinking, drug use, bullying) in comparison to their heterosexual peers? (2) In comparing bisexual girls to heterosexual girls, are there differences in these sexual risk and psychosocial outcomes by identity versus behavior? (3) To what extent does identity moderate the effects of any observed differences in health by sexual behavior? Finally, we also sought (4) to explore whether differences existed by bisexual behavior vs. identity with regard to testing for HIV and STIs as these behaviors may serve as proxies for risk awareness and engagement in preventative health care services.

METHODS

The Massachusetts Youth Risk Behavior Survey (MYRBS) is a survey of public high school students from a scientifically selected random sample of schools across the Commonwealth. The MYRBS is conducted by the Massachusetts Department of Elementary and Secondary

Education (ESE), in conjunction with the Massachusetts Department of Public Health, and with funding from the Centers for Disease Control and Prevention (CDC). The survey monitors risk behaviors related to the leading causes of morbidity and mortality in the US among youth in grades 9–12. The anonymous survey includes questions about sexual behaviors that might lead to unintended pregnancy or sexually transmitted diseases, alcohol and other drug use, dietary behaviors, physical activity, and behaviors associated with intentional or unintentional injuries. Data from the 2007 Massachusetts YRBS were analyzed, as these were the most recent publicly available data.

Details of sampling procedures have been reported previously.⁴⁰ In brief, a probability proportionate to size random sample of public high schools (schools with at least one of grades 9 through 12) was selected. In the sampled schools, 6 classes were randomly selected; 3 were then randomly assigned to receive the MYRBS. Trained survey administrators administered the surveys in the participating schools. In 2007, data were collected from over 3000 high school students within 59 schools for the MYRBS. The overall response rate (student response rate x school response rate) was 73% for the 2007 MYRBS. Data from the MYRBS, using appropriate weighted estimates, provide accurate estimates of the prevalence of risk behaviors among public high school students in the Commonwealth of Massachusetts.⁴¹ Additional documentation concerning weighting procedures have been described in detail elsewhere.⁴⁰

Measures

Demographics—The sex of students was assessed by asking subjects to indicate their sex as male or female. Those who did not report their sex or identified as male were excluded from this analysis. Grade was determined by asking subjects what grade they were in, with response options given as 9th, 10th, 11th, 12th, ungraded, or other grade. A binary variable was created for lower grades (9th/10th) and upper grades (11th/12th). Students who reported being ungraded or in another grade and those who did not respond to this question were excluded from this analysis.

Participants were asked their race/ethnicity and classified as White/Caucasian, Black/African American, Latino/Hispanic, Asian/Pacific Islander, other race/identity (eg, American Indian/Alaskan Native, multiracial, etc.). Students were classified as white or racial/ethnic minority to allow for adequate statistical power to examine differences by sexual orientation and support. Those who did not report their race/ethnicity were excluded from analyses.

Students were asked the following question to assess sexual identity: “Which of the following best describes you?” Response options were heterosexual (straight), gay or lesbian, bisexual, not sure, or missing. Given that the analysis was focused on bisexual vs. heterosexual behavior and identity, female students that indicated a lesbian identity, responded “not sure,” and who did not respond to this question were excluded from the current analysis. Thus, girls were categorized as having either a bisexual or heterosexual identity. Students who had never had sexual intercourse were excluded from the analysis.

To assess sexual behavior, respondents were asked with whom they had sexual contact in their lifetime (ie, girls, boys, both, or neither). Girls indicating lifetime sexual contact with girls only, no history of sexual contact, and who did not respond to this question were excluded from the analysis. Thus, girls were categorized as having had lifetime sexual contact with girls and boys (behaviorally bisexual) or boys only (heterosexual).

Sexual health indicators—Eight domains of sexual health were assessed. [1]

“Unprotected intercourse at last sex” was assessed by asking subjects if they or their partner had used a condom during the most recent sexual intercourse (yes/no). [2] “Alcohol or drug use at last sex” was assessed by asking students reporting prior sexual activity whether they had used alcohol or drugs during their last sexual encounter (yes/no). [3] Students reporting prior sexual activity were asked to indicate the age at which they engaged in sexual intercourse for the first time. Using a cutoff age of 14 years, we assessed “early sexual debut” as those reporting first sexual intercourse at age 14 years or younger (yes) vs. age 15 years older (no). [4] Subjects were asked to indicate the number of lifetime sexual partners. Those indicating “4 or more lifetime partners” were coded as yes, and those with 1 to 3 partners were coded as no. [5] “History of pregnancy” was assessed by asking female students to indicate how many times they had been pregnant in their lifetime. Those who had ever been pregnant were coded as yes and those who had never been pregnant were coded as no. [6] “History of forced or unwanted sex” was assessed by asking subjects whether they had ever had sexual contact with anyone against their will. Students reporting forced or unwanted sex were coded as yes and those who had not experienced forced or unwanted sex were coded as no. [7] “History of HIV or STI diagnosis” was determined by asking students if they had ever been told by a doctor or nurse that they had an HIV infection or any other STI (phrased as “STD” in the question). Responses were coded dichotomously (yes/no). [8] Finally, “history of STI testing” was assessed by asking participants to indicate whether they had been tested for STIs such as genital herpes, chlamydia, syphilis, or genital warts. Responses were coded dichotomously (yes/no).

Psychosocial health indicators—Four psychosocial health indicators were investigated. [1] To assess “depressive distress,” students were asked a single-item screening question: “During the past 12 months, did you ever feel so sad or hopeless almost every day for 2 weeks or more in a row that you stopped doing some usual activities?” Response options were yes or no. [2] “Binge drinking” was measured by asking students the following question, “during the past 30 days, on how many days did you have 5 or more drinks of alcohol in a row, that is, within a couple of hours?” Responses ranged from 0 days to 20 or more days. A binary indicator of binge drinking was created, such that students who reported 5 or more drinks of alcohol in a row on one or more days were classified as having binge drank in the past 30 days, and those who did not were classified as not having binge drank. [3] To assess “drug use,” subjects were asked to report how many times they used illegal drugs during the past 30 days. Students indicating 1 time or more were categorized as having used drugs in the past 30 days. Those reporting 0 times, were categorized as not having used drugs in the past 30 days. [4] Finally experiences of “bullying” were determined by asking subjects to report how many times they had been bullied at school in the past 12

months. Response options ranged from 0 times to 12 or more times. Students were classified as having been bullied (1 or more times) versus not (0 times).

Data Analysis

All statistical analyses were conducted using SAS® version 9.2 statistical software. Using a sample of 3,131 sexually active youth, the data were restricted to girls (N = 1598) who reported being sexually active with a male partner (N = 912). Individuals who did not report or had an invalid response for grade-level (N = 2), race/ethnicity (N = 18), sexual identity (N = 1) were excluded. A complete case analysis was conducted such that respondents were eligible for inclusion in this analysis if they completed MYRBS questions on the sexual and psychosocial health indicators described above (missing ranged from 0.3% for HIV testing history to 1.8% for reports of binge drinking). The final data analytic sample was comprised of 875 high school girls who were sexually active.

For all analyses, statistical significance was pre-determined at the $\alpha = .05$ level. Descriptive statistics were obtained for all variables included in the analysis. Bivariate associations were obtained for all risk factors and covariates by bisexual behavior and bisexual identity (vs. heterosexual). Survey procedures^{42, 43} with appropriate weights were used for all analyses to account for the Massachusetts YRBS survey design and sampling procedures.⁴⁴ Proportional differences were assessed using the Rao-Scott chi-square tests, a version of the Pearson chi-square which adjusts for complex survey sample designs.⁴³

A series of logistic regression models were fit to test the association between bisexual behavior and bisexual identity, separately, and sexual risk outcomes, as well as other risk factors. Additionally, to assess potential effect modification of bisexual behavior and identity, for each outcome, a model was fit with a behavior by identity interaction term in addition to the main effects terms.

RESULTS

Demographics

Table 1 presents demographic characteristics, sexual health-related indicators, and psychosocial health indicators, stratified by behaviorally bisexual and heterosexual girls, and by bisexual and heterosexual identity. Weighted bivariate comparisons are provided for each sexual orientation dimension.

The majority of students in the sample (91.9%) identified as heterosexual, and 8.1% identified as bisexual. Among those having only male sex partners in their lifetime (behaviorally heterosexual: 89.5%), the majority (97.2%) identified as heterosexual and 2.8% identified as bisexual. Among high school girls reporting lifetime sexual behavior with both male and female partners (behaviorally bisexual; 10.5%), 53.2% reported a bisexual identity and 46.8% reported a heterosexual identity ($p < .0001$).

Sexual Health-related Indicators

Table 2 presents weighted multivariable models where the sexual health-related indicators are separately regressed on sexual behavior (comparing behaviorally bisexual and

heterosexual young girls) in Column A and sexual identity (comparing bisexual-identified and heterosexual-identified girls) in Column B. Models testing whether the association between sexual behavior and the sexual health-related indicators are modified by identity are shown in Column C. All models are adjusted for race/ethnicity and grade.

Compared to behaviorally heterosexual girls, behaviorally bisexual girls had higher odds of unprotected intercourse at last sex (aOR=2.22; 95% CI=1.26, 3.92), alcohol and/or drug use at last sex (aOR=2.30; 95% CI=1.33, 3.99), early sexual debut (aOR=3.48; 95% CI=1.54, 7.85), having 4 or more sexual partners in their lifetime (aOR=3.44; 95% CI=1.95, 6.08), history of forced or unwanted sex (aOR=4.04; 95% CI=2.61, 6.26), and having ever been tested for STIs (aOR=2.00; 95% CI=1.13, 3.53). There were no statistically significant differences in history of pregnancy or history or lifetime HIV or STI diagnosis for behaviorally bisexual versus heterosexual adolescent girls (Table 2, Column A).

Relative to girls self-identifying as heterosexual, bisexual-identified girls had higher odds of unprotected intercourse at last sex (aOR=2.89; 95% CI=1.77, 4.71), early sexual debut (aOR=3.37; 95% CI=1.63, 6.97), 4 or more lifetime sexual partners (aOR=3.57; 95% CI=1.80, 7.09), history of forced or unwanted sex (aOR=3.81; 95% CI=2.06, 7.04), lifetime history of HIV or STI diagnosis (aOR=2.96; 95% CI=1.08, 8.11), and having ever been tested for STIs (aOR=2.37; 95% CI=1.35, 4.18). No significant differences by sexual identity were found in alcohol and/or drug use at last sex or lifetime history of pregnancy (Table 2, Column B).

Support for a behavior by identity interaction was found for only 1 of the 8 sexual health-related indicators: having ever been tested for STIs (for interaction: $\beta = -1.20$, $p = .018$) - indicating that while behaviorally bisexual girls had an higher odds of ever having been tested for STIs compared to behaviorally heterosexual girls, also identifying as bisexual decreases the magnitude of this association.

Psychosocial Health Indicators

Table 3 presents weighted multivariable models where the psychosocial health indicators are separately regressed on sexual behavior (comparing behaviorally bisexual and heterosexual girls) in Column A and sexual identity (comparing bisexual-identified and heterosexual-identified girls) in Column B. Models testing whether the association between sexual behavior and the psychosocial health indicators are modified by identity are shown in Column C. All models are adjusted for race/ethnicity and grade.

Relative to behaviorally heterosexual girls, behaviorally bisexual girls had higher odds of depression (aOR=3.59; 95% CI=2.56, 5.05), drug use in the past 30 days (aOR=2.93; 95% CI=1.72, 4.99), and having been bullied in the past 12 months (aOR=2.45; 95% CI=1.61, 3.72). No statistically significant differences were found by sexual behavior and binge drinking in the past 30 days (Table 3, Column A).

Compared to heterosexual-identified girls, those self-identifying as bisexual were at higher odds of depression (aOR=5.21; 95% CI=3.01, 9.04), binge drinking in the past 30 days (aOR=1.62; 95% CI=1.03, 2.56), and past-30 day drug use (aOR=2.28; 95% CI=1.25, 4.17).

Experiences of bullying in the past 12 months did not differ by sexual identity (Table 3, Column B). Additionally, there was no evidence of a sexual behavior by sexual identity interaction for the 4 psychosocial health indicators (Table 3, Column C).

DISCUSSION

Sexual orientation has not been routinely collected by all states and jurisdictions on the national YRBS.^{45, 46} Although the CDC will add questions about same-sex sexual contact and sexual identity to their state, territorial, or local YRBS questionnaires starting in 2015, currently, sexual behavior and identity questions remain optional with only 21 states assessing both same sex behavior and identity in 2011.⁴⁷ Thus, statewide, representative data of high school girls from Massachusetts offer a unique contribution to the female adolescent health literature. The current analysis demonstrates disparities in sexual and psychosocial health indicators by sexual orientation, such that bisexual girls, whether defined by behavior or identity, more frequently experienced poorer sexual and psychosocial health outcomes compared to heterosexual girls. In addition, findings demonstrated several distinct differences in sexual and psychosocial health outcomes by sexual orientation dimensions (ie, identity versus behavior), supporting the need to utilize multiple dimensions of sexual orientation when assessing the health of female adolescents.

Consistent with prior research,^{34, 35, 48} girls who reported sex with both male and female partners more frequently reported being bullied and using substances during sex, compared to girls with only male partners. Moreover, compared to heterosexual behavior and identity, both bisexual behavior and identity were associated with the higher prevalence of depression, drug use, and a history of forced or unwanted sex. Whereas the reasons for such findings are not completely understood due to limited sexual health research involving female adolescents, research among adult bisexual men and women shows that individuals who identify and/or engage in bisexual behavior, while included under the LGBT umbrella, may not have access to the same sense of community as other members of the LGBT community, as the bisexual community is less unified.^{49, 50} Additionally, bisexual people may experience rejection from both their LGBT and heterosexual peers who perceive them as an outsider.^{51–53} Rejection from heterosexual and gay peers may lead to greater minority stress for bisexual people, and in turn, contribute to poorer health outcomes.^{50, 53} Future research should explore the extent to which minority stress contributes to poor health outcomes in female sexual minorities, especially during their formative high school years.

Bisexual behavior and identity were associated with sexual behaviors that may place girls and young women at greater risk for HIV and other STIs. However, both ones with a bisexual identity and those who engaged in bisexual behavior had a higher probability than their heterosexual counterparts of having been tested for STI's in their lifetime - a health behavior important in protecting one's sexual health and that of their sexual partners. Whereas the causal mechanisms behind the higher prevalence of testing among bisexual girls warrant investigation, these findings could suggest that bisexual girls are aware of their risk for HIV and STIs and/or may be more willing to engage in preventative health services, and are, thus, getting tested more frequently than their heterosexual peers. This finding could prove useful to those looking to develop school-based prevention interventions with

at-risk bisexual girls as testing affords those accessing it with the opportunity to engage with healthcare providers who may be able to disseminate sexual health information and assist in building behavioral skills to reduce sexual risk. Point of care interventions with bisexual girls may also benefit from addressing the underlying psychosocial health concerns that many of these girls disproportionately face by providing supportive referrals or triage to mental healthcare and substance abuse treatment. Those engaged in intervention efforts will need to take care to assess both bisexual behavior and identity to ensure that bisexual girls engaged in risky health behaviors are identified and supported.

Girls who reported sexual contact with both male and female partners had significantly greater discordance between identity and behavior than behaviorally heterosexual girls, which is consistent with previous research.^{21, 35, 54} Sexual and psychosocial health outcomes also varied by sexual orientation dimension in some cases. For example, girls with a bisexual identity more frequently reported having received an HIV or STI diagnosis as well as recent binge drinking, compared to heterosexual girls, whereas bisexual behavior was associated with substance use during sex and experiences of bullying. In exploring 2 measures of sexual orientation, we identified a higher frequency of adverse health behaviors and outcomes among bisexual girls relative to their heterosexual peers; differences which may have been obscured had only one measure been utilized or bisexual identity or behavior combined with lesbian identity or behavior. Additional research is needed to understand the specific mechanisms behind these differences, including the contexts in which risk behaviors occur, given prior studies demonstrating that social context is a key determinant of health behaviors.⁵⁵⁻⁵⁸ For example, participants in a qualitative study of girls described the social pressures to conform to the norms of heterosexuality, having a boyfriend, or having sex, and the challenges of meeting these ideals while attempting to make sense of their own desires and attractions.⁵⁸ Longitudinal studies that explore the individual and contextual factors that shape risk behavior may help to better understand the causal pathways for the specific risk differences observed among behaviorally bisexual and identified high school girls relative to their heterosexual peers.

Findings from the current analysis should be interpreted in light of several limitations. The Massachusetts YRBS enrolls school-engaged youth in public schools, which may miss homeless or marginally housed youth,⁵⁹ as well as teens attending private schools. Also, the cross-sectional design does not allow for causal conclusions to be made. Whereas our sample size was large, stratification across measures resulted in small cell sizes for some analyses. Additionally, depressive distress was assessed categorically. Although categorical measurement of mental health disorders do not allow for the collection of all clinically relevant information (eg, severity of symptoms), and can result in the inaccurate measurement of disease prevalence, depressive distress is not a proxy for clinically diagnosed depression in this sample. Data from the 2007 Massachusetts YRBS were analyzed as these were the most recent publicly available data. Although it is possible that the 2007 sample may be different than the 2013 sample, our results are largely consistent with findings from recently published studies,^{21, 45} and should be considered given the unique contribution of our findings using dual measures of sexual orientation. Lastly, given that this is a secondary data analysis, we were limited to the data that were collected and

available for public use. As a result, lack of data on other potential confounders (eg, sexual assault at first sex, childhood sexual abuse) could bias the results.

Overall, these findings document that engaging in bisexual behavior or having a bisexual identity is associated with a variety of adverse sexual and psychosocial health outcomes in high school girls – findings that lend support to including questions using multiple dimensions of sexual orientation when evaluating the risk behavior of sexually active adolescents. Longitudinal research is needed to assess whether and how minority stress contributes to worse health outcomes among bisexual girls and identify the underlying factors that lead to such disparities so that effective interventions may be developed.

IMPLICATIONS FOR SCHOOL HEALTH

Bisexuality, whether defined by identity or behavior, is associated with numerous adverse sexual and psychosocial health outcomes in high school girls. This paper illustrates the importance of identifying at-risk bisexual youth not just by their identity and affiliation with the LGBT community, but also by their behavior. The disparities faced by bisexual female youth must be addressed through comprehensive education and interventions facilitated by school personnel and district leadership.

Targeted, school-based programs addressing the specific health needs of bisexual youth are greatly needed. School settings are an ideal place for such interventions as the materials covered by educators in school-wide sexual health programs could serve to normalize bisexual behavior and identities, thereby improving the social environments of bisexual students and reducing experiences of bullying and other contextual factors. Whereas the discussion of LGBT health can be controversial in some school districts, bisexuality is not historically more difficult to broach with concerned parents than other parts of the LGBT spectrum and should be incorporated into school-based health programs.

Ideal interventions include timely, nonjudgmental, comprehensive school-based sexual education programs to support and reduce the disparities experienced by high school girls. Specifically, school health practices and lessons should elucidate the health needs, risks, and resources directly associated with bisexual behaviors and identities in addition to addressing the needs of other groups within the LGBT spectrum. Additionally, school health practitioners should nurture peer educators to provide individual counseling as well as develop culturally-competent local resources (such as information brochures) to address the sexual and psychosocial health needs of female students engaging in bisexual behavior and/or bisexually identified.

Human Subjects Approval Statement

The Centers for Disease Control and Prevention's institutional review board granted approved the administration of the Youth Risk Behavior Survey MYRBS to be administered nationwide. We obtained administrative approvals from the Massachusetts Department of Elementary and Secondary Education (MDESE), and by the AIDS Advisory and Materials Review Panel, as required by the MDESE Cooperative Agreement with the Centers for

Disease Control and Prevention. MDESE staff members conducted the survey in full compliance with standards for ethical treatment of individuals participating in the project.

Acknowledgments

The authors thank the Massachusetts Department of Elementary and Secondary Education for their permission to use the Massachusetts Youth Risk Behavior Survey data. The authors declare no potential conflicts of interest and received no financial support for the research, authorship, and/or publication of this article. J. White is supported by grants T32MH020031 and P30MH062294 from the National Institute of Mental Health.

References

1. Russell ST, Seif H. Bisexual female adolescents: a critical analysis of past research, and results from a national survey. *J Bisex.* 2010; 2(2–3):73–94.
2. Eaton L, Kalichman S, Cain D, Cherry C, Pope H, Fuhrel A, et al. Perceived prevalence and risks for human papillomavirus (HPV) infection among women who have sex with women. *J Womens Health.* 2008; 17(1):75–84.
3. Fethers K, Marks C, Mindel A, Estcourt C. Sexually transmitted infections and risk behaviors in women who have sex with women. *Sex Transm Infect.* 2000; 76(5):345–349. [PubMed: 11141849]
4. Moore J, Warren D, Zierler S, Schuman P, Solomon L, Schoenbaum E, et al. Characteristics of HIV-infected lesbians and bisexual women in four urban centers. *Womens Health.* 1996; 2(1):49–60.
5. Scheer S, Peterson I, Page-Shafer K, Delgado V, Gleghorn A, Ruiz J, et al. Sexual and drug use behavior among women who have sex with both women and men: results of a population-based survey. *Am J Public Health.* 2002; 92(7):1110–1112. [PubMed: 12084692]
6. Everett BG. Sexual orientation disparities in sexually transmitted infections: examining the intersection between sexual identity and sexual behavior. *Arch Sex Behav.* 2013; 42(2):225–236. [PubMed: 22350122]
7. Stevens S, Korchmaros JD, Miller D. A comparison of victimization and perpetration of intimate partner violence among drug abusing heterosexual and lesbian women. *J Fam Violence.* 2010; 25(7):639–649.
8. Corliss HL, Grella CE, Mays VM, Cochran SD. Drug use, drug severity, and help-seeking behaviors of lesbian and bisexual women. *J Womens Health.* 2006; 15(5):556–568.
9. Koh AS, Ross LK. Mental health issues: A comparison of lesbian, bisexual and heterosexual women. *J Homosex.* 2006; 51(1):33–57. [PubMed: 16893825]
10. Cochran SD, Sullivan JG, Mays VM. Prevalence of mental disorders, psychological distress, and mental services use among lesbian, gay, and bisexual adults in the United States. *J Consult Clin Psychol.* 2003; 71(1):53–61. [PubMed: 12602425]
11. Kecojevic A, Wong CF, Schrager SM, Silva K, Bloom JJ, Iverson E, et al. Initiation into prescription drug misuse: differences between lesbian, gay, bisexual, transgender (LGBT) and heterosexual high-risk young adults in Los Angeles and New York. *Addict Behav.* 2012; 37(11):1289–1293. [PubMed: 22738887]
12. King M, Semlyen J, Tai SS, Killaspy H, Osborn D, Popelyuk D, et al. A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. *BMC Psychiatry.* 2008; 8(1):1–17. [PubMed: 18173833]
13. Hipwell A, Stepp S, Keenan K, Allen A, Hoffmann A, Rottingen L, et al. Examining links between sexual risk behaviors and dating violence involvement as a function of sexual orientation. *J Pediatr Adolesc Gynecol.* 2013; 26(4):212–218. [PubMed: 23726138]
14. Pyra M, Weber K, Wilson TE, Cohen J, Murchison L, Goparaju L, et al. Sexual minority status and violence among HIV infected and at-risk women. *J Gen Intern Med.* 2014; 29(8):1–8.
15. Savin-Williams RC. Verbal and physical abuse as stressors in the lives of lesbian, gay male, and bisexual youths: associations with school problems, running away, substance abuse, prostitution, and suicide. *J Consult Clin Psychol.* 1994; 62(2):261–269. [PubMed: 8201062]

16. Colfax G, Vittinghoff E, Husnik MJ, McKirnan D, Buchbinder S, Koblin B, et al. Substance use and sexual risk: a participant-and episode-level analysis among a cohort of men who have sex with men. *Am J Epidemiol.* 2004; 159(10):1002–1012. [PubMed: 15128613]
17. Stall R, Mills TC, Williamson J, Hart T, Greenwood G, Paul J, et al. Association of co-occurring psychosocial health problems and increased vulnerability to HIV/AIDS among urban men who have sex with men. *Am J Public Health.* 2003; 93(6):939–942. [PubMed: 12773359]
18. Bontempo DE, D’Augelli AR. Effects of at-school victimization and sexual orientation on lesbian, gay, or bisexual youths’ health risk behavior. *J Adolesc Health.* 2002; 30(5):364–374. [PubMed: 11996785]
19. Garofalo R, Wolf RC, Kessel S, Palfrey J, DuRant RH. The association between health risk behaviors and sexual orientation among a school-based sample of adolescents. *Pediatrics.* 1998; 101(5):895–902. [PubMed: 9565422]
20. Faulkner AH, Cranston K. Correlates of same-sex sexual behavior in a random sample of Massachusetts high school students. *Am J Public Health.* 1998; 88(2):262–266. [PubMed: 9491018]
21. Goodenow C, Szalacha LA, Robin LE, Westheimer K. Dimensions of sexual orientation and HIV-related risk among adolescent females: evidence from a statewide survey. *J Info.* 2008; 98(6): 1051–1058.
22. Lindley LL, Nicholson TJ, Kerby MB, Lu N. HIV/STI associated risk behaviors among self-identified lesbian, gay, bisexual, and transgender college students in the United States. *AIDS Educ Prev.* 2003; 15(5):413–429. [PubMed: 14626464]
23. Marshal MP, Friedman MS, Stall R, King KM, Miles J, Gold MA, et al. Sexual orientation and adolescent substance use: a meta-analysis and methodological review*. *Addiction.* 2008; 103(4): 546–556. [PubMed: 18339100]
24. Rosario M, Meyer-Bahlburg H, Hunter J, Gwadz M. Sexual risk behaviors of gay, lesbian, and bisexual youths in New York City: prevalence and correlates. *AIDS Educ Prev.* 1999; 11(6):476–496. [PubMed: 10693645]
25. Saewyc E, Skay C, Richens K, Reis E, Poon C, Murphy A. Sexual orientation, sexual abuse, and HIV-risk behaviors among adolescents in the Pacific Northwest. *J Info.* 2006; 96(6):1104–1110.
26. Chung YB, Katayama M. Assessment of sexual orientation in lesbian/gay/bisexual studies. *J Homosex.* 1996; 30(4):49–62. [PubMed: 8738744]
27. Laumann, EO. *The Social Organization of Sexuality: Sexual Practices in the United States.* Chicago, IL: University of Chicago Press; 1994.
28. Rosario M, Schrimshaw EW, Hunter J, Braun L. Sexual identity development among lesbian, gay, and bisexual youths: consistency and change over time. *J Sex Res.* 2006; 43(1):46–58. [PubMed: 16817067]
29. Tornello SL, Riskind RG, Patterson CJ. Sexual orientation and sexual and reproductive health among adolescent young women in the United States. *J Adolesc Health.* 2014; 54(2):160–168. [PubMed: 24157195]
30. Scheer S, Parks CA, McFarland W, Page-Shafer K, Delgado V, Ruiz JD, et al. Self-reported sexual identity, sexual behaviors and health risks: examples from a population-based survey of young women. *Journal of Lesbian Studies.* 2002; 7(1):69–83. [PubMed: 24815715]
31. Saewyc EM, Bearinger LH, Blum RW, Resnick MD. Sexual intercourse, abuse and pregnancy among adolescent women: does sexual orientation make a difference? *Fam Plann Perspect.* 1999; 31(3):127–131. [PubMed: 10379429]
32. Mojola SA, Everett B. STD and HIV risk factors among US young adults: Variations by gender, race, ethnicity and sexual orientation. *Perspect Sex Reprod Health.* 2012; 44(2):125–133. [PubMed: 22681428]
33. Berlan ED, Corliss HL, Field AE, Goodman E, Bryn Austin S. Sexual orientation and bullying among adolescents in the Growing Up Today Study. *J Adolesc Health.* 2010; 46(4):366–371. [PubMed: 20307826]
34. Russell ST, Driscoll AK, Truong N. Adolescent same-sex romantic attractions and relationships: implications for substance use and abuse. *Am J Public Health.* 2002; 92(2):198–202. [PubMed: 11818291]

35. Lindley LL, Walsemann KM, Carter JW. The association of sexual orientation measures with young adults' health-related outcomes. *Am J Public Health*. 2012; 102(6):1177–1185. [PubMed: 22021310]
36. Marshal MP, Friedman MS, Stall R, Thompson AL. Individual trajectories of substance use in lesbian, gay and bisexual youth and heterosexual youth. *Addiction*. 2009; 104(6):974–981. [PubMed: 19344440]
37. Thoma BC, Huebner DM, Rullo JE. Unseen risks: HIV-related risk behaviors among ethnically diverse sexual minority adolescent females. *AIDS Educ Prev*. 2013; 25(6):535–541. [PubMed: 24245599]
38. Maguen S, Armistead LP, Kalichman S. Predictors of HIV antibody testing among gay, lesbian, and bisexual youth. *J Adolesc Health*. 2000; 26(4):252–257. [PubMed: 10734272]
39. Masters NT, Beadnell B, Morrison DM, Hoppe MJ, Wells EA. Multidimensional characterization of sexual minority adolescents' sexual safety strategies. *J Adolesc*. 2013; 36(5):953–961. [PubMed: 24011111]
40. Centers for Disease Control and Prevention (CDC). Methodology of the youth Risk Behavior Surveillance System - 2013. Atlanta, GA: CDC; 2013.
41. Massachusetts Department of Public Health. Health and Risk Behaviors of Massachusetts Youth, 2007: The Report. Boston, MA: Massachusetts Department of Public Health; 2008. Available at: <http://www.doe.mass.edu/cnp/hprograms/yrbs/> [Accessed June 10, 2014]
42. Lehtonen, R.; Pahkinen, E. Practical Methods for Design and Analysis of Complex Surveys. Chichester, UK: John Wiley & Sons; 2004.
43. Thomas DR, Singh A, Roberts G. Tests of independence on two-way tables under cluster sampling: an evaluation. *Intl Stat Rev*. 1996; 64(3):295–311.
44. Massachusetts Department of Elementary and Secondary Education. Massachusetts Coordinated School Health Program. Mass.Gov; 2013. Nutrition, Health, and Safety: Youth Risk Behavior Survey. Available at: <http://www.doe.mass.edu/cnp/hprograms/yrbs/> [Accessed September June 10, 2014]
45. Mustanski B, Van Wagenen A, Birkett M, Eyster S, Corliss HL. Identifying sexual orientation health disparities in adolescents: analysis of pooled data from the Youth Risk Behavior Survey, 2005 and 2007. *Am J Public Health*. 2014; 104(2):211–217. [PubMed: 24328640]
46. Bradford J, Mustanski B. Health disparities among sexual minority youth: the value of population data. *Am J Public Health*. 2014; 104(2):197–197. [PubMed: 24328630]
47. Centers for Disease Control and Prevention (CDC). National Center for HIV/AIDS VH, STD, and TB Prevention. Lesbian, Gay, Bisexual and Transgender Health. Atlanta, GA: CDC; 2013. LGBTQ youth programs-at-a-glance. Available at: <http://www.cdc.gov/lgbthealth/youth-programs.htm> [Accessed: October 1, 2013]
48. Austin SB, Roberts AL, Corliss HL, Molnar BE. Sexual violence victimization history and sexual risk indicators in a community-based urban cohort of “mostly heterosexual” and heterosexual young women. *Am J Public Health*. 2008; 98(6):1015–1020. [PubMed: 17901440]
49. Alarie M, Gaudet S. “I don't know if she is bisexual or if she just wants to get attention”: analyzing the various mechanisms through which emerging adults invisibilize bisexuality. *J Bisex*. 2013; 13(2):191–214.
50. Dodge B, Schnarrs PW, Reece M, Goncalves G, Martinez O, Nix R, et al. Community involvement among behaviourally bisexual men in the Midwestern USA: experiences and perceptions across communities. *Cult Health Sex*. 2012; 14(9):1095–1110. [PubMed: 22978551]
51. Thorne L. “But I'm attracted to women”: sexuality and sexual identity performance in interactional discourse among bisexual students. *J Lang Sex*. 2013; 2(1):70–100.
52. Scherrer K. Culturally competent practice with bisexual individuals. *Clin Soc Work J*. 2013; 41(3): 238–248.
53. Friedman MR, Dodge B, Schick V, Herbenick D, Hubach RD, Bowling J, et al. From bias to bisexual health disparities: attitudes toward bisexual men and women in the United States. *LGBT Health*. 2014; 1(4):309–318. [PubMed: 25568885]

54. Mustanski B, Birkett M, Greene GJ, Rosario M, Bostwick W, Everett BG. The association between sexual orientation identity and behavior across race/ethnicity, sex, and age in a probability sample of high school students. *Am J Public Health*. 2014; 104(2):237–244. [PubMed: 24328662]
55. Browning CR, Leventhal T, Brooks-Gunn J. Neighborhood context and racial differences in early adolescent sexual activity. *Demography*. 2004; 41(4):697–720. [PubMed: 15622950]
56. Suarez T, Miller J. Negotiating risks in context: a perspective on unprotected anal intercourse and barebacking among men who have sex with men—where do we go from here? *Arch Sex Behav*. 2001; 30(3):287–300. [PubMed: 11330118]
57. Hartel D. Context of HIV risk behavior among female injecting drug users and female sexual partners of injecting drug users. *NIDA Res Monogr*. 1994:41–47. [PubMed: 8742590]
58. Teitelman AM, Bohinski JM, Boente A. The social context of sexual health and sexual risk for urban adolescent girls in the United States. *Issues Ment Health Nurs*. 2009; 30(7):460–469. [PubMed: 19544131]
59. Corliss HL, Goodenow CS, Nichols L, Austin SB. High burden of homelessness among sexual-minority adolescents: findings from a representative Massachusetts high school sample. *Am J Public Health*. 2011; 101(9):1683–1689. [PubMed: 21778481]

Table 1

Sample Characteristics of High School Girls who Completed the 2007 Massachusetts Youth Risk Behavior Surveillance (YRBS) Survey (N=875), Stratified by Lifetime Sexual Behavior (behaviorally bisexual=10.5% vs. behaviorally-heterosexual=89.5%) and Sexual Identity (bisexual=8.1% vs. heterosexual=91.9%)⁺

	SEXUAL BEHAVIOR – Lifetime				SEXUAL IDENTITY – Current				
	Total Sample (N = 875)	Behaviorally Bisexual (N = 91) 10.5 %	Behaviorally Heterosexual (N = 784) 89.5 %	Rao X ²	p-value	Bisexual Identity (N = 72) 8.1 %	Heterosexual Identity (N = 803) 91.9 %	Rao X ²	p-value
DEMOGRAPHICS									
Grade									
Lower Grades (Freshman or Sophomore)	43.4	50.3	42.5	1.29	0.256	55.0	42.4	3.71	0.054
Upper Grades (Junior or Senior)	56.6	49.7	57.5			45.0	57.6		
Race									
White/Caucasian	76.4	73.9	76.7	1.38	0.848	68.9	77.0	4.81	0.307
Black/African American	5.6	3.8	5.8			4.9	5.6		
Latino/Hispanic	12.4	15.6	12.0			15.2	12.1		
Asian/Pacific Islander	2.4	3.0	2.4			5.1	2.2		
Other Race/Ethnicity	3.3	3.7	3.2			5.9	3.0		
SEXUAL IDENTITY & BEHAVIOR									
Bisexual Identity									
Bisexual	8.1	53.2	2.8	238.62	<0.0001	-	-	-	-
Heterosexual	91.9	46.8	97.2			-	-		
SEXUAL HEALTH-RELATED									
Sex in the Past 3 Months	56.9	60.1	56.6	0.27	0.605	71.1	55.7	6.17	0.013
Unprotected Sex at Last Sex ^{a,b}	22.8	35.0	21.4	5.42	0.020	39.2	21.3	11.94	0.001
Alcohol and/or Drug Use During Last Sex ^b	12.1	22.7	10.9	12.00	0.0005	18.8	11.5	5.62	0.018
Early Sexual Debut (Age 14 and Younger)	28.9	40.7	27.6	7.23	0.007	50.0	27.0	14.48	0.0001
Four or More Sex Partners - Lifetime	17.4	37.2	15.2	18.42	<0.0001	38.0	15.5	15.17	<0.0001

	SEXUAL BEHAVIOR – Lifetime				SEXUAL IDENTITY – Current				
	Total Sample (N = 875)	Behaviorally Bisexual (N = 91) 10.5 %	Behaviorally Heterosexual (N = 784) 89.5 %	Rao X ²	p-value	Bisexual Identity (N = 72) 8.1 %	Heterosexual Identity (N = 803) 91.9 %	Rao X ²	p-value
History of Pregnancy - Lifetime	18.5	19.1	18.4	0.02	0.895	19.2	18.4	0.03	0.871
History of Forced/Unwanted Sex - Lifetime	24.7	52.0	21.5	47.34	<0.0001	51.7	22.3	23.88	<0.0001
HIV or STI Diagnosis - Lifetime	2.4	4.9	2.1	2.24	0.134	6.2	2.1	4.07	0.044
Tested for STI - Lifetime	22.1	32.1	20.9	4.09	0.043	35.0	21.0	6.99	0.008
PSYCHOSOCIAL HEALTH RELATED									
Depression – Past 12 Months	38.7	67.0	35.3	59.38	<0.0001	75.3	35.5	51.05	<0.0001
Binge drinking - Past 30 days	39.3	44.1	38.7	0.93	0.334	48.8	38.4	3.53	0.061
Drug use - Past 30 days	31.0	51.9	28.6	13.70	0.0002	46.1	29.7	5.83	0.016
Bullied – Past 12 Months	23.0	39.4	21.0	21.12	<0.0001	30.1	22.3	1.47	0.226

⁺ All frequencies and bivariate estimates are weighted.

^a Unprotected sex = sex without a condom during most recent sexual intercourse act

^b This is for the entire sample (not just those who had sex in past three months)

NE=Non-estimable because there is a zero cell

Table 2

Weighted Multivariable Logistic Regression Models: Regressing Sexual Health Indicators on Behaviorally-Bisexual vs. Behaviorally-Heterosexual and Bisexual Identity vs. Heterosexual Identity (Grade- and Race-Adjusted)

	Sexual Health Indicator 1: Unprotected Intercourse at Last Sex				Sexual Health Indicator 2: Alcohol and/or Drug Use at Last Sex							
	Model 1A		Model 1B		Model 1C		Model 2A		Model 2B		Model 2C	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value
Behaviorally Bisexual	2.22 (1.26, 3.92)	.006	-	-	0.76 (0.37)	.039	2.30 (1.33, 3.99)	.003	-	-	0.78 (0.47)	.094
Bisexual Identity	-	-	2.89 (1.77, 4.71)	<.0001	1.45 (0.48)	.002	-	-	1.68 (0.98, 2.87)	.057	-0.41 (0.80)	.607
Interaction Term	-	-	-	-	-1.27 (0.70)	.070	-	-	-	-	0.49 (1.08)	.648
Covariates:												
Nonwhite Race/Ethnicity	1.46 (0.83, 2.58)	.188	1.44 (0.81, 2.55)	.214	0.38 (0.29)	.196	1.20 (0.74, 1.96)	.464	1.19 (0.73, 1.96)	.487	0.18 (0.25)	.466
Upper Grades*	2.94 (2.11, 4.08)	<.0001	3.04 (2.20, 4.20)	<.0001	1.13 (0.17)	<.0001	1.68 (0.90, 3.15)	.106	1.67 (0.90, 3.09)	.101	0.52 (0.32)	.111

	Sexual Health Indicator 3: Early Sexual Debut (Age 14 and under)				Sexual Health Indicator 4: Four or More Sexual Partners - Lifetime							
	Model 3A		Model 3B		Model 3C		Model 4A		Model 4B		Model 4C	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value
Behaviorally Bisexual	1.67 (1.08, 2.57)	.02	-	-	0.25 (0.36)	.474	3.44 (1.95, 6.08)	<.0001	-	-	1.10 (0.38)	.003
Bisexual Identity	-	-	2.30 (1.38, 3.84)	.001	1.03 (0.49)	.038	-	-	3.57 (1.80, 7.09)	.0003	1.17 (0.72)	.106
Interaction Term	-	-	-	-	-0.52 (0.78)	.505	-	-	-	-	-0.84 (0.80)	.295
Covariates:												
Nonwhite Race/Ethnicity	2.09 (1.44, 3.05)	.0001	2.06 (1.42, 3.01)	.0002	0.73 (0.19)	<.0001	1.20 (0.79, 1.82)	.399	1.17 (0.78, 1.77)	.451	0.17 (0.21)	.421
Upper Grades*	0.49 (0.27, 0.83)	<.0001	0.50 (0.36, 0.68)	<.0001	-0.70 (0.16)	<.0001	2.21 (1.36, 3.60)	.001	2.26 (1.43, 3.55)	.0004	0.83 (0.24)	.0005

	Sexual Health Indicator 5: History of Pregnancy - Lifetime				Sexual Health Indicator 6: History of Forced or Unwanted Sex - Lifetime							
	Model 5A		Model 5B		Model 5C		Model 6A		Model 6B		Model 6C	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value
Behaviorally Bisexual	1.19 (0.63, 2.27)	.589	-	-	0.14 (0.47)	.763	4.04 (2.61, 6.26)	<.0001	-	-	1.30 (0.33)	<.0001
Bisexual Identity	-	-	1.33 (0.72, 2.49)	.365	0.42 (0.64)	.511	-	-	3.81 (2.06, 7.04)	<.0001	1.13 (0.44)	.009

	Sexual Health Indicator 1: Unprotected Intercourse at Last Sex						Sexual Health Indicator 2: Alcohol and/or Drug Use at Last Sex					
	Model 1A		Model 1B		Model 1C		Model 2A		Model 2B		Model 2C	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value
Interaction Term	-	-	-	-	-0.33 (0.80)	.679	-	-	-	-	-0.88 (0.53)	.100
Covariates:												
Nonwhite Race/Ethnicity	0.66 (0.37, 1.20)	.171	0.66 (0.36, 1.20)	.173	-0.41 (0.30)	.173	1.08 (0.78, 1.49)	.660	1.04 (0.75, 1.45)	.806	0.06 (0.17)	.717
Upper Grades*	4.24 (2.73, 6.60)	<.0001	4.27 (2.75, 6.64)	<.0001	1.45 (0.23)	<.0001	0.87 (0.59, 1.29)	.499	0.88 (0.60, 1.30)	.528	-0.11 (0.20)	.568
	Sexual Health Indicator 7: History of HIV or STI Diagnosis – Lifetime						Sexual Health Indicator 8: History of STI Testing – Lifetime					
	Model 7A		Model 7B		Model 7C		Model 8A		Model 8B		Model 8C	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value
Behaviorally Bisexual	2.35 (0.81, 6.81)	.117	-	-	-0.06 (1.07)	.957	2.00 (1.13, 3.53)	0.017	-	-	0.72 (0.35)	.041
Bisexual Identity	-	-	2.96 (1.08, 8.11)	.034	0.38 (0.96)	.693	-	-	2.37 (1.35, 4.18)	.003	1.24 (0.43)	.004
Interaction	-	-	-	-	0.98 (1.66)	.554	-	-	-	-	-1.20 (0.51)	.018
Covariates:												
Nonwhite Race/Ethnicity	2.39 (1.02, 5.61)	.045	2.33 (1.01, 5.36)	.047	0.84 (0.43)	.049	1.69 (1.06, 2.69)	0.028	1.67 (1.05, 2.67)	.032	0.53 (0.24)	.026
Upper Grades*	1.03 (0.33, 3.26)	.954	1.08 (0.34, 3.39)	.901	0.06 (0.58)	.914	2.61 (1.75, 3.89)	<.0001	2.66 (1.79, 3.94)	<.0001	0.99 (0.20)	<.0001

* Upper Grades = 11th and 12th grades (referent = 9th and 10th grades)

Table 3

Weighted Multivariable Logistic Regression Models: Regressing Psychosocial Health Indicators on Behaviorally-Bisexual vs. Behaviorally-Heterosexual and Bisexual Identity vs. Heterosexual Identity (Grade- and Race-Adjusted)

	Psychosocial Indicator 1: Depression – Past 12 Months										Psychosocial Indicator 2: Binge Drinking – Past 30 Days																	
	Model 9A		Model 9B		Model 9C		Model 10A		Model 10B		Model 10C		Model 11A		Model 11B		Model 11C		Model 12A		Model 12B		Model 12C					
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	B (SE)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value		
Behaviorally Bisexual	3.59 (2.56, 5.05)	<.0001	-	-	1.10 (0.34)	.001	1.26 (0.78, 2.03)	.347	-	-	-0.17 (0.32)	.590	2.93 (1.72, 4.99)	<.0001	-	-	0.88 (0.33)	.008	2.45 (1.61, 3.72)	<.0001	-	-	1.07 (0.29)	.0003	2.93 (1.72, 4.99)	<.0001		
Bisexual Identity	-	-	5.21 (3.01, 9.04)	<.0001	2.06 (0.62)	.001	-	-	1.62 (1.03, 2.56)	.039	0.20 (0.41)	.627	-	-	2.28 (1.25, 4.17)	.007	0.0005 (0.53)	.999	-	-	1.47 (0.75, 2.90)	.261	-0.24 (0.62)	.696	-	-	2.28 (1.25, 4.17)	.007
Interaction	-	-	-	-	-1.58 (0.93)	.091	-	-	-	-	0.57 (0.67)	.389	-	-	-	-	0.38 (0.55)	.491	-	-	-	-	-0.11 (0.63)	.865	-	-	-	-
Covariates:																												
Nonwhite Race/Ethnicity	1.33 (0.97, 1.83)	0.082	1.30 (0.93, 1.80)	.122	0.28 (0.17)	.099	0.55 (0.35, 0.84)	.006	0.54 (0.35, 0.83)	.005	-0.62 (0.22)	.004	0.64 (0.48, 0.87)	.004	0.64 (0.48, 0.86)	.003	-0.45 (0.15)	.003	0.63 (0.38, 1.03)	.065	0.63 (0.39, 1.02)	.062	-0.46 (0.25)	.067	0.64 (0.48, 0.87)	.004	0.64 (0.48, 0.86)	.003
Upper Grades*	0.68 (0.49, 0.94)	.020	0.69 (0.50, 0.96)	.027	-0.37 (0.17)	.027	1.41 (1.07, 1.86)	.015	1.43 (1.08, 1.89)	.014	0.35 (0.15)	.015	1.43 (1.03, 1.98)	.033	1.41 (1.02, 1.96)	.039	0.36 (0.17)	.032	0.51 (0.34, 0.77)	.001	0.51 (0.34, 0.76)	.001	-0.68 (0.21)	.001	1.43 (1.03, 1.98)	.033	1.41 (1.02, 1.96)	.039

* Upper Grades = 11th and 12 grades (referent = 9th and 10th grades)