



For reprint orders, please contact: reprints@futuremedicine.com

Overview of current immunotherapeutic strategies for glioma

In the last decade, numerous studies of immunotherapy for malignant glioma (glioblastoma multiforme) have brought new knowledge and new hope for improving the prognosis of this incurable disease. Some clinical trials have reached Phase III, following positive outcomes in Phase I and II, with respect to safety and immunological end points. Results are encouraging especially when considering the promise of sustained efficacy by inducing antitumor immunological memory. Progress in understanding the mechanisms of tumor-induced immune suppression led to the development of drugs targeting immunosuppressive checkpoints, which are used in active clinical trials for glioblastoma multiforme. Insights related to the heterogeneity of the disease bring new challenges for the management of glioma and underscore a likely cause of therapeutic failure. An emerging therapeutic strategy is represented by a combinatorial, personalized approach, including the standard of care: surgery, radiation, chemotherapy with added active immunotherapy and multiagent targeting of immunosuppressive checkpoints.

Keywords: clinical trials • dendritic cells • gene therapy • glioma • immune checkpoints blockade • immunotherapy • vaccination

It has long been known that immunosuppressive treatment regimens or diseases accompanied by an immunosuppressive state are associated with increased incidence of malignancy [1] and that tumors progress more slowly and can even be rejected when an immune response is elicited [2,3]. These observations have led to formulating the concept of immune surveillance [4,5], stipulating that immune mechanisms are responsible for the continuous monitoring and elimination of cells displaying neoplastic mutations. Cancer cells, glioblastoma cells included, activate mechanisms to evade immune surveillance through immunosuppressive cytokines and cells, which act upon all aspects of the immune defense to hinder the recognition and immune eradication of tumor cells.

In the context of the tumor-induced immunosuppressive environment, CD4⁺ T-lymphocytes either do not recognize the tumor antigens, recognize them but become

anergic or differentiate into Tregs, which suppress the immune response further [6]. In patients with glioma, characterized by generalized lymphopenia, the proportion of CD4⁺FoxP3⁺ Tregs has been shown to be markedly increased and to correlate with a decreased proliferative capacity of CD4⁺ T cells. Depletion of Tregs restored proliferation of CD4⁺ T cells and this was accompanied by a decrease in Th-2 (IL-4 and IL-10) and an increase of Th-1 (IL-6, IL-2, TNF- α and IFN- γ) cytokines [7]. CD8⁺ cytotoxic T cells (CTLs) are functionally impaired either directly by tumor cells or indirectly through inflammatory molecules in the tumor microenvironment or through interactions with altered antigen-presenting cells (APCs), Tregs or myeloid-derived suppressor cells [8]. Decreased expression of HLA class I molecules by glioma cells, leading to impaired antigen presentation and lysis by CTLs, has been correlated with increased

Anda-Alexandra Calinescu¹,
Neha Kamran¹, Gregory
Baker¹, Yohei Mineharu²,
Pedro Ricardo Lowenstein^{1,3}
& Maria Graciela Castro^{*1,3}

¹Department of Neurosurgery, University of Michigan School of Medicine, Ann Arbor, MI 48109, USA

²Department of Neurosurgery, Kyoto University, Kyoto, Japan

³Department of Cell & Developmental Biology, University of Michigan, Ann Arbor, MI 48109, USA

*Author for correspondence: mariagcastro2010@gmail.com

Table 1. Summary of preclinical studies of dendritic cell vaccination previously reported in the literature.

Study (year)	Cytokines used for DC differentiation	Antigen loading on DC	Vaccine injection site	Adjuvant therapy
Liau (1999)	GM-CSF, IL-4	Acid-eluted tumor peptide	s.c.	None
Akasaki (2001)	GM-CSF, IL-4	DC chemically fused with irradiated tumor cells	s.c.	rIL-12 (i.p, d5, 7, 9, 11, 13, 15, 17, 19)
Aoki (2001)	GM-CSF	Sonicated tumor cells	i.p.	SUV (small unilaminar vesicle, <i>in vitro</i>)
Ni (2001)	GM-CSF with myc and raf	Irradiated tumor cells (apoptosis)	i.p.	None
Insug (2002)	GM-CSF, IL-4	RNA (transfection)	i.p.	rIL-12 (i.p., d1–5)
Witham (2002)	GM-CSF, IL-4, Flt3L	UCN-01-treated tumor cells (apoptosis)	s.c.	None
Yamanaka (2002)	GM-CSF, IL-4	None	i.t.	SFV-mediated IL-12
Prins (2003)	GM-CSF, IL-4	hgp100/TRP-2 peptides	i.d.	None
Takagi (2003)	GM-CSF, IL-4	DC chemically fused with irradiated tumor cells	s.c.	None
Driessens (2004)	GM-CSF	Irradiated tumor cells (apoptosis)	s.c.	GM-CSF-secreting tumor
Herrlinger (2004)	(GM-CSF, IL-4)	Irradiated, GM-CSF and IL-4 transfected GL261	s.c.	GM-CSF, IL-4
Saito (2004)	GM-CSF, IL-4	Sonicated tumor cells	i.t., s.c.	pSV2mulFN- β (d7)
Zhang (2004)	GM-CSF, IL-4	RNA (coincubation)	s.c.	None
Zhu (2005)	GM-CSF, IL-4	Heat-treated tumor cells (apoptosis)	s.c.	None
Pellegatta (2005)	GM-CSF, IL-4	Freeze-thawed tumor cells (necrosis)	s.c.	None
Kjaergaard (2005)	GM-CSF, IL-4	DC electronically fused with irradiated tumor cells	intrasplenic	OX40R mAb
Kuwashima (2005)	GM-CSF, IL-4	Irradiated tumor cells with IL-4 transfection (apoptosis)	i.t., s.c. (DC, tumor)	IFN- α -producing DC, IL-4-secreting tumor
Clavreul (2006)	(GM-CSF)	Irradiated tumor cells (apoptosis)	s.c.	GM-CSF
Kim (2006)	GM-CSF, IL-4	Freeze-thawed tumor cells (necrosis)	s.c.	rIL-12-producing DC
Jouanneau (2006)	GM-CSF, IL-4, Flt3L; Ribomunyl, IFN- γ	Freeze-thawed tumor cells (necrosis)	i.p.	None
Ciesielski (2006)	GM-CSF	Human survivin (transfection)	s.c.	None
Cho (2007)	GM-CSF, IL-4; TNF- α , PGE2	Survivin fused to protein transduction domain of HIVtat	s.c.	None
Kim (2007)	GM-CSF, IL-4	TAT-survivin	s.c.	TMZ (d3–6, i.p.)
Amano (2007)	GM-CSF, IL-4	RHAMM-mRNA (transfection)	i.p.	None
Kim (2007)	GM-CSF, IL-4	mTERT-mRNA (transfection)	i.m.	None
Grauer (2007)	GM-CSF, IL-4	Freeze-thawed tumor cells (necrosis)	s.c.	Anti-CD25 mAbs
Ciesielski (2008)	GM-CSF with myc and raf	Survivin 53–67 peptide	s.c.	None
Xu (2009)	GM-CSF, IL-4	Irradiated tumor cells (apoptosis)	s.c.	None

DC: Dendritic cell; Flt3L: Fms-like tyrosine kinase 3 ligand; GCV: Gancyclovir; GM-CSF represents granulocyte macrophage colony-stimulating factor; i.d.: Intradermal; IL: Interleukin; i.p.: Intraperitoneal; i.t.: Intratumoral; LPS: Lipopolysaccharide; mAb: Monoclonal antibody; s.c.: Subcutaneous; TK: Thymidine kinase; TMZ: Temozolomide.

Table 1. Summary of preclinical studies of dendritic cell vaccination previously reported in the literature (cont.).

Cell line	Host	Tumor injection site	Treatment initiation	Day of treatment	Survival >60 days	Ref.
9L	Fischer 344 rats	Intracranial	7	7, 14, 21	30%	[55]
SR-B10.A	B10.A mice	Intracranial	5	5, 12	50%	[37]
GL261	C57BL/6 mice	Intracranial	7	7, 14, 21	80% (d28)	[39]
GL261	C57BL/6 mice	Intracranial	3	3, 7, 10, 14	40%	[56]
GL261	C57BL/6 mice	Intracranial	-21	-21, -14, -7	100%	[48]
9L	Fischer 344 rats	Intracranial	0	0, 1, 4	0%	[61]
B16	C57BL/6 mice	Intracranial	7	7, 14, 21	40%	[63]
GL26	C57BL/6 mice	Intracranial	0	0	75%	[58]
SR-B10.A	B10.A mice	Flank	0	0, 7	40% (40d)	[60]
9L	Fischer 344 rats	Intracranial	4	4, 11, 18	60%	[44]
GL261	C57BL/6 mice	Intracranial	3	3, 10	100%	[47]
GL261	C57BL/6 mice	Intracranial	3	3	50%	[59]
G422	Kuming mice	Intracranial	4	4, 11, 18	0%	[64]
C6	Wister rats	Intracranial	ND	Five-times weekly	0%	[65]
GL261	C57BL/6 mice	Intracranial	0	0,7,14	75%	[57]
GL261	C57BL/6 mice	Intracranial	7	7	80%	[53]
GL261	C57BL/6 mice	Intracranial	4	4 (d3, 5, 7 for tumor vaccine)	55%	[54]
F98	Fischer 344 rats	Intracranial	4	4	0% (n.s.)	[43]
GL26	C57BL/6 mice	Intracranial	-21	-21, -14, -7	70%	[50]
GL26	C57BL/6 mice	Intracranial	-14	-14, -7	37.50%	[49]
GL261	C57BL/6 mice	Intracranial	-4	-4, +3, +10	0%	[41]
GL26	C57BL/6 mice	Flank	10	10	75%	[40]
GL26	C57BL/6 mice	Intracranial	13	13	40%	[52]
KR158B	C57BL/6 mice	Intracranial	3	3, 10	15%	[38]
GL26	C57BL/6 mice	Intracranial	-14	-14, -7	85%	[51]
GL261	C57BL/6 mice	Intracranial	-14	-14, -7	80%	[46]
GL261	C57BL/6 mice	Intracranial	4	4, 11, 18	25%	[42]
9L	Fischer 344 rats	Intracranial	7	7, 4, 21	30%	[62]

DC: Dendritic cell; Flt3L: Fms-like tyrosine kinase 3 ligand; GCV: Gancyclovir; GM-CSF represents granulocyte macrophage colony-stimulating factor; i.d.: Intradermal; IL: Interleukin; i.p.: Intraperitoneal; i.t.: Intratumoral; LPS: Lipopolysaccharide; mAb: Monoclonal antibody; s.c.: Subcutaneous; TK: Thymidine kinase; TMZ: Temozolomide.

Table 1. Summary of preclinical studies of dendritic cell vaccination previously reported in the literature (cont.).

Study (year)	Cytokines used for DC differentiation	Antigen loading on DC	Vaccine injection site	Adjuvant therapy
Fujita (2009)	GM-CSF; LPS; poly-ICLC, IFN- γ , IFN- α , IL-4	Garc1 ₇₇₋₈₅ + EphA2 ₆₈₂₋₆₈₉	i.t., s.c.	None
Maes (2009)	GM-CSF	RNA of tumor cells (electroporation)	s.c.	Anti-CD25 mAbs (day-21)
Jiang (2009)	GM-CSF, IL-4	Freeze-thawed tumor cells (necrosis)	s.c.	CXCL10/IP-10 (d5,10,15)
Saka (2010)	GM-CSF, IL-4	IL13ra2-mRNA (transfection)	s.c.	None
Kim (2010)	GM-CSF, IL-4	TMZ and radiation-treated tumor cells (apoptosis)	s.c.	TMZ (d2-6, i.p.)
Xiao (2011)	GM-CSF, IL-4	RNA of tumor stem cells (transfection)	s.c.	None
Mineharu (2011)	hFlt3L, IL-6	Ad-TK/GCV-treated tumor cells (autophagy/apoptosis)	i.t., s.c.	CpG2006 (s.c., d10), Ad-TK/Flt3L (i.t., d10)
Feng (2012)	GM-CSF, IL-4; TNF- α , LPS	Freeze-thawed tumor cells (necrosis)	s.c.	T cell (tail vein, d3)
Zhang (2013)	GM-CSF, IL-4	Freeze-thawed tumor cells (necrosis)	i.p.	COX-2 inhibitor
Peng (2014)	GM-CSF, IL-4	Ad-CPEB4 (transfection)	s.c.	None
Bu (2015)	GM-CSF, IL-4; TNF- α	Exosome from DCs pulsed with chaperon-rich tumor lysate	i.d.	None
Li (2015)	GM-CSF	EphA1-PE38	i.t.	None

DC: Dendritic cell; Flt3L: Fms-like tyrosine kinase 3 ligand; GCV: Gancyclovir; GM-CSF represents granulocyte macrophage colony-stimulating factor; i.d.: Intradermal; IL: Interleukin; i.p.: Intraperitoneal; i.t.: Intratumoral; LPS: Lipopolysaccharide; mAb: Monoclonal antibody; s.c.: Subcutaneous; TK: Thymidine kinase; TMZ: Temozolomide.

grade of malignancy [9]. Also, decreased expression of the costimulatory molecule B7 leads to poor costimulation and T-cell anergy [10], whereas high expression of B7-H1 (PD-L1) on tumor cells inhibits the function of CD4⁺ and CD8⁺ cells [11] and induces apoptosis of CTLs [12]. Glioma cells also express soluble Fas ligand, responsible for the death of antigen-stimulated CTLs [13]. Myeloid-derived suppressor cells, a heterogeneous population of immature myeloid cells, have also been shown to accumulate in the blood of glioma patients and inhibit T-cell function, effect mediated through their production of arginase 1 [14,15]. Natural killer (NK) cells are unable to activate their cytotoxic mechanisms in the absence of activating receptors on tumor cells [16-19], which are downregulated in glioma patients [20]. Also, glioma cells express surface proteins, like galectin-1, which inhibit NK-mediated immune surveillance [21]. Finally, type II NK T cells (NKT cells) contribute to the immunosuppressive tumor microenvironment through secretion of anti-inflammatory cytokines like TGF- β and IL-13 [22]. It is now evident that successful immunotherapy for glioma

needs to address the mechanisms of tumor-induced immune suppression in addition to being mindful of the unique environment of the brain, which, unlike other organs, has minimal tolerance for inflammation.

Glioblastoma (GBM [WHO grade IV]) is the deadliest and most common form of glioma (0.59-3.69 new cases in 100,000 every year) [23]. The current standard of care (SOC): surgery, radiation and temozolomide [24] can only offer patients a median survival of 14.6 months after diagnosis and a 5-year survival rate of 0.05-4.7% [25]. Interestingly, an increasing number of studies (reviewed by [25]) indicate that an overactive immune system, as found in asthma, hay fever, eczema and food allergies, reduces the risk of developing GBM, suggesting that immune therapies for GBM could be a successful avenue to improve patient outcome. Recent accumulating evidence has highlighted the heterogeneous and evolving nature of GBM [26-29], and suggests that this heterogeneity represents a key to treatment failure. Harnessing the power of the dynamic, versatile and continuously adapting immune system to aid in the treatment of a

Table 1. Summary of preclinical studies of dendritic cell vaccination previously reported in the literature (cont.).

Cell line	Host	Tumor injection site	Treatment initiation	Day of treatment	Survival >60 days	Ref.
GL261	C57BL/6 mice	Intracranial	0	0, 10	75%	[45]
GL261	C57BL/6 mice	Intracranial	-7	-7	100%	[66]
GL261	C57BL/6 mice	Intracranial	1	1, 8, 15	60%	[67]
KR158B	C57BL/6 mice	Intracranial	3	3, 10	0%	[68]
GL26	C57BL/6 mice	Intracranial	4	4, 11, 18	70%	[69]
9L	Fischer 344 rats	Intracranial	3	3, 10, 17	0%	[70]
CNS1/9L	Lewis/Fisher rat	Intracranial	10/9	10, 17, 24/ 9, 16, 23	90%/ 83.3%	[71]
C6	Wister rats	Intracranial	3	3, 7, 14	50%	[72]
C6	Wistar rats	Intracranial	3	3, 10, 17	50%	[73]
GL261	C57BL/6 mice	Flank	7	7, 14	Not described	[74]
GL261	C57BL/6 mice	Intracranial	3	3, 6, 9, 12 (every 3 days)	60%	[75]
C6	Wistar rats	Intracranial	10	10,13,16,19 (every 3 days)	Not given	[76]

DC: Dendritic cell; Flt3L: Fms-like tyrosine kinase 3 ligand; GCV: Gancyclovir; GM-CSF represents granulocyte macrophage colony-stimulating factor; i.d.: Intradermal; IL: Interleukin; i.p.: Intraperitoneal; i.t.: Intratumoral; LPS: Lipopolysaccharide; mAb: Monoclonal antibody; s.c.: Subcutaneous; TK: Thymidine kinase; TMZ: Temozolomide.

moving target, like GBM, represents a challenging but worthwhile pursuit.

Numerous preclinical studies have demonstrated that several immunotherapeutic strategies can be successful in animal models of GBM, including: gene therapy [30], passive immunotherapy with antibodies against tumor antigens [31], adoptive T-cell transfer with T cells activated against tumor antigens or engineered to express chimeric antigen receptors (CARs) [32–34], immunomodulatory strategies aimed at inhibiting the immune checkpoints used by tumors to escape from immune surveillance [35,36] as well as active immunotherapy, employing peptide or dendritic cell (DC) vaccines (summarized in **Table 1** [37–76]), to elicit immune reactivity against tumors and induce immunological memory capable of preventing recurrence of the disease.

Taken into the clinic, many Phase I and II clinical trials for GBM using immunotherapy in combination with SOC have come to completion in the last 5 years. Results have shown that immunological approaches are generally safe, with minimal side effects and able

to elicit specific immune responses and in some cases improve progression-free survival (PFS) and overall survival (OS) [77–85]. Studies employed gene therapy [86–93], DC vaccines, which vary primarily in the agents used to prime the DCs for antigen presentation: either GBM-associated antigens (GAA) [81,94], autologous tumor lysates [78,82,94–97] or RNA from GBM stem cells [85], with or without adjuvants aimed to activate Toll-like receptors (TLRs). A few trials tested the effect of antigenic stimulation with peptides or tumor cells [77,79,98–102] and some analyzed the effect of autologous T-cell transfer on eliciting an antiglioma immune response [103,104] or the effect of specific antibodies against GBM receptors or to deplete Tregs [105,106].

A growing interest for immunotherapeutic approaches for GBM is illustrated by the increasing annual number of funded clinical trials worldwide (ClinicalTrials.org database, **Figure 1**). Most of the trials involved active immunotherapy using different vaccination strategies or gene therapy. Following promising results in the treatment metastatic melanoma, where antibodies against CTLA-4 and PD-1

have shown an increase in OS [107], or induce tumor regression [108], an important new addition to the anti-GBM toolbox is represented by drugs targeting immunosuppressive checkpoints. It is currently thought that immunotherapeutic strategies most likely to succeed will entail a combination of active vaccination and immune checkpoint inhibition [109]. In this review, the authors highlight advances in immune treatment strategies for GBM during the past 5 years and present some of the ongoing challenges and future perspectives in the field. The authors apologize to all scientists whose work they were unable to mention due to limitations in space. New insights will soon be brought about by the results of open clinical trials testing immunotherapy regimens for GBM (Table 2).

Targeting immunosuppressive checkpoints

Immune checkpoints are negative-regulatory signaling mechanisms responsible for maintaining self tolerance and preventing autoimmune reactions, which attenuate the strength and duration of normal T-cell-mediated immune responses. It has become apparent that diverse cancers, including GBM, co-opt the physiological function of immune checkpoints to greatly diminish T-cell-mediated antitumor immunity [11,36,110–111].

Two major checkpoints have been identified as immune escape mechanisms in both rodent and human cancers: CTLA-4/CD152 and PD-1/CD279 [110]. Both attenuate T-cell activation and promote T-cell

anergy, however, they differ in their spatial and temporal activity. CTLA-4 is a powerful inhibitory T-cell receptor which preferentially binds to B7.1/CD80 and B7.2/CD86, ligands expressed on the surface of APCs, precluding their binding to the T-cell costimulatory receptor CD28 and thus inhibiting T-cell proliferation and cytokine production [112]. The CTLA-4 immune checkpoint occurs early in the immune response, during the priming phase and acts primarily within secondary lymphoid organs. Conversely, PD-1 signaling takes place directly within the tumor microenvironment, during the effector phase, by interacting with one of two currently identified PD-1 ligands: PD-L1/B7-H1/CD274 [113] or PD-L2/B7-DC/CD273 [112] expressed on the surface of cancer cells. Engagement of PD-1 ligands with the PD-1 T-cell receptor also leads to T-cell inhibition by blocking cell proliferation and inhibiting cytokine production [113,114]. CTLA-4 and PD-1 are both commensurately upregulated at the T-cell surface in response to proinflammatory cues. A diagrammatic summary of the CTLA-4 and PD-1 immune checkpoints is shown in Figure 2.

Cytotoxic T-lymphocyte-associated protein 4

A few human monoclonal antibodies targeting CTLA-4 (e.g., ipilimumab and tremelimumab) have been evaluated for safety and efficacy in human cancer patients: ipilimumab has been shown to improve OS in patients with metastatic melanoma in a Phase III




Figure 1. Timeline of clinical trials for glioma using immunotherapy. A search for ‘immunotherapy’ and ‘glioma’ in the ClinicalTrials.gov database (March 2015) yields a list of 61 clinical trials: 14 with dendritic cell vaccines, 9 testing synthetic peptide vaccines, 8 using autologous T-cell transfer, 6 gene therapy, 4 with tumor cell lysate vaccine combined with T-cell transfer, 2 with autologous NK or NKT cell transfer, 2 with allogeneic T-cell transfer, 4 targeting immunosuppressive checkpoints and 7 using other immune treatment strategies. Limitations of the search engine may not allow a comprehensive listing; nonetheless, the graph illustrates the extensive interest in anti-glioma vaccines and an emerging trend of testing immunosuppressive checkpoints. NK: Natural killer; NKT: NK T cell.

clinical trial [107]; however, another Phase III trial using tremelimumab failed to show significant survival advantage [115]. Several Phase I and II studies of solid tumors using these antibodies show promise of improved PFS [116] but can also elicit severe immune adverse effects [117].

Preclinical investigations using anti-CTLA-4 antibodies against primary brain cancers have demonstrated significant increases in animal survival. For example, Fecci *et al.* have shown that administration of anti-CTLA-4 antibodies results in long-term survival in 80% of immunocompetent mice bearing syngeneic SMA-560 intracranial tumors [111]. The treatment normalized systemic CD4⁺ T-cell counts and decreased the number of Tregs (CD4⁺/CD25⁺/Foxp3⁺/GITR⁺) [111]. Intratumoral administration of IL-12 in combination with Anti-CTLA-4 antibodies leads to the eradication of intracranial GL261 gliomas, increasing the number of CD8⁺ effector cells and reducing Foxp3⁺ Tregs within the tumor microenvironment [118]. Also, vaccination with GM-CSF-expressing glioma cells, when combined with anti-CTLA-4 antibodies, has been shown to be more effective against established murine GL261 intracranial tumors than either treatment alone [119].

Programmed cell death protein 1

Promising results in the treatment of metastatic melanoma and other solid tumors have been demonstrated in many trials using the anti-PD-1 human antibodies: nivolumab and pembrolizumab [108,120,121]. Preclinical glioma studies using anti-PD-1 antibodies have also shown effectiveness. The combination of anti-PD-1 antibodies and radiotherapy has been shown to double median survival and elicit long-term survival in 15–40% of mice bearing GL261 gliomas compared with either treatment alone [122]. The authors show that the tumor microenvironment was infiltrated by CD8⁺/IFN- γ ⁺/TNF- α ⁺ CTLs along with reductions in tumor-infiltrating CD4⁺/Foxp3⁺ Tregs, thus suggesting that the mechanisms targeted by immune checkpoint blocking antibodies in animal studies are similar to those in human clinical trials. Recently, an investigation by Wainwright *et al.* has demonstrated that combinatorial targeting of immune checkpoints in the murine GL261 glioma model is more effective than single agent treatment, a strategy which could carry high potential value for future clinical trials for GBM [123]. The establishment of long-term antitumor immunological memory in many of these preclinical studies as demonstrated by the failure of tumors to grow in response to tumor rechallenge suggests a potential added benefit of immune checkpoint blockade in the prevention of tumor recurrence.

A few clinical trials have recently started to test the safety and efficacy of immune checkpoint blockade in GBM patients. A randomized Phase III study is aimed to test nivolumab versus bevacizumab in adult patients with recurrent GBM (CheckMate 143, ClinicalTrials Identifier: NCT02017717). One of the arms of this trial will test the combination therapy of nivolumab and ipilimumab. Three Phase I/II studies will analyze pembrolizumab with or without bevacizumab (NCT02337491) or pembrolizumab in combination with MRI-guided laser ablation (NCT02311582) in patients with recurrent GBM and will test the effect of anti-PD-1 antibody, pidilizumab against diffuse intrinsic pontine glioma and recurrent GBM (NCT01952769). MEDI4736, a human anti-PD-L1 antibody, is currently being tested in combination with radiotherapy and bevacizumab in the treatment of GBM (NCT02336165). A kinase inhibitor for TGF β RI (galunisertib) will be evaluated in combination with nivolumab in several advanced solid tumors, including GBM in a Phase Ib/II safety study (NCT02423343).

Enthusiasm over the use of immune checkpoint blockade as a powerful immunotherapeutic strategy in the fight against cancer has been undermined by a relatively high frequency of immune-related adverse effects in the form of gastrointestinal, dermatological, hepatic and endocrinological toxicities [124] which, in extreme cases, have led to treatment-related death [125,126]. It is thought that immune-related adverse effects associated with immune checkpoint blockade are due to aberrant infiltration of activated CD4⁺ and CD8⁺ T cells into normal tissues together with elevated levels of proinflammatory cytokines [127]. One study found grade 3–4 toxicities in 41 of 296 patients, with three treatment-related deaths attributed to pneumonitis in response to treatment to the PD-1 inhibitor nivolumab [121]. Newer agents targeting cognate PD-1 ligands (PD-1Ls) have now been tested in NSCLC, renal cell cancer and melanoma (NCT00729664) and show durable tumor regression with less grade 3 or 4 adverse events compared with anti-CTLA-4 and anti-PD-1 blockade [128].

Immune stimulatory gene therapy

Clinical trials targeting gliomas with gene therapy started in the early 1990s [93] and have employed a variety of vehicles to deliver genes, such as viral vectors, synthetic nanoparticles and liposomes, neural and mesenchymal or embryonic stem cells [87,88,92,93,129]. These gene therapeutic approaches have aimed to introduce suicide genes or oncolytic viruses into the tumor cells and/or to induce the expression of immunomodulatory cytokines to aid the antitumor immune response. More recently, genetic techniques have been used to

Table 2. Open clinical trials of immunotherapy in glioblastomas.

NCT number, date first received, category	Phase	Study title	Diagnosis
NCT00045968 September 17 2002 DC vaccine	III	DCVax®-L to treat newly diagnosed GBM	Newly diagnosed GBM
NCT00626483 February 28 2008 DC vaccine	I	Basiliximab in treating patients with newly diagnosed glioblastoma multiforme undergoing targeted immunotherapy and temozolomide-caused lymphopenia	GBM seropositive or seronegative for CMV
NCT00634231 5 March 2008 Gene therapy	I	A Phase I study of AdV-tk + prodrug therapy and radiation therapy for pediatric brain tumors	GBM and ependymoma
NCT01326104 22 July 2010 Autologous T cells w DC vaccine	I/II	Vaccine immunotherapy for recurrent medulloblastoma and primitive neuroectodermal tumor	Medulloblastoma, neuroectodermal tumor
NCT0140067 19 July 2011 Tumor lysate vaccine	I	Imiquimod/brain tumor initiating cell (BTIC) vaccine in brain stem glioma	Diffuse intrinsic pontine glioma
NCT01657734 7 September 2011 Other		Multimodal MR imaging in patients with glioblastoma treated with dendritic cell therapy	GBM following immunotherapy
NCT01454596 6 October 2011 Autologous T cells ahIL-2 and chemotherapy	I	CAR T-cell receptor immunotherapy targeting EGFRvIII for patients with malignant gliomas expressing EGFRvIII	EGFRvIII positive glioma and gliosarcoma with failed prior radiotherapy with or without prior chemotherapy
NCT01522820 25 January 2012 Autologous T cells with DC vaccine	I	A Phase I clinical trial of mTOR inhibition with rapamycin for enhancing intranodal DC vaccine-induced antitumor immunity in patients with NY-ESO-1 expressing solid tumors	NY-ESO-1 expressing solid tumors

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

Table 2. Open clinical trials of immunotherapy in glioblastomas (cont.).

Intervention	Outcome measures			
	Safety Toxicity	Efficacy	Immunologic analyses	Other
DC vaccine made with dendritic cells differentiated from the patients own PBMC primed with autologous tumor lysate	Y	PFS and OS	Analyze immune response	
pp65-lysosomal-associated membrane protein (LAMP) mRNA-loaded DCs with GM-CSF in combination with basiliximab, an anti CD25 monoclonal antibody used to deplete regulatory T cells	Y	PFS	Functional capacity of T-reg cells, vaccine-induced cellular or humoral immune responses development of autoimmunity, phenotype of NK cells	Characterize immunologic infiltrates in recurrent tumors
AdV-tk + valacyclovir in combination with standard of care radiation	Y	PFS, OS	Analyze immunologic function	
One time iv. administration of autologous <i>ex vivo</i> -expanded autologous T-cell transfer (TTRNA-xALT) in combination with 3 weekly doses tumor RNA-loaded DCs (TTRNA-DCs)	Y	PFS, OS	Magnitude and persistence of antitumor humoral and cellular immunity, cytokine production, profile of lymphocytes	Identify potential tumor-specific antigens as vaccine candidates
BTIC (brain tumor initiating cells) lysate vaccine and the TLR7 agonist Imiquimod in combination with radiation therapy	Y	PFS		
Observational	N	N	N	MRI spectroscopy, perfusion imaging and diffusion imaging to characterize inflammatory response, metabolism and tissue structure
Autologous T-cell transfer with retrovirally transduced PBLs with EGFRvIII CAR, combined with IL-2 and the chemotherapeutics: fludarabine and cyclophosphamide	Y	PFS	N	
Intranodal administration of the CDX-1401 vaccine (a fusion protein consisting of a fully human monoclonal antibody with specificity for the dendritic cell receptor DEC-205 linked to the NY-ESO-1 tumor antigen) with or without the mTOR inhibitor: rapamycin	Y	PFS	NY-ESO-1-specific cellular and humoral immunity	

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

Table 2. Open clinical trials of immunotherapy in glioblastomas (cont.).

NCT number, date first received, category	Phase	Study title	Diagnosis
NCT01567202 26 March 2012 DC vaccine	III	A triple-blind randomized clinical study of vaccination with dendritic cells loaded with glioma stem-like cells-associated antigens against brain GBM	GBM
NCT01702792 3 October 2012 Other	I	Vaccination of patients with newly diagnosed GBM using autologous tumor lysate and montanide emulsion for derivation of tumor-specific hybridomas	Newly diagnosed GBM
NCT01678352 30 August 2012 Tumor lysate vaccine	0	Imiquimod and tumor lysate vaccine immunotherapy in adults with high risk or recurrent grade II gliomas	WHO grade II glioma
NCT01795313 12 February 2013 Peptide vaccine		Immunotherapy for recurrent ependymomas in children using HLA-A2-restricted tumor antigen peptides in combination with imiquimod	Recurrent ependymoma in HLA-2A-positive pediatric patients
NCT01811992 13 February 2013 Gene therapy	I	Combined cytotoxic and immune-stimulatory therapy for glioma	Malignant glioma, GBM
NCT01952769 15 September 2013 Immunosuppressive checkpoint blockade	I/II	A Phase I/II clinical trial of CT-011 (pidilizumab) in diffuse intrinsic pontine glioma and relapsed glioblastoma multiforme	DIPG, relapsed GBM
NCT01956734 27 September 2013 Gene therapy	I	Virus DNX2401 and temozolomide in recurrent glioblastoma	GBM recurrent tumor
NCT02017249 7 November 2013 Other	I	Double-blinded randomized placebo-controlled trial analyzing the efficacy of oral arginine to improve immune function in GBM	GBM
NCT02010606 2 December 2013 DC vaccine	I	Phase I trial of vaccination with autologous DCs pulsed with lysate derived from an allogeneic glioblastoma stem-like cell line for patients with newly diagnosed or recurrent glioblastoma	Newly diagnosed or recurrent GBM
NCT02017717 17 December 2013 Immunosuppressive checkpoint blockade	III	A randomized study of nivolumab versus bevacizumab and a safety study of nivolumab in adult subjects with recurrent GBM (CheckMate 143)	Recurrent GBM
NCT02049489 26 January 2014 DC vaccine	I	A study of ICT-121 dendritic cell vaccine in recurrent glioblastoma	GBM

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

Table 2. Open clinical trials of immunotherapy in glioblastomas (cont.).

Intervention	Outcome measures			
	Safety Toxicity	Efficacy	Immunologic analyses	Other
SOC (surgery, radiation, temozolomide) with or without DC vaccine generated from PBMC primed with stem-like cell-associated antigens (SAA) derived from the patients' own tumor	N	PFS, OS	N	
Derivation of tumor-specific hybridomas	Y	N	N	Number of hybridoma clones that produce glioma-specific antibodies
BTIC lysate vaccine and the TLR7 agonist imiquimod	Y	N	Induction of BTIC lysate-specific T-cell response	
Vaccine with HLA-2A restricted peptides and imiquimod as adjuvant	Y		Tumor-associated antigen-specific T-cell responses	
Dose escalation of Ad-hCMV-TK and Ad-hCMV-Flt3L	Y	PFS, postoperative survival	Adenoviral antibodies	MRI and clinical assessment
Administration of CT-011 (pidilizumab, anti PD-L1 antibody) in patients with DIPG or recurrent GBM	Y	PFS, OS, ORR	N	Response on imaging and neurological examination
Stereotaxic, intratumoral injection of the oncolytic virus DNX2401 in combination with temozolomide	Y	PFS, OS12	Immunogenicity	Biomarkers and tumor genetics
Arginine supplement will be administered orally three times per day for 7 days before surgery and 7 days after surgery			Change in immune function (>25% increase in functional response of peripheral T cells)	
SOC with DC vaccine primed with autologous tumor cell lysate and optional bevacizumab	Y		Immune response: T-cell activity	Tumor stem cell antigen expression
Administration of nivolumab (anti PD-L1 antibody) versus bevacizumab and nivolumab or nivolumab in combination with ipilimumab (anti-CTLA antibody) in adult subjects with recurrent GBM	Y	OS, PFS, ORR	N	
Repeated administration of ICT-121 DC vaccine prepared from autologous DCs pulsed with purified CD133 peptides to target GSC	Y	ORR OS, PFS	Cytotoxic T-cell response to the ICT-121 vaccine epitopes	Antigen expression, culture of CD133 ⁺ neurospheres

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

Table 2. Open clinical trials of immunotherapy in glioblastomas (cont.).			
NCT number, date first received, category	Phase	Study title	Diagnosis
NCT02060955 10 February 2014 Autologous T cells	II	Randomized Phase II multicenter study to investigate efficacy of autologous lymphoid effector cells specific against tumor cells (ALECSAT) in patients with GBM measured compared with avastin/irinotecan	GBM
NCT02078648 14 February 2014 Peptide vaccine	I/II	Safety and efficacy study of SL-701, a glioma-associated antigen vaccine to treat recurrent GBM	Adult GBM
NCT02122822 23 April 2014 Tumor lysate vaccine	I	Research for immunotherapy of glioblastoma with autologous heat shock protein gp96	Newly diagnosed supratentorial glioma
NCT02193347 2 July 2014 Peptide vaccine	I	Patients with IDH1-positive recurrent grade II glioma enrolled in a safety and immunogenicity study of tumor-specific peptide vaccine	IDH1-positive recurrent grade II glioma
NCT02197169 20 July 2014 Gene therapy	Ib	DNX-2401 with Interferon gamma (IFN- γ) for recurrent glioblastoma or gliosarcoma brain tumors	Glioblastoma, gliosarcoma
NCT02208362 1 August 2014 Autologous T cells	I	Genetically modified T cells in treating patients with recurrent or refractory malignant glioma	Refractory malignant glioma
NCT02311582 4 December 2014 Immunosuppressive checkpoint blockade	I/II	A Phase I and open label, randomized, controlled Phase II study testing the safety, toxicities and efficacy of MK-3475 in combination with MRI-guided laser ablation in recurrent malignant gliomas	Malignant glioma

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

Intervention	Outcome measures			
	Safety Toxicity	Efficacy	Immunologic analyses	Other
Autologous T-cell transfer with <i>ex vivo</i> antitumor activated and expanded lymphoid effector cells (ALECSAT administered at week 4, 9, 14, 26 and 46) compared with standard bevacizumab/irinotecan treatment	Y	OS, PFS	N	
Multivalent SL-701 synthetic peptide vaccine in combination in combination with GM-CSF and imiquimod (TLR7 agonist)	Y	PFS/OS	N	
Heat shock protein gp96-peptide complex made from a person's tumor cells	Y	PFS/OS	Antigen-specific T-cell responses (ELISPOT)	
PEPIDH1M vaccine (a peptide vaccine derived from the mutated IDH1R132H) with tetanus-toxoid preconditioning and temozolomide	Y	N	IFN ELISPOT on peptide-stimulated lymphocytes	
Modified adenovirus DNX-2401, an oncolytic adenovirus with or without IFN	Y	PFS	Immunological and biological effects after DNX-2401 with interferon gamma	
Intratumoral, intracavitary autologous T-cell transfer with central memory enriched T cells transduced with lentivirus to express an IL13R α 2-specific, hinge-optimized, 41BB-costimulatory chimeric receptor and a truncated CD19	Y	PFS, OS	T-cell detection in tumors	Tumor size, intracerebral inflammation, IL13Ra2 expression on tumor cells
Administration of MK-3475 (anti PD-L1 antibody) in combination with MRI-guided laser ablation	Y	PFS, OS	Antiglioma immune response before and after treatment. Correlate intratumoral expression of PD-L1 with number of glioma cell-specific cytotoxic T cells and OS	Identify PD-1-dependent biomarkers

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

Table 2. Open clinical trials of immunotherapy in glioblastomas (cont.).

NCT number, date first received, category	Phase	Study title	Diagnosis
NCT02149225 16 May 2014 Personalized peptide vaccine	I	Trial of actively personalized peptide vaccinations plus immunomodulators in patients with newly diagnosed GBM concurrent with temozolomide maintenance	GBM
NCT02336165 23 December 2014 Immunosuppressive checkpoint blockade	II	Phase II study to evaluate the clinical efficacy and safety of MEDI4736 in patients with glioblastoma	GBM

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

generate lymphocytes expressing CARs [33,34,130–132], with great selectivity and cytotoxicity against tumor cells. While viral-mediated suicide gene therapy and oncolytic viral therapy are not generally considered ‘classical’ immunotherapeutic approaches, accumulating data demonstrated that, in some cases, viral/gene therapy-mediated tumor cell death qualifies as immunogenic cell death initiating ER stress, expression of calreticulin on the cell surface, release of damage-associated molecular patterns like HMGB1 [133,134] and ATP and of pathogen-associated molecular patterns from the viral vector/oncolytic virus, leading to enhanced antigen presentation and antitumor immune response [135,136].

Suicide gene therapy involves the introduction of viral genes into the tumor cells, most commonly thymidine kinase (TK), resulting in the conversion of a systemically administered prodrug: gancyclovir (GCV) into a toxic metabolite within tumor cells and leading to tumor cell death [93,137]. In addition, vectors for gene therapy can be modified to express immunostimulatory molecules, which will aid in the fight against GBM. A number of studies using replication-deficient viruses have been conducted for GBM [92,138], including a bicistronic system that carries IL-2 and TK [90] and Flt3L and TK [139].

Oncolytic viral therapy utilizes replication-competent viral vectors, able to selectively replicate in tumor

cells, induce tumor cell lysis and spread of viruses to adjacent cells [87,93,140]. Additionally, the nonlytic viruses can express therapeutic genes in target cells. Oncolytic herpes simplex virus (HSV), measles virus, poliovirus, Newcastle disease virus and conditionally replicating adenovirus are all being tested at various stages in GBM therapy [88]. G207 is a conditionally replicating mutant HSV that has an impaired RR enzyme allowing it to replicate only in dividing cells. Since HSV is a human pathogen with neurotropic properties, this genetic manipulation provides tumor selectivity with safety. Phase I and Ib clinical studies using G207 showed no treatment-related toxicity with repeated doses and direct injection into the resected cavity [141,142]. Promising therapeutic responses were also identified in 8 out of 21 patients in the Phase I study and 3 out of 6 patients in the Phase Ib trial. Another genetically modified mutant HSV, HSV1716 was also tested in a Phase I study, demonstrating no toxicity [91]. In a subsequent Phase Ib study, no treatment-related toxicity was observed and evidence of viral replication was seen in tumor biopsies by histological examination. Second-generation oncolytic HSV vectors are also in preclinical development. Such vectors have been engineered to express therapeutic transgenes such as TNF- α , VEGF and IL-4 [93]. The two commonly tested conditionally replicating Ads in glioma are ONYX-015

Table 2. Open clinical trials of immunotherapy in glioblastomas (cont.).

Intervention	Outcome measures			
	Safety Toxicity	Efficacy	Immunologic analyses	Other
Administration of APVAC1 vaccine plus poly-ICLC and GM-CSF or APVAC2 vaccine plus poly-ICLC and GM-CSF	Y	PFS, OS	Frequency of peptide-specific CD8 T cells, No. of T-cell responses per peptides vaccinated No. of patients with at least one and two vaccine-induced T-cell responses Average number of immune responses per patient	Frequency of immune cell populations in the blood. Serum and plasma immunological proteins. Noncellular parameters measured from tumor, plasma or serum. Cellular parameters from PBMCs, leukapheresis samples or isolated TILs
Administration of MEDI4736 (a human monoclonal antibody directed against programmed cell death ligand 1 PD-L1) in combination with radiotherapy and bevacizumab	Y	OS, PFS, ORR	Biological activity of MEDI4736, as assessed by immunologic markers	Pharmacokinetic profile of MEDI4736: half-life, Tmax, Cmax

Worldwide, there are currently (ClinicalTrials.gov, 10 March 2015) 29 open clinical trials assessing immunotherapeutic approaches for the treatment of glioblastoma: therapeutic strategies include: DC vaccines (5 trials), autologous T cell transfer (3), autologous T cell transfer with DC vaccine therapy (1), synthetic peptide vaccines (5), tumor lysate vaccines (3), gene therapy (4) and targeting immunosuppressive checkpoints (4). In addition, one study is looking at generating anti-GBM antibodies with hybridomas, one study is an MRI observational study following DC vaccination and one is looking at arginine as immunostimulant. BTIC: Brain tumor-initiating cell; CAR: Chimeric antigen receptor; CMV: Cytomegalovirus; DC: Dendritic cell; DIPG: Diffuse intrinsic pontine glioma; GBM: Glioblastoma multiforme; ORR: Overall response rate; OS: Overall survival; PBMC: Peripheral blood mononuclear cell; PFS: Progression-free survival.

and Ad5Delta24 [137]. ONYX-015 has a deletion in the early region E1B-55kD protein that normally binds to inactivated p53 in the infected cells preventing them from undergoing apoptosis. Absence of this protein allows the virus to only replicate in p53-deficient tumor cells. Phase I studies with ONYX-015 in patients with recurrent glioma that were injected with various doses of ONYX-015 in the resection cavity showed neither treatment-related toxicity nor clinical benefit [143,144]. The Ad5Delta24 can selectively replicate in glioma cells because of a deletion in the viral protein E1A and preclinical studies have shown therapeutic efficacy against glioma xenografts [145]. However, cancer cells with intact Rb protein are refractory to Ad5Delta24. Ad5Delta24 was further modified (Ad5Delta24-RGD) to increase its targeting to tumor cells and is currently under a Phase I study. Genetically modified variants of measles virus such as MV-Edm, that have a high affinity for cellular CD46 receptors abundantly expressed on tumor cells, have also been tested in preclinical settings with favorable therapeutic success [146,147]. MV-Edm-CEA is also in a Phase I study. Specific targeting to tumor cells has also been achieved by the creation of variants expressing glioma-specific ligands such as IL-13 against IL-13R α 2 [148] and glycoproteins gD and gB to mediate HSV infection exclusively through recognition of EGFR on glioblastoma cells [149].

Immune stimulatory gene therapy serves to generate a robust immune response against glioma-specific antigens. Cytokine-mediated gene therapies also aim to achieve higher local concentrations of cytokines/chemokines and reduce systemic toxicity. Cytokine gene therapy with IFN- β has shown efficacy in preclinical studies with human xenografts and in mouse models of glioma, demonstrating augmentation of T-helper cell response, DC activity, NK cells and prolonged survival [150-152]. IFN- β also induces MHC I expression and therefore enhances the CTL response. A Phase I clinical trial with liposome-mediated IFN- β gene injection into the resection cavity demonstrated minimal toxicity with more than 50% tumor reduction (T1-weighted MRI) in two out of five patients for at least 16 months [153]. Another dose escalation study with Ad-IFN- β in 11 patients with GBM also showed therapeutic efficacy with no adverse effects except in one patient [154].

IFN- γ -mediated gene therapy has not shown efficacy when administered alone [137]; however, in combination with TNF- α , it enhanced the survival of glioma bearing animals along with an increase in T-cell recruitment to the tumor [155]. In other studies, IL-2 or IL-12 gene therapy resulted in growth inhibition of a rat glioma [156]. A combination of IL-2 with TK (GCV) in 12 patients with recurrent glioma using RV (retrovirus)-mediated gene therapy showed 12-month

PFS and OS rates of 14 and 25%, respectively, with minor side effects [86]. Clinical trials using IL-4/TK gene-modified autologous glioma cells or fibroblasts and a replicative oncolytic HSV carrying IL-12 gene therapy are underway [89].

Flt3L was initially characterized as the cytokine that resulted in enhanced myelopoiesis and B lymphopoiesis and subsequently it was shown to have a potent effect on the generation of both myeloid- and lymphoid-derived DC populations in mice [157]. Our group has pioneered the development and efficacy testing of Ad-mediated delivery of recombinant human Flt3L (Ad-Flt3L) in preclinical models of glioma [158].

Administration of Ad-Flt3L significantly inhibited tumor growth and increased survival in a dose-dependent manner. We have also developed a conditionally cytotoxic-immune stimulatory gene therapy that delivers TK and FLT3L using Ads [159,160]. Tumor cells are selectively killed by the TK/GCV administration. Flt3L serves to increase the recruitment of APCs to the tumor microenvironment which take up antigens released by the dying tumor cells and subsequently induce tumor-specific T-cell responses. Dying tumor cells also produce HMGB1, which activates TLR2 on APCs. Our experiments showed that release of HMGB1 and activation of TLR2 were crucial for




Figure 2. CTLA-4 and PD-L1 immune checkpoints in glioma immune escape. The CTLA-4 immune checkpoint (left panel) occurs during the priming phase of the immune response, primarily within secondary lymphoid organs. The inhibitory CTLA-4 T-cell receptor binds with higher affinity to the CD80/86 ligands on the surface of APCs and prevents their binding to and signaling through the costimulatory receptor CD28. This leads to decreased T-cell activation and proliferation in the context of antigen presenting MHC class I. PD-1 signaling (right panel) occurs during the effector phase of the immune response within the tumor microenvironment. The PD-1 receptor on the T-cell surface interacts with one of two PD-1 ligands that are expressed on the surface of tumor cells: PD-L1 or PD-L2. This interaction, in the context of tumor antigen presenting MHC class I, decreases the T-cell tumor lytic capacity and induces T-cell anergy. APC: Antigen-presenting cell.




Figure 3. Gene therapy for glioma with Ad-TK and Ad-Flt3L. (A) Thymidine kinase and Flt3L-expressing adenoviruses are injected directly into the tumor. (B) Following GCV administration, TK will convert GCV to GCV-triphosphate that is incorporated into the DNA of actively proliferating cells, that is, tumor cells, causing them to undergo apoptosis and release tumor antigens, including HMGB1. Flt3L entering systemic circulation will induce the trafficking of DCs into the tumor and their exposure to tumor antigens. (C) DCs exposed to tumor antigens process them into peptides presented on MHC class I/II molecules and increase the expression of the costimulatory molecules CD80/86 on their surface. (D) Activated DCs travel to the draining lymph nodes where they present antigenic peptides in combination with the costimulatory signals to naive CD4⁺ and CD8⁺ T cells, inducing clonal expansion and maturation of glioma-specific T cells, secretion of stimulatory cytokines, trafficking into the tumor and cytolytic killing of glioma cells. DC: Dendritic cell; GCV: Gancyclovir.

the TK-Flt3L-induced antiglioma response [133]. This gene therapy approach has demonstrated tumor regression, long-term survival and immunological memory in several transplantable, orthotopic syngeneic models of GBM in mice and rats [159–162]. Based on the excellent success seen in preclinical testing with the TK-Flt3L gene therapy, a Phase I clinical trial was launched in 2013, using this cytotoxic immune stimulatory approach (NCT01611992). A diagram of this therapeutic mechanism is presented in **Figure 3**.

Active immunotherapy

DC vaccines

DCs, the most effective APCs, can prime both CD4⁺ T helper and CD8⁺ cytotoxic cells [163,164] and also function as strong activators of NK and NKT cells [165,166]. To stimulate T cells and generate a specific and efficient antitumor immune response and induce immunological memory, DCs need to deliver three signals: signal (**Figure 3A**), represented by the by cross-presentation of antigenic peptides via MHC molecules; signal (**Figure 3B**), by the interaction of costimulatory molecules, CD80 and CD86 with the CD28 receptor located on T cells [167,168] and signal (**Figure 3C**) by immunostimulatory cytokines such as IL-12 and IL-2 secreted by DCs and activated CD4⁺ T cells [169].

Preclinical studies

Since 1999, numerous preclinical studies (summarized in **Table 1** [37–76]) have analyzed the efficiency of DC vaccines in the treatment of glioma using rodent syngeneic models. The first step in any vaccination protocol is represented by the generation of sufficient number of DCs, typically from bone marrow cells, induced to differentiate with specific cytokines like GM-CSF and/or Flt3L, pulsed with GAAs (tumor lysate or tumor-specific peptide/mRNA epitopes) and injected (most often intradermally or subcutaneously) into animals either prior to or after tumor inoculation (**Figure 4**). Oftentimes, adjuvants like CpG oligonucleotides or lipopolysaccharide are coinjected in order to increase the expression of maturation markers: CD80 and CD86 on DCs. Preclinical studies have aimed to optimize several factors shown to be critical in the efficacy of DC vaccination: DC differentiation, antigen loading, administration route and adjuvant treatment.

DC differentiation

Bone marrow-derived mononuclear cells can be induced to differentiate into DCs through the actions of two main cytokines, GM-CSF (used in combination with IL-4) and Flt3L (used with IL-6) [170]. It has been shown that alpha-type 1-polarized DCs induce larger numbers of antitumor CTLs, secrete increased amounts

of IL-12 and are resistant to immunosuppression by Tregs [171]. DCs induced by Flt3L + IL-4 are of the α 1 type, whereas DCs generated with GM-CSF are not, but can be converted into α 1-DCs with lipopolysaccharide, IFN- γ , IFN- α IL-4 and poly- ICLC [172].

DC loading with tumor antigens

Most commonly, whole tumor lysates have been used to load DCs with GAAs. These lysates can be generated with methods which induce necrosis or apoptosis: acid elution, freeze-thawing, irradiation, temozolomide or thymidine-kinase + GCV treatment [71,173]. Alternatively, tumor RNA and tumor-specific peptides/mRNA including ephrin (Eph)-A1, EphA2, IL-13 α 2, survivin, gp100 and TRP-2 have been used to pulse DCs, or DCs have been directly fused to tumor cells [45,46,68,76]. Using multiple epitopes to pulse DCs decreases the risk of developing immune tolerance through immunoediting. Genetically engineering DCs to express common glioma antigens permit multiple epitope presentation of GAAs, irrespective of the patient's human leukocyte antigen (HLA) type.

DC vaccination route

The route of DC administration represents a critical aspect of vaccination strategies. It has been shown that increased proximity to the tumor site negatively impacts the efficacy of DC vaccinations due to increased immunosuppression [174]. In addition, intratumoral administration of immunostimulatory cytokines like IL-12, IFN- γ and Flt3L enhances the efficacy of subcutaneous DC administration [45,59,71,175], suggesting that a combinatorial approach, intratumoral, for cytokines and systemic, at the distance from the tumor, for DCs, may represent an optimal strategy for vaccination.

Adjuvant treatment

In animal models, several proinflammatory cytokines have been used to improve the therapeutic efficacy of DC vaccination, including IL-12, IFN- α , IL-4, IFN- β , CXCL10 and Flt3L (**Table 1**). It was thought that chemotherapy negatively impacts immunotherapy for glioma; however, it has been shown that temozolomide does not inhibit but rather enhances the effect of DC vaccination [52] and that it does not impair the efficiency of immunomodulatory gene therapy with Ad-TK+GCV and Ad-Flt3L [134]. Activation by costimulatory molecules is critical to avoid T-cell anergy [176]. OX40 receptor (CD134) is a costimulatory molecule that is expressed on activated CD4⁺ and transiently expressed on activated CD8⁺ T cells. Stimulatory anti-OX40 receptor antibodies have been shown to enhance the therapeutic efficacy of DC vaccination in a mouse glioma model [53].




Figure 4. Dendritic cell vaccination for glioma therapy. Mononuclear cells extracted from bone marrow are cultured with GM-CSF and IL-4 or Flt3L and IL-6 to induce their differentiation into DCs. Tumor cells are killed by irradiation or other cytotoxic stimuli to generate tumor antigens. DCs are then pulsed with tumor antigens by co-culturing DCs with whole tumor lysates, purified and injected subcutaneously or intradermally as a vaccine together with costimulatory agents such as CpG oligonucleotide (CpG-ODN). CpG-ODNs stimulate DCs through signaling via Toll-like receptors. Injected DCs migrate to lymph nodes, where they encounter CD8⁺ and CD4⁺ T cells. MHC-antigen complexes are recognized by T-cell receptors and IL-12 secreted by the DCs further activates CD8⁺ T cells to become antigen-specific CTLs. CTLs migrate into the tumor where they attack and lyse tumor cells. CTL: Cytotoxic T-lymphocyte; DC: Dendritic cell; GCV: Gancyclovir; TK: Thymidine kinase.

Clinical studies

In clinical trials, DCs are induced to differentiate from peripheral blood mononuclear cells using most commonly GM-CSF and IL-4. DCs are then loaded with specific antigens, transfected with tumor RNA or fused with tumor cells and induced to mature using combinations of cytokines like: IL-6, TNF- α , PGE2 and IL-1 β or TNF- α , IL-1 β , IFN- α , IFN- γ and poly[I:C] [95]. Mature DCs can be administered directly to the patient or frozen down for future use. Oftentimes, the number of cells produced is limited and it is critical to harvest the peripheral blood mononuclear cell prior to beginning of any treatment, to avoid collecting blood when the patients become lymphopenic.

The most advanced current clinical trial with DC vaccines for glioblastomas (NCT00045968) using an autologous DC vaccine (DCVax-L), prepared by puls-

ing DCs with proteins from the patient's own tumor, has reached Phase III. In earlier phases of the trial, it has been shown that the vaccine is safe and elicits systemic antitumor CTL response, yet, clinical benefit was very limited. This was attributed to actively progressing tumors and high expression of TGF β 2 in the tumor [96,97]. A study using autologous DC vaccines in combination with TLR agonists as adjuvants: Imiquimod or poly-ICLC has shown that patients who had tumors with a mesenchymal, but not proneural gene expression pattern responded to treatment as demonstrated by increased OS compared with controls with the same genetic signature [82].

Another study of autologous DC vaccine pulsed with six GAA peptides (HER2, TRP-2, gp100, MAGE-1, IL13R α 2 and AIM-2) resulted in response to treatment of 33% of the 21 patients enrolled. PFS and OS

correlated with the quantitative expression of MAGE1 and AIM2 from resected tumors [81]. The tumors of all patients expressed at least three antigens and 75% expressed all six antigens. The median PFS in newly diagnosed patients was 15.9 months and the median OS was 38.4 months. Furthermore, production of IFN- γ and TNF- α in stimulated CD8⁺ T cells was increased and correlated with survival. Expression of CD133 has been shown to increase in recurrent GBM, indicative of resistance to treatment and worse clinical outcome [177,178]. Following the GAA vaccine, there was a decrease in CD133 expression in patients with recurrent GBM, suggesting that the vaccine therapy resulted in a cytotoxic attack on glioma stem cells.

The treatment of 22 patients with recurrent GBM with α -type 1-polarized DCs (α DC1) pulsed with the GAA peptides: EphA2, IL13R α 2, YKL-40 and gp100 combined with administration of poly-ICLC showed the induction of a positive immune response at least against one of the peptides in 58% of patients, an increase in type 1 cytokines (IFN- α 1 and CXCL10) and that production of IL-12 correlated with time to progression [179].

When comparing two Phase I clinical trials with DC vaccines pulsed either with autologous tumor lysate (NCT00068510) or with the GAA: TRP-2, gp100, HER-2/neu and survivin (NCT00612001), there were no significant differences between the trials with respect to frequency of helper, CTLs, B cells or NK cells; however, the GAA trial had a relative increase in activated NK cells (CD3⁺CD4⁺CD16⁺CD25⁺) as well as an increased ratio of Tregs when compared with the tumor lysate-pulsed DC vaccine group [94]. This was associated with decreased survival in the GAA trial. Decreased Treg populations postvaccination correlated with increased survival in both groups, recommending the use of monitoring Treg populations in clinical trials of immunotherapy for GBM.

A randomized Phase II clinical trial using SOC with added antiangiogenic therapy with bevacizumab and DC vaccine (AVO113), generated by priming DCs with autologous tumor antigens, showed an increase in median OS (535 days +/- 155) in the combined bevacizumab and vaccine group compared with the vaccine group (438 days +/- 205) or bevacizumab alone group (406 days +/- 224) [78], suggesting a possible mechanism of decreased immunosuppression induced by the anti-VEGF antibodies.

Targeting GSCs with active immunotherapy represents an attractive therapeutic strategy considering that GSCs are resistant to the conventional approaches of radiotherapy and chemotherapy. Many studies have focused on identifying cellular markers of GSC, notably CD133, but also EGFRvIII, HER2, IL13R α 2 and

L1-CAM, yet these markers are also found on neural stem cells and other nontumor cells [180]. A recently completed Phase I/II trial (NCT00846456) using DCs transfected with autologous GSC mRNA (extracted from glioma neurospheres), demonstrated induction of T-lymphocyte proliferation in response to tumor lysate, hTERT or survivin peptides. The vaccinated patients had significantly longer median PFS than the matched controls (694 days compared with 236 days) and also a longer median OS (759 days) than the controls [85]. This study is encouraging and highlights the benefit of identifying GSCs by their sphere-forming ability rather than by particular surface markers.

Amplifications of the *EGFR* gene with gain of function represent the most common (40%) genetic alteration in GBM [181]. A mutant form of the *EGFR* gene, EGFRvIII, found in ~20–30% of GBM patients, expresses a truncated, constitutively active form of the receptor which results in increased proliferation and survival advantage of GBM tumor cells [182]. Another transforming mutation is EGFRvIV, with a deletion in the C-terminal domain. These mutations are very specific to glioma cells and hence an attractive target for therapy. Numerous preclinical studies demonstrated the efficiency of targeting the EGFRvIII or wild-type EGFR with peptide vaccines [183] or targeted antibodies [184] and led to development of a clinical trial with autologous DC vaccines pulsed with the EGFRvIII keyhole limpet hemocyanin (KLH)-conjugated specific peptide (PEPvIII-KLH/CDX-110), which showed safety and efficacy in eliciting an antitumor immune response and improved survival in GBM patients who express the respective variant [185].

Peptide vaccines

Peptide vaccines offer advantages compared with DC vaccines, as they do not require generation of activated and mature autologous DCs, a process that may not be feasible in all patients. It is important that the peptides are tumor-specific and that immune stimulatory strategies (immune adjuvants, cytokines: IL-2, GM-CSF) are coopted to ensure the proper priming and maturation of the endogenous APCs.

Following promising results with the DC vaccine pulsed with the EGFRvIII peptide, a subsequent Phase II multicenter study (ACTIVATE, ACTII) applied the PEPvIII-KLH/CDX-100 vaccine (Rindopimut/CDX-110) concurrent with temozolomide, without the accompanying DCs, in patients with newly diagnosed EGFRvIII-positive GBM [186]. This study showed that 6 out of 14 patients analyzed developed EGFRvIII-specific antibody responses which correlated positively with OS, the median OS (26.0 months) being higher than in the matched historical control

group (15 months) and that at recurrence 82% of patients lost EGFRvIII expression, demonstrating treatment-induced tumor immunoediting and immune escape [185,186]. A subsequent Phase II multicenter single-arm trial (ACTIII), aimed to confirm previous results using the same therapeutic approach, showed a median OS of 21.8 months, specific anti-EGFRvIII antibody titers in 85% of patients and decrease in EGFRvIII immunoreactivity in 4/6 (67%) tumor samples [84]. A current Phase III multicenter clinical trial (ACTIV, NCT01480479) is testing the efficacy of (CDX-110, Rintega, CellDex therapeutics), GM-CSF, temozolomide and KLH for the treatment of adult patients with EGFRvIII-positive glioblastomas. Another Phase II study is looking at the effects of combining rindopepimut, GM-CSF and bevacizumab for the treatment of relapsed EGFRvIII-positive glioma (NCT01498328).

Given the risk of immunoediting following single-peptide vaccinations, many investigators are aiming to produce effective combinations of GBM-specific peptides to induce robust antitumor immune responses and prevent the induction of immune tolerance. A pilot study of 26 pediatric brain stem and high-grade gliomas used a combination of three GAA peptides: EphA2, IL-13R α 2 and survivin, together with a pan HLA-DR tetanus toxoid peptide and the TLR3 agonist poly[I:C] administered intradermally in HLA-A2-positive children. This study showed that the vaccines were well tolerated, induced specific anti-GAA immune responses (by ELISPOT) and favorable clinical responses [102]. Some patients presented initial pseudoprogression, as evidenced by worsening symptoms and transient increased edema, evidenced on MRI scans, due to tumor infiltration with immune cells following the vaccine. However, patients showing pseudoprogression survived longer, suggesting that this may be a favorable prognostic marker for treatment efficacy.

In adult patients with high-risk low-grade glioma (LGG), a study using vaccinations with eight courses of intramuscular administration of the GAAs: IL13R α 2, EphA2, WT1 and Survivin emulsified with the adjuvant Montanide-ISA-51 demonstrated robust IFN γ ELISPOT responses against at least 3 out of 4 peptides in 14 out of 22 patients and median PFS of 17 months in newly diagnosed patients and 12 months in recurrent LGG. Results are encouraging and warrant further studies using this approach in patients with LGG (WHO grade II), in which the slower course of disease progression allows for repeated vaccinations with improved outcome [101].

Numerous other glioma-specific peptides have been identified (reviewed in [187]) and there is great potential in finding combinations of peptides which may be

used to generate an 'off-the-shelf' vaccine that will be effective in a broad range of glioma patients. Ongoing clinical trials are testing a proprietary combination of 11 HLA-A2-restricted tumor-associated peptides IMA950 alone (NCT020278648) or in combination with GM-CSF (NCT01222221) or poly-ICLC (NCT01920191).

In Europe, the Glioma Actively Personalized Vaccine Consortium aims to rapidly personalize peptide vaccines, within a few months after the initial surgery, using next-generation sequencing, mass spectrometry and computational medicine algorithms. A current Phase I clinical trial in newly diagnosed GBM patients (NCT02149225) will test the safety profile of patient-tailored APVAC vaccines when administered with immunomodulators concurrent with temozolomide. The frequency of antigen-specific CD8⁺ T cells will be monitored, as well as immune cell populations in the blood and tumor together with a panel of serum cytokines/proteins to identify biomarkers of immune response to the vaccine.

Tumor cell vaccines

Results from a recent Phase I/II prospective clinical trial using temozolomide, fractionated radiotherapy and autologous, formalin-fixed tumor cell vaccine in 24 patients with newly diagnosed GBM show promising results with treated patients, exhibiting a median PFS of 8.2 months and OS of 22.2 months [79]. Interestingly, the median PFS in patients with a delayed-type hypersensitivity response at the third vaccination of greater than 10 mm was significantly higher (OS = 29.5 months) compared with patients with smaller delayed-type hypersensitivity, in congruence with the emerging concept that atopic reactions represent a favorable prognostic sign for immunotherapy against GBM.

The heat shock protein chaperone gp96 plays a crucial role in folding, assembly and export of TLRs and in binding to the CD91 receptor on APCs [188,189]. Activation of DCs, with tumor-associated peptides and proteins, is greatly enhanced when coupled to heat-shock protein-peptide gp96. A recent Phase II clinical trial of adult recurrent GBM in which vaccines were made by isolating tumor gp96 complexes from autologous-resected tumors (HSPPC-96) demonstrated their safety and encouraged further studies of efficacy [77]. The median OS was 42.6 weeks, the 6 months survival: 90.2%, and 12 month survival of 29.3%, with poorer outcome among patients with lymphopenia. In the previous Phase I trial, with the same vaccine, significant peripheral immune responses were demonstrated in 11 out of 12 patients treated and also significant infiltration of tumor tissue with CD3⁺, CD8⁺ and CD56⁺ IFN- γ -positive cells. The median survival of

responders was 47 weeks, longer than the 16 weeks of the single nonresponder [98]. A follow-up randomized Phase II open clinical trial (NCT01814813) will compare the efficacy of the HSPPC-96 autologous vaccine with or without bevacizumab therapy in recurrent, resectable GBMs.

Passive immunotherapy

Antibodies

Antibodies against EGFR (nimotuzumab) in combination with radiation and chemotherapy (vinorelbine) have been used in a study of pediatric diffuse intrinsic pontine gliomas, which presents with overexpression and amplification ERBB1/EGFR [105]. Results recently published from this trial show that this treatment increased median OS (15 months) when compared with a study of nimotuzumab and radiation alone (OS = 9.4). To overcome the immunosuppressive effect of tumor-induced Tregs, a randomized, placebo-controlled pilot study combined a selective antibody for the high-affinity IL-2R α (daclizumab) with a vaccine (ZAP IT) targeting EGFRvIII, in temozolomide-treated GBM patients [106]. The study shows that one administration of daclizumab reduced the number of Tregs, without markedly affecting the number or activation of CD4⁺ or CD8⁺ T cells and favorably influenced the production of antibodies against EGFRvIII.

Autologous T-cell transfer & CAR-modified lymphocytes

Adoptive T-cell therapy was first used in the treatment of melanoma patients [32]. Treatment efficacy is, however, limited by the immunosuppressive tumor environment and the fact that not all patients have resectable tumors to be used in the production of tumor-specific lymphocytes [190]. A Phase I clinical trial in patients with recurrent GBM- and cytomegalovirus-positive serology, autologous T-cells transfer with cytomegalovirus-specific peptides-stimulated T cells, four out of ten patients who completed a minimum of three T-cell infusions showed PFS during the extent of the study. The median survival was 403 days and the median PFS was 246 days. The clinical outcome was, however, not correlated with antigen-specific T-cell number or functional phenotype [103], as indicated by expression of CD103, characteristic of recent thymic emigrant status of vaccine-induced CD8⁺ T cells [191].

Genetically altering lymphocytes to express CARs provides a feasible and useful method of passive immunotherapy against tumors. CARs consist of the antigen-binding region of a monoclonal antibody fused with the signal transduction domain of CD3 ζ or Fc ϵ RI γ (gamma subunit of the Fc region of the immunoglobulin

E receptor I) [192]. They permit independence from MHC I expression on tumors and increased penetration and persistence into the tumor microenvironment when compared with monoclonal antibodies.

CARs have been tested in clinical trials for neuroblastoma, renal cell carcinoma and B-cell malignancies [193–195]. While therapies did show therapeutic efficacy, serious adverse effects were also observed in some cases, possibly because of the expression of the targeted antigen on normal tissues. It is therefore essential to select targets that are highly specific to tumor cells. IL-13R α 2 is a cell surface receptor specific for glioma [196] and thus represents a good target for immunotherapy. Preclinical data with IL-13 zetakine CAR to target IL-13R α 2 showed elimination of human xenografts in mice [132]. The clinical trial testing safety and feasibility of this therapeutic approach in patients with recurrent GBM has recently been published [104], showing that the approach is feasible with minimal adverse effects and that two out of three patients who received repeated intracranial infusion of IL13-zetakine+CTLs developed transient anti-glioma responses, suggested by increased regions of tumor necrosis visualized on MRI in one patient and decreased expression of IL-13R α 2 in another.

HER2 is expressed by up to 80% of GBMs and absent from the normal brain [197], and hence a good target for therapy. It has been shown that HER2-positive autologous GBM cells can induce a specific T-cell response with increased production of IFN- γ and IL-2 and result in tumor regression in a xenograft GBM mouse model [130]. Furthermore, these T cells can kill CD133 HER-2-positive glioma stem cells. A Phase I safety/efficacy clinical trial using HER-2-specific CARs will test this treatment in patients with glioblastoma (NCT02442297).

Immune stimulatory adjuvants

Pathogen-associated molecular patterns bind to TLRs and have a high capacity to stimulate cell-mediated immunity by increasing the production of immunostimulatory cytokines and increasing expression of costimulatory molecules on APCs. Compounds most commonly investigated as adjuvants for cancer vaccines are: polyriboinosinic–polyribocytidylic acid (poly[I:C]) and its derivative poly-ICLC, (poly[I:C] stabilized with poly-L-lysine and carboxymethylcellulose) (synthetic analogs of viral dsRNA polymers), TLR3 agonists which have been shown to enhance anti-tumor response by activating NK and T cells [198,199], CpG oligonucleotides (CpG ODN), strong activators of both the native and adaptive immune system [200] and TLR7 agonists, like Imiquimod [201,202].

A prospective Phase II clinical trial of pediatric glioma treated with poly-ICLC has proven to be well

tolerated by children with no observed dose-limiting toxicities, five out of ten children showing long-term stable disease. The authors conclude that the results justify biomarker studies for personalization of poly-ICLC as a single agent or adjuvant. Another Phase II study of poly-ICLC and radiotherapy with concurrent and adjuvant temozolomide in newly diagnosed GBM concluded that poly-ICLC may improve the efficacy of radiotherapy and temozolomide treatment without added toxicity [203].

CpG ODNs (TLR9 agonists) have shown promise in many preclinical studies and also in Phase I clinical studies of glioma. A multicenter Phase II clinical study testing intratumoral administration of CpG-ODN for patients with recurrent GBM, showed little beneficial effect in this use of single-agent CpG-ODN [200,204]. Several trials use CpG-ODNs in combination with other immunotherapeutic agents.

Another immunostimulatory adjuvant, tetanus toxoid, has recently been shown to improve the efficacy of DC vaccination and to prolong the survival of glioblastoma patients, with more than 50% surviving longer than 40 months. In mice, it was shown that tetanus toxoid-enhanced DC migration and suppressed tumor growth in a CCL3-dependent manner [80].

Ongoing challenges

During the last 5 years, much progress has been made in refining immunotherapeutic approaches for the treatment of GBM and it is likely that soon, immunotherapy will be included in the SOC for GBM, next to maximal possible surgical resection, radiation and chemotherapy. It has become clear, however, that many details of immunotherapeutic protocols still need to be carefully assessed. With current approaches, the OS is still very short. A unique challenge of tumors localized in the brain is brought about by the minimal tolerance to inflammation, difficult to avoid when attempting immunostimulatory treatment strategies; hence, a delicate balance needs to be met between enhancing immune-mediated tumor killing and limiting brain inflammation. A great need exists to identify biomarkers with prognostic value and of clinical efficacy to guide the therapeutic intervention. Clinical trials do not have a standardized protocol to analyze the antitumor immune response and this makes it difficult to interpret the results and to compare one trial to the next. Given the tremendous heterogeneity of clinical presentations, the intrinsic heterogeneity of each individual tumor, as well as the immunoeediting following various therapeutic interventions, it is apparent that a single 'magic bullet', a 'one-size-fits-all' approach will not be forthcoming. Personalized medicine with ongoing monitoring of tumor progression and immunological end points using

a dynamically tailored therapeutic approach could bring great promise to the management of GBM.

Conclusion and future perspective

Current open clinical trials of immunotherapy and GBM illustrate a predominance of studies of DC vaccines in various combinatorial treatment strategies and an emerging popularity of studies with antibodies targeting immunosuppressive checkpoints (ClinicalTrials.org, Table 2). Combinatorial approaches in preclinical trials of GBM show benefit when targeting multiple immune checkpoints [123] or when adding cytokine therapy [118] and are likely to result in better outcomes when translated to the clinic. Ongoing clinical trials are testing many combination therapies. Antibodies against CTLA-4 and PD-1 are administered with bevacizumab, and/or radiotherapy or kinase inhibitors (NCT02017717, NCT02337491, NCT02336165, NCT02423343). Suicide gene therapy is combined with radiation therapy or immune-stimulatory therapy (NCT01811992, NCT00634231). Oncolytic viral therapy is used together with temozolomide or IFN- γ (NCT01956734, NCT02197169). Peptide or DC vaccines are tested with GMCSF, temozolomide, KLH or bevacizumab (NCT01920191, NCT01480479, NCT01498328) and autologous T-cell transfer with DC vaccine or IL-2 and chemotherapy (NCT01326104, NCT01522820, NCT01454596). These trials will soon shed light on the most promising avenues to pursue further.

Criteria for radiographic response assessment to immunotherapeutics have been recently defined [205]; however, it has become clear that the neuro-oncology field would benefit from guidelines tailored to the unique characteristics of brain tumors. The immunotherapy Response Assessment in Neuro-Oncology (iRANO) criteria are currently being discussed [206] and are meant to guide therapeutic decisions and prevent premature termination of immunotherapeutic treatment due to pseudoprogression in patients responding to the treatment. An open observational clinical trial (NCT01657734) is using advanced multimodal imaging techniques: MRI spectroscopy, perfusion imaging and diffusion imaging in patients with glioblastoma treated with DC therapy to characterize inflammatory response, metabolism and tissue structures and will be instrumental in further shaping the iRANO criteria.

Increased efforts are dedicated to establish reliable biomarkers to improve the assessment of response to treatment and guide further therapeutic decisions. Advances in technology: quantitative, high-sensitivity and resolution flow cytometry of rare populations, high-throughput microscopy of tumor infiltrating lymphocytes *in situ*, whole transcriptome profiling of

Executive summary

- Immunotherapy for glioblastoma multiforme (GBM) is increasingly viewed as the fourth arm in the standard of care (SOC), as several clinical studies have shown safety, increased progression-free survival (PFS) and the promise of long-term efficacy through eliciting antitumor immunological memory. Successful immunotherapy for glioma needs to address tumor-induced immune suppression in addition to being mindful of the minimal tolerance for inflammation in the brain.

Targeting immunosuppressive checkpoints

- CTLA-4: receptor expressed on T-lymphocytes, inhibits costimulatory molecules on antigen-presenting cells, which results in T-cell anergy. Blocking CTLA-4 with specific antibodies (ipilimumab, tremelimumab) increases the survival of patients with metastatic melanoma and survival of animals in preclinical glioma studies, increasing antitumor CD8 cytotoxicity and decreasing accumulation of Tregs. Current clinical trials are testing these antibodies in glioma patients.
- PD-1 ligands (PD-L1 and PD-L2) expressed by glioma cells bind to PD-1 receptors on T cells, blocking their proliferation and cytokine production. Anti-PD-1 antibodies (nivolumab and pembrolizumab) have shown efficacy in the treatment of metastatic melanoma and xenograft animal models of glioma. In preclinical trials, decreased Treg infiltration, increased cytotoxic antitumor immunity and establishment of immunologic memory have been demonstrated. Clinical trials are underway to test these antibodies in glioma patients.

Immunostimulatory gene therapy

- Viral-mediated gene therapy induces immunogenic cell death of tumor cells which enhances antigen presentation and antitumor immune responses.
- Suicide gene therapy is the most common strategy for gene therapy in glioma. Thymidine kinase introduced into the tumor by viruses (or other vectors) will convert an administered prodrug (ganciclovir), into a toxic analog, which induces apoptosis of dividing tumor cells.
- Oncolytic viral therapy induces tumor cell death through viruses genetically modified to selectively replicate within tumor cells and can also express therapeutic genes like TNF α , IL-4. Viruses most commonly used are oncolytic HSV, RVs, measles virus (MV-Edm) and conditionally replicating adenovirus (ONYX-015, Ad5Delta24).
- Immunomodulatory gene therapy aims to create an immunostimulatory tumor microenvironment within the brain. Clinical trials are testing combinations of suicide gene therapy or oncolytic viral therapy with immunomodulating cytokine gene therapy (IFN β , TNF α , IFN γ , IL-2, IL-12 and Flt3L).

Active immunotherapy

- Dendritic cell (DC) vaccines, generated from autologous peripheral blood mononuclear cells expanded *ex vivo* and induced to differentiate into DCs, pulsed with various tumor antigens and reintroduced into the patient, represent the most common strategy for active immunotherapy and personalized medicine. A Phase III multicenter randomized clinical trial with autologous DC vaccines is currently underway (NCT00045968).
- Peptide vaccines offer advantages over cell-based vaccine with added convenience of an 'off the shelf' product. Many glioma-specific peptides have been identified and tested in combination or alone in many studies. The most commonly tested are EGFRvIII, EphA2, IL13R α 2, survivin, WT2725, gp100, YKL-40, MAGE-1 in various combinations and added immune stimulatory strategies using cytokines (IL-2, GM-CSF) and adjuvants to activate TLR receptors (CpG, poly[I:C]). Limitations are restricted to the HLA type of the patients, most commonly used are HLA-A2-restricted peptides. A Phase I clinical trial for personalized peptide vaccines is conducted in Europe by the Glioma Actively Personalized Vaccine Consortium consortium.
- Tumor cell vaccines using autologous tumor lysates show increased efficacy especially in patients undergoing delayed-type hypersensitivity reactions following vaccination. Glioma peptides coupled to the heat shock protein chaperon gp96 have an increased capacity to activate DCs and elicit-specific peripheral immune responses. Clinical trials using this strategy have shown increased PFS.

Passive immunotherapy

- Antibodies.
- Autologous T-cell transfer with T cells stimulated with cytomegalovirus peptides in patients with CMV-positive serology has shown increased PFS in a Phase I trial.
- Chimeric antigen receptor-modified lymphocytes. This strategy allows the generation of clonally expanded T cells modified to express receptors, which will recognize tumor cells independent of their MHC I expression. This approach is limited to very specific tumor antigens (like EGFRvIII and IL-13R α 2), which are not expressed on normal cells, as to not induce severe autoimmune adverse effects.
- Immune stimulatory adjuvants. These compounds are very useful in nonspecifically stimulating cell-mediated immunity through the activation of TLRs. Compounds used are: poly[I:C], poly-I:LC, CpG-ODNs, imiquimod, tetanus toxoid.

Executive summary**Ongoing challenges**

- The survival of patients treated with SOC and effective immunotherapy is still very short.
- Immunoediting and the intrinsic heterogeneity of GBMs allow tumor cells to escape treatment.
- Clinical trials using combinatorial approaches with SOC and antibodies to target immunosuppressive checkpoints are underway and show promise.
- Care needs to be taken when severe autoimmune adverse reactions develop; however, in many cases delayed hypersensitivity reaction may represent a positive prognostic factor.
- A great need exists to identify biomarkers able to guide therapeutic strategies and monitor efficacy of immunotherapeutic approaches.

Future perspective

- Ongoing clinical trials are testing combinations of immunotherapeutic approaches together with SOC and will soon shed light on the most promising avenues to pursue further.
- The immunotherapy Response Assessment in Neuro-Oncology criteria are currently being discussed and will improve the therapeutic decision making process to prevent premature termination of immunotherapeutic treatment due to pseudoprogression.
- Advances in technology allow for multidimensional profiling of tumors and immunological parameters which will enable a global analysis of response to treatment and will likely improve therapeutic decisions and clinical outcome of GBM through the development of optimal personalized medicine.

tumor and immune genes, high-throughput sequencing of antigen-specific receptors of whole lymphocyte populations (TCRseq and BCRseq) permit nowadays the accumulation of large datasets, which allow for multidimensional profiling of tumors and immunological parameters [207]. This will enable a more in depth analysis of response to treatment and will likely improve therapeutic decisions and clinical outcome of GBM toward the development of optimal personalized medicine.

Acknowledgements

We thank M Dahlgren and M Dzaman for their administrative and technical support.

Financial & competing interests disclosure

This work was supported by NIH/National Institute of Neurological Disorders & Stroke (NIH/NINDS) Grants RO1-

NS094804, R01-NS074387, R01-NS057711, R21-NS091555, University of Michigan U042841 to MG Castro; NIH/NINDS Grants R01-NS054193, R01- NS061107, R01-NS082311 and R21-NS084275 to PR Lowenstein; the Michigan Institute for Clinical and Health Research, NIH 2UL1-TR000433; University of Michigan Cancer Biology Training Grant, NIH/NCI (National Cancer Institute) T32-CA009676; University of Michigan Training in Clinical and Basic Neuroscience, NIH/NINDS T32-NS007222 and the University of Michigan Medical Scientist Training Program, NIH/NIGMS (National Institute of General Medicine Sciences) T32. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

Papers of special note have been highlighted as:

- of interest; •• of considerable interest

- 1 Doll R, Kinlen L. Immunosurveillance and cancer: epidemiological evidence. *Br. Med. J.* 4(5732), 420–422 (1970).
 - 2 Greene HS. Heterotransplantation of tumors. *Ann. NY Acad. Sci.* 69(4), 818–829 (1957).
 - 3 Baldwin RW, Robins RA. Factors interfering with immunological rejection of tumours. *Br. Med. Bull.* 32(2), 118–123 (1976).
 - 4 Burnet FM. Immunological aspects of malignant disease. *Lancet* 1(7501), 1171–1174 (1967).
 - 5 Vesely MD, Kershaw MH, Schreiber RD, Smyth MJ. Natural innate and adaptive immunity to cancer. *Annu. Rev. Immunol.* 29, 235–271 (2011).
 - 6 Facciabene A, Motz GT, Coukos G. T-regulatory cells: key players in tumor immune escape and angiogenesis. *Cancer Res.* 72(9), 2162–2171 (2012).
 - 7 Fecci PE, Mitchell DA, Whitesides JF *et al.* Increased regulatory T-cell fraction amidst a diminished CD4 compartment explains cellular immune defects in patients with malignant glioma. *Cancer Res.* 66(6), 3294–3302 (2006).
- **Demonstrates that glioma patients present with generalized lymphopenia and that the proportion of Tregs (CD4⁺FoxP3⁺) was 2.63-times higher in glioma patients than in normal controls and correlated with decreased proliferating capacity of CD4⁺ T cells. Depletion of Tregs restored the functional impairment of CD4⁺ T cells and this was accompanied by a decrease in Th-2 and an increase of Th-1 cytokines. This is the first study to underscore the role of Tregs in facilitating immune evasion by glioblastoma multiformes (GBMs).**

- 8 Crespo J, Sun H, Welling TH, Tian Z, Zou W. T cell anergy, exhaustion, senescence, and stemness in the tumor microenvironment. *Curr. Opin. Immunol.* 25(2), 214–221 (2013).
- 9 Mehling M, Simon P, Mittelbronn M *et al.* WHO grade associated downregulation of MHC class I antigen-processing machinery components in human astrocytomas: does it reflect a potential immune escape mechanism? *Acta Neuropathol.* 114(2), 111–119 (2007).
- 10 Anderson RC, Anderson DE, Elder JB *et al.* Lack of B7 expression, not human leukocyte antigen expression, facilitates immune evasion by human malignant gliomas. *Neurosurgery* 60(6), 1129–1136; discussion 1136 (2007).
- 11 Wintterle S, Schreiner B, Mitsdoerffer M *et al.* Expression of the B7-related molecule B7-H1 by glioma cells: a potential mechanism of immune paralysis. *Cancer Res.* 63(21), 7462–7467 (2003).
- 12 Dong H, Strome SE, Salomao DR *et al.* Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion. *Nat. Med.* 8(8), 793–800 (2002).
- 13 Didenko VV, Ngo HN, Minchew C, Baskin DS. Apoptosis of T lymphocytes invading glioblastomas multiforme: a possible tumor defense mechanism. *J. Neurosurg.* 96(3), 580–584 (2002).
- **In this study it is demonstrated that apoptotic T cells infiltrated in human GBM express Fas and are localized in close proximity or direct contact to Fas-L expressing tumor cells, suggesting a mechanism of immune evasion by GBM through the Fas/FasL pathway induced killing of T cells.**
- 14 Raychaudhuri B, Rayman P, Ireland J *et al.* Myeloid-derived suppressor cell accumulation and function in patients with newly diagnosed glioblastoma. *Neuro Oncol.* 13(6), 591–599 (2011).
- 15 Sippel TR, White J, Nag K *et al.* Neutrophil degranulation and immunosuppression in patients with GBM: restoration of cellular immune function by targeting arginase I. *Clin. Cancer Res.* 17(22), 6992–7002 (2011).
- 16 Larsen SK, Gao Y, Basse PH. NK cells in the tumor microenvironment. *Crit. Rev. Oncog.* 19(1–2), 91–105 (2014).
- 17 Levy EM, Roberti MP, Mordoh J. Natural killer cells in human cancer: from biological functions to clinical applications. *J. Biomed. Biotechnol.* 2011, 676198 (2011).
- 18 Vitale M, Cantoni C, Pietra G, Mingari MC, Moretta L. Effect of tumor cells and tumor microenvironment on NK-cell function. *Eur. J. Immunol.* 44(6), 1582–1592 (2014).
- 19 Waldhauer I, Steinle A. NK cells and cancer immunosurveillance. *Oncogene* 27(45), 5932–5943 (2008).
- 20 Crane CA, Han SJ, Barry JJ, Ahn BJ, Lanier LL, Parsa AT. TGF-beta downregulates the activating receptor NKG2D on NK cells and CD8⁺ T cells in glioma patients. *Neuro Oncol.* 12(1), 7–13 (2010).
- 21 Baker GJ, Chockley P, Yadav VN *et al.* Natural killer cells eradicate galectin-1-deficient glioma in the absence of adaptive immunity. *Cancer Res.* 74(18), 5079–5090 (2014).
- 22 Robertson FC, Berzofsky JA, Terabe M. NKT cell networks in the regulation of tumor immunity. *Front. Immunol.* 5, 543 (2014).
- 23 Ostrom QT, Gittleman H, Farah P *et al.* CBTRUS statistical report: primary brain and central nervous system tumors diagnosed in the United States in 2006–2010. *Neuro Oncol.* 15(Suppl. 2), ii1–ii56 (2013).
- 24 Stupp R, Hegi ME, Gilbert MR, Chakravarti A. Chemoradiotherapy in malignant glioma: standard of care and future directions. *J. Clin. Oncol.* 25(26), 4127–4136 (2007).
- 25 Ostrom QT, Bauchet L, Davis FG *et al.* The epidemiology of glioma in adults: a “state of the science” review. *Neuro Oncol.* 16(7), 896–913 (2014).
- 26 Burrell RA, McGranahan N, Bartek J, Swanton C. The causes and consequences of genetic heterogeneity in cancer evolution. *Nature* 501(7467), 338–345 (2013).
- 27 Fisher R, Puztai L, Swanton C. Cancer heterogeneity: implications for targeted therapeutics. *Br. J. Cancer* 108(3), 479–485 (2013).
- 28 Patel AP, Tirosh I, Trombetta JJ *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. *Science* 344(6190), 1396–1401 (2014).
- 29 Sottoriva A, Spiteri I, Piccirillo SG *et al.* Intratumor heterogeneity in human glioblastoma reflects cancer evolutionary dynamics. *Proc. Natl Acad. Sci. USA* 110(10), 4009–4014 (2013).
- 30 Kane JR, Miska J, Young JS, Kanojia D, Kim JW, Lesniak MS. Sui generis: gene therapy and delivery systems for the treatment of glioblastoma. *Neuro Oncol.* 17(Suppl. 2), ii24–ii36 (2015).
- 31 Chen KS, Mitchell DA. Monoclonal antibody therapy for malignant glioma. *Adv. Exp. Med. Biol.* 746, 121–141 (2012).
- 32 Morgan RA, Dudley ME, Wunderlich JR *et al.* Cancer regression in patients after transfer of genetically engineered lymphocytes. *Science* 314(5796), 126–129 (2006).
- 33 Morgan RA, Johnson LA, Davis JL *et al.* Recognition of glioma stem cells by genetically modified T cells targeting EGFRvIII and development of adoptive cell therapy for glioma. *Hum. Gene Ther.* 23(10), 1043–1053 (2012).
- 34 Ohno M, Natsume A, Ichiro Iwami K *et al.* Retrovirally engineered T-cell-based immunotherapy targeting type III variant epidermal growth factor receptor, a glioma-associated antigen. *Cancer Sci.* 101(12), 2518–2524 (2010).
- 35 Callahan MK, Postow MA, Wolchok JD. CTLA-4 and PD-1 pathway blockade: combinations in the clinic. *Front. Oncol.* 4, 385 (2014).
- 36 Postow MA, Callahan MK, Wolchok JD. Immune checkpoint blockade in cancer therapy. *J. Clin. Oncol.* 33(17), 1974–1982 (2015).
- 37 Akasaki Y, Kikuchi T, Homma S, Abe T, Kofe D, Ohno T. Antitumor effect of immunizations with fusions of dendritic and glioma cells in a mouse brain tumor model. *J. Immunother.* 24(2), 106–113 (2001).
- 38 Amano T, Kajiwara K, Yoshikawa K *et al.* Antitumor effects of vaccination with dendritic cells transfected with modified receptor for hyaluronan-mediated motility mRNA in a mouse glioma model. *J. Neurosurg.* 106, 638–645 (2007).
- 39 Aoki H, Mizuno M, Natsume A *et al.* Dendritic cells pulsed with tumor extract-cationic liposome complex increase the induction of cytotoxic T lymphocytes in mouse brain tumor. *Cancer Immunol. Immunother.* 50, 463–468 (2001).

- 40 Cho HI, Kim EK, Park SY, Lee SK, Hong YK, Kim TG. Enhanced induction of anti-tumor immunity in human and mouse by dendritic cells pulsed with recombinant TAT fused human survivin protein. *Cancer Lett.* 258, 189–198 (2007).
- 41 Ciesielski MJ, Apfel L, Barone TA, Castro CA, Weiss TC, Fenstermaker RA. Antitumor effects of a xenogeneic survivin bone marrow derived dendritic cell vaccine against murine GL261 gliomas. *Cancer Immunol. Immunother.* 55, 1491–1503 (2006).
- 42 Ciesielski MJ, Kozbor D, Castanaro CA, Barone TA, Fenstermaker RA. Therapeutic effect of a T helper cell supported CTL response induced by a survivin peptide vaccine against murine cerebral glioma. *Cancer Immunol. Immunother.* 57, 1827–1835 (2008).
- 43 Clavreul A, Delhaye M, Jadaud E, Menei P. Effects of syngeneic cellular vaccinations alone or in combination with GM-CSF on the weakly immunogenic F98 glioma model. *J. Neurooncol.* 79, 9–17 (2006).
- 44 Driessens G, Hamdane M, Cool V, Velu T, Bruyns C. Highly successful therapeutic vaccinations combining dendritic cells and tumor cells secreting granulocyte macrophage colony-stimulating factor. *Cancer Res.* 64, 8435–8442 (2004).
- 45 Fujita M, Zhu X, Ueda R *et al.* Effective immunotherapy against murine gliomas using type 1 polarizing dendritic cells—significant roles of CXCL10. *Cancer Res.* 69, 1587–1595 (2009).
- 46 Grauer OM, Suttmuller RPM, Van Maren W *et al.* Elimination of regulatory T cells is essential for an effective vaccination with tumor lysate-pulsed dendritic cells in a murine glioma model. *Int. J. Cancer* 122, 1794–1802 (2008).
- 47 Herrlinger U, Aulwurm S, Strik H, Weit S, Naumann U, Weller M. MIP-1 α antagonizes the effect of a GM-CSF-enhanced subcutaneous vaccine in a mouse glioma model. *J. Neurooncol.* 66, 147–154 (2004).
- 48 Insug O, Ku G, Ertl HCJ, Blaszczyk-Thurin M. A dendritic cell vaccine induces protective immunity to intracranial growth of glioma. *Anticancer Res.* 22(2A), 613–621 (2002).
- 49 Jouanneau E, Poujol D, Gulia S *et al.* Dendritic cells are essential for priming but inefficient for boosting antitumor immune response in an orthotopic murine glioma model. *Cancer Immunol.* 55, 254–267 (2006).
- 50 Kim CH, Hong MJ, Park SD *et al.* Enhancement of anti-tumor immunity specific to murine glioma by vaccination with tumor cell lysate-pulsed dendritic cells engineered to produce interleukin-12. *Cancer Immunol. Immunother.* 55(11), 1309–1319 (2006).
- 51 Kim CH, Yoon JS, Sohn HJ *et al.* Direct vaccination with pseudotype baculovirus expressing murine telomerase induces anti-tumor immunity comparable with RNA-electroporated dendritic cells in a murine glioma model. *Cancer Lett.* 250, 276–283 (2007).
- 52 Kim C-H, Woo S-J, Park J-S *et al.* Enhanced antitumor immunity by combined use of temozolomide and TAT-survivin pulsed dendritic cells in a murine glioma. *Immunology* 122, 615–622 (2007).
- 53 Kjaergaard J, Wang L-X, Kuriyama H, Shu S, Plautz GE. Active immunotherapy for advanced intracranial murine tumors by using dendritic cell-tumor cell fusion vaccines. *J. Neurosurg.* 103, 156–164 (2005).
- 54 Kuwashima N, Nishimura F, Eguchi J *et al.* Delivery of dendritic cells engineered to secrete IFN- α into central nervous system tumors enhances the efficacy of peripheral tumor cell vaccines: dependence on apoptotic pathways. *J. Immunol.* 175, 2730–2740 (2005).
- 55 Liao LM, Black KL, Prins RM *et al.* Treatment of intracranial gliomas with bone marrow-derived dendritic cells pulsed with tumor antigens. *J. Neurosurg.* 90, 1115–1124 (1999).
- 56 Ni HT, Spellman SR, Jean WC, Hall WA, Low WC. Immunization with dendritic cells pulsed with tumor extract increases survival of mice bearing intracranial gliomas. *J. Neurooncol.* 51, 1–9 (2001).
- 57 Pellegatta S, Finocchiaro G. Cell therapies in neuro-oncology. *Neurol. Sci.* 26, 43–45 (2005).
- 58 Prins RM, Odesa SK, Liao LM. Immunotherapeutic targeting of shared melanoma-associated antigens in a murine glioma model. *Cancer Res.* 63, 8487–8491 (2003).
- 59 Saito R, Mizuno M, Nakahara N *et al.* Vaccination with tumor cell lysate-pulsed dendritic cells augments the effect of IFN- β gene therapy for malignant glioma in an experimental mouse intracranial glioma. *Int. J. Cancer* 111, 777–782 (2004).
- 60 Takagi Y, Kikuchi T, Niimura M, Ohno T. Effects of glucocorticoids on antitumor effects of immunizations with fusions of dendritic and tumor cells. *Anticancer Res.* 23, 2553–2558 (2003).
- 61 Witham TF, Erff ML, Okada H *et al.* 7-Hydroxystaurosporine-induced apoptosis in 9L glioma cells provides an effective antigen source for dendritic cells and yields a potent vaccine strategy in an intracranial glioma model. *Neurosurgery* 50, 1327–1335 (2002).
- 62 Xu Q, Liu G, Yuan X *et al.* Antigen-specific T-cell response from dendritic cell vaccination using cancer stem-like cell-associated antigens. *Stem Cells* 27, 1734–1740 (2009).
- 63 Yamanaka R, Zullo S, Ramsey J *et al.* Marked enhancement of antitumor immune responses in mouse brain tumor models by genetically modified dendritic cells producing Semliki Forest virus-mediated interleukin-12. *J. Neurosurg.* 97, 611–618 (2002).
- 64 Zhang Z, Tang L-L, Zhan R-Y, Tong Y, Yao H-P, Du L-A. Immunotherapy of intracranial G422 glioblastoma with dendritic cells pulsed with tumor extract or RNA. *J. Zhejiang Univ. Sci.* 5, 1298–1303 (2004).
- 65 Zhu X, Lu C, Xiao B, Qiao J, Sun Y. An experimental study of dendritic cells-mediated immunotherapy against intracranial gliomas in rats. *J. Neurooncol.* 74, 9–17 (2005).
- 66 Maes W, Rosas GG, Verbinnen B *et al.* DC vaccination with anti-CD25 treatment leads to long-term immunity against experimental glioma. *Neuro Oncol.* 11, 529–542 (2009).
- 67 Jiang X-B, Lu X-L, Hu P, Liu R-E. Improved therapeutic efficacy using vaccination with glioma lysate-pulsed dendritic cells combined with IP-10 in murine glioma. *Vaccine* 27, 6210–6216 (2009).

- 68 Saka M, Amano T, Kajiwara K *et al.* Vaccine therapy with dendritic cells transfected with IL13ra2 mRNA for glioma in mice. *J. Neurosurg.* 113, 270–279 (2010).
- 69 Kim T-G, Kim C-H, Park J-S *et al.* Immunological factors relating to the antitumor effect of temozolomide chemioimmunotherapy in a murine glioma model. *Clin. Vaccine Immunol.* 17, 143–153 (2010).
- 70 Xiao Z-Y, Tang H, Xu Z-M *et al.* An experimental study of dendritic cells transfected with cancer stem-like cells RNA against 9L brain tumors. *Cancer Biol. Ther.* 11, 974–980 (2011).
- 71 Mineharu Y, King GD, Muhammad AKMG *et al.* Engineering the brain tumor microenvironment enhances the efficacy of dendritic cell vaccination: implications for clinical trial design. *Clin. Cancer Res.* 17, 4705–4718 (2011).
- 72 Feng E, Gao H, Su W, Yu C. Immunotherapy of rat glioma without accumulation of CD4(+)CD25(+)FOXP3(+) regulatory T cells. *Neural Regen. Res.* 7, 1498–1506 (2012).
- 73 Zhang H, Tian M, Xiu C, Wang Y, Tang G. Enhancement of antitumor activity by combination of tumor lysate-pulsed dendritic cells and celecoxib in a rat glioma model. *Oncol. Res.* 20, 447–455 (2013).
- 74 Peng W, Nan Z, Liu Y *et al.* Dendritic cells transduced with CPEB4 induced antitumor immune response. *Exp. Mol. Pathol.* 97, 273–278 (2014).
- 75 Bu N, Wu H, Zhang G *et al.* Exosomes from dendritic cells loaded with chaperone-rich cell lysates elicit a potent T cell immune response against intracranial glioma in mice. *J. Mol. Neurosci.* 56(3), 631–643 (2015).
- 76 Li M, Wang B, Wu Z *et al.* A novel recombinant protein of ephrinA1–PE38/GM-CSF activate dendritic cells vaccine in rats with glioma. *Tumor Biol.* 36(7), 5497–5503 (2015).
- 77 Bloch O, Crane CA, Fuks Y *et al.* Heat-shock protein peptide complex-96 vaccination for recurrent glioblastoma: a Phase II, single-arm trial. *Neuro Oncol.* 16(2), 274–279 (2014).
- 78 Buchroithner J, Pichler J, Marosi C *et al.* Vascular endothelial growth factor targeted therapy may improve the effect of dendritic cell-based cancer immune therapy. *Int. J. Clin. Pharmacol. Ther.* 52(1), 76–77 (2014).
- 79 Ishikawa E, Muragaki Y, Yamamoto T *et al.* Phase I/IIa trial of fractionated radiotherapy, temozolomide, and autologous formalin-fixed tumor vaccine for newly diagnosed glioblastoma. *J. Neurosurg.* 121(3), 543–553 (2014).
- 80 Mitchell DA, Batich KA, Gunn MD *et al.* Tetanus toxoid and CCL3 improve dendritic cell vaccines in mice and glioblastoma patients. *Nature* 519, 366–369 (2015).
- **Demonstrates increased efficiency of dendritic cell (DC) vaccination when the vaccine site was preconditioned with tetanus toxoid. This led to more than 50% of patients surviving longer than 40 months. In mice, it was shown that tetanus toxoid-enhanced DC migration to the lymph nodes draining the vaccine site and that this therapy suppressed tumor growth in a CCL3-dependent manner.**
- 81 Phuphanich S, Wheeler CJ, Rudnick JD *et al.* Phase I trial of a multi-epitope-pulsed dendritic cell vaccine for patients with newly diagnosed glioblastoma. *Cancer Immunol. Immunother.* 62(1), 125–135 (2013).
- **Report of a Phase I clinical trial using autologous DC vaccine pulsed with six GAA (HER2, TRP-2, gp100, MAGE-1, IL13Ra2 and AIM-2) demonstrating response to treatment of 33% of the 21 patients enrolled, increased production of IFN- γ and TNF- α in stimulated CD8 $^+$ T cells, which correlated with survival, decreased expression of CD133 in recurrent GBM patients, suggesting that the treatment killed GSC cells. The median progression-free survival was 15.9 months and the median overall survival was 38.4 months, larger than other studies.**
- 82 Prins RM, Soto H, Konkankit V *et al.* Gene expression profile correlates with T-cell infiltration and relative survival in glioblastoma patients vaccinated with dendritic cell immunotherapy. *Clin. Cancer Res.* 17(6), 1603–1615 (2011).
- 83 Riedmann EM. A genus cancer vaccine extends survival of GBM patients. *Hum. Vaccin. Immunother.* 10(2), 252–253 (2014).
- 84 Schuster J, Lai RK, Recht LD *et al.* A Phase II, multicenter trial of rindopepimut (CDX-110) in newly diagnosed glioblastoma: the ACT III study. *Neuro Oncol.* 17(6), 854–861 (2015).
- 85 Vik-Mo EO, Nyakas M, Mikkelsen BV *et al.* Therapeutic vaccination against autologous cancer stem cells with mRNA-transfected dendritic cells in patients with glioblastoma. *Cancer Immunol. Immunother.* 62(9), 1499–1509 (2013).
- 86 Colombo F, Barzon L, Franchin E *et al.* Combined HSV-TK/IL-2 gene therapy in patients with recurrent glioblastoma multiforme: biological and clinical results. *Cancer Gene Ther.* 12(10), 835–848 (2005).
- 87 Kroeger KM, Muhammad AK, Baker GJ *et al.* Gene therapy and virotherapy: novel therapeutic approaches for brain tumors. *Discov. Med.* 10(53), 293–304 (2010).
- 88 Kwiatkowska A, Nandhu MS, Behera P, Chiocca EA, Viapiano MS. Strategies in gene therapy for glioblastoma. *Cancers (Basel)* 5(4), 1271–1305 (2013).
- 89 Okada H, Pollack IF, Lotze MT *et al.* Gene therapy of malignant gliomas: a Phase I study of IL-4-HSV-TK gene-modified autologous tumor to elicit an immune response. *Hum. Gene Ther.* 11(4), 637–653 (2000).
- 90 Palu G, Cavaggioni A, Calvi P *et al.* Gene therapy of glioblastoma multiforme via combined expression of suicide and cytokine genes: a pilot study in humans. *Gene Ther.* 6(3), 330–337 (1999).
- 91 Rampling R, Cruickshank G, Papanastassiou V *et al.* Toxicity evaluation of replication-competent herpes simplex virus (ICP 34.5 null mutant 1716) in patients with recurrent malignant glioma. *Gene Ther.* 7(10), 859–866 (2000).
- 92 Sonabend AM, Ulasov IV, Lesniak MS. Gene therapy trials for the treatment of high-grade gliomas. *Gene Ther. Mol. Biol.* 11(A), 79–92 (2007).
- 93 Tobias A, Ahmed A, Moon KS, Lesniak MS. The art of gene therapy for glioma: a review of the challenging road to the bedside. *J. Neurol. Neurosurg. Psychiatry* 84(2), 213–222 (2013).
- 94 Prins RM, Wang X, Soto H *et al.* Comparison of glioma-associated antigen peptide-loaded versus autologous tumor lysate-loaded dendritic cell vaccination in malignant glioma patients. *J. Immunother.* 36(2), 152–157 (2013).

- 95 Akiyama Y, Oshita C, Kume A *et al.* Alpha-type-1 polarized dendritic cell-based vaccination in recurrent high-grade glioma: a Phase I clinical trial. *BMC Cancer* 12, 623 (2012).
- 96 Liao LM, Prins RM, Kiertscher SM *et al.* Dendritic cell vaccination in glioblastoma patients induces systemic and intracranial T-cell responses modulated by the local central nervous system tumor microenvironment. *Clin. Cancer Res.* 11(15), 5515–5525 (2005).
- 97 Yu JS, Liu G, Ying H, Yong WH, Black KL, Wheeler CJ. Vaccination with tumor lysate-pulsed dendritic cells elicits antigen-specific, cytotoxic T-cells in patients with malignant glioma. *Cancer Res.* 64(14), 4973–4979 (2004).
- 98 Crane CA, Han SJ, Ahn B *et al.* Individual patient-specific immunity against high-grade glioma after vaccination with autologous tumor derived peptides bound to the 96 KD chaperone protein. *Clin. Cancer Res.* 19(1), 205–214 (2013).
- 99 Izumoto S, Tsuboi A, Oka Y *et al.* Phase II clinical trial of Wilms tumor 1 peptide vaccination for patients with recurrent glioblastoma multiforme. *J. Neurosurg.* 108(5), 963–971 (2008).
- 100 Morita S, Oka Y, Tsuboi A *et al.* A Phase I/II trial of a WT1 (Wilms' tumor gene) peptide vaccine in patients with solid malignancy: safety assessment based on the Phase I data. *Jpn J. Clin. Oncol.* 36(4), 231–236 (2006).
- 101 Okada H, Butterfield LH, Hamilton RL *et al.* Induction of robust type-I CD8⁺ T-cell responses in WHO grade 2 low-grade glioma patients receiving peptide-based vaccines in combination with poly-ICLC. *Clin. Cancer Res.* 21(2), 286–294 (2015).
- 102 Pollack IF, Jakacki RI, Butterfield LH *et al.* Antigen-specific immune responses and clinical outcome after vaccination with glioma-associated antigen peptides and polyinosinic-polycytidylic acid stabilized by lysine and carboxymethylcellulose in children with newly diagnosed malignant brainstem and nonbrainstem gliomas. *J. Clin. Oncol.* 32(19), 2050–2058 (2014).
- 103 Schuessler A, Smith C, Beagley L *et al.* Autologous T-cell therapy for cytomegalovirus as a consolidative treatment for recurrent glioblastoma. *Cancer Res.* 74(13), 3466–3476 (2014).
- 104 Brown CE, Badie B, Barish ME *et al.* Bioactivity and safety of IL13Ralpha2-redirected chimeric antigen receptor CD8⁺ T cells in patients with recurrent glioblastoma. *Clin. Cancer Res.* 21(18), 4062–4072 (2015).
- 105 Massimino M, Biassoni V, Miceli R *et al.* Results of nimotuzumab and vinorelbine, radiation and re-irradiation for diffuse pontine glioma in childhood. *J. Neurooncol.* 118(2), 305–312 (2014).
- 106 Sampson JH, Schmittling RJ, Archer GE *et al.* A pilot study of IL-2Ralpha blockade during lymphopenia depletes regulatory T-cells and correlates with enhanced immunity in patients with glioblastoma. *PLoS ONE* 7(2), e31046 (2012).
- 107 Hodi FS, O'Day SJ, McDermott DF *et al.* Improved survival with ipilimumab in patients with metastatic melanoma. *N. Engl. J. Med.* 363(8), 711–723 (2010).
- 108 Hamid O, Robert C, Daud A *et al.* Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. *N. Engl. J. Med.* 369(2), 134–144 (2013).
- 109 Bloch O. Immunotherapy for malignant gliomas. *Cancer Treat. Res.* 163, 143–158 (2015).
- 110 Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat. Rev. Cancer* 12(4), 252–264 (2012).
- 111 Fecci PE, Ochiai H, Mitchell DA *et al.* Systemic CTLA-4 blockade ameliorates glioma-induced changes to the CD4⁺ T cell compartment without affecting regulatory T-cell function. *Clin. Cancer Res.* 13(7), 2158–2167 (2007).
- 112 Chambers CA, Kuhns MS, Egen JG, Allison JP. CTLA-4-mediated inhibition in regulation of T cell responses: mechanisms and manipulation in tumor immunotherapy. *Annu. Rev. Immunol.* 19, 565–594 (2001).
- 113 Freeman GJ, Long AJ, Iwai Y *et al.* Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J. Exp. Med.* 192(7), 1027–1034 (2000).
- 114 Chemnitz JM, Parry RV, Nichols KE, June CH, Riley JL. SHP-1 and SHP-2 associate with immunoreceptor tyrosine-based switch motif of programmed death 1 upon primary human T cell stimulation, but only receptor ligation prevents T cell activation. *J. Immunol.* 173(2), 945–954 (2004).
- 115 Ribas A, Kefford R, Marshall MA *et al.* Phase III randomized clinical trial comparing tremelimumab with standard-of-care chemotherapy in patients with advanced melanoma. *J. Clin. Oncol.* 31(5), 616–622 (2013).
- 116 Lynch TJ, Bondarenko I, Luft A *et al.* Ipilimumab in combination with paclitaxel and carboplatin as first-line treatment in stage IIIB/IV non-small-cell lung cancer: results from a randomized, double-blind, multicenter Phase II study. *J. Clin. Oncol.* 30(17), 2046–2054 (2012).
- 117 Yang JC, Hughes M, Kammula U *et al.* Ipilimumab (anti-CTLA4 antibody) causes regression of metastatic renal cell cancer associated with enteritis and hypophysitis. *J. Immunother.* 30(8), 825–830 (2007).
- 118 Vom Berg J, Vrohling M, Haller S *et al.* Intratumoral IL-12 combined with CTLA-4 blockade elicits T cell-mediated glioma rejection. *J. Exp. Med.* 210(13), 2803–2811 (2013).
- 119 Agarwalla P, Barnard Z, Fecci P, Dranoff G, Curry WT Jr. Sequential immunotherapy by vaccination with GM-CSF-expressing glioma cells and CTLA-4 blockade effectively treats established murine intracranial tumors. *J. Immunother.* 35(5), 385–389 (2012).
- 120 Robert C, Long GV, Brady B *et al.* Nivolumab in previously untreated melanoma without BRAF mutation. *N. Engl. J. Med.* 372(4), 320–330 (2015).
- 121 Topalian SL, Hodi FS, Brahmer JR *et al.* Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N. Engl. J. Med.* 366(26), 2443–2454 (2012).
- 122 Zeng J, See AP, Phallen J *et al.* Anti-PD-1 blockade and stereotactic radiation produce long-term survival in mice with intracranial gliomas. *Int. J. Radiat. Oncol. Biol. Phys.* 86(2), 343–349 (2013).
- 123 Wainwright DA, Chang AL, Dey M *et al.* Durable therapeutic efficacy utilizing combinatorial blockade

- against IDO, CTLA-4, and PD-L1 in mice with brain tumors. *Clin. Cancer Res.* 20(20), 5290–5301 (2014).
- Shows that combinatorial therapeutic inhibition of IDO, CTLA-4 and PD-L1 induces 100% long-term survival in a mouse model of well-established GBMs. This was accompanied by a decrease in tumor-infiltrating Tregs, coincident with a significant increase in CD8⁺IFN γ ⁺ T cells and illustrates the benefits of multiagent targeting of immunosuppressive agents for the treatment of GBM.
- 124 Gelao L, Criscitiello C, Esposito A, Goldhirsch A, Curigliano G. Immune checkpoint blockade in cancer treatment: a double-edged sword cross-targeting the host as an “innocent bystander”. *Toxins (Basel)* 6(3), 914–933 (2014).
- 125 Weber JS, Kahler KC, Hauschild A. Management of immune-related adverse events and kinetics of response with ipilimumab. *J. Clin. Oncol.* 30(21), 2691–2697 (2012).
- 126 Slovin SF, Higanó CS, Hamid O *et al.* Ipilimumab alone or in combination with radiotherapy in metastatic castration-resistant prostate cancer: results from an open-label, multicenter Phase I/II study. *Ann. Oncol.* 24(7), 1813–1821 (2013).
- 127 Kaehler KC, Piel S, Livingstone E, Schilling B, Hauschild A, Schadendorf D. Update on immunologic therapy with anti-CTLA-4 antibodies in melanoma: identification of clinical and biological response patterns, immune-related adverse events, and their management. *Semin. Oncol.* 37(5), 485–498 (2010).
- 128 Brahmer JR, Tykodi SS, Chow LQ *et al.* Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N. Engl. J. Med.* 366(26), 2455–2465 (2012).
- 129 Juratli TA, Schackert G, Krex D. Current status of local therapy in malignant gliomas – a clinical review of three selected approaches. *Pharmacol. Ther.* 139(3), 341–358 (2013).
- 130 Ahmed N, Salsman VS, Kew Y *et al.* HER2-specific T cells target primary glioblastoma stem cells and induce regression of autologous experimental tumors. *Clin. Cancer Res.* 16(2), 474–485 (2010).
- 131 Balyasnikova IV, Ferguson SD, Sengupta S, Han Y, Lesniak MS. Mesenchymal stem cells modified with a single-chain antibody against EGFRvIII successfully inhibit the growth of human xenograft malignant glioma. *PLoS ONE* 5(3), e9750 (2010).
- 132 Kahlon KS, Brown C, Cooper LJ, Raubitschek A, Forman SJ, Jensen MC. Specific recognition and killing of glioblastoma multiforme by interleukin 13-zetakine redirected cytolytic T cells. *Cancer Res.* 64(24), 9160–9166 (2004).
- 133 Curtin JF, Liu N, Candolfi M *et al.* HMGB1 mediates endogenous TLR2 activation and brain tumor regression. *PLoS Med.* 6(1), e10 (2009).
- 134 Candolfi M, Yagiz K, Wibowo M *et al.* Temozolomide does not impair gene therapy-mediated antitumor immunity in syngeneic brain tumor models. *Clin. Cancer Res.* 20(6), 1555–1565 (2014).
- 135 Workenhe ST, Mossman KL. Oncolytic virotherapy and immunogenic cancer cell death: sharpening the sword for improved cancer treatment strategies. *Mol. Ther.* 22(2), 251–256 (2014).
- 136 Krysko DV, Garg AD, Kaczmarek A, Krysko O, Agostinis P, Vandenabeele P. Immunogenic cell death and DAMPs in cancer therapy. *Nat. Rev. Cancer* 12(12), 860–875 (2012).
- 137 Okura H, Smith C, Rutka J. Gene therapy for malignant glioma. *Mol. Cell Ther.* 2(1), 1–19 (2014).
- 138 Oldfield EH, Ram Z, Culver KW, Blaese RM, Devroom HL, Anderson WF. Gene therapy for the treatment of brain tumors using intra-tumoral transduction with the thymidine kinase gene and intravenous ganciclovir. *Hum. Gene Ther.* 4(1), 39–69 (1993).
- 139 Puntel M, Muhammad AK, Candolfi M *et al.* A novel bicistronic high-capacity gutless adenovirus vector that drives constitutive expression of herpes simplex virus type 1 thymidine kinase and tet-inducible expression of Flt3L for glioma therapeutics. *J. Virol.* 84(12), 6007–6017 (2010).
- 140 Grandi P, Peruzzi P, Reinhart B, Cohen JB, Chiocca EA, Glorioso JC. Design and application of oncolytic HSV vectors for glioblastoma therapy. *Expert Rev. Neurother.* 9(4), 505–517 (2009).
- 141 Markert JM, Liechty PG, Wang W *et al.* Phase Ib trial of mutant herpes simplex virus G207 inoculated pre-and post-tumor resection for recurrent GBM. *Mol. Ther. J. Am. Soc. Gene Ther.* 17(1), 199–207 (2009).
- 142 Markert JM, Medlock MD, Rabkin SD *et al.* Conditionally replicating herpes simplex virus mutant, G207 for the treatment of malignant glioma: results of a Phase I trial. *Gene Ther.* 7(10), 867–874 (2000).
- 143 Kirn D, Hermiston T, McCormick F. ONYX-015: clinical data are encouraging. *Nat. Med.* 4(12), 1341–1342 (1998).
- 144 Chiocca EA, Abbed KM, Tatter S *et al.* A Phase I open-label, dose-escalation, multi-institutional trial of injection with an E1B-attenuated adenovirus, ONYX-015, into the peritumoral region of recurrent malignant gliomas, in the adjuvant setting. *Mol. Ther. J. Am. Soc. Gene Ther.* 10(5), 958–966 (2004).
- 145 Fueyo J, Gomez-Manzano C, Alemany R *et al.* A mutant oncolytic adenovirus targeting the Rb pathway produces anti-glioma effect *in vivo*. *Oncogene* 19(1), 2–12 (2000).
- 146 Anderson BD, Nakamura T, Russell SJ, Peng KW. High CD46 receptor density determines preferential killing of tumor cells by oncolytic measles virus. *Cancer Res.* 64(14), 4919–4926 (2004).
- 147 Wollmann G, Ozduman K, van den Pol AN. Oncolytic virus therapy for glioblastoma multiforme: concepts and candidates. *Cancer J.* 18(1), 69–81 (2012).
- 148 Allen C, Paraskevovou G, Iankov I *et al.* Interleukin-13 displaying retargeted oncolytic measles virus strains have significant activity against gliomas with improved specificity. *Mol. Ther. J. Am. Soc. Gene Ther.* 16(9), 1556–1564 (2008).
- 149 Uchida H, Marzulli M, Nakano K *et al.* Effective treatment of an orthotopic xenograft model of human glioblastoma using an EGFR-retargeted oncolytic herpes simplex virus. *Mol. Ther.* 21(3), 561–569 (2013).
- 150 Qin XQ, Beckham C, Brown JL, Lukashev M, Barsoum J. Human and mouse IFN-beta gene therapy exhibits different anti-tumor mechanisms in mouse models. *Mol. Ther.* 4(4), 356–364 (2001).
- 151 Qin XQ, Tao N, Dergay A *et al.* Interferon-beta gene therapy inhibits tumor formation and causes regression of established tumors in immune-deficient mice. *Proc. Natl Acad. Sci. USA* 95(24), 14411–14416 (1998).

- 152 Mizuno M, Yoshida J. Effect of human interferon beta gene transfer upon human glioma, transplanted into nude mouse brain, involves induced natural killer cells. *Cancer Immunol. Immunother.* 47(4), 227–232 (1998).
- 153 Yoshida J, Mizuno M, Fujii M *et al.* Human gene therapy for malignant gliomas (glioblastoma multiforme and anaplastic astrocytoma) by *in vivo* transduction with human interferon beta gene using cationic liposomes. *Hum. Gene Ther.* 15(1), 77–86 (2004).
- 154 Chiocca EA, Smith KM, McKinney B *et al.* A Phase I trial of Ad.hIFN-beta gene therapy for glioma. *Mol. Ther.* 16(3), 618–626 (2008).
- 155 Ehteshami M, Samoto K, Kabos P *et al.* Treatment of intracranial glioma with *in situ* interferon-gamma and tumor necrosis factor-alpha gene transfer. *Cancer Gene Ther.* 9(11), 925–934 (2002).
- 156 Chen B, Timiryasova TM, Andres ML *et al.* Evaluation of combined vaccinia virus-mediated antitumor gene therapy with p53, IL-2, and IL-12 in a glioma model. *Cancer Gene Ther.* 7(11), 1437–1447 (2000).
- 157 Maraskovsky E, Brasel K, Teepe M *et al.* Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J. Exp. Med.* 184(5), 1953–1962 (1996).
- 158 Ali S, Curtin JF, Zirger JM *et al.* Inflammatory and anti-glioma effects of an adenovirus expressing human soluble Fms-like tyrosine kinase 3 ligand (hsFlt3L): treatment with hsFlt3L inhibits intracranial glioma progression. *Mol. Ther.* 10(6), 1071–1084 (2004).
- 159 Ali S, King GD, Curtin JF *et al.* Combined immunostimulation and conditional cytotoxic gene therapy provide long-term survival in a large glioma model. *Cancer Res.* 65(16), 7194–7204 (2005).
- 160 King GD, Muhammad AK, Larocque D *et al.* Combined *Flt3L/TK* gene therapy induces immunological surveillance which mediates an immune response against a surrogate brain tumor neoantigen. *Mol. Ther.* 19(10), 1793–1801 (2011).
- 161 Candolfi M, Curtin JF, Yagiz K *et al.* B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. *Neoplasia* 13(10), 947–960 (2011).
- 162 King GD, Muhammad AK, Curtin JF *et al.* *Flt3L* and *TK* gene therapy eradicate multifocal glioma in a syngeneic glioblastoma model. *Neuro Oncol.* 10(1), 19–31 (2008).
- 163 Rock KL, Clark K. Analysis of the role of MHC class II presentation in the stimulation of cytotoxic T lymphocytes by antigens targeted into the exogenous antigen-MHC class I presentation pathway. *J. Immunol.* 156(10), 3721–3726 (1996).
- 164 Rock KL, Gamble S, Rothstein L. Presentation of exogenous antigen with class I major histocompatibility complex molecules. *Science* 249, 918–921 (1990).
- 165 Gustafsson K, Junevik K, Werlenius O, Holmgren S, Karlsson-Parra A, Andersson P-O. Tumour-loaded α -type 1-polarized dendritic cells from patients with chronic lymphocytic leukaemia produce a superior NK-, NKT- and CD8⁺ T cell-attracting chemokine profile. *Scand. J. Immunol.* 74, 318–326 (2011).
- 166 Shimizu K, Fujii S-I. DC therapy induces long-term NK reactivity to tumors via host DC. *Eur. J. Immunol.* 39, 457–468 (2009).
- 167 Lafferty KJ, Warren HS, Woolnough JA. A mediator acting as a costimulator for the development of cytotoxic responses *in vitro*. *Adv. Exp. Med. Biol.* 114, 497–501 (1979).
- 168 Capece D, Verzella D, Fischietti M, Zazzeroni F, Alesse E. Targeting costimulatory molecules to improve antitumor immunity. *J. Biomed. Biotech.* 2012, 1–17 (2012).
- 169 Joffre O, Nolte MA, Spörri R, Reis E Sousa C. Inflammatory signals in dendritic cell activation and the induction of adaptive immunity. *Immunol. Rev.* 227, 234–247 (2009).
- 170 Cohen PA, Koski GK, Czerniecki BJ *et al.* STAT3- and STAT5-dependent pathways competitively regulate the pan-differentiation of CD34pos cells into tumor-competent dendritic cells. *Blood* 112, 1832–1843 (2008).
- 171 Mailliard RB, Wankowicz-Kalinska A, Cai Q *et al.* Alpha-type-1 polarized dendritic cells: a novel immunization tool with optimized CTL-inducing activity. *Cancer Res.* 64, 5934–5937 (2004).
- 172 Okada H. Brain tumor immunotherapy with type-1 polarizing strategies. *Ann. NY Acad. Sci.* 1174, 18–23 (2009).
- 173 Mineharu Y, Castro MG, Lowenstein PR, Sakai N, Miyamoto S. Dendritic cell-based immunotherapy for glioma: multiple regimens and implications in clinical trials. *Neurol. Med. Chir. (Tokyo)* 53, 741–754 (2013).
- 174 Ohlfest JR, Andersen BM, Litterman AJ *et al.* Vaccine injection site matters: qualitative and quantitative defects in CD8 T cells primed as a function of proximity to the tumor in a murine glioma model. *J. Immunol.* 190(2), 613–620 (2013).
- 175 Yamanaka R, Zullo SA, Tanaka R, Ramsey J, Blaese M, Xanthopoulos KG. Induction of a therapeutic antitumor immunological response by intratumoral injection of genetically engineered Semliki Forest virus to produce interleukin-12. *Neurosurg. Focus* 9, e7 (2000).
- 176 Jenkins M, Schwartz R. Antigen presentation by chemically modified splenocytes induces antigen-specific T cell unresponsiveness *in vitro* and *in vivo*. *J. Exp. Med.* 165, 302–319 (1987).
- 177 Zeppernick F, Ahmadi R, Campos B *et al.* Stem cell marker CD133 affects clinical outcome in glioma patients. *Clin. Cancer Res.* 14(1), 123–129 (2008).
- 178 Pallini R, Ricci-Vitiani L, Montano N *et al.* Expression of the stem cell marker CD133 in recurrent glioblastoma and its value for prognosis. *Cancer* 117(1), 162–174 (2011).
- 179 Okada H, Kalinski P, Ueda R *et al.* Induction of CD8⁺ T-cell responses against novel glioma-associated antigen peptides and clinical activity by vaccinations with α -type 1 polarized dendritic cells and polyinosinic-polycytidylic acid stabilized by lysine and carboxymethylcellulose in patients with recurrent malignant glioma. *J. Clin. Oncol.* 29(3), 330–336 (2011).
- 180 Esparza R, Azad TD, Feroze AH, Mitra SS, Cheshier SH. Glioblastoma stem cells and stem cell-targeting immunotherapies. *J. Neurooncol.* 123(3), 449–457 (2015).
- 181 Gan HK, Kaye AH, Luwor RB. The EGFRvIII variant in glioblastoma multiforme. *J. Clin. Neurosci.* 16(6), 748–754 (2009).

- 182 Heimberger AB, Suki D, Yang D, Shi W, Aldape K. The natural history of EGFR and EGFRvIII in glioblastoma patients. *J. Transl. Med.* 3, 38 (2005).
- 183 Heimberger AB, Crotty LE, Archer GE *et al.* Epidermal growth factor receptor VIII peptide vaccination is efficacious against established intracerebral tumors. *Clin. Cancer Res.* 9(11), 4247–4254 (2003).
- 184 Mishima K, Johns TG, Luwor RB *et al.* Growth suppression of intracranial xenografted glioblastomas overexpressing mutant epidermal growth factor receptors by systemic administration of monoclonal antibody (mAb) 806, a novel monoclonal antibody directed to the receptor. *Cancer Res.* 61(14), 5349–5354 (2001).
- 185 Babu R, Adamson DC. Rindopepimut: an evidence-based review of its therapeutic potential in the treatment of EGFRvIII-positive glioblastoma. *Core Evid.* 7, 93–103 (2012).
- 186 Sampson JH, Heimberger AB, Archer GE *et al.* Immunologic escape after prolonged progression-free survival with epidermal growth factor receptor variant III peptide vaccination in patients with newly diagnosed glioblastoma. *J. Clin. Oncol.* 28(31), 4722–4729 (2010).
- 187 Reardon DA, Wucherpfennig KW, Freeman G *et al.* An update on vaccine therapy and other immunotherapeutic approaches for glioblastoma. *Expert Rev. Vaccines* 12(6), 597–615 (2013).
- 188 Caudill MM, Li Z. HSPPC-96: a personalised cancer vaccine. *Expert Opin. Biol. Ther.* 1(3), 539–547 (2001).
- 189 Binder RJ, Srivastava PK. Essential role of CD91 in representation of gp96-chaperoned peptides. *Proc. Natl Acad. Sci. USA* 101(16), 6128–6133 (2004).
- 190 Chandramohan V, Mitchell DA, Johnson LA, Sampson JH, Bigner DD. Antibody, T-cell and dendritic cell immunotherapy for malignant brain tumors. *Future Oncol.* 9(7), 977–990 (2013).
- 191 Jouanneau E, Black KL, Veiga L *et al.* Intrinsically de-sialylated CD103(+) CD8 T cells mediate beneficial anti-glioma immune responses. *Cancer Immunol. Immunother.* 63(9), 911–924 (2014).
- 192 Ikeda H, Shiku H. Antigen-receptor gene-modified T cells for treatment of glioma. *Adv. Exp. Med. Biol.* 746, 202–215 (2012).
- 193 Chekmasova AA, Rao TD, Nikhamin Y *et al.* Successful eradication of established peritoneal ovarian tumors in SCID-Beige mice following adoptive transfer of T cells genetically targeted to the MUC16 antigen. *Clin. Cancer Res.* 16(14), 3594–3606 (2010).
- 194 Kochenderfer JN, Yu Z, Frasher D, Restifo NP, Rosenberg SA. Adoptive transfer of syngeneic T cells transduced with a chimeric antigen receptor that recognizes murine CD19 can eradicate lymphoma and normal B cells. *Blood* 116(19), 3875–3886 (2010).
- 195 Zhao Y, Wang QJ, Yang S *et al.* A herceptin-based chimeric antigen receptor with modified signaling domains leads to enhanced survival of transduced T lymphocytes and antitumor activity. *J. Immunol.* 183(9), 5563–5574 (2009).
- 196 Mintz A, Gibo DM, Slagle-Webb B, Christensen ND, Debinski W. IL-13Ralpha2 is a glioma-restricted receptor for interleukin-13. *Neoplasia* 4(5), 388–399 (2002).
- 197 Liu G, Ying H, Zeng G, Wheeler CJ, Black KL, Yu JS. HER-2, gp100, and MAG-E-1 are expressed in human glioblastoma and recognized by cytotoxic T cells. *Cancer Res.* 64(14), 4980–4986 (2004).
- 198 Ammi R, de Waele J, Willemsen Y *et al.* Poly(I:C) as cancer vaccine adjuvant: knocking on the door of medical breakthroughs. *Pharmacol. Ther.* 146, 120–131 (2015).
- 199 Black PL, Hartmann D, Pennington R *et al.* Effect of tumor burden and route of administration on the immunotherapeutic properties of polyinosinic-polycytidylic acid stabilized with poly-L-lysine in carboxymethyl cellulose [Poly(I,C)-LC]. *Int. J. Immunopharmacol.* 14(8), 1341–1353 (1992).
- 200 Ursu R, Carpentier AF. Immunotherapeutic approach with oligodeoxynucleotides containing CpG motifs (CpG-ODN) in malignant glioma. *Adv. Exp. Med. Biol.* 746, 95–108 (2012).
- 201 AA Gaspari, Tying SK, Rosen T. Beyond a decade of 5% imiquimod topical therapy. *J. Drugs Dermatol.* 8(5), 467–474 (2009).
- 202 Xiong Z, Ohlfest JR. Topical imiquimod has therapeutic and immunomodulatory effects against intracranial tumors. *J. Immunother.* 34(3), 264–269 (2011).
- 203 Rosenfeld MR, Chamberlain MC, Grossman SA *et al.* A multi-institution Phase II study of poly-ICLC and radiotherapy with concurrent and adjuvant temozolomide in adults with newly diagnosed glioblastoma. *Neuro Oncol.* 12(10), 1071–1077 (2010).
- 204 Carpentier A, Metellus P, Ursu R *et al.* Intracerebral administration of CpG oligonucleotide for patients with recurrent glioblastoma: a Phase II study. *Neuro Oncol.* 12(4), 401–408 (2010).
- 205 Wolchok JD, Hoos A, O'Day S *et al.* Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clin. Cancer Res.* 15(23), 7412–7420 (2009).
- 206 Reardon DA, Okada H. Re-defining response and treatment effects for neuro-oncology immunotherapy clinical trials. *J. Neurooncol.* 123(3), 339–346 (2015).
- 207 Sims JS, Ung TH, Neira JA, Canoll P, Bruce JN. Biomarkers for glioma immunotherapy: the next generation. *J. Neurooncol.* 123(3), 359–372 (2015).