

Location of Initial Visual Field Defects in Glaucoma and Their Modes of Deterioration

Joon Mo Kim,^{1,2} Haksu Kyung,^{2,3} Seong Hee Shim,¹ Parham Azarbod,^{2,4} and Joseph Caprioli²

¹Department of Ophthalmology, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, Republic of Korea

²Stein Eye Institute, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, United States

³Department of Ophthalmology, National Medical Center, Seoul, Korea

⁴Moorfields Eye Hospital, London, United Kingdom

Correspondence: Joseph Caprioli, Department of Ophthalmology, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA 90025; caprioli@ucla.edu.

Submitted: May 19, 2015

Accepted: July 7, 2015

Citation: Kim JM, Kyung H, Shim SH, Azarbod P, Caprioli J. Location of initial visual field defects in glaucoma and their modes of deterioration. *Invest Ophthalmol Vis Sci*. 2015;56:7956-7962. DOI:10.1167/iov.15-17297

PURPOSE. To describe the location of initial visual field defects (VFD) in glaucoma, their modes of deterioration, and those factors associated with different modes of deterioration.

METHODS. Patients with POAG were categorized into four groups based on three consecutive initial VFD: (1) superior paracentral defects (PD), (2) inferior PD, (3) superior nasal defects (ND), and (4) inferior ND. According to the worsening of the VF, four further subgroups were identified: (1) superior central worsening (CW), (2) inferior CW, (3) superior peripheral or nasal worsening (NW), and (4) inferior NW. Systemic and ocular factors were analyzed for each of the subgroups to identify possible associations.

RESULTS. One hundred sixty-two eyes of 162 subjects were analyzed. Superior PD ($n = 40$) were more frequent in females and associated with disc hemorrhage (DH), and were less frequent in patients with systemic hypertension (HT). Inferior PD ($n = 35$) showed a significant association with cup shape measure and axial length. Superior ND ($n = 37$) were more highly associated with HT and diabetes. Inferior ND ($n = 50$) showed a lower incidence of DH. With binary logistic regression analysis, superior PD showed a significant association with both HT and DH. With respect to VF worsening, superior CW showed a significant association with HT and diabetes, whereas superior NW was associated with a high minimum IOP during follow-up, and inferior NW was associated with a high maximum IOP during follow-up.

CONCLUSIONS. The initial location and subsequent direction of worsening of VFD were associated with different systemic and ocular factors.

Keywords: glaucoma, initial visual field defect, visual field progression

Glaucoma is characterized by a chronic progressive optic neuropathy with corresponding and characteristic patterns of visual field (VF) loss. In the majority of patients, VF changes are initially localized and as the disease progresses, these focal areas become wider, deeper, and more numerous. However, there is a great deal of variability in the rate and mode of worsening between patients.¹ Failure to treat early to moderate VF changes could lead to a decline in the quality of a glaucoma patient's life, with various effects depending on the degree and location of the visual loss.^{2,3} Visual field defects (VFD) near fixation have a greater impact on the level of function compared with peripheral changes, even at an early stage of glaucoma.^{4,5} It is generally taught that initial glaucomatous VF abnormalities frequently occur in the periphery with relative preservation of the central field. In actuality, one finds a variety of patterns and in some patients the initial defect may be quite centrally located. There is very little published about the exact nature of the initial VFD and the manner in which it subsequently worsens.

Given that glaucoma is a chronic progressive disease, much effort has been dedicated to the study of the factors relating to visual decay and the relationship between structural and functional changes.⁶⁻⁹ The two key elements in the study of glaucomatous VF deterioration are the direction and the rate of

decay. If deterioration occurs in the central direction (toward fixation), one would expect a greater impact on the patient's function with increasing functional loss in cases of rapidly centrally deteriorating VFs, even in those with relatively small initial defects.

The current study was undertaken to investigate the location of initial glaucomatous VFD, and the possible ocular and systemic factors associated with (1) the location of the initial VFD, and (2) the direction of worsening relative to the initial defect.

MATERIALS AND METHODS

Patient and Visual Field Data

A retrospective case review of open-angle glaucoma (OAG) patients of the University of California at Los Angeles (Los Angeles, CA, USA) Glaucoma Division with 6 or more years follow-up, and who had at least eight VF examinations was conducted. Visual fields were performed with a Humphrey Visual Field Analyzer (Carl Zeiss Ophthalmic Systems, Inc., Dublin, CA, USA) with the 24-2 SITA standard test strategy. Visual fields with acceptable VF reliability scores (according to the Advanced Glaucoma Intervention Study [AGIS] protocol)




FIGURE 1. Initial visual field defect. Paracentral defects were ([A] superior) and ([C] inferior). Nasal defects were ([B] superior) and ([D] inferior).

were included.¹⁰ Within the cohort of subjects whose records were reviewed, only those who matched the additional following criteria were included in the study:

1. Those with single defects, confined to one hemifield only;
2. The record contained all of the following information: a detailed history and complete clinical examination details (slit-lamp biomicroscopy, Goldmann applanation tonometry, gonioscopy, dilated fundus examination), VF examination, optic disc stereophotography, central corneal thickness (CCT) measurements, axial length measurement (IOL master), and confocal scanning laser tomography (HRT, Heidelberg Retina Tomography, Heidelberg Engineering, Heidelberg, Germany); and
3. If both eyes of a patient were eligible, the eye with lower VF mean deviation (MD) was enrolled.

Eyes with systemic or ocular disease known to affect the VF or those with missing data were excluded. The institutional review board of the University of California at Los Angeles approved the present study. All research procedures followed the tenets set forth in the Declaration of Helsinki.

Definitions

Initial VF Defect. These were divided into two groups, paracentral (PD) and nasal (ND) defects based on probability map locations in at least three out of the initial four Humphrey VF print outs. These two groups were then divided into two subgroups (superior and inferior) based on the affected hemifield. This yielded four initial defect arms for the study: (1) superior PD, (2) inferior PD, (3) superior ND, and (4) inferior ND.

Paracentral defects were defined as three or more adjacent points with an abnormal probability score of P less than 2%, or at least one point with a P less than 1% within the two paracentral locations nearest to fixation. Nasal defects were defined as three or more adjacent points with an abnormal probability score of P less than 2%, or one or more points with




FIGURE 2. The direction of extension of VF defects from initial defects. The central progression (CW) groups were 7 and 8 (superior), and 9 and 10 (inferior). The peripheral/nasal progression (NW) groups were 1, 2, and 3 (superior) and 4 and 5, and 6 (inferior).

P less than 1% within the three nasal-most points of the Humphrey VF (Figs. 1, 2). Horizontally, PD was for points lying from 0° to 10° , and ND was for points beyond 20° .

The Direction of VF Deterioration. Deterioration was defined with modified Anderson's criteria^{11,12} as contiguous points on a Humphrey 24-2 SITA standard VF with P less than 2% on the total deviation plot, or the development of new scotoma (at least 1 point of $P < 1\%$ or 3 contiguous points of $P < 2\%$) in the same hemifield.

The direction of worsening was recorded as either central (CW), or peripheral/nasal (NW) depending on the initial location of the VFD (Fig. 2). Deterioration required confirmation in at least three consecutive VFs. Patients were divided into four groups, depending on the direction of the worsening of the defect: superior CW, (2) inferior CW, (3) superior NW, and (4) inferior NW. Cases were excluded from these groups if (1) worsening could not be detected, or (2) the direction of worsening could not be defined such as the occurrence of a new scotoma in the opposite hemifield, a change in the location of a scotoma with repeated VF testing, deepening of a scotoma without spatial expansion, or the development of multiple scotoma, such as from a branched retinal vein occlusion or occurring after a long period of loss to follow up.

Data Analysis

The relationships between both the initial VFD and the direction of subsequent decay with demographic, systemic, and ocular factors were evaluated. Demographic factors included age at the onset of initial VF defect, sex, and family history of glaucoma. Systemic factors included a history of diabetes mellitus (DM), systemic hypertension (HT), and migraine. Ocular factors included disc hemorrhage (DH), CCT, axial length, maximum IOP, minimum IOP, IOP range during follow up, MD, PSD, cup to disc ratio (CDR), retinal nerve fiber layer (RNFL) thickness, cup shape measure (CSM), and disc area. Intraocular pressure range was defined as the difference between maximum IOP and minimum IOP during follow-up. Depending on the sample size, a χ^2 test or a Fisher's

TABLE 1. Comparison Between Initial Paracentral Defect (PD) and Initial Nasal Defect (ND)

	PD	ND	P Value
Subjects	75	87	
Age, y	64 ± 11.54	66.2 ± 9.97	0.201
Sex, male/female	36/39	43/44	0.732
Family history (+)(M/F)	29(13/16)	41(17/24)	0.413
FU duration, y	12.45 ± 3.45	11.64 ± 3.43	0.134
Ocular factors			
Axial length, mm	24.53 ± 0.998	24.15 ± 1.3	0.128
CCT, μm	552.46 ± 36.48	549.64 ± 40.13	0.727
FU minimum IOP, mm Hg	10.1 ± 2.82	9.76 ± 2.28	0.46
FU maximum IOP, mm Hg	19.5 ± 5.05	19.6 ± 5.09	0.706
Disc area, mm	2.068 ± 0.511	2.037 ± 0.581	0.736
CDR	0.5 ± 0.19	0.47 ± 0.17	0.424
RNFL thickness, mm	0.173 ± 0.091	0.175 ± 0.171	0.916
CSM	-0.088 ± 0.093	-0.115 ± 0.089	0.081
Disc hemorrhage	29	18	0.014
PPA	56	69	0.534
Baseline MD, dB	-3.36 ± 2.91	-4.03 ± 4.06	0.062
Baseline PSD, dB	3.84 ± 2.75	3.95 ± 2.81	0.21
Systemic factors			
Hypertension	20	47	0.001
Migraine	9	8	
DM	2	8	

χ^2 test and one-way ANOVA with Bonferroni correction. FU, follow-up; PSD, pattern standard deviation.

exact test was performed to compare categorical variables among groups. Independent *t*-tests and one-way ANOVA (with the Bonferroni correction) were used to compare continuous factors among groups. In order to exclude confounding elements among the factors analyzed, we performed a binary logistic regression analysis for all possible associations. Statistical analyses were performed with SPSS version 20.0 software (SPSS, Inc., Chicago, IL, USA). *P* values of less than 0.05 were considered statistically significant.

RESULTS

Patient and Visual Field Data

One hundred sixty-two eyes of 162 subjects were included. An initial paracentral defect (PD) occurred in 75 and an initial nasal defect (ND) occurred in 88 (Table 1). The baseline characteristics of the subgroups are shown in Tables 2 and 3. The number of eyes in each of the subgroups was 40 in superior PD, 35 in inferior PD, 37 in superior ND, and 50 in inferior ND. The results were compared for each subgroup with all other groups combined and are shown in Table 4. The result of binary logistic regression analysis is shown in Table 5.

The Initial VFD

When analyzing differences in associations between initial PD and initial ND, patients with initial PDs showed more DHs and less HT than patients with ND (*P* = 0.014 and *P* = 0.001, χ^2 test, respectively, Table 1).

Paracentral Defect (PD) Group.

Superior PD subgroup. There was a female preponderance (*P* = 0.018, χ^2 test) in this group but there were no significant

associations with age, family history of glaucoma, DM, or SAS. Among the systemic factors considered, HT (*P* = 0.002, χ^2 test) was negatively associated with the PD superior subgroup compared with the others. There was a significant association with DH (*P* = 0.011, χ^2 test) but no significant association was found with the other ocular factors analyzed.

Inferior PD subgroup. This group had a greater CSM (steeper cup) and longer axial length than other groups (*P* = 0.002 and *P* = 0.038, independent *t*-test, respectively).

Nasal Defect (ND) Group.

Superior ND subgroup. There was a higher incidence of HT (*P* = 0.074, odds ratio [OR] 1.952, χ^2 test) and DM (*P* = 0.05, Fisher exact test) in this group and a more negative MD value (*P* = 0.011, independent *t*-test), higher PSD value (*P* = 0.042, independent *t*-test), lower CSM (less steep cup, *P* = 0.03, independent *t*-test), and shorter axial length (*P* = 0.002, independent *t*-test) compared with the other groups.

Inferior ND subgroup. In these eyes, there was a lower incidence of DH during the follow-up period (*P* = 0.005, χ^2 test).

When binary logistic regression analysis adjusted for sex, HT, and DH was carried out, HT and DH were still significant factors for superior PD (*P* = 0.005 and *P* = 0.048, respectively) (Table 5). No other factors were found to have significant associations with the location of the initial VFD.

The Direction of Worsening of VF Defect

As shown in Table 2, in the group with an initial superior paracentral defect (PD), 14 eyes worsened nasally (superior NW) and three worsened centrally (superior CW). In the group with the initial inferior paracentral defects (PD), two deteriorated centrally (inferior CW) and 14 deteriorated nasally (inferior NW). In the group with initial superior nasal defects (superior ND), 20 eyes deteriorated toward a central direction (superior CW) and six eyes deteriorated toward a nasal direction (superior NW). In the groups with initial inferior nasal defects (inferior ND), 20 eyes deteriorated centrally (inferior CW) and three deteriorated nasally (inferior NW).

Central Worsening Group.

Superior CW subgroup. This group had a significant association with HT (*P* = 0.039, OR 2.65, 95% CI; 1.032 ~ 6.803, χ^2 test) and DM (*P* = 0.048, OR 0.167, 95% CI; 0.029 ~ 0.973, Fisher exact test), and there was a higher minimum IOP (10.88 mm Hg ± 2.33) during the follow up period compared with the other groups (9.44 mm Hg ± 2.07, *P* = 0.017, independent *t*-test).

Inferior CW subgroup. There was a higher maximum IOP (21.55 mm Hg ± 6.25) during the follow-up period compared with the other groups (18.53 mm Hg ± 4.45, *P* = 0.019, independent *t*-test).

Peripheral/Nasal Worsening Group. There were no factors that were significantly associated with either of the subgroups in these patients.

DISCUSSION

The current study was carried out to identify systemic and ocular factors associated with the location of the initial VF defect and its subsequent direction of worsening in glaucoma. The results suggest that factors associated with the location of the initial VF defect are different from those associated with the subsequent direction of worsening. With regard to the initial location of glaucoma damage, the superior paracentral group had a female preponderance, a higher incidence of disc hemorrhages, and was significantly less frequently associated with HT compared with other groups. The inferior paracentral

TABLE 2. Clinical Characteristics of the Subgroups of Initial PD and Initial ND.

	Superior PD	Superior ND	Inferior PD	Inferior ND	P Value
Subjects	40	37	35	50	
Age	64.0 ± 11.25	66.54 ± 9.55	65.29 ± 11.77	65.84 ± 10.44	0.490
Sex, male/female	13/27	15/22	22/13	29/21	0.021
Family history (+)(M/F)	13(6/7)	18(9/9)	16(7/9)	23(10/13)	0.16
FU duration, y	12.21 ± 3.52	10.88 ± 3.54	12.74 ± 3.39	12.18 ± 3.30	0.119
Ocular factors					
Axial length, mm	24.26 ± 0.91	23.68 ± 1.33*	24.80 ± 1.03*	24.55 ± 1.15*	0.009
CCT, μm	548 ± 34.3	574.2 ± 28.66	548.47 ± 39.46	546.66 ± 46.36	0.728
FU minimum IOP, mm Hg	10.07 ± 2.87	9.60 ± 2.41	9.97 ± 2.81	9.90 ± 2.53	0.884
FU maximum IOP, mm Hg	19.31 ± 5.18	19.41 ± 5.05	19.96 ± 4.67	20.1 ± 5.16)	0.885
Disc area	2.045 ± 0.44	2.27 ± 0.572	1.963 ± 0.637	1.99 ± 0.608	0.816
CDR	0.47 ± 0.174	0.48 ± 0.167	0.54 ± 0.206	0.57 ± 0.166	0.330
RNFL thickness, mm	0.18 ± 0.08	0.15 ± 0.071	0.16 ± 0.10	0.19 ± 0.213	0.658
CSM	-0.113 ± 0.086	-0.137 ± 0.09*	-0.057 ± 0.09*	-0.103 ± 0.091	0.007
Disc hemorrhage	18	10	12	7	0.013
PPA	32	27	25	41	0.917
Baseline MD, dB	-3.44 ± 2.99	-4.19 ± 3.97	-3.32 ± 3.23	-3.84 ± 3.61	0.342
Baseline PSD, dB	4.36 ± 2.81	4.67 ± 3.17	3.46 ± 2.71	3.84 ± 2.21	0.201
Systemic factors					
Hypertension	32	17	22	24	0.006
Migraine	5	3	4	5	
DM	2	5	0	3	
Worsening direction					
Superior CW	3	20			
Inferior CW			2	20	
Superior NW	14	6			
Inferior NW			14	3	

χ^2 test and one-way ANOVA with Bonferroni correction.

* Highlight statistical significant differences following Bonferroni correction.

group was significantly associated with CSM and axial length. Thus, initial superior paracentral defects were generally associated with systemic factors, while initial inferior paracentral defects were generally associated with ocular factors.

With respect to the direction of worsening, the superior paracentral group exhibited a stronger association with hypertension (HT) compared with the other groups. Central worsening showed a significant association with IOP during the follow-up period; however, peripheral worsening did not because most centrally worsening patients had an initial peripheral VFD. A portion of this result simply reflects the natural course of the disease. Werner and Berashow reported that early VFD often begin with a nasal step.¹³ Another study suggested that nasal steps appear to be a common defect in acute and early chronic angle-closure glaucoma.^{14,15} Park et al.¹⁶ previously reported that the development of an initial paracentral scotoma showed a stronger association with IOP-independent risk factors compared with initial peripheral scotomas, and that high IOP may be a risk factor for peripheral VFD.

Previous studies have suggested an association between systemic HT and glaucoma.¹⁷⁻²³ In this study, while two of the subgroups (superior ND and superior CW) had a positive association with HT, the superior PD group showed a negative association. The relationship between systemic HT and glaucoma is not entirely clear. By increasing the ocular perfusion pressure, systemic HT may in fact be a protective factor in glaucoma by reducing the effects of low ocular perfusion pressure, which is a known risk factor for glaucoma.^{21-22,24-26} It has previously been shown that a 10-mm Hg increase in systolic pressure is associated with 0.24- to 0.3-mm Hg increase in IOP, and a 10-mm Hg increase in

diastolic pressure is associated with 0.4- to 0.6-mm Hg increase in IOP.^{25,27} This occurs due to increasing aqueous humor formation and an increasing choroidal volume.^{28,29} Hypertension may induce obliteration of small vessels that can harm the perfusion of the optic nerve.³⁰ Small vessel changes with longstanding HT can be a risk. Leung et al.³⁰ suggested that retinal arterioles were narrowed in HT patients due to microvascular damage from increased blood pressure. However, systemic HT was determined based solely on history; we cannot analyze the effect of treatment. Patients with a history of systemic HT present the possibility of low BP with treatment. Episodes of hypotension could affect the location of initial VFD.³¹

Disc hemorrhage is an important risk factor for glaucoma and a sign of active disease.³²⁻³⁶ Jeoung et al.³⁷ have shown that the RNFL thickness is decreased by the time a DH is detected. Park et al.¹⁶ have suggested that DH is closely associated with the presence of a central visual field defect. In our study we found that the superior PD group was associated with DH. The inferior PD showed significant associations with axial length and CSM. High axial myopia is a risk factor of glaucoma development and worsening.^{38,39} Structural changes associated with myopia, such as a longer axial length, a larger or tilted optic disc, and peripapillary atrophy (PPA) may make the maculopapillary bundle more susceptible to glaucomatous injury.³⁸ Mayama et al.³⁹ suggested that higher myopia threatens the remaining lower cecocentral VF, and that it may also negatively affect the quality of vision in eyes with advanced OAG.

Cup shape measure (as measured by the HRT) is a reflection of the overall 3-dimensional shape of the optic nerve head. Steeper cupping may be due to the effect of thin RNFL

TABLE 3. Clinical Characteristics of the Study Population of Progression Direction

	Superior NW	Superior CW	Inferior NW	Inferior CW	P Value
Subjects	17	17	23	26	
Age	62.23 ± 10.41	69.53 ± 7.50	62.70 ± 13.45	64.60 ± 11.12	0.242
Sex, male/female	5/12	9/8	15/8	15/11	0.144
Family history(+)(M/F)	6(2/4)	6(3/3)	12(7/9)	15(10/13)	0.418
Ocular factors					
Axial length	24.07 ± 0.87	23.55 ± 1.34*	25.05 ± 1.13*	24.39 ± 1.01	0.008
CCT	559.3 ± 39.4	552.9 ± 30.9	562.9 ± 39.10	544.5 ± 40.0	0.599
FU minimum IOP, mm Hg	11.00 ± 2.28	9.15 ± 2.75	9.86 ± 2.27	10.24 ± 2.05	0.143
FU maximum IOP, mm Hg	21.56 ± 5.87	18.47 ± 5.74	20.27 ± 5.33	21.46 ± 5.22	0.301
Disc area, mm	1.97 ± 0.60	2.13 ± 0.51	2.07 ± 0.57	2.03 ± 0.74	0.912
CDR	0.42 ± 0.22	0.50 ± 0.15	0.46 ± 0.18	0.47 ± 0.19	0.754
RNFL thickness, mm	0.19 ± 0.97	0.17 ± 0.19	0.19 ± 0.09	0.21 ± 0.28	0.914
CSM	-0.127 ± 0.094*	-0.134 ± 0.093*	-0.092 ± 0.098	-0.056 ± 0.074*	0.037
Disc hemorrhage	6	4	7	5	0.731
PPA	12	13	15	21	0.803
Baseline MD, dB	-2.79 ± 2.4	-6.15 ± 6.1	-3.82 ± 3.2	-3.86 ± 2.94	0.072
Baseline PSD, dB	3.87 ± 2.32	4.48 ± 2.55	4.42 ± 3.06	4.10 ± 2.49	0.69
Systemic factors					
Hypertension	13	6	16	13	0.048
Migraine	2	0	4	2	
DM	0	3	0	2	

χ² test and one-way ANOVA with Bonferroni correction.

* Highlight statistical significant differences following Bonferroni correction.

thickness or the reflection of the natural variation of optic disc structure.⁴⁰ In our results, the inferior PD subgroup had a greater CSM (steeper cup) and superior ND subgroup had a lower CSM (less steep cup). The amount of VFD of both groups was comparable. Inferior PD showed longer axial length and superior ND showed shorter axial length than others. With elongation of axial length, RNFL thickness decreases⁴¹; this may be one of the factors which affects our results.

With respect to worsening, inferior NW was associated with a significantly higher maximum IOP during follow up than other worsening groups. Considering that inferior PD were associated with CSM and longer axial length and worsening of this group (inferior CW) showed higher maximum IOP during follow-up, inferior PD may be more affected by ocular factors rather than systemic factors.

TABLE 4. Comparing Each Subgroup in Turn With all the Other Groups Combined Together

	Positive Association	Negative Association
Superior PD	Female (<i>P</i> = 0.011), DH (<i>P</i> = 0.018)	HT (<i>P</i> = 0.002)
Inferior PD	csm (<i>P</i> = 0.002), axial length (<i>P</i> = 0.038)	None
Superior ND	HT (<i>P</i> = 0.074), DM (<i>P</i> = 0.05)	CSM (<i>P</i> = 0.03), axial length (<i>P</i> = 0.002)
Inferior ND	None	None
Superior CW	HT (<i>P</i> = 0.039), DM (<i>P</i> = 0.048), higher minimum IOP (<i>P</i> = 0.017)	None
Inferior CW	Higher maximum IOP (<i>P</i> = 0.019)	None
Superior NC	None	None
Inferior NC	None	None

χ² test or Fisher exact test and independent *t*-test.

According to a recently presented study that analyzed initial VFD as either central and peripheral, the central cases showed a higher association with systemic risk factors and with DH than the peripheral ones.¹⁶ The differences between those results and those of this study may largely be due to the study design. Their subjects mainly included high IOP glaucoma patients (including chronic angle closure glaucoma); however, our study was in OAG patients (including normal-tension patients), and those with chronic angle closure glaucoma were excluded. Furthermore, our patients were subdivided into four groups rather than two. Considering that IOP-independent factors may preferentially affect the neuroretinal rim or RNFL closer to the papillomacular bundle in the inferior half of optic nerve head or retina compared with high IOP,^{16,42,43} the negative association between HT and the superior PD group is interesting. The glaucoma patients with HT had superior nasal defects and that tended to progress centrally. Generally, most of initial VF defects, particularly initial parafoveal scotomas, developed in the superior hemifield.^{16,44,45} Systemic hypotension has been reported as a risk factor for central scotomas.^{16,26,46} In the early stages of systemic HT, blood flow to the eye may be enhanced. With longstanding systemic HT, smaller arterioles may be affected earlier than larger arterioles. We hypothesize that although there may be a high ocular perfusion pressure of the large arteries, decreased perfusion in the smaller, diseased vessels may produce peripheral VFD earlier than central VFD. As the effects of systemic HT or its treatment evolve, even central retinal ganglion cells may become vulnerable to decreased perfusion, and the VFD progress centrally. According to our findings, the central VF in patients

TABLE 5. Binary Logistic Regression Analysis for Superior PD

	OR	95% CI	P Value
Sex, male	0.535	0.236 ~ 1.211	0.134
Disc hemorrhage	0.443	0.199 ~ 0.984	0.045
Hypertension	3.761	1.498 ~ 9.443	0.005

with systemic HT remains relatively preserved during the initial stage of the disease as compared with those without systemic HT.

There are some limitations of this study. The data were collected retrospectively and information such as the initial history and the systemic risk factors were dependent on patient reporting and medical record documentation during clinic visits. All study patients were recruited from a tertiary care glaucoma clinic, which could create a sampling bias. Some patients were excluded from analysis because of missing data, such as axial length, CCT, and records at the initial visit. The effect of this selection of patients on our results is not known. A confounding factor in the design of the study is that the central most locations of the VF may preferentially deteriorate toward the periphery and the peripheral most points may do so centrally, and this could have introduced a bias into the results.

In our study, patients with an initial paracentral VFD had different characteristics than those with other patterns of defects. In particular, systemic factors (more migraine and less HT) were closely associated with superior paracentral defects. However, ocular factors (steeper cup and longer axial length) were more closely associated with inferior paracentral defects. Glaucoma patients with normal blood pressure appear to show a greater loss of superior central VF as compared with their hypertensive counterparts. These findings may help clinicians identify glaucoma phenotypes that may help anticipate the course of worsening patients and guide appropriate treatment.

Acknowledgments

Supported by grants from Pfizer, Inc. (New York, NY, USA), Allergan, Inc. (Irvine, CA, USA), Alcon, Inc. (Fort Worth, TX, USA), an unrestricted grant from Research to Prevent Blindness, Inc. (New York, NY, USA), and the National Eye Institute/National Institutes of Health (Bethesda, MD, USA) Grant R01-EY-018644 (all JC).

Disclosure: **J.M. Kim**, None; **H. Kyung**, None; **S.H. Shim**, None; **P. Azarhod**, None; **J. Caprioli**, Pfizer, Inc. (F), Allergan, Inc. (F, C), Alcon, Inc. (F)

References

- Caprioli J, Spaeth GL. Comparison of visual field defects in the low-tension glaucomas with those in the high-tension glaucomas. *Am J Ophthalmol*. 1984;97:730-737.
- McKean-Cowdin R, Wang Y, Wu J, Azen SP, Varma R; for the Los Angeles Latino Eye Study Group. Impact of visual field loss on health-related quality of life in glaucoma: the Los Angeles Latino Eye Study. *Ophthalmology*. 2008;115:941-948.
- Yanagisawa M, Kato S, Kobayashi M, Watanabe M, Ochiai M. Relationship between vision-related quality of life and different types of existing visual fields in Japanese patients. *Int Ophthalmol*. 2012;32:523-529.
- Fujita K, Yasuda N, Oda K, Yuzawa M. Reading performance in patients with central visual field disturbance due to glaucoma [in Japanese]. *Nippon Ganka Gakkai Zasshi*. 2006;110:914-918.
- Coexkelbergh TR, Brouwer WH, Cornelissen FW, et al. The effect of visual field defects on driving performance: a driving simulator study. *Arch Ophthalmol*. 2002;120:1509-1516.
- Drance S, Anderson DR, Schulzer M. Collaborative normal-tension glaucoma study group. Risk factors for progression of visual field abnormalities in normal tension glaucoma. *Am J Ophthalmol*. 2001;131:699-708.
- Lee EJ, Kim T-W, Weinreb RN, Park KH, Kim SH, Kim DM. Trend-based analysis of retinal nerve fiber layer thickness measured by optical coherence tomography in eyes with localized nerve fiber layer defects. *Invest Ophthalmol Vis Sci*. 2011;52:1138-1144.
- De Moraes CGV, Juthani VJ, Liebmann JM, et al. Risk factors for visual field progression in treated glaucoma. *Arch Ophthalmol*. 2011;129:562-568.
- Caprioli J, Mock D, Bitrian E, et al. A method to measure and predict rates of regional visual field decay in glaucoma. *Invest Ophthalmol Vis Sci*. 2011;52:4765-4773.
- Thissen D, Steinberg L, Kaung D. Quick and easy implementation of the Benjamini-Hockberg procedure for controlling the false positive rate in multiple comparison. *J Educ Behav Stat*. 2002;27:77-83.
- Anderson DR. *Automated Static Perimetry 1992*. St. Louis: CV Mosby; 1992.
- Thomas R, Paul P, Muliyl J. Use of pattern standard deviation instead of corrected pattern standard deviation in Anderson's criteria. *J Glaucoma*. 2000;9:480-482.
- Werner EB, Berashow J. Peripheral nasal field defect in glaucoma. *Ophthalmology*. 1979;86:1875-1878.
- Lau LI, Liu CJ, Chou JC, et al. Patterns of visual field defects in chronic angle-closure glaucoma with different disease severity. *Ophthalmology*. 2003;110:1890-1894.
- Bonomi L, Marraffa M, Marchini G, et al. Perimetric defects after a single acute angle-closure glaucoma attack. *Graefes Arch Clin Exp Ophthalmol*. 1999;237:908-914.
- Park SC, De Moraes CG, Teng CCW, Tello C, Liebmann JM, Ritch R. Initial parafoveal versus peripheral scotomas in glaucoma: risk factors and visual field characteristics. *Ophthalmology*. 2011;118:1782-1789.
- Sun J, Zhou X, Kang Y, et al. Prevalence and risk factors for primary open-angle glaucoma in a rural northeast China population: a population-based survey in Bin County, Harbin. *Eye*. 2012;26:283-291.
- Hulsman CA, Vingerling JR, Hofman A, Witteman JC, de Jong PT. Blood pressure, arterial stiffness, and open-angle glaucoma: the Rotterdam study. *Arch Ophthalmol*. 2007;125:805-812.
- Mitchell P, Lee AJ, Rohtchina E, Wang JJ. Open-angle glaucoma and systemic hypertension: the Blue Mountains Eye Study. *J Glaucoma*. 2004;13:319-326.
- Memarzadeh F, Ying-Lai M, Chung J, Azen SP, Varma R. Blood pressure, perfusion pressure, and open-angle glaucoma: the Los Angeles Latino Eye Study. *Invest Ophthalmol Vis Sci*. 2010;51:2872-2877.
- Leske MC, Heijl A, Hyman L, Bengtsson B, Dong L, Yang Z. Predictors of long-term progression in the early manifest glaucoma trial. *Ophthalmology*. 2007;114:1965-1972.
- Leske MC, Wu SY, Hennis A, Honkanen R, Nemesure B. Risk factors for incident open-angle glaucoma: the Barbados Eye Studies. *Ophthalmology*. 2008;115:85-93.
- Zhao D, Cho J, Kim MH, Guallar E. The association of blood pressure and primary open-angle glaucoma: a meta-analysis. *Am J Ophthalmol*. 2014;158:615-627.
- Leske MC, Connell AM, Wu SY, et al. Risk factors for open-angle glaucoma. The Barbados Eye Study. *Arch Ophthalmol*. 1995;113:918-924.
- Bonomi L, Marchini G, Marraffa M, et al. Vascular risk factors for primary open angle glaucoma: the Egna-Neumarkt Study. *Ophthalmology*. 2000;107:1287-1293.
- Sung KR, Cho JW, Lee S, et al. Characteristics of visual field progression in medically treated normal-tension glaucoma patients with unstable ocular perfusion pressure. *Invest Ophthalmol Vis Sci*. 2011;52:737-743.
- Klein BE, Klein R, Knudtson MD. Intraocular pressure and systemic blood pressure: longitudinal perspective: the Beaver Dam Eye Study. *Br J Ophthalmol*. 2005;89:284-287.

28. Xu L, Wang H, Wang Y, et al. Intraocular pressure correlated with arterial blood pressure: the Beijing eye study. *Am J Ophthalmol*. 2007;144:461-462.
29. Hvidberg A, Kessing SV, Fernandes A. Effect of changes in CWO2 and body positions on intraocular pressure during general anaesthesia. *Acta Ophthalmol (Copenh)*. 1981;59:465-475.
30. Leung H, Wang JJ, Rochtchina E, et al. Relationships between age, blood pressure, and retinal vessel diameters in older population. *Invest Ophthalmol Vis Sci*. 2003;44:2900-2904.
31. Kang JW, Park B, Cho BJ. Comparison of risk factors for initial central scotoma versus initial peripheral scotoma in normal-tension glaucoma. *Korean J Ophthalmol*. 2015;29:102-108.
32. Leske MC, Heijl A, Hyman L, et al; for the EMGT Group. Predictors of long-term progression in the Early Manifest Glaucoma Trial. *Ophthalmology*. 2007;114:1965-1972.
33. Miglior S, Torri V, Zeyen T, et al; for the European Glaucoma Prevention Study (EGPS) Group. Intercurrent factors associated with the development of open-angle glaucoma in the European Glaucoma Prevention Study. *Am J Ophthalmol*. 2007;144:266-275.
34. Suh MH, Park KH, Kim HJ, et al. Glaucoma progression after the first-detected optic disc hemorrhage by optical coherence tomography. *J Glaucoma*. 2012;21:358-366.
35. Siegnier SW, Netland PA. Optic disc hemorrhages and progression of glaucoma. *Ophthalmology*. 1996;103:1014-1024.
36. Ishida K, Yamamoto T, Sugiyama K et al. Disc hemorrhage is a significantly negative, prognostic factor in normal tension glaucoma. *Am J Ophthalmology*. 2000;129:707-714.
37. Jeung JW, Park KH, Kim DM, et al. Optic disc hemorrhage may be associated with retinal nerve fiber loss in otherwise normal eyes. *Ophthalmology*. 2008;115:2132-2140.
38. Quigley HA, Addicks EM, Green WR, Maumenee AE. Optic nerve damage in human glaucoma. II. The site of injury and susceptibility to damage. *Arch Ophthalmol*. 1981;99:635-649.
39. Mayama C, Suzuki Y, Araie M, et al. Myopia and advanced-stage open-angle glaucoma. *Ophthalmology*. 2002;109:2072-2077.
40. Kim JM, Jeoung JW, Bitrian E, et al. Comparison of clinical characteristics between Korean and western normal-tension glaucoma patients. *Am J Ophthalmol*. 2013;155:852-857.
41. Park SH, Park KH, Kim JM, Choi CY. Relation between axial length and ocular parameters. *Ophthalmologica*. 2010;224:188-193.
42. Ferreras A, Pablo LE, Garway-Heath DF, Fogagnolo P, Garcia-Feijoo J. Mapping standard automated perimetry to the peripheral retinal nerve fiber layer in glaucoma. *Invest Ophthalmol Vis Sci*. 2008;49:3018-3025.
43. Horn FK, Mardin CY, Laemmer R, et al. Correlation between local glaucomatous visual field defects and loss of nerve fiber layer thickness measured with polarimetry and spectral domain OCT. *Invest Ophthalmol Vis Sci*. 2009;50:1971-1977.
44. Jung KI, Park HY, Park CK. Characteristics of optic disc morphology in glaucoma patients with parafoveal scotoma compared to peripheral scotoma. *Invest Ophthalmol Vis Sci*. 2012;20:53:4813-4820.
45. Hood DC, Raza AS, de Moraes CG, et al. Initial arcuate defects within the central 10 degrees in glaucoma. *Invest Ophthalmol Vis Sci*. 2011;16:52:940-946.
46. Lee M, Cho EH, Lew HM, Ahn J. Relationship between ocular pulse amplitude and glaucomatous central visual field defect in normal-tension glaucoma. *J Glaucoma*. 2012;21:596-600.