

Expression Divergence of Chemosensory Genes between *Drosophila sechellia* and Its Sibling Species and Its Implications for Host Shift

Meng-Shin Shiao^{1,2,†}, Jia-Ming Chang^{3,4,5,†}, Wen-Lang Fan^{1,6}, Mei-Yeh Jade Lu¹, Cedric Notredame^{3,4}, Shu Fang¹, Rumi Kondo^{7,*}, and Wen-Hsiung Li^{1,8,*}

¹Biodiversity Research Center, Academia Sinica, Taipei, Taiwan

²Research Center, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

³Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Dr. Aiguader 88, 08003, Barcelona, Spain

⁴Universitat Pompeu Fabra (UPF), Barcelona, Spain

⁵Institute of Human Genetics (IGH), UPR 1142, CNRS, Montpellier, France

⁶Whole-Genome Research Core Laboratory of Human Diseases, Chang Gung Memorial Hospital, Keelung, Taiwan

⁷Graduate School of Humanities and Sciences, Ochanomizu University, Tokyo, Japan

⁸Department of Ecology and Evolution, University of Chicago

*Corresponding author: E-mail: kondo.rumi@ocha.ac.jp; whli@sinica.edu.tw.

†These authors contributed equally to this work.

Accepted: September 8, 2015

Data deposition: The data have been deposited at NCBI GEO under the accession numbers GSE67587, GSE67861, and GSE67862.

Abstract

Drosophila sechellia relies exclusively on the fruits of *Morinda citrifolia*, which are toxic to most insects, including its sibling species *Drosophila melanogaster* and *Drosophila simulans*. Although several odorant binding protein (*Obp*) genes and olfactory receptor (*Or*) genes have been suggested to be associated with the *D. sechellia* host shift, a broad view of how chemosensory genes have contributed to this shift is still lacking. We therefore studied the transcriptomes of antennae, the main organ responsible for detecting food resource and oviposition, of *D. sechellia* and its two sibling species. We wanted to know whether gene expression, particularly chemosensory genes, has diverged between *D. sechellia* and its two sibling species. Using a very stringent definition of differential gene expression, we found a higher percentage of chemosensory genes differentially expressed in the *D. sechellia* lineage (7.8%) than in the *D. simulans* lineage (5.4%); for upregulated chemosensory genes, the percentages were 8.8% in *D. sechellia* and 5.2% in *D. simulans*. Interestingly, *Obp50a* exhibited the highest upregulation, an approximately 100-fold increase, and *Or85c*—previously reported to be a larva-specific gene—showed approximately 20-fold upregulation in *D. sechellia*. Furthermore, *Ir84a* (*ionotropic receptor 84a*), which has been proposed to be associated with male courtship behavior, was significantly upregulated in *D. sechellia*. We also found expression divergence in most of the chemosensory gene families between *D. sechellia* and the two sibling species. Our observations suggest that the host shift of *D. sechellia* was associated with the enrichment of differentially expressed, particularly upregulated, chemosensory genes.

Key words: host shift, *Drosophila sechellia*, chemosensory genes, RNA-seq, antennal transcriptome.

Introduction

Most *Drosophila* species utilize a wide variety of food sources (Rio et al. 1983; Louis and David 1986), but *Drosophila sechellia*, which is endemic to Seychelles (Tsacas and Bachli 1981), exclusively uses the fruit of *Morinda citrifolia*, commonly

known as Tahitian Noni fruit, as its food source and for oviposition. Interestingly, the fruit of *M. citrifolia* is toxic to other insects, including most of the *Drosophila* species (Rkha et al. 1991). How *D. sechellia* evolved the ability to sense its specific host plant is not well understood.

© The Author(s) 2015. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Chemosensory genes are responsible for sensing odors and thus are essential for survival (finding food sources) and reproduction (finding oviposition) of most animal species. In hexapods (insects), the chemosensory gene superfamily comprises the odorant binding protein (*Obp*) genes, the chemosensory protein (*Csp* and *CheA/B*) genes, and three transmembrane receptor gene families: the olfactory receptor (*Or*) genes, the gustatory receptor (*Gr*) genes, and the ionotropic receptor (*Ir*) genes (Vosshall et al. 1999; Clyne et al. 1999, 2000; Galindo and Smith 2001; Xu et al. 2002; Benton et al. 2009; Vieira and Rozas 2011).

OBPs transport hydrophobic odorant molecules across the aqueous lymph surrounding the olfactory sensory neurons (OSNs) on the sensilla of fly antennae and lymph of other chemosensory sensilla. To be noted, CSPs and CHEA/B shared the same protein description, that is, chemosensory proteins, in the previous studies (Xu et al. 2002; Starostina et al. 2009; Vieira and Rozas 2011). Vieira and Rozas (2011) identified four CSPs and proposed that CSPs are homologous to OBPs. CHEA/B were found in the front legs of mature male flies, and thought to interact directly with lipid-like pheromones; they therefore are thought to be involved in male-specific mating behaviors (Xu et al. 2002; Starostina et al. 2009).

ORs and GRs were responsible for “smelling” and “tasting,” respectively, whereas IRs, which were identified on the basis of their structural similarity with ionotropic glutamate receptors (*iGluRs*), were recently found to be responsible for salt detection in *Drosophila* (Zhang et al. 2013). *Or* genes and *Ir* genes are expressed mostly in OSNs on different types of sensilla in antennae, and *Gr* genes are expressed in other body parts including proboscis, wing margins, legs, and ovipositors. As *Ir* genes show sequence homology to *iGluRs*, they are evolutionarily unrelated to *Or* and *Gr* genes (Benton et al. 2009). *Ir* genes are expressed in OSNs, different from those expressing *Or* or *Gr* genes, and are generally thought to mediate responses to acids and amines.

Previous studies suggested that the host shift of *D. sechellia* was associated with changes of several chemosensory genes either at the DNA sequence level or at the messenger RNA expression level. Examining interspecies hybrids between *D. sechellia* and *Drosophila melanogaster* deficiency strains, Matsuo et al. (2007) suggested that *Obp57d* and *Obp57e* (*Obp57d/e*) are responsible for the host preference in *D. sechellia*. They further showed that transferring *Obp57d/e* of *D. sechellia* to the *Obp57d/e* knockout *D. melanogaster* changed their preference to higher concentrations of hexanoic acid (HA) and octanoic acid (OA), the toxins contained in the ripe fruit of *M. citrifolia*. They proposed that the functional alteration in *D. sechellia* was due to a 4-bp insertion upstream of *Obp57e*. Dworkin and Jones (2009) showed that the *D. sechellia* food specialization was related to several cases of loss-of-function in genes involved in detoxification, metabolic pathways, and olfaction. They found that *D. sechellia* *Obp56e* harbors a premature stop codon, which might play

a role in *D. sechellia*'s preference of *M. citrifolia* because a knockdown of this gene in *D. melanogaster* reduced the avoidance of *M. citrifolia*. Hungate et al. (2013) identified an approximately 170-kb region related to the tolerance of OA by testing the *D. sechellia* introgression lines under the *Drosophila simulans* genetic background. This locus comprises 18 genes belonging to the *Obp* and *Osiri* (*Osi*) gene families.

The first large-scale microarray analysis of *D. sechellia* antennal transcriptome found that *Or22a*, *Or22b*, and *Or85b* have markedly increased expression in *D. sechellia* (Kopp et al. 2008). However, *Or22a* and *Or22b* also differed significantly in expression level between *D. simulans* and *D. melanogaster* and *Or22a* showed significant expression differences between two strains of *D. melanogaster* as well. Microarray analysis might not be suitable for studying lowly expressed genes, such as *Or* genes, which are lowly expressed because one OSN expresses only 1–3 *Or* genes, including the universally expressed olfactory receptor co-receptor gene (*Orco*).

Another study suggested a 5-fold acceleration of gene loss in the *D. sechellia* genome by comparing genomes of five closely related *Drosophila* species (McBride and Arguello 2007). However, the gene-loss phenomenon may only explain the process for a species not to avoid a plant host, but is insufficient to explain the adaptation to a new host as it may require a novel receptor or changes in the expression of certain existing genes. Interestingly, no novel chemosensory genes have been discovered in the *D. sechellia* genome so far. Furthermore, most studies focused on the well-defined chemosensory gene families, such as *Obp*, *Or*, and *Gr* genes. How the other chemosensory gene families are expressed in antennae and whether they are involved in the host shift of *D. sechellia* are unknown.

Dekker et al. (2006) proposed that the host shift in *D. sechellia* was related to the large differences in the numbers of sensilla in antennae. In comparison to *D. melanogaster*, *D. sechellia* has many more ab3 sensilla at the cost of ab1 and especially ab2 sensilla (Stensmyr et al. 2003; Dekker et al. 2006). The total number of ab3 sensilla on the *D. sechellia* antennae is about 2.5- to 3-fold higher than that of *D. melanogaster* (Dekker et al. 2006). Dekker et al. (2006) also suggested that the increased sensitivity to *M. citrifolia*, particularly to the chemical MeHex, was achieved by increasing the expression of *Or* genes in ab3 antennae, including *Or22a*, *Or22b*, and *Or85b*.

In this study, we hypothesized that the adaptation of *D. sechellia* to the new host was achieved by changing expression levels of existing genes, and we expected enrichment of differentially expressed chemosensory genes in the *D. sechellia* lineage. We would like to know whether the adaptation involved expression divergence in all chemosensory gene families, including recently identified *Ir*, *Csp*, and *Che* genes. We would also like to study whether the expression levels of chemosensory genes are associated with the differences in the number of sensilla in *D. sechellia*.

To answer these questions, we isolated total RNA from more than 300 pairs of antennae detached from the head and we used the Next Generation RNA-seq technology to identify differentially expressed genes (DEGs) between *D. sechellia* and its two sibling species: *D. melanogaster* and *D. simulans*. Our data revealed many highly differentially expressed chemosensory genes in *D. sechellia* that had not been identified in previous studies. We discussed the implications of these observations for the host preference of *D. sechellia*.

Materials and Methods

Sample Collection and RNA Isolation

Drosophila simulans clade split from *D. melanogaster* 3 Ma (Lachaise et al. 1988), and two island endemic species, *D. sechellia* and *D. mauritiana*, independently evolved from the *D. simulans*-like ancestor approximately 250,000 years ago (Kliman et al. 2000; McDermott and Kliman 2008; Garrigan et al. 2012). So, *D. simulans* is a better candidate than *D. mauritiana* as *D. simulans* clade reference for studying the host shift of *D. sechellia*. Therefore, to infer the evolution of chemosensory genes in the host specialization of *D. sechellia*, we chose the most closely related generalist species, *D. simulans*, as a comparison and *D. melanogaster* as the out-group generalist species.

We collected antennae from *D. melanogaster* (Canton-S, Bloomington *Drosophila* Stock Center at Indiana University: #1; collected from Canton, OH in 1920s) and *D. sechellia* (genome strain, UC San Diego *Drosophila* Stock Center: #14021-0248.25; collected from Cousin Island, Seychelles in 1980) at Academia Sinica, Taiwan and from *D. simulans* (genome strain, Kyorin-Fly: k-s05; originally from #14021-0251.194, collected from Wolfskill orchard, Winters, CA in 1995) and *D. sechellia* (Kyorin-Fly: k-s10; originally from #14021-0248.25) at Ochanomizu University, Japan. The flies in Taiwan and Japan were reared on standard cornmeal fly food under similar environmental conditions, a 12/12 h light/dark cycle at 25 °C.

About 300–700 pairs of fly antennae of each sex from each species were collected for total RNA isolation. The antennae resected each day were preserved in approximately 50–100 µl Trizol (Life Technology) and stored at –80 °C before further processing. Right before the RNA isolation, we spun down antennae preserved in Trizol at the max speed (13,000 rpm) for 3 min at 4 °C and transferred each sample into one MagNA Lyser Green Beads tube (Roche).

The RNA isolation protocol we developed combined steps from conventional Trizol extraction and RNeasy Micro Elute Kit (QIAGEN, Inc.) with some modifications to obtain high yield and quality of total RNA. Fly antennae were disrupted with MagNA Lyser (Roche) at 7,000 rpm for 15 s each time and repeated for 4–5 times until the tissue was almost invisible.

After homogenization, we cooled down the lysate on ice for 10 s to prevent RNA from degradation. Tissue lysate was transferred to a new RNase free Eppendorf tube. We used about 400 µl Trizol to rinse the beads of each sample and combine this 400 µl Trizol with the tissue lysate for the RNA isolation. Based on the Trizol standard protocol, 200 µl of chloroform per 1 ml of tissue lysate was added to the lysate and mixed well by shaking vigorously for 15 s and set 2–3 min at room temperature. The aqueous phase was separated by centrifuging the lysate at the maximum speed (13,000 rpm) at 4 °C for 15 min and carefully transferred to a new Eppendorf tube after centrifuging.

We added 1 volume of 5 µl Carrier RNA (QIAGEN, Inc.) and 1 ml of 70% EtOH to the aqueous phase and mixed well by inverting the tubes carefully. To obtain greater amount of RNA, we kept the samples at –80 °C for overnight to enhance RNA precipitation for greater yield. RNA isolation followed manufacturer's protocol of the RNeasy Micro Elute Kit with increased volumes (700 µl) of RPE buffer and 80% ethanol to wash RNA. Genomic DNA was removed by applying on-column DNase I treatment at room temperature for 15 min.

Paired-End mRNA-seq

For paired-end mRNA-seq library preparation, we used the TruSeq v2 kits from Illumina. Input of 4 µg of total RNA was used for mRNA enrichment by oligo-dT beads followed by cation-catalyzed fragmentation for 7.5 min at 94 °C. The mRNA fragments were then converted into double-stranded cDNA by random priming followed by end repair and A-tailing. The fragments were then ligated to the barcoded paired-end adaptors and subjected to ten cycles of polymerase chain reaction (PCR) amplification and purified by Ampure XP beads (Beckman Agencourt). The absolute concentrations of the libraries were determined by Qubit fluorometer (Invitrogen) and profiled by BioAnalyzer 2100 with High Sensitivity DNA Kit (Agilent). The six barcoded cDNA libraries were pooled together at equal molar ratio after quantitative PCR normalization (KAPA Library Quantification Kits) and loaded into three lanes of flow cell, and paired-end 2*100 nt multiplexed sequencing was conducted on Illumina HiSeq2000, yielding an average of 0.5 lanes of sequencing reads in total. The raw sequencing data reported in this work have been deposited in the NCBI GEO with accession numbers GSE67587, GSE67861 and GSE67862 for *D. sechellia* (Tuson, #14021-0248.25), *D. sechellia* (k-s10; #14021-0248.25) and *D. simulans* (k-s05; #14021-0251.194), respectively.

The raw reads data of *D. melanogaster* from Shiao et al. (2013) were included for analysis. Reference genomes version r5.48 (*D. melanogaster*), r1.3 (*D. simulans*), and r1.3 (*D. sechellia*) were used. For RNA-seq analysis, the 100-bp paired-end sequencing reads were mapped to the reference genome using *TopHat* 2.0.6 (Trapnell et al. 2009), with

allowance of two mismatches in read mapping to the reference genome. The expression levels of genes were measured in Fragments Per Kilobase of exon per Million fragments mapped (FPKM) by *Cufflinks* with bias correction (“–multi-read-correct” and “–frag-bias-correct”).

Generating Orthologous Gene Set and Identifying DEGs

Gene annotation tables of the three species were retrieved from FlyBase (version 2012_06) with individual versions 5.48, 1.3 and 1.3 for *D. melanogaster*, *D. simulans* and *D. sechellia*, respectively. The orthologous gene set (OGS) was manually curated by excluding genes duplicated in any of the species and only single copy genes in all three species were selected.

The potential bias from sequencing between different samples was normalized by upper quartile implemented in NOISeq (Tarazona et al. 2012). NOISeq requires technical replicates to perform the calculation. To identify DEGs, we used sequencing data from three lanes as technical repeats. As Dmel_TW had RNA-seq data from only a single sequencing lane for each sex, we generated technical replicates of Dmel_TW artificially in the software for comparison, using one function of NOISeq, NOISeq-sim.

Quantitative Reverse Transcription Polymerase Chain Reaction

We used NanoString quantitative reverse transcription polymerase chain reaction (qRT-PCR) technology to verify the gene expression levels estimated with RNA-seq. Hundred nanograms of total RNA from the same pool of total RNA for RNA-seq was used to determine the expression levels of 43 candidate genes simultaneously and three technical replicates of each gene were performed. Targeting sequences are shown in [supplementary table S1, Supplementary Material online](#). Five genes, *Actin 42A* (*Act42A*), *Cyclophilin-33* (*cyp33*), *mitochondrial ribosomal protein L20* (*mRpl20*), *Ribosomal protein L32* (*Rpl32*) and *Succinate dehydrogenase A* (*SdhA*), that were found to have relatively consistent normalized FPKM values across all the samples (no significant differential expression was detected) and were used as commercialized products by some companies (such as Qiagen) were selected as endogenous genes for qRT-PCR experiments. *Lush* (*Obp76a*) was used as a positive control.

Identification of Gene Sequences of *Or22a* and *Or22b*

Or22a and *Or22b* were not included in the OGS table because of the presence of duplicate gene IDs. In the gene annotation files we used for analyses (version FB2012_06), *Or22a* has two distinct gene IDs in *D. simulans* (FBgn0068650 and FBgn0194492) and in *D. sechellia* (FBgn0171736 and FBgn0259897), whereas *Or22b* has two FlyBase IDs in *D. simulans* (FBgn0068649 and FBgn0194493). This resulted in the removal of *Or22a* and *Or22b* from the OGS in our data

set. In the latest version of the gene annotation file (FB2015_02), FBgn0194492, FBgn0171736 and FBgn0194493 were assigned to other genes. We therefore manually curated the data set, and assigned FBgn0068650 to *Or22a* and FBgn0068649 to *Or22b* in *D. simulans*, and FBgn0259897 to *Or22a* in *D. sechellia*. Furthermore, we aligned DNA sequences of these two IDs to confirm that *Or22a* and *Or22b* of the three species form a monophyletic clade in the phylogenetic analysis (data not shown).

Results

Expression Profiles of Orthologous Genes in the Three Species

To obtain total RNA fully representing the transcriptomes of fly antennae, we detached antennae from heads under a microscope within one hour of eclosion. Antennal RNA of the three species (~300–700 pairs of antennae from each species) was collected independently in two laboratories. First, antennae of *D. sechellia* and *D. melanogaster* for both sexes were collected in Academia Sinica, Taiwan. Second, antennae of *D. sechellia* and *D. simulans* for both sexes were collected at Ochanomizu University, Japan.

The transcriptomes of *D. melanogaster* (~110 million paired-end reads) collected in Taiwan were published earlier in Shiao et al. (2013). To obtain similar sequencing amounts as in Shiao et al. (2013), we barcoded the six samples and pooled them to perform sequencing using three lanes on Illumina HiSeq2000. We obtained about 88–116 million paired-end sequencing reads for each ([supplementary table S2, Supplementary Material online](#)).

To compare the expression levels between species, we first generated an OGS without duplicate genes or gene loss in any of the three species, and obtained 10,034 annotated genes in OGS, including 51 *Obp* genes, 48 *Or* genes, 45 *Gr* genes, 34 *Ir* genes, 4 *Csp* genes, and 20 *CheA/B* genes ([supplementary tables S3 and S4, Supplementary Material online](#)). Eight chemosensory genes were excluded from OGS, including 5 *Or* genes (*Or22a*, *Or22b*, *Or46a*, *Or65c*, and *Or67c*), 1 *Gr* gene (*Gr59b*), and 2 *Ir* genes (*Ir40a* and *Ir93a*); most of these *Or* and *Gr* genes were proposed to be gene losses in *D. sechellia* (McBride and Arguello 2007). There were two copies of *Ir40a* and *Ir93a* in *D. sechellia* and *D. simulans* but only one copy in *D. melanogaster* based on gene annotation. We therefore removed these two *Ir* genes because of the uncertainty of orthology.

The expression levels of genes in OGS were then calculated in terms of FPKM. We used the three lanes as technical replicates, thus obtaining three FPKM values for each gene. The FPKM values of each gene from any pair of sequencing lanes were highly correlated (Pearson correlation $r > 0.99$), indicating no technical bias between different sequencing lanes (data not shown).

As observed in Shiao et al. (2013), the chemosensory genes that were expressed most abundantly in the antennae of *D. melanogaster* included the *Obp* gene family and a universal *Or* gene, *Orco* (also named *Or83b*). In total, 13 *Obp* genes, 2 *Or* genes, 1 *Ir* gene, and 1 *Gr* gene were consistently expressed at moderately high levels (FPKM > 50) in all eight samples, including *D. melanogaster* (supplementary table S4, Supplementary Material online).

There were ten genes consistently expressed at FPKM > 1,000 in all three species: *a10*, *PebIII*, *Obp19a*, *Obp19d*, *Obp28a*, *Obp56d*, *Obp69a*, *Obp83a*, *Obp83b*, and *lush*. The most highly expressed genes in antennae were all ligand binding protein genes. The gene *lush* is known to respond to 11-*cis*-vaccenyl acetate and acts as a sex and aggregation pheromone in *D. melanogaster* (Kim et al. 1998; Xu et al. 2005; Ha and Smith 2006). *Obp19a* and *Obp19d* belong to an *Obp* gene cluster that includes *Obp19a*, *Obp19b*, *Obp19c* and *Obp19d*, spanning an approximately 50-kb region on the *X* chromosome, according to the annotation of the *D. melanogaster* genome. The absence of *Obp19b* in our OGS was due to the missing annotation of *D. simulans* in the gene annotation file; it was expressed at moderate levels (FPKMs between 40.75 and 70.75) in *D. melanogaster* and *D. sechellia*. *Obp19c* was expressed at much lower levels in all three species, compared with *Obp19a* and *Obp19d*. A previous study found that in *Obp19d*, one nucleotide variation in the 5'-untranslated region and a synonymous substitution in the fourth exon were associated with variation in life span (Arya et al. 2010). *Obp28a* was proposed to be responsible for bitter taste in *D. melanogaster*, as suggested by the effect of *Obp28a* knockdown in *D. melanogaster* that resulted in an increased intake of bitter tartans, particularly quinine (Swarup et al. 2014). *Obp69a*, *Obp83a*, and *Obp83b* were proposed to be involved in "hearing" courtship song of males in female *D. melanogaster* (Immonen and Ritchie 2012).

Among the *Or* genes, *Orco* was expressed at the highest level in all samples, at FPKM > 1,000 in both *D. melanogaster* and *D. simulans* and > 500 in *D. sechellia*. The high expression level of *Orco* is expected because it is universally expressed in OSN cells (Stengl and Funk 2013), including in the trichoid and basiconic sensilla of antennae (Larsson et al. 2004). Other highly expressed *Or* genes include *Or42b*, *Or43b*, *Or47b*, *Or59b*, and *Or92a*.

For the other chemosensory gene families, only *Gr63a* and *Ir76b* were expressed at moderate levels in all species. These two genes have been found to detect carbon dioxide and low salt, respectively, in *D. melanogaster* (Jones et al. 2007; Zhang et al. 2013).

DEGs in *D. sechellia* Antennae

We looked for genes involved in dietary shift, assuming that these genes should show differential expression between

D. sechellia and the other two species, but not between male and female flies within a species because there is no sex difference in sensing the fruits of *M. citrifolia*. Under this hypothesis, genes related to the host shift in *D. sechellia* should show differential expression patterns in all of the following four comparisons: 1) Male *D. melanogaster* and male *D. sechellia*, 2) female *D. melanogaster* and female *D. sechellia*, 3) male *D. simulans* and male *D. sechellia*, and 4) female *D. simulans* and female *D. sechellia*.

The probability of significantly differential expression suggested by the authors of NOISeq software is $q \geq 0.8$. A gene with $q = 0.8$ means it is four times more likely differentially expressed than nondifferentially expressed. Using this criterion, we obtained approximately 5,500 nonredundant single copy DEGs either between *D. sechellia* and *D. melanogaster* or between *D. sechellia* and *D. simulans* (supplementary tables S5–S8, Supplementary Material online). Among these genes, 1,532 genes are differentially expressed in *D. sechellia* compared with both sibling species. They contain 41 chemosensory genes (out of 177 chemosensory genes in OGS), including 14 *Obp* genes, 19 *Or* genes (including *Orco*), 3 *Gr* genes, and 5 *Ir* genes (supplementary table S9, Supplementary Material online).

As *Orco* is a universally expressed gene, the significantly lower expression level of *Orco* in *D. sechellia* might have resulted from the normalization process. We therefore conducted qRT-PCR by the NanoString technology to verify the differential expression of *Orco* between species (see "Materials and Methods"). We did not find a significant fold change in the expression levels of *Orco* in any of the pairwise comparisons (supplementary table S10, Supplementary Material online): -0.46 (in log₂ scale) for *D. sechellia* versus *D. melanogaster* males, -0.76 for *D. sechellia* versus *D. melanogaster* females, 0.07 for *D. sechellia* versus *D. simulans* males, and -0.17 for *D. sechellia* versus *D. simulans* females. This observation suggested that the criterion of $q \geq 0.8$ might not be high enough to define the truly DEGs in *D. sechellia*. In addition to *Orco* gene, we conducted qRT-PCR for some randomly selected chemosensory genes with q values between 0.8 and 0.95 or lower than 0.8. The log₂-ratios of fold changes did not show consistent expression levels among species and most of the log₂-ratios were not significant (supplementary table S10, Supplementary Material online). Furthermore, if no technical replicates were available, the authors of NOISeq recommended to use a higher threshold such as $q = 0.9$. For these reasons, we decided to use $q = 0.95$ in our study.

As mentioned above, we chose $q \geq 0.95$ to define DEGs. In total, we identified 986 and 1,156 nonredundant single copy genes up- and downregulated, respectively, in *D. sechellia* antennae from the four independent comparisons: 1) 985 DEGs between males of *D. melanogaster* and *D. sechellia*, including 516 up- and 469 downregulated genes; 2) 1,142 DEGs between females of *D. melanogaster* and *D. sechellia*,

including 591 up- and 551 downregulated; 3) 1,254 DEGs between males of *D. simulans* and *D. sechellia*, including 570 and 684 genes up- and downregulated; and 4) 751 DEGs between females of *D. simulans* and *D. sechellia*, including 384 up- and 367 downregulated. There were 147 genes upregulated and 84 genes downregulated in *D. sechellia* compared with *D. melanogaster* and *D. simulans*, including 13 and 5 chemosensory genes, respectively (fig. 1, table 1, supplementary tables S11 and S12, Supplementary Material online).

The chemosensory genes with the greatest log₂-ratio are *CheA75a* and *CheA87a*. *CheA75a* was expressed at a very low level, almost no expression, in *D. simulans*, resulting in a very high log₂-ratio between *D. sechellia* and *D. simulans* (5.61 and 18.39 in males and females, respectively). More interestingly, *CheA87a* was expressed at a high level in *D. sechellia* but at a low level in the two sibling species. *CheA/B* was proposed to be involved in male-specific pheromone sense in the front legs. Our data suggest a possible association of the *CheA/B* gene family with the host-plant shift of *D. sechellia*.

For the other chemosensory genes, the log₂-ratios of FPKM values for *Obp50a* between *D. sechellia* and the two sibling species ranged from 3.74 to 7.04, indicating that the expression level of *Obp50a* was more than 100-fold higher in *D. sechellia* than in *D. melanogaster* (FPKM < 1) and *D. simulans* (FPKM = 1.79 and 1.65 in the two sexes, respectively). *Obp50a* belongs to a gene cluster that includes *Obp50a/b/c/d/e*, but only *Obp50a* was expressed at an intermediately high level in *D. sechellia*. Among the *Or* genes, *Or85c* showed the greatest differences in expression level between species: Greater than 20-fold higher in *D. sechellia* than in *D. melanogaster* and greater than 10-fold higher in *D. sechellia* than in *D. simulans*. *Or85c* was reported to be a larva-specific receptor in *D. melanogaster* (Couto et al. 2005; Kreher et al. 2005) and was proposed to be the receptor of 3-octanol (Mathew et al. 2013). More recent data indicated that it was expressed at a very low level in *D. melanogaster* antennae (Shiao et al. 2013). However, as there is no evidence that 3-octanol is one of the chemicals emitted by Noni fruit; the potential role of *Or85c* in sensing *M. citrifolia* by *D. sechellia* is unclear.

DEGs Validated by qRT-PCR

To verify DEGs, we conducted qRT-PCR using the NanoString technology. A total of 18 differentially expressed chemosensory genes were analyzed, along with 20 other genes to be discussed later in this section. The same total RNA used for RNA-seq was used for qRT-PCR. Targeting probes were designed in the consensus regions of the three species for all the genes selected. The *Act42A*, *Cyp33*, *mRpL20*, *RpL32*, and *SdhA* genes were selected as endogenous controls to normalize the expression levels. In addition, *lush* and *Orco* were included as positive controls. All the endogenous

genes selected for qRT-PCR had similar expression levels (FPKMs) in the three species.

For most of these genes, the qRT-PCR data corresponded very well to the RNA-seq data except for *Or9a*, *Or42b*, *Or67b*, and *CheA75a* (supplementary table S13, Supplementary Material online). The qRT-PCR data for *Or9a* showed that the expression level was not significantly different between *D. sechellia* and the two other species because the log₂-ratios ranged only from -0.13 to 0.08. On the other hand, the RNA-seq data suggested a much lower expression level for *Or9a* in *D. sechellia*. We checked the genomic location and found that a putative gene, FBgn0166364, overlapped with *Or9a* in the gene annotation table. We then calculated FPKMs by using a different version of annotation table (gene_orthologs_fb_2013_01.tsv.gz) that did not include FBgn166364 in the table (data not shown). The log₂-ratios of FPKM values then suggested no significant differences in expression levels between species, consistent with the qRT-PCR results. Thus, some of the incongruences between RNA-seq and qRT-PCR analysis might be due to annotation artifacts (e.g., overlapping genes) and associated mapping issues. However, the reason why the qRT-PCR and the RNA-seq data for *Or42b* and *Or67b* differed is unknown.

Based on gene sequence analysis, *CheA75a* in *D. simulans* has diverged greatly from those in the other two species (~68% divergence in peptide sequence between *D. simulans* and the two other species). We therefore could not find any probe-targeting consensus region in *D. simulans* *CheA75a* and expected very low or no amplification of *CheA75a* in *D. simulans*. Interestingly, we still detected expression of *CheA75a* in *D. simulans* by qRT-PCR and the expression level was almost the same as that in *D. melanogaster*. We performed a BLAST (Basic Local Alignment Search Tool) search using the 100-bp probe sequence of *CheA75a* against the *D. simulans* genome assembly and annotated genes in FlyBase, but no identical sequence segment longer than 20 bp was found. However, *CheA75a* had a sequence identity of 98% (gi190067942) and 99% (gi190008972) with two clones from a full-length cDNA in the cDNA library of *D. simulans* adult males in GenBank. Thus, the expression level of *CheA75a* in *D. simulans* remains unclear.

In addition to chemosensory genes, we also studied the expression levels of 18 genes with conspicuous differential expression patterns between *D. sechellia* and the other two species (*CG16799*, *CG4797*, and *Listericin*) with potentially interesting functions (supplementary table S13, Supplementary Material online), such as sense perception (*nompC*), olfactory learning (*vsg*), protein binding (*Nup50*), hormone catabolic process (*Jheh1*, *Jheh2*, *Jheh3*, and *to*), and insecticide metabolism (*Cyp308a1* and *GstE9*). For most of these genes, the qRT-PCR data corresponded very well to the RNA-seq data.

For *D. melanogaster*, we generated artificial technical replicates, as recommended by NOISeq (see Materials and




FIG. 1.—Venn diagrams showing genes expressed differentially in *D. sechellia* (Dsec) in comparing with *D. melanogaster* (Dmel) or *D. simulans* (Dsim). A total of 147 genes were significantly upregulated (A) and a total of 84 genes were significantly downregulated (B) in *D. sechellia*. Males and females are presented in blue and red, respectively. The 147 upregulated genes include 3 *Obp* genes, 6 *Or* genes, 1 *Gr* gene, 1 *Ir* gene, 2 *CheA/B* genes and 134 other genes in the transcriptome (A). The 84 downregulated genes include 3 *Obp* genes, 2 *Or* genes, and 79 other genes in the transcriptome (B).

Methods). This procedure ignores the potential stochastic variation in the library preparation or sequencing reaction. Our qRT-PCR data were in agreement with the RNA-seq data.

Enrichment of Differentially Expressed Chemosensory Genes in *D. sechellia*

Based on our analysis, there were a total of 18 chemosensory genes among the 231 genes that showed up- or downregulation in the *D. sechellia* lineage (7.8%). Among the genes in OGS, there are a total of 9,844 genes including 177 chemosensory genes expressed in antennae. Note that we set the threshold of $FPKM \geq 0.05$ to define that a gene is expressed in a species, following our previous papers (Shiao et al. 2012, 2013). All genes with $FPKM \geq 0.05$ in at least one species were included in the analysis. In other words, a gene with $FPKM < 0.05$ in all three species was considered as “not expressed” in the antennae. We observed enrichment of chemosensory genes among the DEGs (18 [chemosensory DEGs]/213 [nonchemosensory DEGs] versus 159 [non-DEG chemosensory genes]/9,454 [non-DEG nonchemosensory genes], $\chi^2 = 48.13$, $P < 0.00001$). For comparison, we also investigated the DEGs in the *D. simulans* lineage (supplementary fig. S1, Supplementary Material online). We found that in

D. simulans, there were 17 chemosensory genes among a total of 316 DEGs (5.4%), and compared with a total of 177 chemosensory genes among the 9,844 genes expressed in antennae, there was enrichment of chemosensory genes (17 [chemosensory DEGs]/299 [nonchemosensory DEGs] versus 160 [non-DEG chemosensory genes]/9,368 [non-DEG nonchemosensory genes], $\chi^2 = 23.72$, $P < 0.00001$). The analysis showed that chemosensory genes were enriched in the DEGs in the antennae of both *D. sechellia* and *D. simulans*, but there was a higher percentage of chemosensory genes differentially expressed in *D. sechellia*.

We further found that in *D. sechellia*, 13 and 5 chemosensory genes were up- or downregulated among a total of 147 upregulated and 84 downregulated genes, respectively, showing a higher percentage of upregulated chemosensory genes (8.8%) than downregulated chemosensory genes (6.0%). In contrast, the percentages of up- and downregulated chemosensory genes were quite similar in *D. simulans*; 10 (5.2%) and 7 (5.6%) among 191 upregulated and 125 downregulated genes, respectively. In summary, we proposed that some of these chemosensory genes have undergone Darwinian selection in the *D. sechellia* lineage.

Table 1

Expression Levels (FPKMs) and Log₂-Ratios of Chemosensory Genes Differentially Expressed in *Drosophila sechellia* (Dsec) Compared with *Drosophila melanogaster* (Dmel) or *Drosophila simulans* (Dsim)

Gene	Male			Female			Male			Female		
	Dsec	Dmel	Log ₂ -Ratio	Dsec	Dmel	Log ₂ -Ratio	Dsec	Dsim	Log ₂ -Ratio	Dsec	Dsim	Log ₂ -Ratio
Upregulated												
<i>Obp19a</i>	8,457.39	3,030.22	1.48	14,437.39	4,482.95	1.69	6,512.19	3,393.15	0.94	9,217.77	4,944.62	0.90
<i>Obp50a</i>	45.16	0.35	7.02	47.24	0.75	5.98	23.85	1.79	3.74	22.26	1.65	3.75
<i>Obp56d</i>	18,266.52	2,260.36	3.01	16,892.33	2,426.42	2.80	13,876.18	4,499.63	1.62	10,827.04	4,812.76	1.17
<i>CheA75a^a</i>	61.82	20.36	1.60	59.02	22.74	1.38	29.27	0.60	5.61	17.26	0.00	18.39
<i>CheA87a</i>	226.03	6.97	5.02	279.17	7.25	5.27	149.12	16.98	3.13	153.23	18.24	3.07
<i>Or23a</i>	44.99	4.42	3.35	44.62	4.16	3.42	40.34	14.91	1.44	34.92	13.72	1.35
<i>Or35a</i>	88.84	28.26	1.65	117.30	32.10	1.87	75.51	11.84	2.67	85.61	13.15	2.70
<i>Or56a</i>	98.88	40.70	1.28	109.11	48.20	1.18	111.82	29.27	1.93	109.05	33.82	1.69
<i>Or67b^a</i>	44.83	17.05	1.39	44.11	20.50	1.11	36.12	11.17	1.69	31.19	11.95	1.38
<i>Or85b</i>	156.59	42.97	1.87	151.82	50.11	1.60	128.78	17.20	2.90	107.25	20.00	2.42
<i>Or85c</i>	95.97	4.53	4.41	161.15	5.80	4.80	68.71	4.28	4.00	103.45	8.54	3.60
<i>Gr64f</i>	43.67	18.36	1.25	57.43	25.85	1.15	29.77	3.67	3.02	35.23	5.85	2.59
<i>Ir84a</i>	30.10	5.72	2.40	22.87	5.74	1.99	25.92	2.62	3.30	19.13	2.75	2.80
Downregulated												
<i>Obp83a</i>	7,520.39	16,673.69	-1.15	5,927.25	14,355.04	-1.28	6,196.39	10,887.62	-0.81	4,466.68	9,469.02	-1.08
<i>Obp99c</i>	22.62	170.49	-2.91	32.12	183.01	-2.51	54.44	487.17	-3.16	72.02	543.59	-2.92
<i>Obp99d</i>	6.02	60.59	-3.33	6.04	57.55	-3.25	2.55	36.81	-3.85	2.97	34.06	-3.52
<i>Or9a^a</i>	1.69	55.75	-5.05	6.36	86.09	-3.76	0.91	19.77	-4.44	3.09	48.11	-3.96
<i>Or42b^a</i>	115.98	214.92	-0.89	116.81	265.63	-1.19	110.75	245.24	-1.15	104.88	285.36	-1.44

NOTE.—FPKMs shown in this table were upper-quartile normalized. Differential expression was defined with $q \geq 0.95$ in the NOISeq test (Tarazona et al. 2012).

^aThose four genes do not show consistent results between qRT-PCR and RNA-seq.

Differentially Expressed Nonchemosensory Genes

Among the upregulated nonchemosensory genes, those most enriched in *D. sechellia* were mostly those with uncertain functions such as genes predicted to be in the subfamily of choline kinases (CG10513, CG10514, CG10553, CG32195, CG9497, and CG9498) and genes predicted to be involved in the following metabolism pathways (KEGG): Propanoate metabolism (succinyl-CoA synthetase [*Sucb*], CG10932, CG31075, and CG8778); butanoate and fatty acid metabolism (CG10932, CG31075, Arc42, and CG8778); tryptophan metabolism and lysine degradation (CG10932, CG31075, and CG8778) (supplementary table S14, Supplementary Material online). These genes tended to have higher fold-changes in expression between *D. sechellia* and its sibling species (log₂-ratios ranging from 1.13 to 2.83), suggesting that these genes might be associated with new functions related to the host-plant adaptation of *D. sechellia*.


The nonchemosensory genes downregulated in *D. sechellia* were enriched in the GO category of toxin response and in gene families putatively involved in odorant degradation (Younus et al. 2014). These genes included glutathione S transferase E9 and D8 (*GstE9* and *GstD8*), juvenile hormone eposide hydrolase 1, 2, and 3 (*Jheh1*, *Jheh2*, and *Jheh3*), and cytochrome P450 308a1 and 4d8 (*Cyp308a1* and *Cyp4d8*) (supplementary table S15, Supplementary Material online). None of the above genes and gene families was significantly

upregulated in *D. sechellia*, suggesting the need for further investigation of odorant degradation genes in *D. sechellia* antennae.

Expression Divergence of Chemosensory Genes between *D. sechellia* and Its Sibling Species

We generated heat maps for the *Or*, *Ir*, *Obp*, and *Gr* gene families and also the whole antennal transcriptome for each sex in each species (figs. 2–4 and supplementary figs. S2–S3, Supplementary Material online). We calculated the average expression level (without upper quartile normalization) of each gene estimated from three sequencing lanes followed by ranking within each of the above four gene families. Duplicate genes excluded in the OGS were now included for expression profile analysis. Genes with lowest expression levels (FPKM = 0 in all the samples) were ranked as 1 (colored yellow in the heat maps), whereas those with highest expression levels were colored red in the heat maps. Hierarchical agglomerative clustering was applied to cluster species as well as genes for each gene family. The significance of clustering was tested by multiscale bootstrap resampling implemented in the *pvclust* R package (Suzuki and Shimodaira 2006).

The results showed that the expression profiles of *Or* and *Ir* genes in *D. sechellia* antennae were significantly divergent from the two sibling species (figs. 2 and 3, approximately unbiased [*au*] *P* value > 0.95). The *Obp* genes also showed an



B Cluster dendrogram with AU/BP values (%)




FIG. 2.—Heat maps of expression profiles of *Or* genes. (A) Heat maps of ranked expression levels of *Or* genes. The red color indicates the highest expression level, whereas the yellow color indicates a low expression level. Clustering results of genes is shown on the left of the figure, and the gene names

expression profile divergent from the two sibling species, although *D. sechellia* was closer to *D. melanogaster* than to *D. simulans* (fig. 4). We did not observe any significant divergence in *Gr* genes or the rest of the transcriptome (supplementary figs. S2–S3, Supplementary Material online). The nonsignificant expression divergence in the *Gr* genes was likely due to the very low expression level of the gene family in the antennae.

The heat maps showed divergent expression between *D. sechellia* and the two sibling species for eight *Obp* and *Or* genes: *Obp99a*, *Or13a*, *Or22a*, *Or22b*, *Or45a*, *Or65c*, *Or67a*, and *Or83c*. Five of them (*Obp99a*, *Or22a*, *Or22b*, *Or65c*, and *Or67a*) were not included in OGS because they had undergone recent duplications or deletions in at least one of the three species. The *Or13a*, *Or45a*, and *Or83c* genes were not identified based on the criterion of $q \geq 0.95$.

In addition, some of the genes showed differential expression in terms of the FPKM values, but were not seen in the color patterns of heat maps. For instance, *Or56a* was ranked as one of the highest expressed *Or* genes in all samples, but the FPKM values were actually significantly different among the three species. We therefore were not able to observe differential expression of this gene from heat map directly. A similar situation was observed in *Obp* and *Ir* genes. Five of them (*Obp19a*, *Obp56d*, *Obp83a*, *Obp99c*, and *Ir84a*) found to be significantly differentiated were all expressed at very high levels, so they were denoted by the same color patterns.

Discussion

We speculate that the adaptation of *D. sechellia* to the new host was under positive selection. This view was supported by the enrichment of differentially expressed chemosensory genes, particularly upregulated chemosensory genes, in the *D. sechellia* lineage. We also showed that chemosensory genes expressed in different developmental stages or responsible for courtship behavior in one of the sibling species might have evolved new function and contributed to the host shift of *D. sechellia*. In addition, *Ir*, *Csp*, and *Che* genes might also have played a role in the host shift of *D. sechellia*.

Dekker et al. (2006) proposed that the host shift in *D. sechellia* resulted from great differences in the numbers of sensilla in antennae. According to their results, the number of ab3 sensilla increased 2.5- to 3-fold in *D. sechellia* and a reduction of 93% to 100% of ab2 sensilla. Based on this observation, we expected to see upregulation of ab3-specific genes but downregulation of ab2-specific genes

in *D. sechellia*. However, among ab3-specific genes, we found upregulation of only *Or85b* but not *Or22a* and *Or22b* in *D. sechellia*. Moreover, genes expressed on the ab2 sensilla, such as *Or59b* and *Or85a*, were not expressed consistently lower in *D. sechellia* than in the other two species as *D. sechellia* has fewer ab2 sensilla than the other two species. Therefore, the host shift may not be related to different numbers of sensilla in the antennae among the three species.

In this study, we included two strains of *D. sechellia* from two different labs, which were cultured in similar conditions. The two strains were derived from the same strain that has been widely used in fly laboratories. We did an analysis to compare expression levels of genes between the two strains of *D. sechellia*. There were only 131 genes differentially expressed (with the criterion of $q \geq 0.95$) between the two strains. Thus, our data from the two *D. sechellia* strains may be taken as two biological replicates.

Csp Genes

Compared with other chemosensory gene families, the *Csp* gene family has the smallest number of genes: Only four genes in the *Drosophila* genome including *a10* (also known as *Os-D*), *CG30172*, *PebIII*, and *Phk-3* (Zhou et al. 2006). Among these four genes, only *a10* is antenna-specific and expressed 6-fold higher than the other *Csp* genes in *D. melanogaster* (Zhou et al. 2006; Foret et al. 2007). Our data confirmed that *a10* was expressed much higher than other *Csp* genes in all three species.

If $q = 0.8$ was used to determine DEGs (supplementary tables S5–S8, Supplementary Material online), then *Phk-3* was expressed significantly higher in *D. sechellia* than in the two sibling species ($q = 0.85–1.00$ in the comparisons). Gene *a10* and *PebIII* were expressed higher in *D. sechellia* than in *D. melanogaster* but lower in *D. sechellia* than in *D. simulans*. *CG30172* showed higher expression in *D. simulans* than in *D. sechellia*, but not higher than in *D. melanogaster*. Based on these observations, we suggest that only *Phk-3* might be associated with the host shift.

Ir Genes

Benton et al. (2009) identified 61 *Ir* genes in *Drosophila*. Using qRT-PCR or RNA in situ hybridization, the authors showed that 15 *Ir* genes were expressed in fly antennae. No expression was detected for the other *Ir* genes in any of the tissues studied. It remains unclear whether these genes were not expressed, expressed at different life stages, or expressed at levels below the detection threshold of the two techniques. In

FIG. 2.—Continued

and expressed sensilla (*Or* genes) are shown on the right. Clustering of species is shown at the top of each figure and the information of each sample is shown at the bottom of the figure. Dmel, *D. melanogaster*; Dsim, *D. simulans*; Dsec, *D. sechellia*; M, males; F, females; TW, Taiwan; JP, Japan. (B) Statistical significance of clustering results among three species. The approximately unbiased (*au*) *P* value (% in red) and bootstrap probability (*bp*) value (% in green) are shown on each branch (in gray color).





Fig. 3.—Heat maps of expression profiles of *Ir* genes. (A) Heat maps of ranked expression levels of *Ir* genes and (B) statistical significance of clustering results among three species. Color patterns and sample names are as described in figure 2.




Fig. 4.—Heat maps of expression profiles of *Obp* genes. (A) Heat maps of ranked expression levels of *Obp* genes and (B) statistical significance of clustering results among three species. Color patterns and sample names are as described in figure 2.

Table 2

Expression Levels of Genes Proposed to be Associated with Host-Plant Adaptation in *Drosophila sechellia* in Literature

Gene Name	Function/Sensilla Types	References	Sex	Taiwan		Japan	
				<i>Drosophila melanogaster</i>	<i>Drosophila sechellia</i>	<i>Drosophila simulans</i>	<i>Drosophila sechellia</i>
<i>Orco</i>	Universal OR		Males	1,860.84	1,418.23	1,587.59	1,156.52
			Females	1,996.82	1,400.83	1,856.26	1,100.28
<i>Or19a</i>	Sensing citrus	Dweck et al. (2013)	Males	2.00	14.07	7.64	13.32
			Females	2.16	16.01	9.73	14.84
<i>Or22a</i>	Ab3 sensilla	Hallem and Carlson (2006) Kopp et al. (2008) Dekker et al. (2006)	Males	11.67	1.06	75.07	1.09
			Females	15.04	0.29	108.11	0.74
<i>Or22b</i>	Ab3 sensilla	Kopp et al. (2008) Dekker et al. (2006)	Males	71.97	0.00	61.53	0.00
			Females	82.33	0.00	85.56	0.00
<i>Or59b</i>	Ab2 sesilla	Dekker et al. (2006)	Males	73.95	106.80	192.70	104.11
			Females	120.07	151.79	303.92	142.07
<i>Or85a</i>	Ab2 sesilla	Dekker et al. (2006)	Males	16.32	45.68	31.47	22.87
			Females	21.92	74.15	51.18	35.98
<i>Or98a</i>	Sensing MeHex	Hallem and Carlson (2006)	Males	34.96	50.92	5.62	29.05
			Females	39.94	58.93	7.96	29.32
<i>Or85b</i>	Sensing MeHex Ab3 sensilla	Hallem and Carlson (2006) Kopp et al. (2008) Dekker et al. (2006)	Males	42.97	156.59	17.20	128.78
			Females	50.11	151.82	20.00	107.25
<i>Or47b</i>	Sensing MeHex	Hallem and Carlson (2006)	Males	89.18	66.00	62.48	119.29
			Females	51.51	39.30	41.11	65.91
<i>Obp57d</i>	Host-plant preference	Higa and Fuyama (1993) Matsuo et al. (2007)	Males	0.00	0.53	0.00	0.56
			Females	0.00	0.37	0.00	0.63
<i>Obp57e</i>	Host-plant preference	Higa and Fuyama (1993) Matsuo et al. (2007)	Males	6.45	2.43	2.66	0.98
			Females	3.27	1.11	3.79	0.74

NOTE.—Expression levels (FPKMs) in bold indicate that the gene was differentially expressed between the two species. The FPKM values for some of the genes, such as *Or22a* and *Or22b*, were not normalized because they were not included in the OGS. *Obp57d* overlapped with *Cpr57A* based on gene annotation file, resulting in no read mapped to *Obp57d* in *D. melanogaster* and *D. simulans*.

addition, 13 of the 15 *Ir* genes did not overlap in expression with the *Or* genes or *Orco* in adult antennae. The RNA-seq data in this study confirmed the expression of 10 of the 15 *Ir* genes reported in Benton et al. (2009) (supplementary table S4, Supplementary Material online). The other five genes (*Ir8a*, *Ir31a*, *Ir40a*, *Ir75b*, and *Ir93a*) were not included in the OGS, possibly due to incomplete gene annotation in the *D. simulans* and *D. sechellia* genomes.

Grosjean et al. (2011) proposed that *Ir84a*, a food-derived receptor, promotes male courtship in *D. melanogaster*. The authors showed that *Ir84a* was activated by food-derived aromatic odors, phenylacetic acid and phenylacetaldehyde, but not by fly-derived chemicals. The two chemicals are common in plants and fruits used as food sources and oviposition sites by *Drosophila*. This receptor was expressed in the OSNs that also expressed *Ir75d*, or *Ir76a* and *Ir76b* (Benton et al. 2009). Interestingly, we found that *Ir84a* and *Ir76a* were differentially expressed in *D. sechellia*. *Ir84a* was overexpressed in *D. sechellia* compared with the two sibling species. *Ir76a* was expressed significantly higher in *D. sechellia* than in *D. melanogaster* in both sexes, and was expressed significantly higher in females, though significantly lower in

males compared with *D. simulans* (supplementary table S8, Supplementary Material online). Thus, *Ir84a* and *Ir76a* could have been associated with the host shift of *D. sechellia*.

Other Genes

We now discuss the genes previously proposed to be associated with the host shift in *D. sechellia*. The FPKM values of these genes are summarized in table 2.

The major volatile odors of *M. citrifolia* fruits are contributed by a variety of compounds, including HA, OA, methyl octanoate (MeOct), methyl hexanoate (MeHex), and 3-methyl-3-butenyl octanoate (Legal et al. 1994; Dekker et al. 2006; Wei et al. 2011). Although the acids HA and OA comprise the vast majority of the volatiles in *M. citrifolia*, a previous study showed that *D. sechellia* is more sensitive to the ester hexanoate than to acids, and is particularly sensitive to the hexanoate MeHex (Dekker et al. 2006). Hallem and Carlson (2006) showed that the two ester compounds, MeHex and MeOct, mostly stimulate *Or22a*, *Or85b* and *Or98a*, whereas they mostly inhibit *Or47b* in *D. melanogaster* (fig. 2). Using microarray expression analysis, Kopp et al.

(2008) showed that *Or22a*, *Or22b*, and *Or85b* were significantly overexpressed in *D. sechellia*. However, *Or22a* and *Or22b* were also differentially expressed between *D. simulans* and *D. melanogaster*, and *Or22a* showed significant differences between two strains of *D. melanogaster*. For these two genes, our data are inconclusive as there are duplicated genes in one of the genomes, so that *Or22a* and *Or22b* were not included in OGS. In our data, *Or85b* was expressed at a significantly higher level in *D. sechellia* than in *D. melanogaster* and *D. simulans* (all the q values > 0.97). Also, *Or98a* was expressed at significant higher levels in both sexes in *D. sechellia*. *Or47b*, a receptor inhibited in *D. melanogaster*, showed no significant difference between *D. melanogaster* and *D. sechellia*, although a significantly higher expression level was detected in *D. sechellia* than in *D. simulans*. Therefore, only *Or85b* might be involved in the host shift of *D. sechellia*.

Dweck et al. (2013) proposed that *Or19a* is very important for the preference of citrus fruits as egg-laying substrate in flies. Our data are consistent with this hypothesis, because we found *Or19a* significantly upregulated in *D. sechellia* compared with the other two species.

Matsuo and his coworkers suggested that *Obp57d* and *Obp57e* are associated with the host-plant preference of *D. sechellia* (Higa and Fuyama 1993; Matsuo et al. 2007). Knocking out *Obp57d* and *Obp57e* in *D. melanogaster* led to a preference to the odorants emitted by the host-plant of *D. sechellia*. Using qRT-PCR, the authors found that *Obp57d* and *Obp57e* were expressed mainly in the legs of all three species. *Obp57d* was not included in our OGS because it completely overlaps with a cuticular protein 57A (*Cpr57A*) gene in *D. melanogaster*, making it impossible to do an informative differential expression analysis with *D. melanogaster*. Moreover, *Obp57d* was expressed at a very low level in the antennae of *D. sechellia* (FPKM < 1 in all *D. sechellia* samples). As for *Obp57e*, when lowering the criterion to $q=0.8$, we found it to be expressed at a significantly lower level in *D. sechellia* (FPKM = 0.74 in Taiwan and 2.43 in Japan) than in *D. melanogaster* (FPKM = 6.45, $P=0.82$) and *D. simulans* (FPKM = 3.79, $P=0.84$).

Hungate et al. (2013) identified a genomic locus spanning approximately 170 KB comprising 18 genes, which are related to the OA tolerance in *D. sechellia*. These 18 genes included two major gene clusters belonging to the *Obp* gene family and the *Osi* gene family. The *Obp* gene cluster includes three *Obp* genes: *Obp83cd*, *Obp83ef*, and *Obp83g*. In our data, *Obp83ef* was upregulated in *D. sechellia* compared with *D. melanogaster*, but not *D. simulans*. The reason is that *Obp83ef* was expressed at very low levels in *D. melanogaster* compared with the other two species. Thus, it is unlikely that *Obp83ef* was involved in the host shift of *D. sechellia*. *Obp83cd* was not included in OGS and *Obp83g* showed an expression level of FPKM < 10 across all the data sets. All the genes (nine genes) in the *Osi* gene cluster with the exception

of *Osi8* and *Gasp* were expressed at very low levels in the antennae of all three species. Taken together, these observations suggest that this genomic locus probably does not play an important role in antenna-mediated adaptation, though they may be important in OA tolerance in other body parts.

In conclusion, our study suggests that all chemosensory gene families might have contributed to the host shift of *D. sechellia*, likely through upregulation of some chemosensory genes. That is, besides the genes proposed by previous studies, some additional chemosensory genes might have contributed to the host shift of *D. sechellia*.

Although we have shown that there was enrichment of differentially expressed chemosensory genes in *D. sechellia*, functional tests are needed to check whether any of the candidate genes contributed to the host shift of *D. sechellia*. Two approaches could be used to study the function of candidate chemosensory genes in the future. First, we could apply the *Escherichia coli*-based cell free expression system developed by Tegler et al. (2015) to express olfactory receptor genes of *D. melanogaster* to test the ligand binding affinity. This system would be able to help us confirm the binding affinity of candidate genes to the chemicals emitted by the host plant. Second, we could apply the CRISPR/Cas9 system to generate loss-of-function mutations in a specific gene or introduce a desired mutation by homologous repair in *Drosophila* (the review by Bassett and Liu 2014). With this technique, plasmids carrying guiding DNA of candidate chemosensory genes could be injected into embryos of *D. sechellia* and crossed to Cas9 strains. We can evaluate the function of mutated candidate genes by observing the behavior of mutated *D. sechellia* in terms of moving toward or avoiding chemicals emitted by the host plant.

Data Accessibility

The raw sequencing data reported in this work have been deposited in the NCBI GEO with accession numbers GSE67587, GSE67861 and GSE67862 for *D. sechellia* (Taiwan), *D. sechellia* (Japan) and *D. simulans* (Japan), respectively.

Supplementary Material

Supplementary figures S1–S3 and tables S1–S15 are available at *Genome Biology and Evolution* online (<http://www.gbe.oxfordjournals.org/>).

Acknowledgments

This work was supported by Postdoctoral Fellows Program, Academia Sinica, Taiwan (to M.-S. S.); Plan Nacional (BFU2011–28575 to C.N. and J.-M.C.); Spanish Ministry of Economy and Competitiveness, “Centro de Excelencia Severo Ochoa 2013–2017” (SEV- 2012–0208 to C.N. and J.-M.C.); “Fundacio’ Obra Social la Caixa” (to J.-M.C.); and

European Research Council (ERC- 2008-AdG No 232947 to J.-M.C.). The authors thank Dr Carol Lee, University of Wisconsin-Madison for valuable comments on the manuscript.

Literature Cited

- Arya GH, et al. 2010. Natural variation, functional pleiotropy and transcriptional contexts of odorant binding protein genes in *Drosophila melanogaster*. *Genetics* 186:1475–1485.
- Bassett AR, Liu J-LL. 2014. CRISPR/Cas9 and genome editing in *Drosophila*. *J Genet Genomics*. 41:7–19.
- Benton R, Vannice KS, Gomez-Diaz C, Vosshall LB. 2009. Variant ionotropic glutamate receptors as chemosensory receptors in *Drosophila*. *Cell* 136:149–162.
- Clyne PJ, et al. 1999. A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila*. *Neuron* 22:327–338.
- Clyne PJ, Warr CG, Carlson JR. 2000. Candidate taste receptors in *Drosophila*. *Science* 287:1830–1834.
- Couto A, Alenius M, Dickson BJ. 2005. Molecular, anatomical, and functional organization of the *Drosophila* olfactory system. *Curr Biol*. 15:1535–1547.
- Dekker T, Ibba I, Siju KP, Stensmyr MC, Hansson BS. 2006. Olfactory shifts parallel superspecialism for toxic fruit in *Drosophila melanogaster* sibling, *D. sechellia*. *Curr Biol*. 16:101–109.
- Dweck HK, et al. 2013. Olfactory preference for egg laying on citrus substrates in *Drosophila*. *Curr. Biol*. 23:2472–2480.
- Dworkin I, Jones CD. 2009. Genetic changes accompanying the evolution of host specialization in *Drosophila sechellia*. *Genetics* 181:721–736.
- Foret S, Wanner KW, Maleszka R. 2007. Chemosensory proteins in the honey bee: insights from the annotated genome, comparative analyses and expressional profiling. *Insect Biochem Mol Biol*. 37:19–28.
- Galindo K, Smith DP. 2001. A large family of divergent *Drosophila* odorant-binding proteins expressed in gustatory and olfactory sensilla. *Genetics* 159:1059–1072.
- Garrigan D, et al. 2012. Genome sequencing reveals complex speciation in the *Drosophila simulans* clade. *Genome Res*. 22:1499–1511.
- Grosjean Y, et al. 2011. An olfactory receptor for food-derived odours promotes male courtship in *Drosophila*. *Nature* 478:236–240.
- Ha TS, Smith DP. 2006. A pheromone receptor mediates 11-cis-vaccenyl acetate-induced responses in *Drosophila*. *J Neurosci*. 26:8727–8733.
- Hallem EA, Carlson JR. 2006. Coding of odors by a receptor repertoire. *Cell* 125:143–160.
- Higa I, Fuyama Y. 1993. Genetics of food preference in *Drosophila sechellia*. *Genetica* 88:129–136.
- Hungate EA, et al. 2013. A locus in *Drosophila sechellia* affecting tolerance of a host plant toxin. *Genetics* 195:1063–1075.
- Immonen E, Ritchie MG. 2012. The genomic response to courtship song stimulation in female *Drosophila melanogaster*. *Proc Biol Sci*. 279:1359–1365.
- Jones WD, Cayirlioglu P, Kadow IG, Vosshall LB. 2007. Two chemosensory receptors together mediate carbon dioxide detection in *Drosophila*. *Nature* 445:86–90.
- Kim MS, Repp A, Smith DP. 1998. LUSH odorant-binding protein mediates chemosensory responses to alcohols in *Drosophila melanogaster*. *Genetics* 150:711–721.
- Kliman RM, et al. 2000. The population genetics of the origin and divergence of the *Drosophila simulans* complex species. *Genetics* 156:1913–1931.
- Kopp A, et al. 2008. Evolution of gene expression in the *Drosophila* olfactory system. *Mol Biol Evol*. 25:1081–1092.
- Kreher SA, Kwon JY, Carlson JR. 2005. The molecular basis of odor coding in the *Drosophila* larva. *Neuron* 46:445–456.
- Lachaise D, et al. 1988. Historical biogeography of the *Drosophila melanogaster* species subgroup. *Evol Biol*. 22:159–225.
- Larsson MC, et al. 2004. Or83b encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron* 43:703–714.
- Legal L, Chappe B, Jallon JM. 1994. Molecular-basis of *Morinda citrifolia* (L)—toxicity on *Drosophila*. *J Chem Ecol*. 20:1931–1943.
- Louis J, David JR. 1986. Ecological specialization in the *Drosophila melanogaster* species subgroup—a case-study of *Drosophila sechellia*. *Acta Oecol Oecol Generalis*. 7:215–229.
- Mathew D, et al. 2013. Functional diversity among sensory receptors in a *Drosophila* olfactory circuit. *Proc Natl Acad Sci U S A*. 110:E2134–E2143.
- Matsuo T, Sugaya S, Yasukawa J, Aigaki T, Fuyama Y. 2007. Odorant-binding proteins OBP57d and OBP57e affect taste perception and host-plant preference in *Drosophila sechellia*. *PLoS Biol*. 5:e118.
- McBride CS, Arguello JR. 2007. Five *Drosophila* genomes reveal nonneutral evolution and the signature of host specialization in the chemoreceptor superfamily. *Genetics* 177:1395–1416.
- McDermott SR, Kliman RM. 2008. Estimation of isolation times of the island species in the *Drosophila simulans* complex from multilocus DNA sequence data. *PLoS one* 3:e2442.
- Rio B, Couturier G, Lemeunier F, Lachaise D. 1983. Evolution d'une specialisation saisonniere chez *Drosophila erecta* (Diptera, Drosophilidae). *Ann Entomol Soc Fr*. 19:235–248.
- Rkha S, Capy P, David JR. 1991. Host plant specialization in the *Drosophila melanogaster* species complex—a physiological, behavioral, and genetic-analysis. *Proc Natl Acad Sci U S A*. 88:1835–1839.
- Shiao MS, et al. 2012. Transcriptomes of mouse olfactory epithelium reveal sexual differences in odorant detection. *Genome Biol Evol*. 4:703–712.
- Shiao MS, et al. 2013. Transcriptional profiling of adult *Drosophila* antennae by high-throughput sequencing. *Zool Stud*. 52:42–51.
- Starostina E, Xu A, Lin H, Pikielny CW. 2009. A *Drosophila* protein family implicated in pheromone perception is related to Tay-Sachs GM2-activator protein. *J Biol Chem*. 284:585–594.
- Stengl M, Funk NW. 2013. The role of the coreceptor Orco in insect olfactory transduction. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*. 199:897–909.
- Stensmyr MC, Dekker T, Hansson BS. 2003. Evolution of the olfactory code in the *Drosophila melanogaster* subgroup. *Proc Biol Sci*. 270:2333–2340.
- Suzuki R, Shimodaira H. 2006. Pvcust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics* 22:1540–1542.
- Swarup S, Morozova TV, Sridhar S, Nokes M, Anholt RR. 2014. Modulation of feeding behavior by odorant-binding proteins in *Drosophila melanogaster*. *Chem Senses*. 39:125–132.
- Tarazona S, et al. 2012. NOseq: a RNA-seq differential expression method robust for sequencing depth biases. *EMBnet* 17:18–19.
- Tegler LT, et al. 2015. Cell-free expression, purification, and ligand-binding analysis of *Drosophila melanogaster* olfactory receptors DmOR67a, DmOR85b and DmORCO. *Sci Rep*. 5:7867.
- Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 25:1105–1111.
- Tsacas L, Bachli G. 1981. *Drosophila sechellia* n. sp., huitieme espece du sous-group melanogaster des ilees Sechelles (Diptera, Drosophilidae). *Rev Fr Entomol*. 3:146–150.
- Vieira FG, Rozas J. 2011. Comparative genomics of the odorant-binding and chemosensory protein gene families across the Arthropoda: origin and evolutionary history of the chemosensory system. *Genome Biol Evol*. 3:476–490.
- Vosshall LB, Amrein H, Morozov PS, Rzhetsky A, Axel R. 1999. A spatial map of olfactory receptor expression in the *Drosophila* antenna. *Cell* 96:725–736.

- Wei GJ, Ho CT, Huang AS. 2011. Analysis of volatile compounds in noni fruit (*Morinda citrifolia* L.) juice by steam distillation-extraction and solid phase microextraction coupled with GC/AED and GC/MS. *J Food Drug Anal.* 19:33–39.
- Xu A, et al. 2002. Novel genes expressed in subsets of chemosensory sensilla on the front legs of male *Drosophila melanogaster*. *Cell Tissue Res.* 307:381–392.
- Xu P, Atkinson R, Jones DN, Smith DP. 2005. *Drosophila* OBP LUSH is required for activity of pheromone-sensitive neurons. *Neuron* 45:193–200.
- Younus F, et al. 2014. Identification of candidate odorant degrading gene/enzyme systems in the antennal transcriptome of *Drosophila melanogaster*. *Insect Biochem Mol Biol.* 53:30–43.
- Zhang YV, Ni J, Montell C. 2013. The molecular basis for attractive salt-taste coding in *Drosophila*. *Science* 340:1334–1338.
- Zhou JJ, Kan Y, Antoniw J, Pickett JA, Field LM. 2006. Genome and EST analyses and expression of a gene family with putative functions in insect chemoreception. *Chem Senses.* 31:453–465.

Associate editor: Soojin Yi