



HHS Public Access

Author manuscript

Nat Rev Genet. Author manuscript; available in PMC 2015 December 22.

Published in final edited form as:

Nat Rev Genet. 2009 September ; 10(9): 639–650. doi:10.1038/nrg2611.

Genetics in geographically structured populations: defining, estimating and interpreting F_{ST}

Kent E. Holsinger* and Bruce S. Weir†

Kent E. Holsinger: kent@darwin.eeb.uconn.edu; Bruce S. Weir: bsweir@u.washington.edu

*Department of Ecology and Evolutionary Biology, U-3043, University of Connecticut, Storrs, Connecticut 06269-3043, USA

†Department of Biostatistics, University of Washington, Box 357232, Seattle, Washington 98195, USA

Abstract

Wright's F -statistics, and especially F_{ST} , provide important insights into the evolutionary processes that influence the structure of genetic variation within and among populations, and they are among the most widely used descriptive statistics in population and evolutionary genetics. Estimates of F_{ST} can identify regions of the genome that have been the target of selection, and comparisons of F_{ST} from different parts of the genome can provide insights into the demographic history of populations. For these reasons and others, F_{ST} has a central role in population and evolutionary genetics and has wide applications in fields that range from disease association mapping to forensic science. This Review clarifies how F_{ST} is defined, how it should be estimated, how it is related to similar statistics and how estimates of F_{ST} should be interpreted.

Nearly every plant or animal species includes many partially isolated populations. As a result of genetic drift or divergent natural selection, such populations become genetically differentiated over time. For example, recent analyses based on more than 370

FURTHER INFORMATION

Kent E. Holsinger's homepage: <http://darwin.eeb.uconn.edu>

1,000 Genomes project: <http://www.1000genomes.org>

ABC4F (approximate Bayesian computation for F -statistics): <http://www-leca.ujf-grenoble.fr/logiciels.htm>

Arlequin (an integrated software application for population genetics data analysis): <http://cmpg.unibe.ch/software/arlequin3>

BayeScan (BAYesian genome SCAN for outliers): <http://www-leca.ujf-grenoble.fr/logiciels.htm>

Bayesian population genetic data analysis: <http://darwin.eeb.uconn.edu/summer-institute/summer-institute.html>

GDA (Genetic Data Analysis): <http://www.eeb.uconn.edu/people/plewis/software.php>

GenAIEx (integrated software for analysis of genetic data with an interface to Excel): http://www.anu.edu.au/BoZo/GenAIEx/genalex_download_6_1.php

Genepop: <http://kimura.univ-montp2.fr/~rousset/Genepop.htm>

GESTE (GENetic STRucture inference based on genetic and Environmental data): <http://www-leca.ujf-grenoble.fr/logiciels.htm>

Hickory (software for the analysis of geographic structure in genetic data): <http://darwin.eeb.uconn.edu/hickory/hickory.html>

Hierfstat (Weir & Cockerham F -statistics for any number of levels in a hierarchy): <http://www2.unil.ch/popgen/softwares/hierfstat.htm>

International HapMap Project: <http://www.hapmap.org>

Nature Reviews Genetics series on Fundamental Concepts in Genetics: <http://www.nature.com/nrg/series/fundamental/index.html>

The genetic structure of populations: http://darwin.eeb.uconn.edu/eeb348/lecture.php?rl_id=402

The genetic structure of populations: a Bayesian approach: http://darwin.eeb.uconn.edu/eeb348/lecture.php?rl_id=403

The Wahlund effect and Wright's F -statistics: http://darwin.eeb.uconn.edu/eeb348/lecture.php?rl_id=445

ALL LINKS ARE ACTIVE IN THE ONLINE PDF

short tandem repeat loci¹ (microsatellites) and 600,000 SNPs² suggest that only 5–10% of human genetic diversity is accounted for by genetic differences among populations from major geographical regions. These results indicate that there are far more similarities among geographically distinct human populations than differences. But what does it mean to say that 5–10% of diversity is accounted for by differences among populations, and how is this figure derived? The short answer is that the estimate of F_{ST} among human populations sampled from these regions is 0.05 for the microsatellite data and 0.10 for the SNP data. However, this answer helps only if one understands what F_{ST} is, how it is estimated from data and what it means to get two different estimates for the same set of populations when different genetic markers are used.

Working independently in the 1940s and 1950s, Sewall Wright³ and Gustave Malécot⁴ introduced F -statistics as a tool for describing the partitioning of genetic diversity within and among populations. In a paper published in 1931 (REF. 5), Wright had already provided a comprehensive account of the processes that cause genetic differentiation among populations. He showed that the amount of genetic differentiation among populations has a predictable relationship to the rates of important evolutionary processes (migration, mutation and drift). For example, large populations among which there is much migration tend to show little differentiation, whereas small populations among which there is little migration tend to be highly differentiated. F_{ST} is a convenient measure of this differentiation, and as a result F_{ST} and related statistics are among the most widely used descriptive statistics in population and evolutionary genetics.

But F_{ST} is more than a descriptive statistic and a measure of genetic differentiation. F_{ST} is directly related to the variance in allele frequency among populations and, conversely, to the degree of resemblance among individuals within populations. If F_{ST} is small, it means that the allele frequencies within each population are similar; if it is large, it means that the allele frequencies are different. If natural selection favours one allele over others at a particular locus in some populations, the F_{ST} at that locus will be larger than at loci in which among-population differences are purely a result of genetic drift. Genome scans that compare single-locus estimates of F_{ST} with the genome-wide background might therefore identify regions of the genome that have been subjected to diversifying selection^{6–8}. Alternatively, if the demographic history of populations affects the genetic variation on sex chromosomes in a different way from the genetic variation on autosomes, the estimates of F_{ST} derived from sex chromosome markers might be different from those derived from autosomal markers⁹.

Estimates of F_{ST} are also important in association mapping of human disease genes and in forensic science. The same evolutionary processes that increase differentiation among populations also increase the similarity among individuals within populations. Therefore, F_{ST} must be considered when allele frequencies are compared between cases and controls to ensure that the differences between them are greater than expected by chance. Similarly, the probability of a match between a suspect and a crime scene sample is specific to the set of people who might reasonably be expected to be sources of the sample. However, defining this set is difficult, so a ‘ θ correction’ is applied to population frequencies to accommodate variation among subpopulations. The θ correction depends on the value of F_{ST} .

In this review, we discuss how F_{ST} is defined, describe approaches for estimating it from data and illustrate several ways in which analysis of F_{ST} can provide insights into the genetic structure and evolutionary dynamics of populations. In addition, we discuss four statistics that are related to F_{ST} (G_{ST} , R_{ST} , Φ_{ST} and Q_{ST}), clarify the differences among them and recommend when each should be used.

These additional statistics partition genetic diversity into within- and among-population components. of the four, G_{ST} is most closely related to F_{ST} , and it has been widely used as a measure of genetic differentiation among populations. However, as we describe below, G_{ST} is an appropriate measure of genetic differentiation only when the contribution of genetic drift to among-population differences is not of interest. As a result, the contexts in which it is useful are limited. By contrast, R_{ST} (for microsatellite data) and Φ_{ST} (for molecular sequence data) are useful in a wide range of contexts in which it is important to account for the mutational ‘distances’ among alleles, and Q_{ST} is useful in the analysis of continuously varying traits.

Definitions

Wright introduced F_{ST} as one of three interrelated parameters to describe the genetic structure of diploid populations³. These parameters are: F_{IT} , the correlation between gametes within an individual relative to the entire population; F_{IS} , the correlation between gametes within an individual relative to the subpopulation to which that individual belongs; and F_{ST} , the correlation between gametes chosen randomly from within the same subpopulation relative to the entire population. We describe here how these parameters are defined in terms of the departure of genotype frequencies from Hardy-Weinberg proportions.

Deriving measures of genetic diversity

As an example of how to calculate genetic diversity, consider two populations that are segregating for two alleles at a single locus. The frequency of allele A_1 in the first population is labelled as p_1 and its frequency in the second population is labelled as p_2 . The frequency of genotype A_1A_1 in the first population is labelled as $x_{11,1}$, the frequency of genotype A_1A_2 in the first population is labelled as $x_{12,1}$, and so on. The genotype frequencies in the two populations are given by the following set of equations:

$$x_{11,1} = p_1^2 + f_1 p_1 (1 - p_1)$$

$$x_{12,1} = 2p_1(1 - p_1)(1 - f_1)$$

$$x_{22,1} = (1 - p_1)^2 + f_1 p_1 (1 - p_1)$$

$$x_{11,2} = p_2^2 + f_2 p_2 (1 - p_2)$$

$$x_{12,2} = 2p_2(1 - p_2)(1 - f_2)$$

$$x_{22,2} = (1 - p_2)^2 + f_2 p_2 (1 - p_2) \quad (1)$$

In this context, f_1 and f_2 are often referred to as the within-population inbreeding coefficients, but this term can be misleading. In practice, f is a measure of the frequency of heterozygotes compared with that expected when genotypes are in Hardy–Weinberg proportions. Inbreeding leads to a deficiency of heterozygotes relative to Hardy–Weinberg expectations, so when there is inbreeding in both populations, f_1 and f_2 will have positive values. But if individuals avoid inbreeding or if there is heterozygote advantage, then heterozygotes will be more common than expected under Hardy–Weinberg expectations, and f_1 and f_2 will be negative. In short, f_1 and f_2 are measures of how different the genotype proportions within populations are from Hardy–Weinberg expectations, and positive values of f indicate a deficiency of heterozygotes, whereas negative values indicate an excess.

Now consider the genotype frequencies in a combined sample that consists of a proportion c of individuals from the first population and a proportion $1 - c$ of individuals from the second population. Similar to the way in which the genotype frequencies in each population differ from Hardy–Weinberg expectations based on the allele frequency in each population, genotype frequencies in the combined sample differ from Hardy–Weinberg expectations based on the average allele frequency. The allele frequencies are given by:

$$x_{11} = \pi^2 + F\pi(1 - \pi)$$

$$x_{12} = 2\pi(1 - \pi)(1 - F)$$

$$x_{22} = (1 - \pi)^2 + F\pi(1 - \pi) \quad (2)$$

in which $\pi = cp_1 + (1 - c)p_2$ is the average allele frequency for A_1 in the combined sample and F is the total inbreeding coefficient¹⁰. F can be expressed as:

$$(1 - F) = (1 - f)(1 - \theta) \quad (3)$$

in which $f = cf_1 + (1 - c)f_2$ is the average within-population departure from Hardy–Weinberg expectations and θ is a measure of allele frequency differentiation among populations (see BOX 1 for a summary of the mathematical notation used in this review). We can define θ as:

$$\theta = \frac{\sigma_{\pi}^2}{\pi(1 - \pi)} \quad (4)$$

in which σ_{π}^2 is the variance in allele frequency among populations. $\pi(1 - \pi)$ is the variance in the allelic state for an allele chosen randomly from the entire population, so it can be regarded as a measure of genetic diversity in the entire population. θ can therefore be interpreted as the proportion of genetic diversity that is due to the differences in allele frequency among populations.

Wright first developed these ideas in the context of a model of discrete populations, in which each population is the same size and receives immigrants from all other populations at the same rate⁵. However, the same statistical argument can be applied to any partitioning of genetic diversity in which the populations differ in allele frequency, whether or not those populations are discrete¹¹. Therefore, when we use θ as a purely descriptive statistic that describes the partitioning of genetic diversity among ‘populations’, we do not need to make assumptions about whether the ‘populations’ we sample are discrete or about the evolutionary processes that might have led to differences among them. Nonetheless, other methods of analysis could be more informative in continuously distributed populations^{12–14}.

Linking f , θ and F to Wright’s F -statistics

Using a different approach, Cockerham^{10,15} showed that f , θ and F can also represent intraclass correlation coefficients. He showed that f is the correlation between alleles within individuals relative to the population to which they belong, θ is the correlation between alleles within populations relative to the combined population and F is the correlation between alleles within individuals relative to the combined population. These are the definitions that Wright gave for F_{IS} , F_{ST} and F_{IT} , respectively. In short, f and F_{IS} can be thought of either as the average within-population departure from Hardy–Weinberg expectations or as the correlation between alleles within individuals relative to the population to which they belong. θ and F_{ST} can be thought of either as the proportion of genetic diversity due to allele frequency differences among populations or as the correlations between alleles within populations relative to the entire population. F and F_{IT} can be thought of either as the departure of genotype frequencies in the combined sample from Hardy–Weinberg expectations or as the correlation between alleles within individuals relative to the combined sample.

In Wright’s notation, subscripts refer to a comparison between levels in a hierarchy: IS refers to ‘individuals within subpopulations’, ST to ‘subpopulations within the total population’ and IT to ‘individuals within the total population’¹⁶. The hierarchy in equation 1 can be extended indefinitely to accommodate such structures. For example, Wright¹⁶ describes variation in the frequency of the Standard chromosome in *Drosophila pseudoobscura* in the western United States at the level of demes (D; local populations), regions (R; groups of several demes), subdivisions (S; groups of several regions) and the total range (T). The corresponding F -statistics are related in the same multiplicative way as f , θ and F :

$$(1 - F_{DT}) = (1 - F_{DR})(1 - F_{RS})(1 - F_{ST}) \quad (5)$$

In this scheme, F_{DR} measures the differentiation among demes within a region, F_{RS} measures the differentiation among regions within subdivisions and F_{ST} measures the differentiation among subdivisions within the total range.

If we return to the examples of genetic differentiation among human populations that were mentioned at the beginning of this review, we can now see that an estimate for F_{ST} or θ of 0.05 (from microsatellites) and 0.10 (from SNPs) suggests that only 5–10% of human genetic diversity is a result of genetic differentiation among human populations. What might be surprising is that the two estimates are derived from the same set of populations — this indicates that the amount of genetic differentiation among human populations is greater at SNP loci than at microsatellites.

Estimation

Statistical sampling

When Wright and Malécot introduced F -statistics, they did not distinguish between the parameters defined in the preceding section and the estimates of those parameters that we make from data. Not making this distinction is similar to confusing the mean height of the human population with an estimate of the mean height calculated from a sample of the population. Estimates of height must account for the variation associated with taking a finite sample from a population. New samples from the same population will have different characteristics. We refer to this variation as statistical sampling¹⁷ (BOX 2). In the context of F -statistics, statistical sampling refers to the variation associated with collecting genetic samples from a fixed set of populations that have fixed but unknown genotype frequencies. The magnitude of variation associated with statistical sampling can be reduced by increasing the size of within-population samples.

Genetic sampling

There is an important difference between estimates made by F -statistics and estimates of height. In addition to accounting for statistical sampling, F -statistics must account for differences among the sets of populations that have been sampled. These differences might arise either because the populations that are sampled are only a subset of all of the populations that could be sampled (statistical sampling of populations rather than statistical sampling of genotypes within populations) or because the populations that are sampled represent only one possible outcome of an underlying stochastic evolutionary process. even if we could take the set of sampled populations back to a previous point in time and re-run the evolutionary process under all of the same conditions (the same population sizes, mutation rates, migration rates and selection coefficients), the genotype frequencies in the new set of populations would differ from those in the populations that were actually sampled¹⁸. This genetic sampling¹⁷ is an unavoidable consequence of genetic drift. The magnitude of variation associated with genetic sampling cannot be reduced by increasing either the number of individuals sampled within populations or the number of populations

sampled. Indeed, the characteristics of genetic sampling are shown by estimates of F -statistics.

In simple cases, it might make sense to estimate statistical parameters using simple functions of the data, such as the sample mean. In more complicated cases, such as those presented by F -statistics, it is useful to have well-defined approaches for constructing estimates. Statisticians have developed several different approaches for estimating parameters from data¹⁹. Three widely used approaches are the method of moments, the method of maximum likelihood and Bayesian methods.

Approaches to estimating F_{ST} : method-of-moments estimates

The method of moments produces an estimate by finding an algebraic expression that makes the expected value of certain sample statistics equal to simple functions of the parameters that are being estimated (as explained in more detail below)¹⁹. Method-of-moments estimates are designed to have low bias in the sense that if samples are taken repeatedly from the same population, the average of the corresponding sample estimates will be close to the unknown population parameter. These estimates have the additional advantages that they are easy to calculate and do not require any assumptions about the shape of the distribution from which the sample is drawn, other than that it has a mean and variance.

For F -statistics, method-of-moments estimates^{17,20,21} are based on an analysis of variance (ANOVA) of allele frequencies. ANOVA is a statistical method that tests whether the means of two or more groups are equal and can therefore be used to assess the degree of differentiation between populations. Briefly, if the variance among populations is the same as the variance within populations, there is no population substructure. ANOVA calculations are framed in terms of mean squares. Therefore, in practice, one calculates the expected mean square among populations (that is, the variance of sample allele frequencies around the mean allele frequency over all populations) and the expected mean square within populations (that is, the heterozygosity within populations when genotypes are in Hardy–Weinberg proportions) averaged over all possible samples (statistical sampling) from all possible populations with the same evolutionary history (genetic sampling). These expected values are then equated to the observed mean squares that are calculated from a sample, and the resulting set of equations is solved for the corresponding variance components. Following the work of Cockerham^{10,22}, F -statistics are defined in terms of these variance components (BOX 3).

Approaches to estimating F_{ST} : maximum-likelihood and Bayesian estimates

In contrast to method-of-moments estimates, likelihood and Bayesian estimates are difficult to calculate and require the specification of the probability distribution from which the sample was drawn. Once this probability distribution is specified, we can calculate a quantity called the likelihood, which is proportional to the probability of our observed data given those parameters. A maximum-likelihood estimate for the parameters is obtained by finding the values of the unknown parameters that maximize that likelihood¹⁹. In most cases, maximum-likelihood estimates are biased. Nonetheless, they typically have a smaller variance and deviate less from the unknown population parameter than the corresponding

method-of-moments estimates¹⁹. For these and other reasons, the method of maximum likelihood is the most widely used technique for deriving statistical estimators^{23,24}.

Bayesian estimates share many of the advantages associated with maximum-likelihood estimates because they use the same likelihood to relate the data to unknown parameters. However, they differ from maximum-likelihood estimates because the likelihood is modified by placing prior distributions on unknown parameters, and estimates are based on the posterior distribution, which is proportional to the product of the likelihood and the prior distributions. Both maximum-likelihood and Bayesian methods suffer the disadvantage that simple algebraic expressions for the estimates are rarely available. Instead, the estimates are obtained through computational methods. Because the Markov chain Monte Carlo methods (MCMC methods) used for analysis of Bayesian models do not require a unique point of maximum likelihood to be identified, Bayesian estimates can be obtained even in complex models with thousands or tens of thousands of parameters, for which numerical maximization of the likelihood would be difficult or impossible²⁶.

For F -statistics, the likelihood approach^{27,28} specifies a probability distribution that describes the variation in allele frequencies among populations and a multinomial distribution that describes genotype samples within populations. θ is related to the variance of the probability distribution that describes the among-population distribution of allele frequencies, and the genotype frequencies are determined by the allele frequencies in each population and f . Estimates are obtained by maximizing the likelihood function with respect to θ , f and the allele frequencies. The Bayesian approach uses the same likelihood function, and after placing appropriate prior distributions on f , θ and allele frequencies, MCMC methods are used to sample from the posterior distributions of f and θ .

Comparing the methods

With more than 5,000 citations, the moments method described by Weir and Cockerham²⁰ has been widely used, partly because of its robustness and partly because it is simple to implement. The maximum-likelihood methods also give simple equations when the distribution of allele frequencies among populations is assumed to be normal²⁷, but only if the sample sizes are equal²⁹. Bayesian methods allow probability statements to be made about F -statistics, and extensions of these methods allow the relationship between F -statistics and demographic or environmental covariates to be explored in the context of a single model³⁰. However, implementations of Bayesian methods may be computationally demanding.

A simple data set is used in BOX 3 to illustrate the slightly different estimates obtained from each approach. Estimates of F_{ST} using moments and Bayesian methods have not been extensively compared, but our experience suggests that the differences in estimates are small when the average number of individuals per population is moderate to large (>20), when the number of populations is moderate to large ($>10-15$) and when most populations are polymorphic. When differences arise, they reflect differences in the treatment of allele frequency estimates when alleles are rare or sample sizes are small. The Bayesian approach 'smooths' population allele frequencies towards the mean²⁴ and does so more aggressively

when alleles are rare or sample sizes are small. The moments approach treats the sample frequencies as fixed quantities without such smoothing. The simulation results in REF. 31 are consistent with this interpretation, although they compare Bayesian estimates with estimates of G_{ST} ³², which does not account for genetic sampling.

Related statistics

Population geneticists have proposed several statistical measures that are related to F_{ST} . Here, we describe four of them: G_{ST} , R_{ST} , Φ_{ST} and Q_{ST} . Nei³³ introduced G_{ST} as a measure of population differentiation. We discuss its relationship to F_{ST} in BOX 2. Haplotype and microsatellite data contain information not only about the frequency with which particular alleles occur but also on the evolutionary distance between them. Statistics such as Φ_{ST} (for haplotype data) and R_{ST} (for microsatellite data) are intended to take advantage of this additional information and to provide greater insight into the patterns of relationships among populations. Whereas F_{ST} , Φ_{ST} and R_{ST} all apply to discrete genetic data, Q_{ST} is an analogous statistic for continuously varying traits. If the markers used to estimate F_{ST} can be presumed to be selectively neutral, comparing an estimate of Q_{ST} with an estimate of F_{ST} can provide investigators with evidence that natural selection has shaped the pattern of variation in the quantitative trait.

R_{ST} , Φ_{ST} and AMOVA

The methods for estimating f , θ and F described above are appropriate for multi-allelic data when the alleles are regarded as equivalent to one another. However, when the data consist of variation at microsatellite loci or of nucleotide sequence (haplotype) information, related methods that allow mutation rates to differ between different pairs of alleles might be more appropriate. Excoffier *et al.*³⁴ introduced analysis of molecular variance (AMOVA) for analysis of haplotype variation. AMOVA is based on an analysis-of-variance framework that is analogous to the one developed by Weir and Cockerham²⁰. The mean squares in an AMOVA analysis are based on a user-specified measure of the evolutionary distance between haplotypes, and AMOVA leads to quantities that are analogous to classical F -statistics (BOX 1). Similarly, the mean squares used to calculate R_{ST} ^{35,36} are based on differences in the number of repeats between alleles at each microsatellite locus. Although the result of both analyses is a partitioning of genetic variance into within- and among-population components analogous to F_{ST} , neither has a direct interpretation as a parameter of a statistical distribution. Instead, they estimate an index that is derived from two different statistical distributions: the distribution of allele (haplotype or microsatellite) frequencies among populations and the distribution of evolutionary distances among alleles.

Nonetheless, such measures may be thought of as estimating the additional time since the common ancestry of randomly chosen alleles that accrues as a result of populations being subdivided^{37,38}, provided that the measure of evolutionary distance between any two alleles is proportional to the time since their most recent common ancestor. Extensive simulation studies have shown that estimates of R_{ST} may be unreliable unless many loci are used^{39–41}, but unlike F_{ST} the expected value of R_{ST} does not depend on the rate of mutation. Estimates of Φ_{ST} or R_{ST} may be useful when mutations have contributed substantially to allelic differences among populations, but their usefulness may be limited by the extent to which

the mutational model underlying the statistics matches the actual mutational processes occurring in the system³⁹.

Q_{ST} and polygenic variation

Spitze⁴² noted that another quantity analogous to θ can be estimated for continuously varying traits. Specifically, we can define:

$$Q_{ST} = \frac{\sigma_{GP}^2}{\sigma_{GP}^2 + 2\sigma_{GI}^2} \quad (6)$$

in which σ_{GP}^2 is the additive genetic variance among populations and σ_{GI}^2 is the additive genetic variance within populations. σ_{GP}^2 can be estimated from between-population crosses, and σ_{GI}^2 can be estimated from within-population crosses. Because the total variance in between-population crosses is $\sigma_{GP}^2 + \sigma_{GI}^2$, Q_{ST} is the proportion of additive genetic variance in a trait that is due to among-population differences. If the trait is selectively neutral, if all genetic variation is additive and if the mutation rates at loci contributing to the trait are the same as those at other loci, we expect Q_{ST} and F_{ST} to be equal^{43,44}. Comparing the magnitude of Q_{ST} and F_{ST} may therefore indicate whether a particular trait has been subject to stabilizing selection ($Q_{ST} < F_{ST}$) or diversifying selection ($Q_{ST} > F_{ST}$). However, because of the uncertainties associated with estimates of Q_{ST} and F_{ST} , such comparisons are likely to be useful only when they are available for a moderately large number of populations (>20)⁴⁵. Furthermore, caution is necessary when suggesting that a comparison of Q_{ST} and F_{ST} provides evidence for stabilizing selection because non-additive genetic variation tends to change Q_{ST} , even for a neutral trait⁴⁶.

Applications

F -statistics include both F_{ST} , which measures the amount of genetic differentiation among populations (and simultaneously the extent to which individuals within populations are similar to one another), and F_{IS} , which measures the departure of genotype frequencies within populations from Hardy–Weinberg proportions. Here, we focus on the applications of F_{ST} for several reasons (BOX 4).

Estimating migration rates

Wright⁵ showed that if all populations in a species are equally likely to exchange migrants and if migration is rare, then:

$$F_{ST} \approx \frac{1}{4N_e m + 1} \quad (7)$$

in which m is the fraction of each population composed of migrants (the backward migration rate)⁴⁷ and N_e is the effective population size of local populations⁴⁸. Because of this simple relationship, it is tempting to use estimates of F_{ST} from population data to estimate $N_e m$.

Unfortunately, it has been recognized for many years that this simple approach to estimating migration rates might fail⁴⁹. The most obvious reason for this failure is that populations are rarely structured so that all populations exchange migrants at the same rate, which causes some populations to resemble one another more than others. If differentiation between populations is solely a result of isolation by distance⁵⁰, for example, then the slope of the regression of $F_{ST}/(1 - F_{ST})$ on either the logarithm of between-population distance (for populations distributed in two dimensions) or the between-population distance alone (for populations in a linear habitat) is proportional to $D_e\delta^2$, in which D_e is the effective density of the population ($D_e = N_e/\text{area}$) and δ^2 is the mean squared dispersal distance⁵¹. However, if differentiation is the result not only of isolation by distance but also of natural selection or if the drift–migration process has not reached a stationary point, the slope of this relationship cannot be interpreted as an estimate of migration. Moreover, a pure migration–drift process, a pure drift–divergence process or a combination of the two could produce the same distribution of allele frequencies. Indeed, migration–drift, drift–divergence or a combination of the two can account for any pattern of allele frequency differences among populations⁵². Therefore, although pairwise estimates of F_{ST} (or Φ_{ST} or R_{ST}) provide some insight into the degree to which populations are historically connected^{37,38}, they do not allow us to determine whether that connection is a result of ongoing migration or of recent common ancestry.

There are additional difficulties with interpreting estimates of F_{ST} . Different genetic markers may give different estimates of F_{ST} for many reasons, and to derive an estimate of migration rates from F_{ST} , one must assume that the particular set of markers that are chosen have the expected relationship with $N_e m$. This may often be problematic. For example, differences between F_{ST} estimates from human microsatellites (0.05) and SNPs (0.10) cannot reflect differences in migration rate because both estimates are derived from the same set of individuals and the same set of populations — the Human Genome Diversity Project–Centre d’Étude du Polymorphisme Humain sample^{1,2,53}. The use of coalescent-based approaches (see later section) that incorporate models of the mutational process is one method of overcoming this difficulty^{54–56}.

Inferring demographic history

Population-specific or pairwise estimates of F_{ST} may provide insights into the demographic history of populations when estimates are available from many loci. For example, Keinan *et al.*⁹ reported pairwise estimates of F_{ST} for 13,600–62,830 autosomal SNP loci and 1,100–2,700 X chromosome SNP loci in human population samples from northern Europe, East Asia and West Africa. Because there are four copies of each autosome in the human population for every three copies of the X chromosome, one would expect there to be greater differentiation at X chromosome loci than at autosomal loci. Specifically, for two populations that diverged t generations ago, one might expect:

$$1 - F_{ST} = \left(1 - \frac{1}{2N_e}\right)^t \quad (8)$$

in which N_e is the effective size of the local populations. Therefore, if Q is defined as:

$$\ln(1 - F_{ST}^{\text{auto}})/\ln(1 - F_{ST}^X) \quad (9)$$

Q is approximately:

$$N_e^X/N_e^{\text{auto}}=0.75 \quad (10)$$

Q is approximately 0.75 for comparisons between East Asians and northern Europeans ($Q = 0.72 \pm 0.05$), but it is substantially smaller for comparisons between West Africans and other populations in the sample ($Q = 0.58 \pm 0.03$ for the comparison with northern Europeans and $Q = 0.62 \pm 0.03$ for the comparison with East Asians). These results suggest either sex-biased dispersal (long-range immigration of males from Africa after non-African populations were initially established) or selection on X chromosome loci after the divergence of African and non-African populations.

Identifying genomic regions under selection

Similarly, locus-specific estimates of F_{ST} may identify genomic regions that have been subject to selection. The logic is straightforward; the pattern of genetic differentiation at a neutral locus is completely determined by the demographic history of the populations (that is, the history of population expansions and contractions), the mutation rates at the loci concerned and the rates and patterns of migration among the populations^{6,57–60}. In a typical multilocus sample, it is reasonable to assume that all autosomal loci have experienced the same demographic history and the same rates and patterns of migration. If the loci also have similar mutation rates and if the variation at each locus is selectively neutral, the allelic variation at each locus represents a separate sample from the same underlying stochastic evolutionary process. loci showing unusually large amounts of differentiation may indicate regions of the genome that have been subject to diversifying selection, whereas loci showing unusually small amounts of differentiation may indicate regions of the genome that have been subject to stabilizing selection⁵⁸. Several groups have used such genome scans to examine patterns of differentiation in the human genome.

By comparing locus-specific estimates of F_{ST} with the genome-wide distribution, Akey *et al.*⁶ identified 174 regions (out of the 26,530 examined) that showed what they called ‘signatures of selection’ in the human genome. of these loci, 156 showed unusually large amounts of differentiation (suggesting diversifying selection) and 18 showed unusually small amounts of differentiation (suggesting stabilizing selection). By contrast, when Weir *et al.*⁷ examined the high-resolution Perlegen (~1 million SNPs) and phase I HapMap (~0.6 million SNPs) data sets in humans to examine locus-specific estimates of F_{ST} , they also found large differences in F_{ST} among loci, but their analyses suggested that the very high variance associated with single-locus estimates of F_{ST} precluded using these estimates to detect selection. Both sets of investigators noted a particular problem with single-locus estimates when using high-resolution SNP maps: the high correlation between F_{ST} estimates when loci are in strong gametic disequilibrium makes it difficult to determine whether the F_{ST} at any particular SNP is markedly different from expectation.

Although single-locus estimates of F_{ST} are highly uncertain, simulation studies suggest that when loci are inherited independently, background information about a few hundred loci is sufficient to allow the reliable identification of loci that are subject to selection when a suitable criterion for detecting ‘outliers’ is used^{8,58,61}. Although few loci are falsely identified as being subject to selection when they are neutral, genome scans using F_{ST} may often fail to detect selection when it is present. For example, when a single allele is strongly favoured in all populations, not only is F_{ST} expected to be nearly zero but variation is also expected to be nearly non-existent, rendering estimates of F_{ST} either highly unreliable or unobtainable. Similarly, when selection is weak, data from many loci are needed to recognize that the estimate of F_{ST} at the locus involved is unusual. More importantly, as mentioned above, high-resolution genome scans must account for the statistical association between closely linked loci. Guo *et al.*⁸ used a conditional autoregressive scheme to identify 57 loci that showed unusually large amounts of among-population differentiation in a sample of 3,000 SNP loci on human chromosome 7 separated by only 860 nucleotides on average. Sixteen of these markers are associated with *LEP*, a gene encoding a leptin precursor that is associated with behaviours that influence the balance between food intake and energy expenditure⁶² (FIG. 1). Moreover, association studies in one French population had previously suggested a relationship between one of the SNPs identified as an outlier in this study and obesity⁶³.

Forensic science and association mapping

In forensic science, matching a genetic profile taken from a suspect with a profile taken from a stain left at a crime scene serves as evidence linking the suspect to the crime. To quantify the strength of this evidence, it is useful to determine the probability of a random match — that is, the probability that the genetic profile at the crime scene matches that of the suspect if the suspect was not the source of the stain. In some cases, two people, the suspect and the person who left the crime sample, may belong to a subpopulation for which there is no specific allele frequency information. In such a case, we can use a θ correction⁶⁴ to calculate the probability of a match based on allele frequency information from a larger population of which the subpopulation is a part. The probability of a random match takes into account the allele frequency variation among subpopulations within the wider population for which allele frequencies are available. For example, if the matching profile consisted of a homozygote AA at a single locus and if p_A is the population frequency of allele A, the probability that the crime profile is AA given that the suspect is AA and the suspect is not the source of the stain is (REF. 65):

$$P(AA|AA) = \frac{(3\theta + (1 - \theta)p_A)(2\theta + (1 - \theta)p_A)}{(1 + \theta)(1 + 2\theta)} \quad (11)$$

There is a similar equation for heterozygotes, and these θ -correction results are multiplied over loci. The 1996 National research Council report⁶⁶ recommended using $\theta = 0.01$ except for small isolated subpopulations, for which they suggested that a value of $\theta = 0.03$ was more appropriate. The practical effect of the θ correction is that the numerical strength of the evidence against a suspect is reduced. If $p_A = 0.01$, for example, the uncorrected probability of a match is 0.0001. However, with $\theta = 0.01$, the probability of a match is an order of

magnitude larger — 0.0012. With $\theta = 0.03$, it is even larger — 0.0064. Therefore, it is much less surprising to see a match when we take account of the population substructure than when we ignore it.

In association mapping, case–control studies compare the allele frequencies at genetic markers (generally SNPs) between groups of people with a disease and groups who do not have the disease. When frequencies at a marker locus differ between the groups, it is interpreted as evidence for gametic disequilibrium between the marker and a disease-related gene. This in turn suggests that the marker and disease-related genes are in close proximity on the same chromosome. However, as many authors have pointed out, population substructure unrelated to disease status could cause the same kind of allele frequency difference^{67–70}. The genomic control method is one way to account for population substructure. It uses background estimates of F_{ST} to control for subpopulation differences that are unrelated to disease status^{67,68}. If cases and controls have different marker allele frequencies for reasons unconnected with the disease, as would be shown by frequency differences across the whole genome, an uncorrected case–control test would give spurious indications of marker–disease associations.

Relationship to coalescent-based methods

When Kingman introduced the coalescent process to population genetics just over 25 years ago^{71,72}, it revolutionized the field. Many approaches to the analysis of molecular data, particularly molecular sequence and SNP data, now take advantage of the conceptual, computational and analytical framework that coalescent-based methods provide^{73–79}. For example, whereas F -statistics provide only limited insight into the rates and patterns of migration, statistics based on the coalescent process can provide insights into the rates of mutation, migration and other evolutionary processes. Coalescent analysis is based on maximizing the likelihood of a given sample configuration or sampling from the corresponding Bayesian posterior distribution. The likelihood is constructed from the genealogical histories for the sample that are consistent with the unknown evolutionary parameters of interest; for example, the size of the population or populations from which the sample was taken, or the history of population size changes, mutation rates, recombination rates or migration rates^{55,80–86}. Coalescent analyses are likely to provide precise estimates of effective population size, mutation rates and migration rates when certain conditions are met — that is, when the model used for analysis is consistent with the demographic history of populations from which samples are collected, with the migration patterns among populations in the sample and with the mutational processes that generated allelic differences in the sample, and also when it is reasonable to presume that the drift–mutation–migration process has reached an evolutionary equilibrium^{54,73}. When these assumptions are not met it may not be reasonable to estimate the related evolutionary parameters, and the examples presented above show that analyses based on F -statistics may still provide substantial insights.

Conclusions

Sewall Wright⁵ provided a comprehensive account of the processes leading to genetic differentiation among populations nearly 80 years ago, but he did not provide the tools that empirical population geneticists needed to apply his insights to understanding variation in wild populations. During his work on isolation by distance in the plant *Linanthus parryae* in the 1940s^{50,87}, the theory of F -statistics that he and Gustave Malécot later developed^{3,4,16,88} began to emerge. Because of the insights that F -statistics can provide about the processes of differentiation among populations, over the past 50 years they have become the most widely used descriptive statistics in population and evolutionary genetics. From the time population geneticists first began to collect data on allozyme variation^{89–94} to recent analyses of SNP variation in the human genome^{2,9,95–97}, F -statistics, and F_{ST} in particular, have been used to investigate processes that influence the distribution of genetic variation within and among populations. Unfortunately, neither Wright nor Malécot distinguished carefully between the definition of F -statistics and the estimation of F -statistics. In particular, until Cockerham introduced his indicator formalism^{10,22}, few if any population geneticists understood that estimators of F -statistics must take into account both statistical sampling and genetic sampling.

The statistical methodology for estimating F -statistics is now well established. With the availability of methods to estimate locus- and population-specific effects on F_{ST} ^{7,8,27,58,61,98}, geneticists now have a set of tools for identifying genomic regions or populations with unusual evolutionary histories. Through further extensions of this approach, it is even possible to determine the relationship between the recent evolutionary history of populations and environmental or demographic variables⁹⁹. The basic principles of how population size, mutation rate and migration are related to the genetic structures of populations have been well understood for nearly 80 years. Analyses of F -statistics in populations of plants, animals and microorganisms have broadened and deepened this understanding, but these analyses have mostly been applied to data sets that contain a small number of loci. The age of population genomics is now upon us^{100,101}. The 1,000 Genomes project and the International HapMap Project give a hint of what is to come. Despite the scale of these projects, much of the data can be understood fundamentally as allelic variation at individual loci. As a result, we expect F -statistics to be at least as useful in understanding these massive data sets as they have been in population and evolutionary genetics for most of the past century.

Acknowledgements

We thank R. Prunier and K. Theiss for their helpful comments on earlier versions of this Review. The work in the laboratories of the authors was supported in part by grants from the US National Institutes of Health (1 R01 GM 068449-01A1 to K.E.H; 1 R01 GM 075091 to B.S.W).

Glossary

Genetic drift The random fluctuations in allele frequencies over time that are due to chance alone.

Short tandem repeat loci	Loci consisting of short sequences (2–6 nucleotides) that are repeated multiple times. Alleles at short tandem repeat loci differ from one another in their number of repeats.
Variance	A measure of the amount of variation around a mean value.
Diversifying selection	Selection in which different alleles are favoured in different populations. It is often a consequence of local adaptation (in which genotypes from different populations have higher fitness in their home environments owing to historical natural selection).
Hardy–Weinberg proportions	When the frequency of each diploid genotype at a locus equals that expected from the random union of alleles. That is, the genotypes AA, Aa and aa will be at frequencies p^2 , $2pq$ and q^2 , respectively.
Heterozygote advantage	A pattern of natural selection in which heterozygotes are more likely to survive than homozygotes.
Likelihood	A mathematical function that describes the relationship between the unknown parameters of a statistical distribution — for example, the mean and variance of the allele frequency distribution among populations or the allele frequency in a particular population — and the data. It is directly proportional to the probability of the data given the unknown parameters.
Prior distribution	A statistical distribution used in Bayesian analysis to describe the probability that parameters take on a particular value before examining any data. It expresses the level of uncertainty about those parameters before the data have been analysed.
Posterior distribution	A statistical distribution used in Bayesian analysis to describe the probability that parameters take a particular value after the data have been analysed. It reflects both the likelihood of the data given particular parameters and the prior probability that parameters take particular values.
Markov chain Monte Carlo methods	Methods that implement a computational technique that is widely used for approximating complex integrals and other functions. In this context, these methods are used to approximate the posterior distribution of a Bayesian model.
Multinomial distribution	A statistical distribution that describes the probability of obtaining a sample with a specified number of objects in each of several categories. The probability is determined by the total sample size and the probability of drawing an object from each category. The binomial distribution is a special case of the multinomial distribution in which there are two categories.
Additive genetic variance	The part of the total genetic variation that is due to the main (or additive) effects of alleles on a phenotype. The additive variance

	determines the degree of resemblance between relatives and therefore the response to selection.
Stabilizing selection	Selection in which either the same allele or the same genotype is favoured in different populations.
Effective population size	Formulated by Wright in 1931, the effective population size reflects the size of an idealized population that would experience drift in the same way as the actual (census) population. The effective population size can be lower than the census population size owing to various factors, including a history of population bottlenecks and reduced recombination.
Coalescent-based approaches	Approaches that use statistical properties of the genealogical relationship among alleles under particular demographic and mutational models to make inferences about the effective size of populations and about rates of mutation and migration.
Conditional autoregressive scheme	A statistical approach developed for analysis of data in which a random effect is associated with the spatial location of each observation. The magnitude of the random effect is determined by a weighted average of the random effects of nearby positions. In most applications, the weights of the averages are inversely related to the spatial distance between two sample points.

References

1. Rosenberg NA, et al. Genetic structure of human populations. *Science*. 2002; 298:2381–2385. [PubMed: 12493913]
2. Li JZ, et al. Worldwide human relationships inferred from genome-wide patterns of variation. *Science*. 2008; 319:1100–1104. [PubMed: 18292342]
3. Wright S. The genetical structure of populations. *Ann. Eugen.* 1951; 15:323–354. [PubMed: 24540312] This paper develops the explicit framework for the analysis and interpretation of *F*-statistics in an evolutionary context.
4. Malecot G. *Les Mathématiques de l'Hérédité*. 1948 Masson, Paris This book develops a framework — equivalent to Wright's *F*-statistics — for the analysis of genetic diversity in hierarchically structured populations.
5. Wright S. Evolution in Mendelian populations. *Genetics*. 1931; 16:97–159. [PubMed: 17246615] A landmark paper in population genetics in which the effect of population size, mutation and migration on the abundance and distribution of genetic variation in populations is first quantitatively described.
6. Akey JM, Zhang G, Khang K, Jin L, Shriver MD. Interrogating a high-density SNP map for signatures of natural selection. *Genome Res*. 2002; 12:1805–1814. [PubMed: 12466284]
7. Weir BS, Cardon LR, Anderson AD, Nielsen DM, Hill WG. Measures of human population structure show heterogeneity among genomic regions. *Genome Res*. 2005; 15:1468–1476. [PubMed: 16251456]
8. Guo F, Dey DK, Holsinger KE. A Bayesian hierarchical model for analysis of SNP diversity in multilocus, multipopulation models. *J. Am. Stat. Assoc.* 2009; 164:142–154. [PubMed: 19623271]
9. Keinan A, Mullikin JC, Patterson N, Reich D. Accelerated genetic drift on chromosome X during the human dispersal out of Africa. *Nature Genet.* 2009; 41:66–70. [PubMed: 19098910]

10. Cockerham CC. Variance of gene frequencies. *Evolution*. 1969; 23:72–84. This paper develops the first approach for the analysis of F -statistics that recognizes the effect of genetic sampling on estimates of F -statistics from population data.
11. Wahlund S. Zusammensetzung von Population und Korrelationserscheinung vom Standpunkt der Vererbungslehre aus betrachtet. *Hereditas*. 1928; 11:65–106.
12. Sokal RR, Oden NL, Thomson BA. A simulation study of microevolutionary inferences by spatial autocorrelation analysis. *Biol. J. Linn. Soc.* 1997; 60:73–93.
13. Sokal RR, Oden NL. Spatial autocorrelation analysis as an inferential tool in population genetics. *Am. Nat.* 1991; 138:518–521.
14. Epperson, BK. *Geographical Genetics*. Princeton Univ. Press; 2003.
15. Weir BS, Cockerham CC. Mixed self- and random-mating at two loci. *Genet. Res.* 1973; 21:247–262. [PubMed: 4731639]
16. Wright, S. *Evolution and the Genetics of Populations*. Vol. 4. Univ. Chicago Press; 1978.
17. Weir, BS. *Genetic Data Analysis II. Methods for Discrete Population Genetic Data*. Sunderland, USA: Sinauer Associates; 1996.
18. Rousset F. Inbreeding and relatedness coefficients: what do they measure? *Heredity*. 2002; 88:371–380. [PubMed: 11986874]
19. Casella, G.; Berger, RL. *Statistical Inference*. Duxbury, Pacific Grove: 2002.
20. Weir BS, Cockerham CC. Estimating F -statistics for the analysis of population structure. *Evolution*. 1984; 38:1358–1370. This paper develops the ANOVA framework to apply Cockerham's approach to F -statistics and provides method-of-moments estimates for F -statistics.
21. Excoffier, L. *Handbook of Statistical Genetics*. Balding, DJ.; Bishop, M.; Cannings, V., editors. Chichester: John Wiley & Sons; 2001. p. 271-307.
22. Cockerham CC. Analyses of gene frequencies. *Genetics*. 1973; 74:679–700. [PubMed: 17248636]
23. Berger, JO. *Statistical Decision Theory and Bayesian Analysis*. New York: Springer; 1985.
24. Robert, CP. *The Bayesian Choice: From Decision-Theoretic Foundations to Computational Implementation*. New York: Springer; 2001.
25. Lee, PM. *Bayesian Statistics: An Introduction*. London: Edward Arnold; 1989.
26. Gelfand AE, Smith AFM. Sampling-based approaches to calculating marginal densities. *J. Am. Stat. Assoc.* 1990; 85:398–409.
27. Weir BS, Hill WG. Estimating F -statistics. *Annu. Rev. Genet.* 2002; 36:721–750. [PubMed: 12359738]
28. Wehrhahn C. Proceedings of the ecological genetics workshop. *Genome*. 1989; 31:1098–1099.
29. Samanta S, Li YJ, Weir BS. Drawing inferences about the coancestry coefficient. *Theor. Popul. Biol.* 2009; 75:312–319. [PubMed: 19345237]
30. Gaggiotti OE, et al. Patterns of colonization in a metapopulation of grey seals. *Nature*. 2002; 13:424–427. [PubMed: 11919630]
31. Levens ND, Crawford DJ, Archibald JK, Santos-Geurra A, Mort ME. Nei's to Bayes': comparing computational methods and genetic markers to estimate patterns of genetic variation in *Tolpis* (Asteraceae). *Am. J. Bot.* 2008; 95:1466–1474. [PubMed: 21628154]
32. Nei M, Chesser RK. Estimation of fixation indices and gene diversities. *Ann. Hum. Genet.* 1983; 47:253–259. [PubMed: 6614868]
33. Nei M. Analysis of gene diversity in subdivided populations. *Proc. Natl Acad. Sci. USA.* 1973; 70:3321–3323. [PubMed: 4519626] This article introduces G_{ST} as a measure of genetic differentiation among populations.
34. Excoffier L, Smouse PE, Quattro JM. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics*. 1992; 131:479–491. [PubMed: 1644282] This paper introduces Φ_{ST} and AMOVA for the analysis of haplotype data.
35. Slatkin M. A measure of population subdivision based on microsatellite allele frequencies. *Genetics*. 1995; 139:457–462. [PubMed: 7705646] This article introduces R_{ST} for the analysis of microsatellite data.

36. Rousset F. Equilibrium values of measures of population subdivision for stepwise mutation processes. *Genetics*. 1996; 142:1357–1362. [PubMed: 8846911]
37. Slatkin M. Inbreeding coefficients and coalescence times. *Genet. Res.* 1991; 58:167–175. [PubMed: 1765264]
38. Holsinger KE, Mason-Gamer RJ. Hierarchical analysis of nucleotide diversity in geographically structured populations. *Genetics*. 1996; 142:629–639. [PubMed: 8852859]
39. Balloux F, Lugon-Molin N. The estimation of population differentiation with microsatellite markers. *Mol. Ecol.* 2002; 11:155–165. [PubMed: 11856418]
40. Balloux F, Brunner F, Goudet J. Microsatellites can be misleading: an empirical and simulation study. *Evolution*. 2000; 54:1414–1422. [PubMed: 11005307]
41. Gaggiotti OE, Lange O, Rassman K, Gliddon C. A comparison of two indirect methods for estimating average levels of gene flow using microsatellite data. *Mol. Ecol.* 1999; 8:1513–1520. [PubMed: 10564457]
42. Spitze K. Population structure in *Daphnia obtusa*: quantitative genetic and allozymic variation. *Genetics*. 1993; 135:467–374. This paper introduces Q_{ST} for the analysis of continuously varying trait data.
43. Lande R. Neutral theory of quantitative genetic variance in an island model with local extinction and colonization. *Evolution*. 1992; 46:381–389.
44. McKay JK, Latta RG. Adaptive population divergence: markers, QTL and traits. *Trends Ecol. Evol.* 2002; 17:285–291.
45. O'Hara RB, Merila J. Bias and precision in Q_{ST} estimates: problems and some solutions. *Genetics*. 2005; 171:1331–1339. [PubMed: 16085700]
46. Goudet J, Martin G. Under neutrality $Q_{ST} = F_{ST}$ when there is dominance in an island model. *Genetics*. 2007; 176:1371–1374. [PubMed: 17339207]
47. Notohara M. The coalescent and the genealogical process in geographically structured population. *J. Math. Biol.* 1990; 29:59–75. [PubMed: 2277236]
48. Charlesworth B. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. *Nature Rev. Genet.* 2009; 10:195–205. [PubMed: 19204717]
49. McCauley DE, Whitlock MC. Indirect measures of gene flow and migration: $F_{ST} = 1/(4Nm + 1)$. *Heredity*. 1999; 82:117–125. [PubMed: 10098262]
50. Wright S. Isolation by distance. *Genetics*. 1943; 28:114–138. [PubMed: 17247074]
51. Rousset F. Genetic differentiation and estimation of gene flow from F -statistics under isolation by distance. *Genetics*. 1997; 145:1219–1228. [PubMed: 9093870]
52. Felsenstein J. How can we infer geography and history from gene frequencies? *J. Theor. Biol.* 1982; 96:9–20. [PubMed: 7109659]
53. Cann HM, et al. A human genome diversity cell line panel. *Science*. 2002; 296:261–262. [PubMed: 11954565]
54. Beerli P. Comparison of Bayesian and maximum-likelihood estimation of population genetic parameters. *Bioinformatics*. 2006; 22:341–345. [PubMed: 16317072]
55. Kuhner MK. Coalescent genealogy samplers: windows into population history. *Trends Ecol. Evol.* 2009; 24:86–93. [PubMed: 19101058]
56. Kuhner MK. LAMARC 2.0: maximum likelihood and Bayesian estimation of population parameters. *Bioinformatics*. 2006; 22:768–770. [PubMed: 16410317]
57. Fu R, Gelfand A, Holsinger KE. Exact moment calculations for genetic models with migration, mutation, and drift. *Theor. Popul. Biol.* 2003; 63:231–243. [PubMed: 12689794]
58. Beaumont MA, Balding DJ. Identifying adaptive genetic divergence among populations from genome scans. *Mol. Ecol.* 2004; 13:969–980. [PubMed: 15012769]
59. Vitalis R, Dawson K, Boursot P. Interpretation of variation across marker loci as evidence of selection. *Genetics*. 2001; 158:1811–1823. [PubMed: 11514464]
60. Beaumont MA, Nichols RA. Evaluating loci for use in the genetic analysis of population structure. *Proc. R. Soc. Lond. B.* 1996; 263:1619–1626.

61. Foll M, Gaggiotti O. A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: a Bayesian perspective. *Genetics*. 2008; 180:977–993. [PubMed: 18780740]
62. Zhang Y, et al. Positional cloning of the mouse obese gene and its human homologue. *Nature*. 1994; 372:425–432. [PubMed: 7984236]
63. Mammès O, et al. Association of the G2548A polymorphism in the 5' region of the *LEP* gene with overweight. *Ann. Hum. Genet.* 2000; 64:391–394. [PubMed: 11281277]
64. Balding DJ, Donnelly P. How convincing is DNA evidence? *Nature*. 1994; 368:285–286. [PubMed: 8127359]
65. Balding DJ, Nichols RA. DNA match probability calculation: how to allow for population stratification, relatedness, database selection, and single bands. *Forensic Sci. Int.* 1994; 64:125–140. [PubMed: 8175083]
66. Council, NR. *The Evaluation of Forensic DNA Evidence*. Washington DC: National Academy Press; 1996.
67. Devlin B, Roeder K, Wasserman L. Genomic control, a new approach to genetic-based association studies. *Theor. Popul. Biol.* 2001; 60:155–166. [PubMed: 11855950]
68. Devlin B, Roeder K. Genomic control for association studies. *Biometrics*. 1999; 55:997–1004. [PubMed: 11315092]
69. Pritchard JK, Donnelly P. Case-control studies of association in structured or admixed populations. *Theor. Popul. Biol.* 2001; 60:227–237. [PubMed: 11855957]
70. Pritchard JK, Rosenberg NA. Use of unlinked genetic markers to detect population stratification in association studies. *Am. J. Hum. Genet.* 1999; 65:220–228. [PubMed: 10364535]
71. Kingman JFC. On the genealogy of large populations. *J. Appl. Prob.* 1982; 19A:27–43.
72. Kingman JFC. The coalescent. *Stoch. Proc. Appl.* 1982; 13:235–248.
73. Kuhner MK, Smith LP. Comparing likelihood and Bayesian coalescent estimation of population parameters. *Genetics*. 2007; 175:155–165. [PubMed: 16510781]
74. Wang J. A coalescent-based estimator of admixture from DNA sequences. *Genetics*. 2006; 173:1679–1692. [PubMed: 16624918]
75. Innan H, Zhang K, Marjoram P, Tavaré S, Rosenberg NA. Statistical tests of the coalescent model based on the haplotype frequency distribution and the number of segregating sites. *Genetics*. 2005; 169:1763–1777. [PubMed: 15654103]
76. Wall JD, Hudson RR. Coalescent simulations and statistical tests of neutrality. *Mol. Biol. Evol.* 2001; 18:1134–1135. [PubMed: 11371601]
77. Nordborg M. Structured coalescent processes on different time scales. *Genetics*. 1997; 146:1501–1514. [PubMed: 9258691]
78. Donnelly P, Tavaré S. Coalescents and genealogical structure under neutrality. *Annu. Rev. Genet.* 1995; 29:401–421. [PubMed: 8825481]
79. Griffiths RC, Tavaré S. Simulating probability distributions in the coalescent. *Theor. Popul. Biol.* 1994; 46:131–159.
80. Fearnhead P, Donnelly P. Estimating recombination rates from population genetic data. *Genetics*. 2001; 159:1299–1318. [PubMed: 11729171]
81. Kuhner MK, Beerli P, Yamato J, Felsenstein J. Usefulness of single nucleotide polymorphism data for estimating population parameters. *Genetics*. 2000; 156:439–447. [PubMed: 10978306]
82. Kuhner MK, Yamato J, Felsenstein J. Maximum likelihood estimation of recombination rates from population data. *Genetics*. 2000; 156:1393–1401. [PubMed: 11063710]
83. Kuhner MK, Felsenstein J. Sampling among haplotype resolutions in a coalescent-based genealogy sampler. *Genet. Epidemiol.* 2000; 19(Suppl. 1):15–21.
84. Kuhner MK, Yamato J, Felsenstein J. Maximum likelihood estimation of population growth rates based on the coalescent. *Genetics*. 1998; 149:429–434. [PubMed: 9584114]
85. Beerli P, Felsenstein J. Maximum-likelihood estimation of migration rates and effective population numbers in two populations using a coalescent approach. *Genetics*. 1999; 152:763–773. [PubMed: 10353916]

86. Drummond AJ, Nicholls GK, Rodrigo AG, Solomon W. Estimating mutation parameters, population history and genealogy simultaneously from temporally spaced sequence data. *Genetics*. 2002; 161:1307–1320. [PubMed: 12136032]
87. Wright S. An analysis of local variability of flower color in *Linanthus parryae*. *Genetics*. 1943; 28:139–156. [PubMed: 17247075]
88. Malécot, G. *The Mathematics of Heredity*. San Francisco: W. H. Freeman; 1969.
89. Hamrick JL, Godt MJW. Effects of life history traits on genetic diversity in plant species. *Philos. Trans. R. Soc. Lond. B*. 1996; 351:1291–1298.
90. Hamrick, JL. *Isozymes in Plant Biology*. Soltis, DE.; Soltis, PS., editors. Portland: Dioscorides; 1989. p. 87-105.
91. Loveless MD, Hamrick JL. Ecological determinants of genetic structure in plant populations. *Annu. Rev. Ecol. Syst.* 1984; 15:65–95.
92. Hamrick JL, Linhart YB, Mitton JB. Relationships between life history characteristics and electrophoretically detectable genetic variation in plants. *Annu. Rev. Ecol. Syst.* 1979; 10:173–200.
93. Gottlieb, LD. *Progress in Phytochemistry*. Reinhold, L.; Harborne, JB.; Swain, T., editors. Vol. 7. Oxford: Pergamon; 1981. p. 1-46.
94. Brown AHD. Enzyme polymorphism in plant populations. *Theor. Popul. Biol.* 1979; 15:1–42.
95. International HapMap Consortium. A second generation human haplotype map of over 3.1 million SNPs. *Nature*. 2007; 449:851–861. [PubMed: 17943122]
96. International HapMap Consortium. A haplotype map of the human genome. *Nature*. 2005; 437:1299–1320. [PubMed: 16255080]
97. He M, et al. Geographical affinities of the HapMap samples. *PLoS ONE*. 2009; 4:e4684. [PubMed: 19259268]
98. Balding DJ. Likelihood-based inference for genetic correlation coefficients. *Theor. Popul. Biol.* 2003; 63:221–230. [PubMed: 12689793]
99. Foll M, Gaggiotti O. Identifying the environmental factors that determine the genetic structure of populations. *Genetics*. 2006; 174:875–891. [PubMed: 16951078]
100. Begun DJ, et al. Population genomics: whole-genome analysis of polymorphism and divergence in *Drosophila simulans*. *PLoS Biol.* 2007; 5:e310. [PubMed: 17988176]
101. Luikart G, England PR, Tallmon D, Jordan S, Taberlet P. The power and promise of population genomics: from genotyping to genome typing. *Nature Rev. Genet.* 2003; 4:981–994. [PubMed: 14631358]
102. Goudet J, Raymond M, de Meeus T, Rousset F. Testing differentiation in diploid populations. *Genetics*. 1996; 144:1933–1940. [PubMed: 8978076]
103. Workman PL, Niswander JD. Population studies on southwest Indian tribes. II. Local genetic differentiation in the Papago. *Am. J. Hum. Genet.* 1970; 22:24–49. [PubMed: 5411647]
104. Holsinger, KE. *Hierarchical Modeling for the Environmental Sciences*. Clark, JS.; Gelfand, AE., editors. Oxford Univ. Press; 2006. p. 25-37.
105. Holsinger KE. Analysis of genetic diversity in hierarchically structured populations: a Bayesian perspective. *Heredity*. 1999; 130:245–255.
106. Weir BS. The rarity of DNA profiles. *Ann. Appl. Stat.* 2007; 1:358–370. [PubMed: 19030117]
107. Ritland KR. Joint maximum-likelihood estimation of genetic and mating system structure using open-pollinated progenies. *Biometrics*. 1986; 42:25–43.
108. Thompson SL, Ritland K. A novel mating system analysis for modes of self-oriented mating applied to diploid and polyploid arctic Easter daisies (*Townsendia hookeri*). *Heredity*. 2006; 97:119–126. [PubMed: 16721390]

Box 1

Mathematical notation

In this box, we provide definitions for the mathematical symbols used throughout the Review.

Parameter	Definition
<i>Among-population allele frequency distribution</i>	
π	Mean allele frequency
σ_{π}^2	Variance in allele frequency
Wright's F -statistics and Cockerham's θ -statistics	
F_{IS}	Correlation of alleles within an individual relative to the subpopulation in which it occurs; equivalently, the average departure of genotype frequencies from Hardy–Weinberg expectations within populations
F_{ST}	Correlation of randomly chosen alleles within the same subpopulation relative to the entire population; equivalently, the proportion of genetic diversity due to allele frequency differences among populations
F_{IT}	Correlation of alleles within an individual relative to the entire population; equivalently, the departure of genotype frequencies from Hardy–Weinberg expectations relative to the entire population
f	Co-ancestry for alleles within an individual relative to the subpopulation in which it occurs; equivalent to F_{IS}
θ	Co-ancestry for randomly chosen alleles within the same subpopulation relative to the entire population; equivalent to F_{ST}
F	Co-ancestry for alleles within an individual relative to the entire population; equivalent to F_{IT}
Φ -statistics and R_{ST} *	
Φ_{IS}	Excess similarity of alleles within an individual relative to the subpopulation in which it occurs; analogous to F_{IS}
Φ_{ST}	Excess similarity among randomly chosen alleles within the same subpopulation relative to the entire population; equivalently, the proportion of genetic diversity (measured as the expected squared evolutionary distance between alleles) due to differences among populations; analogous to F_{ST}
Φ_{IT}	Excess similarity of alleles within an individual relative to the entire population; analogous to F_{IT}
R_{ST}	Excess similarity among randomly chosen alleles within the same subpopulation relative to the entire population; equivalently, the proportion of genetic diversity (measured as the expected squared difference in repeat numbers between alleles) due to differences among populations; analogous to F_{ST}
<i>Measuring genetic differentiation among populations in quantitative traits</i>	
σ_{GI}^2	Additive genetic variance within populations
σ_{GP}^2	Additive genetic variance among populations
Q_{ST}	Proportion of additive genetic variation in the entire population due to differences among populations; analogous to F_{ST}

* Φ_{ST} from analysis of molecular variance (AMOVA) is used for haplotype data (for example, nucleotide sequence data or mapped restriction site data) and requires a measure of evolutionary distance among all pairs of haplotypes. R_{ST} is used for microsatellite data and requires that alleles are labelled according to the number of repeat units that they contain.

Box 2**Genetic sampling versus statistical sampling**

Genetic drift leads to differences among populations that are described by the distribution of allele frequencies among those populations. The variance of this distribution is directly related to F_{ST} (see equation 2), but in a typical study only a subset of populations is sampled. Therefore, in addition to accounting for the variation associated with sampling from populations, estimates of F -statistics must account for the variation associated with sampling sets of populations from the allele frequency distribution.


Genetic (or evolutionary) sampling

Part **a** of the figure shows the distribution of allele frequencies among populations corresponding to a mean allele frequency of $\pi = 0.5$ and $F_{ST} = \theta = 0.1$. If two sets of populations (represented by dark and light circles) are sampled from this distribution, the allele frequencies in the first set of populations (light circles) will differ from those in the second set (dark circles). Part **b** provides an example in which two different sets of five population frequencies are drawn randomly from the distribution of allele frequencies shown in part **a**.

The variation in allele frequencies illustrated in part **a** reflects the effect of genetic or evolutionary sampling. The differences between the sets of samples in part **b** reflect the effect of sampling particular populations from the distribution of allele frequencies in part **a** and are analogous to the results that would be expected in an empirical study if it were repeated on a different set of populations.

Statistical sampling

Part **c** illustrates the more familiar idea of statistical sampling. It shows the distribution of sample allele frequencies obtained in 1,000 samples of 20 individuals from the population with the largest allele frequency in the population sample on the left in part **b**. Statistical sampling refers to the variation in sample composition that is expected when alleles are repeatedly sampled from a population with a particular allele frequency.



Investigators can control the amount of variation associated with statistical sampling by increasing the number of individuals sampled within populations: the larger the number of individuals sampled, the less that the sample allele frequencies will differ from the underlying population frequencies. By contrast, investigators cannot control the amount of variation associated with genetic sampling: the variation associated with genetic sampling is an intrinsic property of the underlying stochastic evolutionary process that contributes to the differentiation among populations.

The relationship between F_{ST} and G_{ST}

Nei introduced the statistic G_{ST} as a measure of genetic differentiation among populations³³. It is defined in terms of the population frequencies in part **b**, not the allele frequency distribution in part **a**. By contrast, estimates of F_{ST} account for genetic sampling and they are intended to reflect the properties of the allele frequency distribution in part **a**. As a result, F_{ST} and G_{ST} measure different properties. Therefore, G_{ST} will be an appropriate measure only when interest focuses on characteristics of the particular samples illustrated in part **b**. In a typical population study, θ will be a more appropriate measure of differentiation.

It might seem that similar arguments should apply to exact tests of population differentiation¹⁰² because they also use permutations of sample configurations to determine whether populations are differentiated from one another. However, the permutation test is equivalent to determining whether the allele frequency distribution in part **a** has a variance greater than zero, so exact tests implicitly consider both statistical and genetic sampling effects.

Box 3**Comparing methods for estimating F_{ST}**

To illustrate the differences among calculating method-of-moments, maximum-likelihood and Bayesian estimates of F -statistics, we use data from a classic study on human populations that investigated the allele frequency differences at blood group loci (see the table). We use a subset of the data that were originally reported by Workman and Niswander¹⁰³. Their data consist of genotype counts at several loci in Native American Papago and were collected from ten political districts in south-western Arizona. Estimates of F_{IS} , F_{ST} and F_{IT} derived from the MN blood group locus suggest that there is little departure of the genotype frequencies from Hardy–Weinberg expectations within each district and little genetic differentiation among the districts.

Method-of-moments analysis

Analysis of variance on the indicator variable $y_{ij,k}$, in which $y_{ij,k} = 1$ if allele i in individual j of population k is M , gives moment estimates for the variance components of $\sigma_G^2 = 0.16$, $\sigma_I^2 = 0.00511$, and $\sigma_P^2 = 0.0000667$, in which G stands for genotypes (alleles within individuals), I stands for individuals (individuals within populations) and P stands for populations (among populations). Following Cockerham¹⁰:

$$F = \frac{\sigma_P^2 + \sigma_I^2}{\sigma_P^2 + \sigma_I^2 + \sigma_G^2}$$

$$\theta = \frac{\sigma_P^2}{\sigma_P^2 + \sigma_I^2 + \sigma_G^2}$$

$$f = \frac{\sigma_I^2}{\sigma_I^2 + \sigma_G^2}$$

Therefore, the moment estimates are $F = 0.0348$, $\theta = 0.00402$ and $f = 0.0309$. As expected for human populations, there is little evidence that the genotype proportions within each political district differ from Hardy–Weinberg expectations ($f \approx 0$). Similarly, there is little evidence of genetic differentiation among political districts ($\theta \approx 0$).

Bayesian and likelihood analysis

By contrast, current implementations of a Bayesian approach to analysing these data typically assume independent uniform (0,1) prior distributions for both f and θ . The posterior mean of f and θ for these data are 0.0503 and 0.0189, respectively. The posterior distribution of f has a mode near 0 but is broad (with a 95% credible interval of 0.0033–0.123), which causes the posterior mean of f to be larger than the method-of-moments estimate. Similarly, the estimates of allele frequencies within each population

are uncertain and the estimate of θ takes this uncertainty into account, suggesting that there is slightly more among-population differentiation than detected with moment estimates. For comparison, the maximum-likelihood estimates are $F = 0.0408$, $\theta = 0.00640$ and $f = 0.0346$ (obtained by estimating the variance components in a Gaussian mixed model applied to the indicator variables and by using Cockerham's definitions of F , f and θ in terms of the variance components).

Parameter	Method of moments	Maximum likelihood	Bayesian
f	0.0309	0.0346	0.0503
θ	0.00402	0.00640	0.0189
F	0.0348	0.0408	0.0683

To extend the method-of-moments approach to multiple alleles and multiple loci, calculations are done separately for every allele at every locus and the sums of squares are combined^{17,27}. To extend the likelihood or Bayesian approaches, we make the assumption that f and θ have the same value at every locus and that genotype counts are sampled independently across loci and populations^{104,105}.

Box 4**Why focus on F_{ST} ?**

We focus here on F_{ST} for several reasons. First, F_{IS} is easier to interpret. It is defined with respect to the populations that are included in the sample, either through population-specific estimates or through the average of those estimates. By contrast, F_{ST} is defined and interpreted with respect to the distribution of allele frequencies among all populations that could have been sampled, not merely those that have been included in the sample. As a result, estimates of F_{ST} must account for genetic sampling, which introduces a level of complexity and subtlety that requires extra attention.

Second, the application of F -statistics to problems in population and evolutionary genetics often centres on estimates of F_{ST} . For example, when interpreting aspects of demographic history, such as sex-biased dispersal out of Africa in human populations⁹, detecting regions of the genome that might have been subject to stabilizing or diversifying selection^{8,58,61} or correcting the probabilities of obtaining a match in a forensic application for genetic substructure within populations¹⁰⁶, estimates of F_{ST} often play a crucial part in interpretations of genetic data. Estimates of F_{IS} reveal important properties of the mating system within populations, but estimates of F_{ST} reveal properties of the evolutionary processes that lead to divergence among populations.

Finally, in many populations of animals, and in human populations in particular, within-population departures from Hardy–Weinberg proportions are small. Where they are present, such departures may reveal more about genetic substructuring within populations than about departures from random mating. Moreover, although estimates of F_{IS} may provide insights into the patterns of mating in inbred populations of plants or animals, the direct analysis of mother–offspring genotype combinations is usually more informative and reliable^{107,108}.




Figure 1. Locus-specific estimates of F_{ST} on human chromosome 7

Estimates are as inferred from the phase II HapMap data set⁹⁵. Horizontal bars indicate the locations of known genes. The red circles are posterior means for SNPs with estimates that are detectably different from the genomic background (purple circles). All ‘outliers’ show significantly more differentiation among the four populations in the sample than is consistent with the level of differentiation seen in the genomic background. The excess differentiation suggests that these SNPs are associated with genomic regions in which loci have been subject to diversifying selection among populations. *CALU*, calumenin; *FSCN3*, ascasin homolog 3; *GCC1*, GRIP and coiled-coil domain containing 1; *GRM8*, glutamate receptor, metabotropic 8; *LEP*, leptin; *SND1*, staphylococcal nuclease and tudor domain containing 1. Figure is modified, with permission, from REF. 8 © (2009) American Statistical Association.

Parameter	Definition
<i>Among-population allele frequency distribution</i>	
π	Mean allele frequency
σ_{π}^2	Variance in allele frequency
Wright's F -statistics and Cockerham's θ -statistics	
F_{IS}	Correlation of alleles within an individual relative to the subpopulation in which it occurs; equivalently, the average departure of genotype frequencies from Hardy–Weinberg expectations within populations
F_{ST}	Correlation of randomly chosen alleles within the same subpopulation relative to the entire population; equivalently, the proportion of genetic diversity due to allele frequency differences among populations
F_{IT}	Correlation of alleles within an individual relative to the entire population; equivalently, the departure of genotype frequencies from Hardy–Weinberg expectations relative to the entire population
f	Co-ancestry for alleles within an individual relative to the subpopulation in which it occurs; equivalent to F_{IS}
θ	Co-ancestry for randomly chosen alleles within the same subpopulation relative to the entire population; equivalent to F_{ST}
F	Co-ancestry for alleles within an individual relative to the entire population; equivalent to F_{IT}
Φ -statistics and R_{ST} *	
Φ_{IS}	Excess similarity of alleles within an individual relative to the subpopulation in which it occurs; analogous to F_{IS}
Φ_{ST}	Excess similarity among randomly chosen alleles within the same subpopulation relative to the entire population; equivalently, the proportion of genetic diversity (measured as the expected squared evolutionary distance between alleles) due to differences among populations; analogous to F_{ST}
Φ_{IT}	Excess similarity of alleles within an individual relative to the entire population; analogous to F_{IT}
R_{ST}	Excess similarity among randomly chosen alleles within the same subpopulation relative to the entire population; equivalently, the proportion of genetic diversity (measured as the expected squared difference in repeat numbers between alleles) due to differences among populations; analogous to F_{ST}
<i>Measuring genetic differentiation among populations in quantitative traits</i>	
σ_{GI}^2	Additive genetic variance within populations
σ_{GP}^2	Additive genetic variance among populations
Q_{ST}	Proportion of additive genetic variation in the entire population due to differences among populations; analogous to F_{ST}

* Φ_{ST} from analysis of molecular variance (AMOVA) is used for haplotype data (for example, nucleotide sequence data or mapped restriction site data) and requires a measure of evolutionary distance among all pairs of haplotypes. R_{ST} is used for microsatellite data and requires that alleles are labelled according to the number of repeat units that they contain.

Parameter	Method of moments	Maximum likelihood	Bayesian
f	0.0309	0.0346	0.0503
θ	0.00402	0.00640	0.0189
F	0.0348	0.0408	0.0683

To extend the method-of-moments approach to multiple alleles and multiple loci, calculations are done separately for every allele at every locus and the sums of squares are combined^{17,27}. To extend the likelihood or Bayesian approaches, we make the assumption that f and θ have the same value at every locus and that genotype counts are sampled independently across loci and populations^{104,105}.