



HHS Public Access

Author manuscript

J Neurosci Nurs. Author manuscript; available in PMC 2017 February 01.

Published in final edited form as:

J Neurosci Nurs. 2016 February ; 48(1): 15–27. doi:10.1097/JNN.0000000000000176.

Pathophysiology and Treatment of Severe Traumatic Brain Injuries in Children

Kimberly A Allen, PhD, RN

Senior Research Fellow, University of Washington, Biobehavioral Nursing and Health Systems.

Work Address: 1959 NE Pacific St, Box 357266, Seattle, WA 98195. Home Address: 523

Broadway E, Apt 549, Seattle, WA 98102, Phone: 706-951-3432

Kimberly A Allen: Kimberly.allen2@gmail.com

Traumatic brain injuries in children are a major cause of morbidity and mortality worldwide (Bramlett & Dietrich, 2004; CDC, 2010; Feigin et al., 2013; Shao et al., 2012). Traumatic brain injuries are an insult to the brain due to direct trauma from an external force, a penetrating object, blast waves, or a jolt to the head (Faul, Xu, Wald, & Coronado, 2010; Klimo, Ragel, Scott, & McCafferty, 2010). In the United States, motor vehicle collisions and falls are the most frequent precipitating events resulting in over 500,000 pediatric traumatic brain injuries (TBI) that lead to hospitalization and deaths each year (Faul et al., 2010). Severe TBIs account for 15,000 admissions annually and a mortality rate of 24% in children in the United States (Piatt & Neff, 2012). Severe pediatric traumatic brain injuries are defined as children less than 18 years of age with a Glasgow Coma Scale score of 3 to 8 (Adelson et al., 2003). An appreciation of the pathophysiologic effects of this injury allow nurses to understand the role of key therapeutic management principles. The purpose of this paper is to explore pathophysiologic events, examine monitoring techniques, and explain current treatment modalities and management related to caring for children with severe TBI.

Pathophysiology

By understanding the pathophysiology associated with a TBI, pediatric nurses may be better able to manage and prevent injury progression. Injury progression may be subdivided into primary and secondary injuries, which are associated with major pathways of dysfunction in TBI.

Primary Injury

Primary injury comes about as a result of linear and rotational forces to brain tissue at the time of impact. Linear forces from acceleration trauma that transverse the skull result in coup and contracoup contusions (Hirsch & Kaufman, 1975). Coup contusions occur when the brain impacts the side of the skull; whereas, contracoup contusions occur when the brain hits the side of the skull and then bounces back to the other side of the skull (Stewart, 1944). The two most common types of hemorrhages following TBI in children are epidural and subdural (Case, 2008; Jamous et al., 2009). Epidural hemorrhages are often associated with skull fractures because the skull fragments cause laceration of an artery leading to accumulation of blood between the skull and the dura (Black, 1956). Subdural hemorrhages are a result from the rupture of one or more of the bridging veins within the space between

the dura mater and arachnoid membrane. The clinical presentation evolves over days to weeks both in the physical skills and electroencephalographic changes (Cohn, 1948).

Rotational forces are a result of the brain moving at a different angular velocity than the skull because the head is not held in a fixed position during the injury (Holbourn & Edin, 1943). Rotational forces can lead to shearing and twisting injury in critical areas of the brain and are often manifested clinically as concussions and diffuse axonal injury (DAI) (Blumbergs et al., 1994; Mihalik et al., 2010; Moen et al., 2012). Concussions are a traumatic injury to the head that can result in a loss of consciousness and traumatic amnesia (Ward, 1964). However, diagnostic imaging does not show any damage to the brain despite symptom continuation for several weeks post-injury (Eisenberg, Meehan, & Mannix, 2014). Diffuse axonal injury is characterized by widespread damage to axons from the force and shearing sustained during the primary injury (Adams, Graham, Scott, Parker, & Doyle, 1980). These injuries and outcomes vary from child to child based on several factors because the primary injury plays a major part in the extent of the injury (Kondo et al., 2010; Moen et al., 2012).

Anatomic considerations confound the primary injury process for young children with TBI when compared to adults. Compared to adults, young children have proportionately large, heavy heads, weaker cervical ligaments and muscles, and thin, pliable skulls; all of which can result in increased severity of injury (Calder, Hill, & Scholtz, 1984). When young children sustain a blunt injury to the head, their large, heavy heads and weaker cervical ligaments and muscles cause the mechanical force to be transferred to the cervical spine region, which can result in cervical spinal fractures from the occiput to C2 and cervical cord damage (Nitecki & Moir, 1994). In children < 8 years of age, the fulcrum of movement is located at C2–3 compared to C5–6 in adults, which may explain why children experience cervical fractures in different locations than adults (Finch & Barnes, 1998). The thin, pliable skull provides less protection to the underlying brain, leaving it more vulnerable to shearing from linear and rotational forces (Leventhal, 1960). The long-term clinical impact of the initial structural changes in the brain from the primary injury is difficult to discern because the early effects of the primary injury are quickly compounded by the secondary injury, which activates complex biomolecular and physiologic reactions.

Secondary Injury

Secondary injury occurs because of alterations in cerebral blood flow (CBF) and the development of cerebral edema, which may lead to neuronal cellular death. If the CBF and cerebral metabolism continues to be compromised (Kilbaugh et al., 2011; Mandera, Larysz, & Wojtacha, 2002; Stiefel, Tomita, & Marmarou, 2005) additional cellular damage and death will occur and more toxic substances will be released (Pasvogel, Miketova, & Moore, 2010; Pun, Lu, & Moochhala, 2009). The longer the secondary injury phase continues, the more likely the child will sustain long-term disabilities (Casey, McKenna, Fiskum, Saraswati, & Robertson, 2008).

Cerebral Blood Flow—CBF is the mechanism that supplies the cerebral tissue with oxygenated blood. To maintain a consistent cerebral flood flow, cerebral arteries and

arterioles dilate and contract in response to hemodynamic changes called cerebral autoregulation (Hekmatpanah, 1970). During the first 0–9 days cerebral autoregulation is often impaired following a TBI, this results in CBF becoming dependent on cerebral perfusion pressures (CPP) and ICP (Tontisirin et al., 2007). An increased risk for impaired autoregulation has been associated with increased ICP, hyperemia, decreased hematocrit, decreased arterial oxygen (PaO₂), increased cerebral lesion size, and are less than 4 years of age (Freeman, Udomphorn, Armstead, Fisk, & Vavilala, 2008; Tontisirin et al., 2007). Clinically, children with impaired cerebral autoregulation have increased odds of poor outcomes (e.g., functional impairment, vegetative state, or death) (Tontisirin et al., 2007).

Cerebral Metabolism—Cerebral metabolism is affected following a severe TBI because the tissue is deprived of glucose and oxygen forcing cells into anaerobic metabolism (Holbach, Schroder, & Koster, 1972). Anaerobic metabolism produces less energy and increases lactate production, ultimately making it difficult for cells to maintain normal functional processes. Failure to maintain cellular functioning due to lack of energy creates an acidotic environment, which leads to serious consequences (Bouzat et al., 2014). When the sodium/potassium pumps (Na/K) pumps fail, an excessive influx of the positively charged sodium ions precipitate massive depolarization of neurons. This depolarization leads to the release of glutamate, an excitatory neurotransmitter that leads to excitotoxicity, as well causing more damage (Chamoun, Suki, Gopinath, Goodman, & Robertson, 2010).

The excess extracellular glutamate activates cell receptors that allow a massive influx of calcium and sodium into the cells leading to clinically significant consequences. The increased intracellular calcium leads to a) activation of the nitric oxide synthetase (NO) leading to cerebral edema and ischemia; b) activation of casplases (destructive enzymes) that damage the DNA of the cell leading to apoptosis; c) activation of the proteases family of Calpains, which cause the breakdown of the neuronal cytoskeleton leading to the loss of structural integrity of the neuron; d) damage to the sodium/potassium pump, leading to an influx of additional sodium followed by water with resultant cerebral edema; and e) mitochondrial injury leading to reactive oxygen species (ROS) generation, microvascular damage and cerebral edema (Huh, Franklin, Widing, & Raghupathi, 2006; Robertson, Bucci, & Fiskum, 2004; Robertson, Saraswati, & Fiskum, 2007). The influx of sodium can lead to increased permeability of the calcium ion channels resulting in cell swelling and/or neuronal death (Staal et al., 2010). Glutamate, calcium, and sodium can cause significant damage to the structures and functions throughout the nervous system.

If cerebral blood flow has been disrupted for a period of time, reperfusion injury may occur when adequate blood flow is restored. The areas of ischemia will likely demonstrate the sequelae associated with poor cerebral blood flow (Robertson, Scafidi, McKenna, & Fiskum, 2009). Upon restoration of perfusion, leukocytes attempt to repair the ischemic tissue, which leads to further generation of free radicals. The free radicals likely play a role in cellular death during reperfusion injury (Reddy & Labhasetwar, 2009).

Blood Brain Barrier and Edema—The blood brain barrier (BBB) is responsible for restricting movement of certain molecules, including medications and white blood cells into the central nervous system. This barrier consists of astrocyte endfeet, brain microvascular

endothelial cells, and capillary basement membrane. Shearing damage sustained in the primary injury can cause structural damage to the blood brain barrier increasing the membrane permeability. During the secondary injury phase, research suggest that excitotoxicity from the excess glutamate leads to an influx of calcium and sodium into the cells (Pun et al., 2009).

Brain edema is the increase of fluid in the brain, which is likely a result of changes in the cellular, molecular, structural, and functional properties of the BBB that usually prevent excess fluid from entering the brain (Unterberg, Stover, Kress, & Kiening, 2004). There are two types of cerebral edema after TBI: vasogenic (interstitial) and cytotoxic (intracellular) (Greve & Zink, 2009). Vasogenic cerebral edema is a result of the increased permeability of the BBB due to necrosis of capillary endothelial cells (Unterberg et al., 2004). The damage to capillary endothelial cells allows the leakage of proteins into the interstitial fluid creating an osmotic gradient. This osmotic gradient allows excess water and proteins to accumulate in the interstitial space of the brain (Donkin & Vink, 2010).

Cytotoxic edema occurs due to energy failure and the inability to maintain the sodium/potassium pump. The influx of sodium ions into the intracellular space is followed by an influx of water from the extracellular space in an attempt to create homeostasis (Donkin & Vink, 2010). The interstitial space within the brain is reduced in cytotoxic edema because the cells swell (Unterberg et al., 2004).

Importantly, both types of edema may lead to increased intracranial pressure (ICP) because of the brain's inability to compensate for the increased volume of the brain tissues or increased volume in the interstitial space of the brain. Based on the Monroe-Kellie hypothesis, the brain will initially attempt to compensate for the edema by displacing CSF into the spinal canal and venous blood into the jugular veins; however, if these mechanisms fail to decrease the volume within the cranial cavity, the ICP will increase (Weed, 1929). The cerebral edema can be monitored clinically through intracranial pressures, visualized on radiographic images, and observed through changes in the child's behavior.

Necrosis and Apoptosis—Primary and secondary injury can lead to two types of cellular death: necrosis and apoptosis. Necrosis is cellular death that occurs in conditions of very severe hypoxia and ischemia due to lack of oxygen (Johnston et al., 2009). Pathologically the cell membrane ruptures, the mitochondria are severely damaged and all the toxic contents (e.g., phospholipases, proteases, and lipid peroxidase) are leaked into the extracellular space leading to inflammation (Minambres et al., 2008). The leakage from the cells can activate the inflammatory cascade and lead to additional consequences. This is why apoptosis would be the preferred method of cellular death.

If the insult is less severe, the cells may recover or may progress to programmed cell death, known as apoptosis (Johnston et al., 2009). Apoptosis is a caspase-dependent pathway that ultimately results in cell shrinkage and general preservation of the cellular membranes without associated inflammation that may begin on the first day of injury (Minambres et al., 2008; Pasvogel et al., 2010). Both necrosis and apoptosis can lead to clinical changes such as decreased cognitive and physical disabilities because of losses in neurons through cellular

death (Brown, Elovic, Kothari, Flanagan, & Kwasnica, 2008). In an attempt to mitigate the damage sustained following severe TBI, advanced monitoring techniques are utilized to monitor the progression of the secondary injury or to evaluate the responsiveness to treatments.

Assessing and Monitoring Children with Severe TBI

Initial Assessment and Stabilization

In the emergency department, the specially-trained trauma team is responsible for the initial assessment and stabilization of a child with a severe TBI. The primary assessment begins with addressing all, potential life-threatening injuries (e.g., tension pneumothorax) including the head injury. The team must determine whether or not the child has a patent airway, can maintain an adequate respiratory drive (breathing), has adequate circulation, and initial neurologic status (disability). This primary assessment is often referred to as the ABCDs (ENA, 2007).

The assessment of the airway is used to determine if air can pass through without obstruction to the lungs (McFadyen, Ramaiah, & Bhananker, 2012). The airway must be cleared of any debris by suctioning or other methods, if obstructed. The team may choose to elicit a gag reflex with the suction catheter while suctioning the child to determine if the child is able to manage his oral secretions and protect his airway from potential aspiration of stomach contents (Marcoux, 2005). If the child has a depressed level of consciousness, the child may be unable to protect his airway and endotracheal intubation may be necessary, while maintaining inline stabilization of the cervical spine (C-spine) (McFadyen et al., 2012).

The assessment of breathing is used to determine which interventions are necessary to help the child maintain adequate oxygenation and ventilation (McFadyen et al., 2012). Children who present with abnormal breathing patterns (e.g., central neurogenic hyperventilation, cluster breathing, ataxic breathing, and apnea) may have sustained a brainstem lesion/injury or a C-spine injury with resultant phrenic nerve failure requiring initial respiratory support (e.g., endotracheal intubation and placement on a ventilator) (Easter, Barkin, Rosen, & Ban, 2010; Marcoux, 2005). C-spine precautions (e.g., rigid cervical collar, log roll turning method) should be maintained because head injuries are associated with spinal cord injuries (Easter et al., 2010). Maintaining systemic oxygenation and ventilation parameters of PaO₂ 301–500 mmHg (40–67 kPa) and PaCO₂ of 36–45 mmHg (5–6 kPa) are associated with higher rates of survival for children (Ramaiah et al., 2013).

Initial assessment and stabilization of circulation begins with checking for a pulse and beginning cardiopulmonary resuscitation, if necessary (Chameides, Ralston, Pediatrics, & Association, 2011). Stopping any external hemorrhage is also critical to minimize blood loss. Maintaining adequate circulation in children with head injuries is important because cerebral perfusion pressure (CPP) depends on the ICP and mean arterial pressure (MAP) (Hekmatpanah, 1970). The MAP is a reflection of the average arterial system pressure during a cardiac cycle and can be calculated based on this equation: $MAP = (1/3 \text{ systolic BP}) + (2/3 \text{ diastolic BP})$. To ensure adequate perfusion the guidelines from Pediatric

Advanced Life Support (PALS), the recommendation is to maintain a pediatric (1–10 years of age) systolic blood pressure $> 70 \text{ mmHg} + (2 \times \text{age in years})$ to avoid a hypotensive state (Kleinman et al., 2010). See Table 1 for recommended MAPS based on age (Hazinski, 2013; Top, Tasker, & Ince, 2011).

The initial neurological status (disability) is assessed through the pediatric Glasgow Coma Scale (GCS), pupil response, and overall responsiveness (ENA, 2007). The pediatric GCS is a modified version of the adult GCS that accounts for the changing developmental status of children (scored 3–15). If the child has an initial pediatric GCS < 8 (classified as a severe head injury) or is unarousable, the trauma/emergency team considers emergent endotracheal intubation and placement on a ventilator (Falk, 2012; Marcoux, 2005). The initial assessment and stabilization should take less than 10 minutes.

Neurological Assessment

Assessing Cerebral Edema and Posttraumatic Cerebrovascular Injuries—

Cranial imaging through computed tomography (CT) and magnetic resonance imaging (MRI) are used to identify cerebral edema and cerebrovascular injuries. This information is used to determine what neurosurgical interventions are necessary and to provide prognostic information about long-term developmental outcomes (Beauchamp et al., 2011; Suskauer & Huisman, 2009). Cranial CT scans are useful in identifying large hematomas, midline shift, skull fractures, excess CSF accumulation in the ventricles, cerebral edema, and brain herniation (Steinborn et al., 2010; Suskauer & Huisman, 2009). The results of the initial CT scans are used to determine if emergent neurosurgical interventions (e.g., decompression craniotomy) are needed (Suskauer & Huisman, 2009). The major advantage of using CT scan is the scan takes less than 10 minutes to perform and the child is easily accessible to the nurse throughout the scan. The major disadvantage of using CT is the potential for exposing the child to ionizing radiation (Duhaime, Holshouser, Hunter, & Tong, 2012).

Empirical evidence suggests diagnostic radiation both *in utero* and in childhood is associated with an increased risk of childhood cancer (IARC, 2012). The overall cancer incidence is 24% greater for those exposed to CT scans than for those who are not exposed in childhood (Mathews et al., 2013). Children needing a CT scan of the brain are at a greater risk of exposure because the skull is less dense than adult skulls and offer less protection to the underlying cerebral tissue (Morton et al., 2013). The American College of Radiology and the *Image Gently* campaign have made specific suggestions for radiologists and radiologic technologists and medical physicist to lower CT doses of radiation and use alternative testing when possible (Strauss et al., 2010). Nurses can assist the clinicians in meeting these new goals by ensuring that only areas that need to be scanned are scanned.

MRI scans are helpful in identifying extra-axial hemorrhage (e.g., subdural hemorrhage), early areas of ischemia and detecting edema in subacute moderate-to-severe injury (Bigler et al., 2013; Hunter, Wilde, Tong, & Holshouser, 2011). The major disadvantage of using MRI scan is that it takes longer than a CT scan and the equipment taken into the room to monitor the child must be MRI-safe. The nurse needs to review MRI safety with parents who may accompany the child into the scan, and the child or parent does not have any MR-contraindicated implantable devices (e.g., deep brain stimulators) (Duhaime et al., 2012).

Assessing Intracranial Pressures—The pressure in the cranial vault is made up of the cerebral spinal fluid (CSF), blood, and brain. According to the Monro-Kellie hypothesis, ICP is relatively constant because the cranial cavity represents a fixed volume (Weed, 1929). If an increase in the volume of CSF, blood, or brain compartments occurs, the brain will compensate initially by displacing one of the two other compartments. However, when displacement does not decrease the pressure within the cranial cavity, the ICP increases. In the case of TBI, one or more of these components can become increased, leading to increased ICP (Weed, 1929).

ICP can be assessed through clinical observations (e.g., headache, irritability), non-invasive examinations (e.g., transcranial Doppler), and direct invasive measures (e.g., intraparenchymal monitors). Careful and frequent assessment of children with severe TBI may allow the clinician to observe and manage the signs and symptoms of early increased ICP before the child progresses to the late signs. Early signs of increased ICP may be subtle such as irritability and cranial nerve dysfunction. The most ominous late sign of increased ICP is Cushing's triad. Cushing's triad is the result of cerebral ischemia. Clinically, it manifests as increased systolic blood pressure (also widened pulse pressure) to increase cerebral perfusion, bradycardia due to a vagal response triggered by the cardiac baroreceptors, and abnormal or irregular respirations (Cushing, 1903). See Table 2 for signs and symptoms of increasing intracranial pressure.

Transcranial Doppler (TCD) is a non-invasive examination of the blood flow velocity of the middle cerebral arteries through the transtemporal window (temple anterior to the ear) to determine cerebral hemodynamics and indirectly assess intracranial hypertension (Trabold, Meyer, Blanot, Carli, & Orliaguet, 2004; Verlhac, 2011). The use of TCD to evaluate increased ICP is limited in children (Melo et al., 2011; Meyer et al., 2005) and has previously been used to evaluate cerebral vasospasms (Trabold et al., 2004) and cerebral autoregulation (Vavilala et al., 2004). This non-invasive method may be useful as screening tool in early resuscitation because it can be used in children with coagulation disorders and those with critical injuries prior to computed tomography (CT) scan (Melo et al., 2011). Nursing responsibilities for the procedure include positioning the child supine throughout the procedure.

The two most common types of direct invasive intracranial monitoring devices utilized for monitoring in pediatric TBI are the external ventricular device (EVD) and intraparenchymal (IP) monitors (Wiegand & Richards, 2007). The EVD is the gold-standard for measuring ICP. An EVD is a fluid-coupled device with the catheter placed directly in the lateral ventricles that allows access to CSF, making it ideal for drainage of CSF for increased ICP and sampling of CSF (Padayachy, Figaji, & Bullock, 2010). Nursing considerations when caring for a child with an EVD involve assessing for complications: monitoring for infection including systemic temperature (Geyer, Meller, Kulpan, & Mowery, 2013); assessing the dressing for drainage (Ngo et al., 2009), assessing CSF drainage for color, clarity, and amount (Geyer et al., 2013); and monitoring for signs and symptoms of over-drainage of CSF (e.g., dehydration, hyponatremia) and under-drainage of CSF (Ngo et al., 2009). Nurses must also ensure each time the child is repositioned the EVD is zeroed with the foramen of Monro (external auditory meatus) at the prescribed level (Geyer et al., 2013).

Intraparenchymal devices such as the Codman microsensor (Codman, Raynham, MA, USA) and the Camino (Integra Neurosciences, Plainsboro, NJ, USA) are also utilized to directly monitor ICP. Generally the transducer/catheter is placed within the subarachnoid space and then is secured to the scalp (AANN, 2011). During the insertion process, the ICP monitor and the intracranial transducers are zeroed and do not require further zeroing (Marcoux, 2005). The extracranial transducers are recalibrated per institutional protocol (usually every 12 hours). The nursing responsibilities involve monitoring for complications associated with IP devices including assessing for hemorrhage and fractures, and monitoring for infection (Anderson et al., 2004). With both the EVD and IP monitor, the nurse must also try to keep the child from removing the device and pulling the equipment.

When ICP monitors are available CPP can be calculated by the following equation: $CPP = MAP - ICP$. CPP represents an indirect measure of cerebral blood flow. The ICP should be treated to maintain adequate CPP between 40–50 mmHg for children (Kochanek et al., 2012).

Assessing Cerebral Oxygenation—The main methods of measuring alterations in cerebral oxygenation are jugular venous oxygen saturation ($SjvO_2$), brain tissue oxygen partial pressure ($PbtO_2$), and near-infrared spectroscopy (NIRS). Jugular venous oxygen saturation measures the balance of global cerebral oxygen delivery and the rate of cerebral metabolic oxygen consumption (Perez et al., 2003). The catheter to measure $SjvO_2$ is positioned in the bulb of the internal jugular vein (Blissitt, 2009) with the tip of the catheter above the level of C1/C2 disk to minimize contamination from the facial vein (Rohlwink & Figaji, 2010). Generally, normal values of $SjvO_2$ range from 55 to 75% (Perez et al., 2003). Decreased $SjvO_2$ can occur with decreased systemic oxygen supply, local or systemic hypoperfusion, and increased cerebral metabolism or oxygen extraction (e.g., pyrexia and seizures). Children with severe TBI who experienced two or more $SjvO_2$ measurements 55% had an increased risk of poor neurologic outcome (severe disability, coma or persistent vegetative state, or brain death) (Perez et al., 2003). Increased $SjvO_2$ may occur with decreased cerebral metabolism, increased systemic oxygen supply, restricted oxygen diffusion, and hyperemia (Rohlwink & Figaji, 2010). The limitation to monitoring $SjvO_2$ is the inability to measure focal ischemia, as the measurement only provides an assessment of global cerebral oxygenation (Rohlwink & Figaji, 2010).

Careful assessment and management of $SjvO_2$ catheters is necessary to ensure accurate results and prevent complications. When sampling blood from the catheter, withdrawn at a rate of 2 milliliters/minute to decrease the risk of extracerebral contamination. Recalibrate in vivo using a blood sample at least every 24 hours. Maintaining a pressurized saline bag with continuous flush may prevent clots and facilitate proper functioning of the catheter (Blissitt, 2009). Nurses need to monitor for complications that can occur during insertion or while the catheter is indwelling: carotid artery puncture, hematoma formation, infection, thrombosis, and raised ICP (Rohlwink & Figaji, 2010).

Brain tissue oxygenation ($PbtO_2$) monitors (e.g., Licox [Integra Neurosciences, Plainsboro, NJ, USA]) provide an estimate of the balance between cellular oxygen consumption and regional oxygen supply (Purins, Lewen, Hillered, Howells, & Enblad, 2014). $PbtO_2$ is

thought to be reflective of CBF, arterial oxygen tension, or a product of CBF and arteriovenous tension difference in oxygen tension (De Georgia, 2014; Rosenthal et al., 2008). Normal parameters for PbtO₂ values are > 30 mmHg (millimeters mercury) (Maloney-Wilensky & Le Roux, 2010; Stippler et al., 2012). PbtO₂ values < 20 mmHg are usually considered to be approaching ischemia (De Georgia, 2014). When PbtO₂ values are between 0 and 5 mmHg for greater than 1 hour, there is a high risk of severe disability or death (Figaji & Kent, 2010; Figaji et al., 2009). Leading causes of decreased PbtO₂ levels include decreased arterial partial pressure of oxygen, poor cerebral blood flow, tissue barriers to diffusion (e.g., cytotoxic edema), increased arterial CO₂, increased intracranial pressure, and low cerebral perfusion pressure can decrease PbtO₂ and lead to ischemia (Rohlwink et al., 2012).

The Licox system, the major PbtO₂ monitor utilized, can also monitor brain temperature, and ICP simultaneously by probes inserted through a triple-lumen bolt (Stevens, 2004). The triple-lumen bolt is usually placed in the frontal lobe in non-contused tissue identified on CT scan (Narotam, Burjonrappa, Raynor, Rao, & Taylon, 2006; Stiefel et al., 2006; Ushewokunze & Sgouros, 2009) and at least 14 mm (millimeters) from other probes (e.g. EVD); (Stevens, 2004). Due to localized microtrauma at the insertion site, the PbtO₂ sensor readings are not considered accurate for the first 30–120 minutes after insertion. The current recommendation is to maintain PbtO₂ > 10 mmHg (Kochanek et al., 2012). Following insertion, the nurse must ensure the Licox cables are secured per institutional protocol, usually at two points of tensions, one directly on the patient's head where the bolt is anchored to the skin and the other at the patient's shoulder (Wilensky et al., 2005). The PbtO₂ monitor does not require daily recalibration (Stevens, 2004). The dressing at the site is changed every 48 hours or when saturated and replaced with a dry, sterile, occlusive dressing (Wilensky et al., 2005). The nurse also needs to monitor for possible complications associated with PbtO₂ monitors: infection (Stiefel et al., 2006), hemorrhage (Stiefel et al., 2006), and catheter dislodgement (Stevens, 2004).

Near-infrared spectroscopy (NIRS) is a non-invasive measure of regional oxygen saturation (rSO₂) that provides continuous measurements. This measurement describes the balance of oxygen supply and oxygen demand of the regional cerebral tissue (Drayna, Abramo, & Estrada, 2011). The normal parameter of rSO₂ for children is about 70% (Yaron et al., 2003; Yoxall, Weindling, Dawani, & Peart, 1995). Preliminary data from 30 children suggests cerebral rSO₂ is impacted by changes in end tidal CO₂, heart rate, and hematocrit changed cerebral rSO₂, but is not effected by changes in ICP or CPP (Zuluaga, Esch, Cvijanovich, Gupta, & McQuillen, 2010). NIRS has not been used extensively in children with TBI (Amigoni et al., 2011).

Current Treatments for Children with Severe TBI

Many of the treatments for severe TBI are determined based on assessment findings of increased ICP, impaired cerebral oxygenation, cerebral edema, and cerebrovascular injuries. The treatments aim to restore adequate CBF and cerebral metabolism, reduce cerebral edema, and minimize cellular death to ameliorate secondary injury. The Guidelines for the Acute Medical Management of Severe Traumatic Brain Injury in Infants, Children, and

Adolescents (2012) provides treatment recommendations based level: I, II, III. The treatment recommendation based levels are determined by the strength of the study designs available for each treatment with level I having the strongest evidence and level III having the least evidence (Kochanek et al., 2012). See Table 3 for a summary of the evidence for treatment of severe TBI. The main treatment modalities include hyperosmolar therapy, temperature control, consideration of etomidate to control severe increased ICP and thiopental to control ICP, and consideration of prophylactic treatment to reduce incidence of early posttraumatic seizures.

ICP monitoring is a mainstay in the treatment of severe TBI (Exo et al., 2011) with most guidelines considering values of > 20–25 mmHg requiring treatment (Kochanek et al., 2012). Current research also suggests maintaining PbtO₂ at a minimum of 10 mmHg (Kochanek et al., 2012; Narotam et al., 2006). When the monitoring parameters are not within the acceptable ranges, treatments are necessary to prevent further injury to the cerebral tissue.

Hyperosmolar Therapy—Hyperosmolar therapy creates an increased serum osmolality by pulling fluid from interstitial space back into the cerebral vessel and into systemic circulation, lowering the volume in the cranial vault (Knapp, 2005). Hyperosmolar therapy with 3% hypertonic saline solution should be considered for intracranial hypertension because it alters cerebral volume (Weed, 1929). The effective acute dose of hypertonic saline is 6.5 to 10 mL/kg (milliliter per kilogram) and the continuous dose is 0.1 mL to 1.0 mL/kg/hr (milliliter per kilogram per hour) to maintain ICP < 20 mmHg (Kochanek et al., 2012). Nurses need to monitor serum osmolality every 4 to 6 hours to ensure it is maintained below 360 mOsm/L (milliosmole per liter) for 3% hypertonic saline (Kochanek et al., 2012).

Temperature Control—Systemic hyperthermia from inflammation following TBI can lead to additional secondary injury because pyrexia exacerbates many of the biochemical reactions (e.g., excitotoxicity, production of ROS) (Puccio et al., 2009). Animal studies suggest that hyperthermia is associated with poor outcomes and should be avoided and thus maintaining normothermia is recommended in severe TBI (Kochanek et al., 2012). Nurses should monitor the child's temperature and adjust the environment to ensure the child is not hyperthermic, also alerting the healthcare team if the child becomes hyperthermic.

Cerebrospinal fluid drainage—Drainage of cerebrospinal fluid is to reduce the volume of intracranial fluid, which in turn decreases the volume within the cranial vault and thus reduces the intracranial pressure. Drainage of CSF can occur through the EVD (Kochanek et al., 2012). The EVD can also be used for monitoring ICP, which is helpful in the management of children with severe TBI. Normal output of CSF is 3–5 mL/hour for infants; 5–10 mL/hr for children; and 10–15 mL/hr for adolescents (Vernon-Levett, 2006).

Barbiturate Therapy—Very limited evidence exists for the use of barbiturates in severe TBI. The use of barbiturates may be necessary, when CSF drainage, hyperosmolar therapy, and sedation and analgesia have not effectively lowered the ICP to maintain an adequate CPP (Marshall et al., 2010). Barbiturates decrease ICP by suppressing metabolism and

altering vascular tone (Kochanek et al., 2012). Pentobarbital or thiopental are administered as the medications of choice (Glick, Ksendzovsky, Greesh, & Raksin, 2011; Marshall et al., 2010; Mellion et al., 2013). Nurses generally monitor patients with TBI who receive barbiturates for medically-induced comas closely. Vital signs are measured at least hourly, if not more frequently depending on the child's condition (Mazzola & Adelson, 2002). Continuous arterial blood pressure monitoring and cardiovascular support to maintain adequate CPP is usually required (Kochanek et al., 2012). Nurses also monitor for electrographic burst suppression, which is evaluated through the use of continuous electroencephalography (EEG) (Mellion et al., 2013). These children require critical care by vigilant specially trained nurses.

Decompressive craniectomy—Decompressive craniectomy (DC) is a surgical procedure to remove a portion of the skull and debride the necrotic tissue and allow unimpeded swelling of the brain and improved CBF (Morrow & Pearson, 2010). DC may be considered for children showing early signs of neurologic deterioration or herniation in the early stages of ICP, or evaluation of a mass lesion refractory to medical treatment (Bowers, Riva-Cambrin, Hertzler, & Walker, 2013). The outcomes of children who undergo DC are generally positive; however, few studies exist examining pediatric patients (Thomale, Graetz, Vajkoczy, & Sarrafzadeh, 2010; Weintraub, Williams, & Jane, 2012). In studies with children who underwent DC due to sudden increases in ICP, long-term outcomes showed moderate disability to good recovery of the majority of the children (Figaji, Fieggen, & Peter, 2003; Josan & Sgouros, 2006; Ruf et al., 2003). Currently, a large international randomized-controlled trial, Randomised Evaluation of Surgery with Craniectomy for Uncontrollable Elevation of Intra-Cranial Pressure (RESCUEicp), is underway to better understand if DC is an effective form of treatment for traumatic brain injuries for individuals ages 10–65 years of age (Bohman & Schuster, 2013; Hutchinson et al., 2006). Nurses caring for these children need to monitor for signs and symptoms of infection (Hockenberry & Wilson, 2007), maintain precautions to prevent physical pressure around the area without an intact skull, and assess and manage pain (Vernon-Levett, 2006)

Analgesia, sedation, and neuromuscular blockade—The purpose of administering analgesia and sedation medication is to decrease pain and stress, which increase cerebral metabolic demands and increases cerebral blood volume, in turn raising ICP (Kochanek et al., 2012). Different analgesics and sedation medications can effect the cerebral vasculature, metabolism, autoregulation, ICP, and CPP. Unfortunately because most analgesic and sedation medications have not been evaluated, in regard, to the effect on ICP, they cannot be recommended as a treatment for intracranial hypertension. Etomidate and thiopental can be considered to treat intracranial hypertension. If etomidate is utilized, the risks of adrenal suppression must be considered (Kochanek et al., 2012). The use of neuromuscular blockade agents is mentioned in the guidelines as a consideration, but no recommendations are offered. Due to the paucity of evidence, additional medications frequently utilized in the pediatric intensive care are also reviewed. The recommendation is to allow the treating clinician to choose which medication is best for the patient.

Opioids (e.g., morphine, fentanyl), classified as analgesics agents, can directly affect the respiratory centers in the medulla, leading to a decreased respiratory rate and subsequent increased PaCO₂ (Kilbaugh, Friess, Raghupathi, & Huh, 2010). The increased PaCO₂ can cause vasodilatation of the cerebral vessels and increase ICP and decrease CPP. The nurse should monitor for common side effects of opioids: constipation, urinary retention, sedation, nausea, vomiting, respiratory depression, bradycardia, hypotension, and pruritis (Kilbaugh et al., 2010).

Analgesia is often administered in conjunction with sedation. The classification of medications generally used for sedation is benzodiazepines (e.g., midazolam, lorazepam, diazepam). Benzodiazepines have several advantages for children with severe TBI: sedation, anxiolysis, muscle relaxation, anterograde amnesia, decreased cerebral metabolic rate, and anticonvulsant properties (Kilbaugh et al., 2010). However, clinically important side effects exist that could be detrimental: decreased blood pressure and depressed ventilation (M. J. Meyer et al., 2010). Initially, nurses should monitor vital signs each hour or more frequently depending on the child's condition and continue to monitor the child's pain level (Hockenberry & Wilson, 2007)

Antiseizure prophylaxis—Prophylactic treatment with phenytoin may be considered to reduce the incidence of early posttraumatic seizures (PTS) (Kochanek et al., 2012). Early PTS are witnessed clinical seizures or abnormal trace recordings on EEG within the first 7 days following the traumatic event (Arango et al., 2012) with about 44% of children having at least one seizure (Arndt et al., 2013). Risk factors associated with any type of PTS are young age, type of trauma (abusive head trauma), epidural hematoma (Arango et al., 2012), and skull fracture (Arndt et al., 2013). Risk factors associated with subclinical PTS are young age, abusive head trauma, subdural hematoma, and intraaxial hemorrhage (Arndt et al., 2013). Additional research is needed to guide the treatment of antiseizure prophylaxis.

Conclusion

The pathophysiologic effects of primary and secondary injury play a key role in the overall outcomes for children with TBI. Trauma prevention is the only way to combat primary injury. Nurses can help educate parents and children about seat belt safety, falls preventions, and bicycle safety. The secondary injury phase that occurs when a complex biochemical and physiologic process is activated requires vigilant nursing care. Nurses must monitor for alterations in arterial oxygenation, cerebral metabolism, cerebral autoregulation, and cerebral blood flow through changes in the child's behavior and highly technical biometric equipment. The care and monitoring of children with severe TBI continues for days to weeks because the cerebral ischemic leads to increased permeability of the blood brain barrier, and cellular death. Treatments exist to combat the effects of the secondary injury, but these treatments cannot 'cure' TBI. The children who sustain a TBI may have no residual impairments but other may suffer catastrophic social, emotional, cognitive, and functional impairments.

References

- Thompson, H., editor. AANN. Care of a Patient Undergoing Intracranial Pressure Monitoring/External Ventricular Drainage or Lumbar Drainage: AANN Clinical Practice Guideline Series. Glenview, IL: 2011. p. 38
- Adams JH, Graham DI, Scott G, Parker LS, Doyle D. Brain damage in fatal non-missile head injury. *J Clin Pathol.* 1980; 33(12):1132–1145. [PubMed: 7451661]
- Adelson PD, Bratton SL, Carney NA, Chesnut RM, du Coudray HE, Goldstein B, Wright DW. Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents. Chapter 1: Introduction. *Pediatric Critical Care Medicine.* 2003; 4(3 Suppl):S2–4.10.1097/01.CCM.0000066600.71233.01 [PubMed: 12847337]
- Amigoni A, Mozzo E, Brugnaro L, Tiberio I, Pittarello D, Stellan G, Bonato R. Four-side near-infrared spectroscopy measured in a paediatric population during surgery for congenital heart disease. *Interactive Cardiovascular Thoracic Surgery.* 2011; 12(5):707–712.10.1510/ievts.2010.253328 [PubMed: 21335618]
- Anderson RC, Kan P, Klimo P, Brockmeyer DL, Walker ML, Kestle JR. Complications of intracranial pressure monitoring in children with head trauma. *Journal of Neurosurgery.* 2004; 101(1 Suppl):53–58.10.3171/ped.2004.101.2.0053 [PubMed: 16206972]
- Arango JI, Deibert CP, Brown D, Bell M, Dvorchik I, Adelson PD. Posttraumatic seizures in children with severe traumatic brain injury. *Childs Nerv Syst.* 2012; 28(11):1925–1929.10.1007/s00381-012-1863-0 [PubMed: 22843174]
- Arndt DH, Lerner JT, Matsumoto JH, Madikians A, Yudovin S, Valino H, Giza CC. Subclinical early posttraumatic seizures detected by continuous EEG monitoring in a consecutive pediatric cohort. *Epilepsia.* 2013; 54(10):1780–1788.10.1111/epi.12369 [PubMed: 24032982]
- Beauchamp MH, Ditchfield M, Babl FE, Kean M, Catroppa C, Yeates KO, Anderson V. Detecting traumatic brain lesions in children: CT versus MRI versus susceptibility weighted imaging (SWI). *Journal of Neurotrauma.* 2011; 28(6):915–927.10.1089/neu.2010.1712 [PubMed: 21501069]
- Bigler ED, Abildskov TJ, Petrie J, Farrer TJ, Dennis M, Simic N, Owen Yeates K. Heterogeneity of brain lesions in pediatric traumatic brain injury. *Neuropsychology.* 2013; 27(4):438–451.10.1037/a0032837 [PubMed: 23876117]
- Black SP. Epidural hematoma; several clinical variations. *Mo Med.* 1956; 53(10):849–852. [PubMed: 13369395]
- Blissitt, P. Brain Oxygen Monitoring. In: Littlejohns, LR.; Bader, MK., editors. AACN-AANN Protocols for Practice: Monitoring Technologies in Critically Ill Neuroscience Patients. Ontario: Jones and Bartlett Publishers, LLC; 2009. p. 103-145.
- Blumbergs PC, Scott G, Manavis J, Wainwright H, Simpson DA, McLean AJ. Staining of amyloid precursor protein to study axonal damage in mild head injury. *Lancet.* 1994; 344(8929):1055–1056. [PubMed: 7523810]
- Bohman LE, Schuster JM. Decompressive craniectomy for management of traumatic brain injury: an update. *Curr Neurol Neurosci Rep.* 2013; 13(11):392.10.1007/s11910-013-0392-x [PubMed: 24101348]
- Bouzat P, Sala N, Suys T, Zerlauth JB, Marques-Vidal P, Feihl F, Oddo M. Cerebral metabolic effects of exogenous lactate supplementation on the injured human brain. *Intensive Care Med.* 2014; 40(3):412–421.10.1007/s00134-013-3203-6 [PubMed: 24477453]
- Bowers CA, Riva-Cambrin J, Hertzler DA 2nd, Walker ML. Risk factors and rates of bone flap resorption in pediatric patients after decompressive craniectomy for traumatic brain injury. *J Neurosurg Pediatr.* 2013; 11(5):526–532.10.3171/2013.1.PEDS12483 [PubMed: 23473303]
- Bramlett HM, Dietrich WD. Pathophysiology of cerebral ischemia and brain trauma: similarities and differences. *Journal of Cerebral Blood Flow and Metabolism.* 2004; 24(2):133–150.10.1097/01.WCB.0000111614.19196.04 [PubMed: 14747740]
- Brown AW, Elovic EP, Kothari S, Flanagan SR, Kwasnica C. Congenital and acquired brain injury. 1. Epidemiology, pathophysiology, prognostication, innovative treatments, and prevention. *Archives of Physical Medicine and Rehabilitation.* 2008; 89(3 Suppl 1):S3–8.10.1016/j.apmr.2007.12.001

- Calder IM, Hill I, Scholtz CL. Primary brain trauma in non-accidental injury. *J Clin Pathol*. 1984; 37(10):1095–1100. [PubMed: 6490948]
- Case ME. Accidental traumatic head injury in infants and young children. *Brain Pathology*. 2008; 18(4):583–589.10.1111/j.1750-3639.2008.00203.x [PubMed: 18782170]
- Casey PA, McKenna MC, Fiskum G, Saraswati M, Robertson CL. Early and sustained alterations in cerebral metabolism after traumatic brain injury in immature rats. *Journal of Neurotrauma*. 2008; 25(6):603–614.10.1089/neu.2007.0481 [PubMed: 18454682]
- CDC. Ten leading causes of death and injury. 2010. Retrieved March 7, 2014, from <http://www.cdc.gov/injury/wisqars/LeadingCauses.html>
- Chameides, L.; Ralston, M. Pediatrics, A. A. o., & Association, A. H. Pediatric advanced life support. Dallas, TX: American Heart Association; 2011.
- Chamoun R, Suki D, Gopinath SP, Goodman JC, Robertson C. Role of extracellular glutamate measured by cerebral microdialysis in severe traumatic brain injury. *Journal of Neurosurgery*. 2010; 113(3):564–570.10.3171/2009.12.JNS09689 [PubMed: 20113156]
- Cohn R. Subdural hematoma; an experimental study. *Arch Neurol Psychiatry*. 1948; 59(3):360–367.
- Cushing H. The blood pressure reaction of acute cerebral compression, illustrated by cases of intracranial hemorrhage. *American Journal of Science*. 1903; 125:1017–1044.
- De Georgia MA. Brain Tissue Oxygen Monitoring in Neurocritical Care. *J Intensive Care Med*. 2014.10.1177/0885066614529254
- Donkin JJ, Vink R. Mechanisms of cerebral edema in traumatic brain injury: therapeutic developments. *Current Opinion in Neurology*. 2010; 23(3):293–299.10.1097/WCO.0b013e328337f451 [PubMed: 20168229]
- Drayna PC, Abramo TJ, Estrada C. Near-infrared spectroscopy in the critical setting. *Pediatric Emergency Care*. 2011; 27(5):432–439. quiz 440–432. 10.1097/PEC.0b013e3182188442 [PubMed: 21546812]
- Duhaime AC, Holshouser B, Hunter JV, Tong K. Common data elements for neuroimaging of traumatic brain injury: pediatric considerations. *J Neurotrauma*. 2012; 29(4):629–633.10.1089/neu.2011.1927 [PubMed: 21671798]
- Easter JS, Barkin R, Rosen CL, Ban K. Cervical Spine Injuries in Children, Part I: Mechanism of Injury, Clinical Presentation, and Imaging. *Journal of Emergency Medicine*. 2010.10.1016/j.jemermed.2009.11.034
- Eisenberg MA, Meehan WP 3rd, Mannix R. Duration and Course of Post-Concussive Symptoms. *Pediatrics*. 2014.10.1542/peds.2014-0158
- ENA. Trauma Nursing Core Course (TNCC). 6. Des Plaines, IL: Emergency Nurses Association; 2007.
- Exo J, Kochanek PM, Adelson PD, Greene S, Clark RS, Bayir H, Bell MJ. Intracranial pressure-monitoring systems in children with traumatic brain injury: Combining therapeutic and diagnostic tools. *Pediatr Crit Care Med*. 2011.10.1097/PCC.0b013e3181e8b3ee
- Falk AC. Impact of elevated ICP on outcome after paediatric traumatic brain injury requiring intensive care. *Childs Nerv Syst*. 2012; 28(7):1069–1075.10.1007/s00381-012-1772-2 [PubMed: 22573139]
- Faul, M.; Xu, L.; Wald, MM.; Coronado, VG. Traumatic brain injury in the United States: Emergency department visits, hospitalizations and deaths 2002–2006. Atlanta: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2010.
- Feigin VL, Theadom A, Barker-Collo S, Starkey NJ, McPherson K, Kahan M, Group BS. Incidence of traumatic brain injury in New Zealand: a population-based study. *Lancet Neurol*. 2013; 12(1):53–64.10.1016/S1474-4422(12)70262-4 [PubMed: 23177532]
- Figaji AA, Fieggen AG, Peter JC. Early decompressive craniotomy in children with severe traumatic brain injury. *Childs Nerv Syst*. 2003; 19(9):666–673.10.1007/s00381-003-0804-3 [PubMed: 12908115]
- Figaji AA, Kent SJ. Brain tissue oxygenation in children diagnosed with brain death. *Neurocrit Care*. 2010; 12(1):56–61.10.1007/s12028-009-9298-5 [PubMed: 19847675]
- Figaji AA, Zwane E, Thompson C, Fieggen AG, Argent AC, Le Roux PD, Peter JC. Brain tissue oxygen tension monitoring in pediatric severe traumatic brain injury. Part 1: Relationship with outcome. *Childs Nervous System*. 2009; 25(10):1325–1333.10.1007/s00381-009-0822-x

- Finch GD, Barnes MJ. Major cervical spine injuries in children and adolescents. *J Pediatr Orthop*. 1998; 18(6):811–814. [PubMed: 9821142]
- Freeman SS, Udomphorn Y, Armstead WM, Fisk DM, Vavilala MS. Young age as a risk factor for impaired cerebral autoregulation after moderate to severe pediatric traumatic brain injury. *Anesthesiology*. 2008; 108(4):588–595.10.1097/ALN.0b013e31816725d7 [PubMed: 18362589]
- Geyer K, Meller K, Kulpan C, Mowery BD. Traumatic brain injury in children: acute care management. *Pediatr Nurs*. 2013; 39(6):283–289. [PubMed: 24640314]
- Glick RP, Ksendzovsky A, Greesh J, Raksin P. Initial observations of combination barbiturate coma and decompressive craniectomy for the management of severe pediatric traumatic brain injury. *Pediatr Neurosurg*. 2011; 47(2):152–157.10.1159/000330709 [PubMed: 21934277]
- Greve MW, Zink BJ. Pathophysiology of traumatic brain injury. *Mount Sinai Journal of Medicine*. 2009; 76(2):97–104.10.1002/msj.20104 [PubMed: 19306379]
- Hazinski, MF., editor. *Nursing Care of the Critically Ill Child*. St. Louis, Missouri: Elsevier; 2013.
- Hekmatpanah J. Cerebral circulation and perfusion in experimental increased intracranial pressure. *J Neurosurg*. 1970; 32(1):21–29.10.3171/jns.1970.32.1.0021 [PubMed: 5410794]
- Hirsch CS, Kaufman B. Contrecoup skull fractures. *J Neurosurg*. 1975; 42(5):530–534.10.3171/jns.1975.42.5.0530 [PubMed: 1151449]
- Hockenberry, M.; Wilson, D., editors. *Wong's Nursing Care of Infants and Children*. 8. St. Louis: Mosby; 2007.
- Holbach KH, Schroder FK, Koster S. Alterations of cerebral metabolism in cases with acute brain injuries during spontaneous respiration of air, oxygen and hyperbaric oxygen. *Eur Neurol*. 1972; 8(1):158–160. [PubMed: 5052137]
- Holbourn A, Edin D. Mechanics of head injuries. *Lancet*. 1943; 242(6267):438–441.10.1016/S0140-6736(00)87453-X
- Huh JW, Franklin MA, Widing AG, Raghupathi R. Regionally distinct patterns of calpain activation and traumatic axonal injury following contusive brain injury in immature rats. *Developmental Neuroscience*. 2006; 28(4–5):466–476.10.1159/000094172 [PubMed: 16943669]
- Hunter JV, Wilde EA, Tong K, Holshouser B. Emerging Imaging Tools for Use with Traumatic Brain Injury Research. *Journal of Neurotrauma*. 2011;10.1089/neu.2011.1906
- Hutchinson PJ, Corteen E, Czosnyka M, Mendelow AD, Menon DK, Mitchell P, Kirkpatrick PJ. Decompressive craniectomy in traumatic brain injury: the randomized multicenter RESCUEicp study (www.RESCUEicp.com). *Acta Neurochir Suppl*. 2006; 96:17–20. [PubMed: 16671415]
- IARC. A review of human carcinogens: radiation: IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 100D. Lyon, France: 2012.
- Jamous MA, Abdel Aziz H, Al Kaisy F, Eloqayli H, Azab M, Al-Jarrah M. Conservative management of acute epidural hematoma in a pediatric age group. *Pediatric Neurosurgery*. 2009; 45(3):181–184.10.1159/000218200 [PubMed: 19440005]
- Johnston MV, Ishida A, Ishida WN, Matsushita HB, Nishimura A, Tsuji M. Plasticity and injury in the developing brain. *Brain & Development*. 2009; 31(1):1–10.10.1016/j.braindev.2008.03.014 [PubMed: 18490122]
- Josan VA, Sgouros S. Early decompressive craniectomy may be effective in the treatment of refractory intracranial hypertension after traumatic brain injury. *Childs Nerv Syst*. 2006; 22(10):1268–1274.10.1007/s00381-006-0064-0 [PubMed: 16496158]
- Kilbaugh TJ, Bhandare S, Lorom DH, Saraswati M, Robertson CL, Margulies SS. Cyclosporin A Preserves Mitochondrial Function after Traumatic Brain Injury in the Immature Rat and Piglet. *Journal of Neurotrauma*. 2011;10.1089/neu.2010.1635
- Kilbaugh TJ, Friess SH, Raghupathi R, Huh JW. Sedation and analgesia in children with developmental disabilities and neurologic disorders. *Int J Pediatr*. 2010; 201010.1155/2010/189142
- Klimo P Jr, Ragel BT, Scott WH Jr, McCafferty R. Pediatric neurosurgery during Operation Enduring Freedom. *J Neurosurg Pediatr*. 2010; 6(2):107–114.10.3171/2010.3.PEDS109 [PubMed: 20672929]
- Knapp JM. Hyperosmolar therapy in the treatment of severe head injury in children: mannitol and hypertonic saline. *AACN Clinical Issues*. 2005; 16(2):199–211. [PubMed: 15876888]

- Kochanek PM, Carney N, Adelson PD, Ashwal S, Bell MJ, Bratton S, Critical Care S. Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents--second edition. *Pediatr Crit Care Med*. 2012; 13(Suppl 1):S1–82.10.1097/PCC.0b013e31823f435c [PubMed: 22217782]
- Kondo K, Maruishi M, Ueno H, Sawada K, Hashimoto Y, Ohshita T, Matsumoto M. The pathophysiology of prospective memory failure after diffuse axonal injury--lesion-symptom analysis using diffusion tensor imaging. *BMC Neurosci*. 2010; 11:147.10.1186/1471-2202-11-147 [PubMed: 21092119]
- Leventhal HR. Birth injuries of the spinal cord. *J Pediatr*. 1960; 56:447–453. [PubMed: 14416159]
- Maloney-Wilensky E, Le Roux P. The physiology behind direct brain oxygen monitors and practical aspects of their use. *Childs Nervous System*. 2010; 26(4):419–430.10.1007/s00381-009-1037-x
- Mandera M, Larysz D, Wojtacha M. Changes in cerebral hemodynamics assessed by transcranial Doppler ultrasonography in children after head injury. *Childs Nervous System*. 2002; 18(3–4): 124–128.10.1007/s00381-002-0572-5
- Marcoux KK. Management of increased intracranial pressure in the critically ill child with an acute neurological injury. *AACN Clinical Issues*. 2005; 16(2):212–231. quiz 270–211. [PubMed: 15876889]
- Marshall GT, James RF, Landman MP, O'Neill PJ, Cotton BA, Hansen EN, May AK. Pentobarbital coma for refractory intra-cranial hypertension after severe traumatic brain injury: mortality predictions and one-year outcomes in 55 patients. *J Trauma*. 2010; 69(2):275–283.10.1097/TA.0b013e3181de74c7 [PubMed: 20699736]
- Mathews JD, Forsythe AV, Brady Z, Butler MW, Goergen SK, Byrnes GB, Darby SC. Cancer risk in 680,000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ*. 2013; 346:f2360.10.1136/bmj.f2360 [PubMed: 23694687]
- Mazzola CA, Adelson PD. Critical care management of head trauma in children. *Critical Care Medicine*. 2002; 30(11 Suppl):S393–401. [PubMed: 12528780]
- McFadyen JG, Ramaiah R, Bhananker SM. Initial assessment and management of pediatric trauma patients. *Int J Crit Illn Inj Sci*. 2012; 2(3):121–127.10.4103/2229-5151.100888 [PubMed: 23181205]
- Mellion SA, Bennett KS, Ellsworth GL, Moore K, Riva-Cambrin J, Metzger RR, Bratton SL. High-dose barbiturates for refractory intracranial hypertension in children with severe traumatic brain injury. *Pediatr Crit Care Med*. 2013; 14(3):239–247.10.1097/PCC.0b013e318271c3b2 [PubMed: 23392360]
- Melo JR, Di Rocco F, Blanot S, Cuttaree H, Sainte-Rose C, Oliveira-Filho J, Meyer PG. Transcranial Doppler can predict intracranial hypertension in children with severe traumatic brain injuries. *Childs Nervous System*. 2011; 27(6):979–984.10.1007/s00381-010-1367-8
- Meyer MJ, Megyesi J, Meythaler J, Murie-Fernandez M, Aubut JA, Foley N, Teasell R. Acute management of acquired brain injury part II: an evidence-based review of pharmacological interventions. *Brain Injury*. 2010; 24(5):706–721. [PubMed: 20376996]
- Meyer PG, Ducrocq S, Rackelbom T, Orliaguet G, Renier D, Carli P. Surgical evacuation of acute subdural hematoma improves cerebral hemodynamics in children: a transcranial Doppler evaluation. *Childs Nervous System*. 2005; 21(2):133–137.10.1007/s00381-004-1016-1
- Mihalik JP, Blackburn JT, Greenwald RM, Cantu RC, Marshall SW, Guskiewicz KM. Collision type and player anticipation affect head impact severity among youth ice hockey players. *Pediatrics*. 2010; 125(6):e1394–1401.10.1542/peds.2009-2849 [PubMed: 20478933]
- Minambres E, Ballesteros MA, Mayorga M, Marin MJ, Munoz P, Figols J, Lopez-Hoyos M. Cerebral apoptosis in severe traumatic brain injury patients: an in vitro, in vivo, and postmortem study. *Journal of Neurotrauma*. 2008; 25(6):581–591.10.1089/neu.2007.0398 [PubMed: 18363508]
- Moen KG, Skandsen T, Folvik M, Brezova V, Kvistad KA, Rydland J, Vik A. A longitudinal MRI study of traumatic axonal injury in patients with moderate and severe traumatic brain injury. *J Neurol Neurosurg Psychiatry*. 2012; 83(12):1193–1200.10.1136/jnnp-2012-302644 [PubMed: 22933813]
- Morrow SE, Pearson M. Management strategies for severe closed head injuries in children. *Seminars in Pediatric Surgery*. 2010; 19(4):279–285. S1055-8586(10)00044-2. [PubMed: 20889084]

- Morton RP, Reynolds RM, Ramakrishna R, Levitt MR, Hopper RA, Lee A, Browd SR. Low-dose head computed tomography in children: a single institutional experience in pediatric radiation risk reduction: clinical article. *J Neurosurg Pediatr.* 2013; 12(4):406–410.10.3171/2013.7.PEDS12631 [PubMed: 23971634]
- Narotam PK, Burjonrappa SC, Raynor SC, Rao M, Taylon C. Cerebral oxygenation in major pediatric trauma: its relevance to trauma severity and outcome. *Journal of Pediatric Surgery.* 2006; 41(3): 505–513.10.1016/j.jpedsurg.2005.11.069 [PubMed: 16516625]
- Ngo QN, Ranger A, Singh RN, Kornecki A, Seabrook JA, Fraser DD. External ventricular drains in pediatric patients. *Pediatr Crit Care Med.* 2009; 10(3):346–351.10.1097/PCC.0b013e3181a320cd [PubMed: 19325503]
- Nitecki S, Moir CR. Predictive factors of the outcome of traumatic cervical spine fracture in children. *J Pediatr Surg.* 1994; 29(11):1409–1411. [PubMed: 7844708]
- Padayachy LC, Figaji AA, Bullock MR. Intracranial pressure monitoring for traumatic brain injury in the modern era. *Childs Nervous System.* 2010; 26(4):441–452.10.1007/s00381-009-1034-0
- Pasvogel AE, Miketova P, Moore IM. Differences in CSF phospholipid concentration by traumatic brain injury outcome. *Biological Research Nursing.* 2010; 11(4):325–331.10.1177/1099800409346056
- Perez A, Minces PG, Schnitzler EJ, Agosta GE, Medina SA, Ciraolo CA. Jugular venous oxygen saturation or arteriovenous difference of lactate content and outcome in children with severe traumatic brain injury. *Pediatric Critical Care Medicine.* 2003; 4(1):33–38.10.1097/01.PCC.0000043309.10452.35 [PubMed: 12656539]
- Piatt JH Jr, Neff DA. Hospital care of childhood traumatic brain injury in the United States, 1997–2009: a neurosurgical perspective. *J Neurosurg Pediatr.* 2012; 10(4):257–267.10.3171/2012.7.PEDS11532 [PubMed: 22900487]
- Puccio AM, Fischer MR, Jankowitz BT, Yonas H, Darby JM, Okonkwo DO. Induced normothermia attenuates intracranial hypertension and reduces fever burden after severe traumatic brain injury. *Neurocrit Care.* 2009; 11(1):82–87.10.1007/s12028-009-9213-0 [PubMed: 19337864]
- Pun PB, Lu J, Moochhala S. Involvement of ROS in BBB dysfunction. *Free Radical Research.* 2009; 43(4):348–364.10.1080/10715760902751902 [PubMed: 19241241]
- Purins K, Lewen A, Hillered L, Howells T, Enblad P. Brain tissue oxygenation and cerebral metabolic patterns in focal and diffuse traumatic brain injury. *Front Neurol.* 2014; 5:64.10.3389/fneur.2014.00064 [PubMed: 24817863]
- Ramaiah VK, Sharma D, Ma L, Prathep S, Hoffman NG, Vavilala MS. Admission oxygenation and ventilation parameters associated with discharge survival in severe pediatric traumatic brain injury. *Childs Nerv Syst.* 2013; 29(4):629–634.10.1007/s00381-012-1984-5 [PubMed: 23207977]
- Reddy MK, Labhsetwar V. Nanoparticle-mediated delivery of superoxide dismutase to the brain: an effective strategy to reduce ischemia-reperfusion injury. *FASEB Journal.* 2009; 23(5):1384–1395.10.1096/fj.08-116947 [PubMed: 19124559]
- Robertson CL, Bucci CJ, Fiskum G. Mitochondrial response to calcium in the developing brain. *Brain Research Developmental Brain Research.* 2004; 151(1–2):141–148.10.1016/j.devbrainres.2004.04.007 [PubMed: 15246700]
- Robertson CL, Saraswati M, Fiskum G. Mitochondrial dysfunction early after traumatic brain injury in immature rats. *Journal of Neurochemistry.* 2007; 101(5):1248–1257.10.1111/j.1471-4159.2007.04489.x [PubMed: 17403141]
- Robertson CL, Scafidi S, McKenna MC, Fiskum G. Mitochondrial mechanisms of cell death and neuroprotection in pediatric ischemic and traumatic brain injury. *Experimental Neurology.* 2009; 218(2):371–380.10.1016/j.expneurol.2009.04.030 [PubMed: 19427308]
- Rohlwink UK, Figaji AA. Methods of monitoring brain oxygenation. *Childs Nervous System.* 2010; 26(4):453–464.10.1007/s00381-009-1033-1
- Rohlwink UK, Zwane E, Fieggen AG, Argent AC, le Roux PD, Figaji AA. The relationship between intracranial pressure and brain oxygenation in children with severe traumatic brain injury. *Neurosurgery.* 2012; 70(5):1220–1230. discussion 1231. 10.1227/NEU.0b013e318243fc59 [PubMed: 22134142]

- Rosenthal G, Hemphill JC 3rd, Sorani M, Martin C, Morabito D, Obrist WD, Manley GT. Brain tissue oxygen tension is more indicative of oxygen diffusion than oxygen delivery and metabolism in patients with traumatic brain injury. *Critical Care Medicine*. 2008; 36(6):1917–1924.10.1097/CCM.0b013e3181743d77 [PubMed: 18496376]
- Ruf B, Heckmann M, Schroth I, Hugens-Penzel M, Reiss I, Borkhardt A, Jodicke A. Early decompressive craniectomy and duraplasty for refractory intracranial hypertension in children: results of a pilot study. *Crit Care*. 2003; 7(6):R133–138.10.1186/cc2361 [PubMed: 14624687]
- Shao J, Zhu H, Yao H, Stallones L, Yeates K, Wheeler K, Xiang H. Characteristics and trends of pediatric traumatic brain injuries treated at a large pediatric medical center in China, 2002–2011. *PLoS One*. 2012; 7(12):e51634.10.1371/journal.pone.0051634 [PubMed: 23251600]
- Staal JA, Dickson TC, Gasperini R, Liu Y, Foa L, Vickers JC. Initial calcium release from intracellular stores followed by calcium dysregulation is linked to secondary axotomy following transient axonal stretch injury. *Journal of Neurochemistry*. 2010; 112(5):1147–1155.10.1111/j.1471-4159.2009.06531.x [PubMed: 19968758]
- Steinborn M, Schaffeler C, Kabs C, Kraus V, Rudisser K, Hahn H. CT and MR imaging of primary cerebrovascular complications in pediatric head trauma. *Emergency Radiology*. 2010; 17(4):309–315.10.1007/s10140-010-0860-4 [PubMed: 20127265]
- Stevens WJ. Multimodal monitoring: head injury management using SjvO₂ and LICOX. *Journal of Neuroscience Nursing*. 2004; 36(6):332–339. [PubMed: 15673209]
- Stewart OW. Contrecoup Lesions in the Relatively Severe Craniocerebral Injuries. *Proc R Soc Med*. 1944; 37(7):381–383. [PubMed: 19992856]
- Stiefel MF, Tomita Y, Marmarou A. Secondary ischemia impairing the restoration of ion homeostasis following traumatic brain injury. *Journal of Neurosurgery*. 2005; 103(4):707–714.10.3171/jns.2005.103.4.0707 [PubMed: 16266054]
- Stiefel MF, Udoetuk JD, Storm PB, Sutton LN, Kim H, Dominguez TE, Huh JW. Brain tissue oxygen monitoring in pediatric patients with severe traumatic brain injury. *Journal of Neurosurgery*. 2006; 105(4 Suppl):281–286.10.3171/ped.2006.105.4.281 [PubMed: 17328278]
- Stippler M, Ortiz V, Adelson PD, Chang YF, Tyler-Kabara EC, Wisniewski SR, Bell MJ. Brain tissue oxygen monitoring after severe traumatic brain injury in children: relationship to outcome and association with other clinical parameters. *J Neurosurg Pediatr*. 2012; 10(5):383–391.10.3171/2012.8.PEDS12165 [PubMed: 22978637]
- Strauss KJ, Goske MJ, Kaste SC, Bulas D, Frush DP, Butler P, Applegate KE. Image gently: Ten steps you can take to optimize image quality and lower CT dose for pediatric patients. *AJR Am J Roentgenol*. 2010; 194(4):868–873.10.2214/AJR.09.4091 [PubMed: 20308484]
- Suskauer SJ, Huisman TA. Neuroimaging in pediatric traumatic brain injury: current and future predictors of functional outcome. *Developmental Disabilities Research Reviews*. 2009; 15(2):117–123.10.1002/ddr.62 [PubMed: 19489082]
- Thomale UW, Graetz D, Vajkoczy P, Sarrafzadeh AS. Severe traumatic brain injury in children—a single center experience regarding therapy and long-term outcome. *Childs Nerv Syst*. 2010; 26(11):1563–1573.10.1007/s00381-010-1103-4 [PubMed: 20177687]
- Tontisirin N, Armstead W, Waitayawinyu P, Moore A, Udomphorn Y, Zimmerman JJ, Vavilala MS. Change in cerebral autoregulation as a function of time in children after severe traumatic brain injury: a case series. *Childs Nervous System*. 2007; 23(10):1163–1169.10.1007/s00381-007-0339-0
- Top AP, Tasker RC, Ince C. The microcirculation of the critically ill pediatric patient. *Crit Care*. 2011; 15(2):213.10.1186/cc9995 [PubMed: 21457503]
- Trabold F, Meyer PG, Blanot S, Carli PA, Orliaguet GA. The prognostic value of transcranial Doppler studies in children with moderate and severe head injury. *Intensive Care Medicine*. 2004; 30(1):108–112.10.1007/s00134-003-2057-8 [PubMed: 14600812]
- Unterberg AW, Stover J, Kress B, Kiening KL. Edema and brain trauma. *Neuroscience*. 2004; 129(4):1021–1029.10.1016/j.neuroscience.2004.06.046 [PubMed: 15561417]
- Ushewokunze S, Sgouros S. Brain tissue oxygenation changes in children during the first 24 h following head injury. *Childs Nervous System*. 2009; 25(3):341–345.10.1007/s00381-008-0752-z

- Vavilala MS, Lee LA, Boddu K, Visco E, Newell DW, Zimmerman JJ, Lam AM. Cerebral autoregulation in pediatric traumatic brain injury. *Pediatr Crit Care Med*. 2004; 5(3):257–263.10.1097/01.PCC.0000123545.69133.C3 [PubMed: 15115564]
- Verlhac S. Transcranial Doppler in children. *Pediatric Radiology*. 2011; 41(Suppl 1):S153–165.10.1007/s00247-011-2038-y [PubMed: 21523592]
- Vernon-Levett, P. Neurologic System. In: Slota, M., editor. *Core Curriculum for Pediatric Critical Care Nursing*. 2. Pittsburgh: Saunders; 2006.
- Ward AA Jr. The physiology of concussion. *Clin Neurosurg*. 1964; 12:95–111. [PubMed: 5865053]
- Weed L. Some limitations of the Monro-Kellie hypothesis. *Archives of Surgery*. 1929; 18(4):1049–1068.
- Weintraub D, Williams BJ, Jane J Jr. Decompressive craniectomy in pediatric traumatic brain injury: a review of the literature. *NeuroRehabilitation*. 2012; 30(3):219–223.10.3233/NRE-2012-0748 [PubMed: 22635127]
- Wiegand C, Richards P. Measurement of intracranial pressure in children: a critical review of current methods. *Developmental Medicine and Child Neurology*. 2007; 49(12):935–941.10.1111/j.1469-8749.2007.00935.x [PubMed: 18039242]
- Wilensky EM, Bloom S, Leichter D, Verdiramo AM, Ledwith M, Stiefel M, Grady MS. Brain tissue oxygen practice guidelines using the LICOX CMP monitoring system. *Journal of Neuroscience Nursing*. 2005; 37(5):278–288. [PubMed: 16379135]
- Yaron M, Niermeyer S, Lindgren KN, Honigman B, Strain JD, Cairns CB. Physiologic response to moderate altitude exposure among infants and young children. *High Alt Med Biol*. 2003; 4(1): 53–59.10.1089/152702903321488988 [PubMed: 12713712]
- Yoxall CW, Weindling AM, Dawani NH, Peart I. Measurement of cerebral venous oxyhemoglobin saturation in children by near-infrared spectroscopy and partial jugular venous occlusion. *Pediatric Research*. 1995; 38(3):319–323. [PubMed: 7494653]
- Zuluaga MT, Esch ME, Cvijanovich NZ, Gupta N, McQuillen PS. Diagnosis influences response of cerebral near infrared spectroscopy to intracranial hypertension in children. *Pediatr Crit Care Med*. 2010; 11(4):514–522.10.1097/PCC.0b013e3181c51cf1 [PubMed: 19935441]

Table 1

Recommended Average Mean Arterial Pressures by Age

Age in years	MAP (mmHg)
1–2 years	50–70
3–4 years	60–75
5–6 years	65–75
7–8 years	70–75
9–10 years	70–75
11–12 years	70–80
13–14 years	80–90

Hazinski, 2013; Top, Tasker, & Ince, 2011

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Signs and Symptoms of Increasing Intracranial Pressure

Early signs and symptoms
Headache
Vomiting
Change in alertness
Irritability
Decreased eye contact
Cranial nerve dysfunction
Seizures
Decrease in Glasgow Coma Scale score

Late signs and symptoms
Further deterioration of consciousness
Bulging fontanel
Decreased spontaneous movement
Posturing
Papilledema
Pupil dilation with decreased or no response to light
Increased blood pressure
Irregular respirations
Cushing's triad

Marcoux, 2005

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Treatments for Children with Severe TBI

Treatment	Level of Evidence	Recommendation
Hyperosmolar Therapy	Level II	Hypertonic saline acute bolus dose 6.5 – 10 ml/kg for increased ICP
	Level III	Continuous 3% saline dose 0.1 – 1.0 ml/kg/hr to maintain ICP < 20 mmHg and serum osmolality <360 mOsm/L
Temperature Control	Level III	Avoid hyperthermia
CSF Drainage	Level III	CSF drainage through an EVD may be considered.
Barbiturates	Level III	High-dose barbiturate therapy in cases of refractory intracranial hypertension.
Decompressive craniectomy	Level III	Consider in early signs of neurologic deterioration or herniation or are developing intracranial hypertension refractory to medical management during the early stages of treatment.
Analgesics, sedatives, and neuromuscular blockade	Level III	Etomidate may be considered to control severe intracranial hypertension. Thiopental may be considered to control intracranial hypertension.
Antiseizure prophylaxis	Level III	Prophylactic treatment with phenytoin may be considered to reduce the incidence of early PTS.

ICP = intracranial pressure; CSF = cerebrospinal fluid; EVD = external ventricular device; mL = milliliter; kg = kilogram; hr = hour; PaCO₂ = arterial carbon dioxide; PTS = posttraumatic seizures

Adapted from (Kochanek et al., 2012)