

SomamiR 2.0: a database of cancer somatic mutations altering microRNA–ceRNA interactions

Anindya Bhattacharya^{1,*} and Yan Cui^{2,3,*}

¹Machine Intelligence Unit, Indian Statistical Institute, Kolkata, WB 700108, India, ²Department of Microbiology, Immunology and Biochemistry, University of Tennessee Health Science Center, Memphis, TN 38163, USA and ³Center for Integrative and Translational Genomics, University of Tennessee Health Science Center, Memphis, TN 38163, USA

Received September 15, 2015; Revised October 23, 2015; Accepted October 28, 2015

ABSTRACT

SomamiR 2.0 (<http://compbio.uthsc.edu/SomamiR>) is a database of cancer somatic mutations in microRNAs (miRNA) and their target sites that potentially alter the interactions between miRNAs and competing endogenous RNAs (ceRNA) including mRNAs, circular RNAs (circRNA) and long noncoding RNAs (lncRNA). Here, we describe the recent major updates to the SomamiR database. We expanded the scope of the database by including somatic mutations that impact the interactions between miRNAs and two classes of non-coding RNAs, circRNAs and lncRNAs. Recently, a large number of miRNA target sites have been discovered by newly emerged high-throughput technologies for mapping the miRNA interactome. We have mapped 388 247 somatic mutations to the experimentally identified miRNA target sites. The updated database also includes a list of somatic mutations in the miRNA seed regions, which contain the most important guiding information for miRNA target recognition. A recently developed web-server, miR2GO, was integrated with the database to provide a seamless pipeline for assessing functional impacts of somatic mutations in miRNA seed regions. Data and functions from multiple sources including biological pathways and genome-wide association studies were updated and integrated with SomamiR 2.0 to make it a better platform for functional analysis of somatic mutations altering miRNA–ceRNA interactions.

INTRODUCTION

SomamiR is a database of cancer somatic mutations that potentially alter microRNA (miRNA) targeting and function (1). The miRNAs are small non-coding RNAs, known

for their role as post-transcriptional regulators of protein-coding mRNAs (2,3). Many genetic and somatic mutations impacting miRNA–mRNA interactions have been associated with various diseases including cancers (4–15). Recent studies have shown that miRNAs also interact with non-coding RNAs such as circular RNAs (circRNA) and long non-coding RNAs (lncRNA) (16–18). The lncRNAs, circRNAs and mRNAs with common miRNA target sites compete for miRNA binding and form a complex network of interaction and regulation, commonly known as the competing endogenous RNA (ceRNA) network (19–21). The dysregulation of crosstalk between ceRNAs in the network plays important roles in cancer pathogenesis (22–27). The miRNA target recognition is largely dependent on sequence complementarity between the seed region (nucleotides 2–7 in the mature miRNA sequence) of an miRNA and its target sites on ceRNAs. Mutations in miRNAs (especially the seed regions) and their target sites may alter miRNA–ceRNA interactions and rewire the ceRNA network (28–30). The updated SomamiR database contains somatic mutations in miRNAs and their target sites on three types of ceRNAs: lncRNAs, circRNAs and mRNAs.

The information guiding miRNA recognition is mainly encoded in the seed regions. Mutations that alter the seed region of an miRNA may have large functional effects because they may disrupt the interactions between the miRNA and many of its original targets and may create interactions with new targets (1,11,31–33). We recently developed a web-server, miR2GO, to assess the functional impacts of mutations in miRNA seed regions (34). The updated SomamiR database exploits miR2GO as a tool for functional analysis of miRNA seed mutations.

When we created the SomamiR database in 2012, there were only a small number of experimentally identified miRNA target sites and no somatic mutation was mapped to those sites. However, two important latest advances have made it possible to map hundreds of thousands of somatic mutations to experimentally identified miRNA target sites on various ceRNAs. The first advance is that

*To whom correspondence should be addressed. Tel: +1 901 448 3240; Fax: +1 901 448 7360; Email: ycui2@uthsc.edu
Correspondence may also be addressed to Anindya Bhattacharya. Email: anindyamail123@gmail.com

a large number of miRNA target sites have been identified by the newly emerging high-throughput technologies such as PAR-CLIP (photoactivatable-ribonucleoside-enhanced crosslinking and immunoprecipitation) (35,36), HITS-CLIP (high-throughput sequencing of RNA isolated by crosslinking immunoprecipitation) (37–39) and CLASH (cross linking, ligation and sequencing of hybrids) (40–42). The second is the very rapid growth of somatic mutations discovered by whole-genome sequencing of many types of cancers.

In summary, the large amount of newly available data for miRNA and ceRNA sequences, the miRNA interactome, and cancer genome sequences have enabled us to perform a major update of the SomamiR database and make it a more useful and complete resource for analyzing the functional impacts of cancer somatic mutations in miRNAs and their target sites.

NEW FEATURES

The SomamiR database has been enhanced by several new features:

1. *Experimentally identified miRNA target sites from CLASH, PAR-CLIP and HITS-CLIP data*: In the past few years, high throughput technologies such as CLASH, PAR-CLIP and HITS-CLIP have been developed to map the miRNA interactome (35–42). The number of experimentally identified miRNA target sites have grown exponentially. There have also been a very rapid increase of non-coding somatic mutation data from the cancer genome sequencing projects such as The Cancer Genome Atlas (TCGA; <http://cancergenome.nih.gov>), the International Cancer Genome Consortium (ICGC; <https://dcc.icgc.org>) (43) and the Pediatric Cancer Genome Project (PCGP; <http://www.pediatriccancergenomeproject.org>) (44). As a result, now we can map over 388 000 somatic mutations in experimentally identified miRNA target sites (Table 1).
2. *Somatic mutations in miRNA seeds*: The seed regions of miRNAs are functionally important because they provide guiding information for target recognition. Mutations in the seed regions of miRNAs may cause big changes in their target gene sets (31,45). SomamiR 2.0 contains 181 somatic mutations mapped to miRNA seed regions.
3. *Integration with the miR2GO webserver* (34): Recently we developed a web-based tool, miR2GO (<http://compbio.uthsc.edu/miR2GO>), for the analysis of functional impacts of miRNA seed mutations. We have integrated miR2GO with SomamiR to allow easy assessment of the functional impacts of miRNA seed mutations.
4. *Somatic mutations in lncRNA–miRNA and circRNA–miRNA binding sites*: Recent studies have shown that lncRNAs and circRNAs play important roles in cancer-related biological processes (46–50). Somatic mutations may alter their interactions with miRNAs. The updated SomamiR database contains somatic mutations in lncRNA–miRNA and circRNA–miRNA binding sites (Table 2).

DATA COLLECTION AND DATABASE CONTENT

Somatic mutations in miRNA sequences

Genomic locations of pre-miRNAs and mature miRNAs were downloaded from miRBase (51,52). We found 987 miRNAs containing 2423 somatic mutations in 50 different types of cancers. Among these somatic mutations, 644 were in mature miRNAs and 1779 were in pre-miRNAs. The somatic mutations mapped to miRNA seed regions are expected to be the most consequential as the seed complementarity was found in most of the functional miRNA target bindings (53). We found 181 somatic mutations in miRNA seeds. We created links that automatically send the somatic mutations as queries to the miR2GO webserver, to allow users to easily perform functional analysis of the seed mutations (Figure 1).

Somatic mutations in experimentally identified miRNA target sites: CLASH

We downloaded the miRNA target sites identified by CLASH experiments from starBase (54). The genomic coordinates of the target sites were mapped to human genome build hg38 by applying the liftOver utility from the UCSC Genome Browser (55,56). Somatic mutations in the genomic locations of the target sites were then collected from the COSMIC database (57). We found 31 863 somatic mutations located in the miRNA target sites on 4553 mRNAs and 9 lncRNAs.

Somatic mutations in experimentally identified miRNA target sites: PAR-CLIP and HITS-CLIP

We downloaded the miRNA target sites identified in 21 PAR-CLIP and 13 HITS-CLIP experiments from starBase (54). We then used liftOver to convert the genomic coordinates of the target sites to the current human genome build (hg38). We searched for the genomic locations of target sites for somatic mutations from the COSMIC database (57). Somatic mutations were found in the miRNA target sites on each of the three types of ceRNAs (Table 2):

1. *Target sites on mRNAs*: The transcription start and end locations for all the RefSeq genes were downloaded from ‘refGene’ table in ‘RefSeq Genes’ track of UCSC table browser. The start and end locations of target sites were then compared against the genomic coordinates to identify the targets sites in mRNAs.
2. *Target sites on lncRNAs*: For all the reported lncRNAs in LNCipedia (58), the transcript locations were computed from the genomic intervals, block starts and block sizes that were downloaded as a BED file. The transcript locations were mapped to the human genome build hg38 by applying liftOver. For finding the target sites in lncRNAs, the target sites locations were then compared against the lncRNA transcript locations.
3. *Target sites on circRNAs*: The genomic coordinates of the circRNAs, transcript block starts and block sizes were downloaded from circBase (59). After applying liftOver, the transcript locations were then compared against the target site locations to identify the target sites on circRNAs.

Table 1. Summary of SomamiR 2.0 contents*

Database Contents	Version/Counts
Genome build	GRCH 38
Somatic mutations in pre-miRNA sequences	1779
Somatic mutations in mature miRNA sequences	644
Somatic mutations in miRNA seeds	181
Somatic mutations (records) in experimentally identified miRNA target sites	388 247 (512 047)
Somatic mutations (records) in predicted miRNA target sites	96 181 (2 868 677)
Biological pathways impacted by somatic mutation in miRNA target sites	199
Cancer-associated genes with somatic mutations in their miRNA target sites	3049
Types of cancers	620

*The numbers in the parentheses are the counts of database records for somatic mutations in miRNA target sites. A record consists of a miRNA, a target site of the miRNA and a somatic mutation in the target site. Multiple miRNAs may share same target sites or have overlapping target sites.

Table 2. Somatic mutations in miRNA target sites on the three types of ceRNAs*

	miRNA target site identification method	ceRNA type		
		mRNA	lncRNA	circRNA
Somatic mutations (records) in experimentally identified miRNA target sites	CLASH	31 841 (36 151)	22 (24)	0 (0)
	PAR-CLIP and HITS-CLIP	226 343 (280 665)	48 511 (77 612)	81 530 (117 595)
Somatic mutations (records) in predicted miRNA target sites	TargetScan	67 366 (458 952)	962 (22 806)	18 075 (85 218)
	Seed match	75 583 (2 313 417)	1049 (127 024)	19 549 (428 236)

*The numbers in the parentheses are the counts of database records for somatic mutations in miRNA target sites. A record consists of a miRNA, a target site of the miRNA and a somatic mutation in the target site. Multiple miRNAs may share same target sites or have overlapping target sites.




Figure 1. Overview of data integration in SomamiR 2.0.

Somatic mutations in predicted miRNA target sites

Somatic mutations in predicted miRNA target sites were also identified on the three classes of ceRNAs (Table 2):

1. *Target sites on mRNAs*: The genomic coordinates of the 3' UTRs for all the RefSeq genes were downloaded from the UCSC table browser (55). Somatic mutations in the 3' UTRs were then collected from COSMIC by comparing the genome locations. Mature miRNA sequences were downloaded from miRBase (release 21) and 3' UTR sequences were downloaded from UCSC table browser. For each somatic mutation in a 3' UTR, a reference and a mutated sequence were scanned for perfect sequence complementarity with the six classes of miRNA seeds described by Ellwanger *et al.* (60). The TargetScan context+ score (53) and the PITA score (61) were provided for assessing the impacts of somatic mutations on miRNA binding.
2. *Target sites on lncRNAs*: We downloaded all the lncRNA transcript sequences in FASTA format from LNCipedia (58). Transcript locations in human genome (hg38) were then compared against COSMIC data to identify somatic mutations in lncRNA transcripts. We then determined the effects of somatic mutations on lncRNA–miRNA target sites by applying the six seed matches (60) and TargetScan on lncRNA sequences.
3. *Target sites on circRNAs*: circRNA sequences were downloaded from circBase (59). Somatic mutations were compared against the transcript locations of circRNAs in human genome build hg38. The alterations of target site binding for somatic mutations were determined by using the six seed matches (60) and TargetScan.

Biological pathways impacted by somatic mutations in miRNA target sites

The KEGG pathways (62) were downloaded from the 'keggPathway' table of the UCSC table browser. We found 20 020 genes in 199 pathways that contain somatic mutations in their miRNA target sites. The KEGG API interface is used to display the biological pathways. The genes with somatic mutations in miRNA target sites are highlighted in the pathways (Figure 1).

Genes associated with cancer risk that contain miRNA related somatic mutations

There has been a rapid growth of genome-wide association studies (GWAS) and candidate gene association studies (CGAS) in the past few years. Newly available GWAS results were processed for the update and 2500 new gene–phenotype associations were added to the database. High-scoring markers associated with cancer phenotypes were collected from the UCSC Table Browser (55), NHGRI GWAS Catalog (63) and the Cancer GAMAdb (64).

DATABASE ACCESS AND USAGE

The content of the SomamiR database is accessible through its browse and search interfaces. Six browsing options are provided on the database homepage. Users can browse

the somatic mutations in miRNA sequences, the somatic mutations in the miRNA target sites identified from the CLASH, PAR-CLIP and HITS-CLIP experiments, the somatic mutations in predicted miRNA target sites and the KEGG biological pathways (62) in which genes with somatic mutations in their miRNA target sites are highlighted. Users can also browse the genes associated with cancer phenotypes in genome-wide association studies and candidate gene association studies. The SomamiR database can be searched by chromosome location, miRNA, ceRNA ID and gene symbol. Filtering options such as target ceRNA type and cancer type are made available for efficient browsing and searching. The database contents are shown in sortable and downloadable tables. Each 'Transcript ID' in the tables are linked to a data page showing all the somatic mutations in the miRNA target sites on the selected transcript. In the data page of each transcript, a UCSC Genome Browser window is used to display the somatic mutations and miRNA target sites in the genomic context. Currently, the SomamiR database includes three types of ceRNAs: mRNA, lncRNA and circRNA. To define cancer types, we adopted the hierarchical classification system from COSMIC cancer browser (57). This classification system has four levels: tissue selection, sub-tissue selection, histology selection and sub-histology selection. For example, the cancer type [haematopoietic_and_lymphoid_tissue][lymph_node][lymphoid_neoplasm][Hodgkin_lymphoma] indicates that the tissue is 'haematopoietic and lymphoid', the sub-tissue is 'lymph node', and the histology and sub-histology are 'lymphoid neoplasm' and 'Hodgkin lymphoma', respectively. The database contains somatic mutations from 620 types of cancers (Supplementary Table S1). The entire database contents are downloadable as spreadsheet files from the download link in the database homepage. Detailed information about the SomamiR database is available in the help page of the database.

DISCUSSION

At the time we developed the first version of the SomamiR database, only a few thousand somatic mutations were identified in the predicted miRNA–mRNA binding sites. The rapid drop of sequencing cost led to a very fast growth of somatic mutations from cancer genome sequencing projects in the past few years. Moreover, the emergence of high-throughput miRNA interactome mapping technologies such as CLASH, PAR-CLIP and HITS-CLIP, enabled large-scale experimental identification of miRNA target sites and thereby provided a source of highly reliable miRNA targets for the updated SomamiR database. We expect that the somatic mutations mapped to experimentally identified miRNA target sites will continue to increase rapidly in the future releases of the SomamiR database.

Our knowledge of miRNA regulatory mechanisms has been greatly expanded by recent findings of ceRNA network and crosstalk. Somatic mutations can disrupt and alter the ceRNA crosstalk and thereby contribute to the pathogenesis of cancers. It is very likely that the full scope and depth of miRNA regulation has yet to be discovered. This will provide the opportunity to further expand the

scope of the SomamiR database by including the somatic mutations impacting new types of interactions involving miRNAs.

Both cancer genomics data and miRNA interactome data are growing very rapidly. It becomes increasingly important to automate the data processing and database update. We developed a semi-automatic data curation pipeline for updating the database contents, which will make it easier to keep the database contents up-to-date.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the editorial assistance of Dr Martha M. Howe provided through the Writing Assistance Program of the College of Graduate Health Sciences.

FUNDING

A.B. was supported by the Ramalingaswami Re-entry fellowship, Department of Biotechnology, India [Project Title: Functional Annotation of Somatic Mutations in Cancer, Project Number: BT/RLF/Re-entry/09/2013]. Funding for open access charge: Ramalingaswami Re-entry fellowship from Department of Biotechnology, India [Project Title: Functional Annotation of Somatic Mutations in Cancer, Project Number: BT/RLF/Re-entry/09/2013].
Conflict of interest statement. None declared.

REFERENCES

- Bhattacharya,A., Ziebarth,J.D. and Cui,Y. (2013) SomamiR: a database for somatic mutations impacting microRNA function in cancer. *Nucleic Acids Res.*, **41**, D977–D982.
- Ambros,V. (2004) The functions of animal microRNAs. *Nature*, **431**, 350–355.
- Bartel,D.P. (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, **116**, 281–297.
- Esquela-Kerscher,A. and Slack,F.J. (2006) Oncomirs - MicroRNAs with a role in cancer. *Nat. Rev. Cancer*, **6**, 259–269.
- Mishra,P.J. and Bertino,J.R. (2009) MicroRNA polymorphisms: The future of pharmacogenomics, molecular epidemiology and individualized medicine. *Pharmacogenomics*, **10**, 399–416.
- Landi,D., Gemignani,F. and Landi,S. (2012) Role of variations within microRNA-binding sites in cancer. *Mutagenesis*, **27**, 205–210.
- Ryan,B.M., Robles,A.I. and Harris,C.C. (2010) Genetic variation in microRNA networks: the implications for cancer research. *Nat. Rev. Cancer*, **10**, 389–402.
- Bao,L., Zhou,M., Wu,L., Lu,L., Goldowitz,D., Williams,R.W. and Cui,Y. (2007) PolymiRTS Database: linking polymorphisms in microRNA target sites with complex traits. *Nucleic Acids Res.*, **35**, D51–D54.
- Ziebarth,J.D., Bhattacharya,A., Chen,A. and Cui,Y. (2012) PolymiRTS Database 2.0: linking polymorphisms in microRNA target sites with human diseases and complex traits. *Nucleic Acids Res.*, **40**, D216–D221.
- Bhattacharya,A., Ziebarth,J.D. and Cui,Y. (2014) PolymiRTS Database 3.0: linking polymorphisms in microRNAs and their target sites with human diseases and biological pathways. *Nucleic Acids Res.*, **42**, D86–D91.
- Ziebarth,J.D., Bhattacharya,A. and Cui,Y. (2012) Integrative analysis of somatic mutations altering microRNA targeting in cancer genomes. *PLoS One*, **7**, e47137.
- Cui,Y. (2014) In silico mapping of polymorphic miRNA–mRNA interactions in autoimmune thyroid diseases. *Autoimmunity*, **47**, 327–333.
- Bhattacharya,A., Ziebarth,J.D. and Cui,Y. (2012) Systematic analysis of microRNA targeting impacted by small insertions and deletions in human genome. *PLoS One*, **7**, e46176.
- Bulik-Sullivan,B., Selitsky,S. and Sethupathy,P. (2013) Prioritization of genetic variants in the microRNA regulome as functional candidates in genome-wide association studies. *Hum. Mutat.*, **34**, 1049–1056.
- Gamazon,E.R., Ziliak,D., Im,H.K., LaCroix,B., Park,D.S., Cox,N.J. and Huang,R.S. (2012) Genetic architecture of microRNA expression: Implications for the transcriptome and complex traits. *Am. J. Hum. Genet.*, **90**, 1046–1063.
- Hansen,T.B., Jensen,T.I., Clausen,B.H., Bramsen,J.B., Finsen,B., Damgaard,C.K. and Kjems,J. (2013) Natural RNA circles function as efficient microRNA sponges. *Nature*, **495**, 384–388.
- Wang,K. and Chang,H. (2011) Molecular mechanisms of long noncoding RNAs. *Mol. Cell*, **43**, 904–914.
- Memczak,S., Jens,M., Elefsinioti,A., Torti,F., Krueger,J., Rybak,A., Maier,L., Mackowiak,S.D., Gregersen,L.H., Munschauer,M. et al. (2013) Circular RNAs are a large class of animal RNAs with regulatory potency. *Nature*, **495**, 333–338.
- Salmena,L., Poliseno,L., Tay,Y., Kats,L. and Pandolfi,P.P. (2011) A ceRNA hypothesis: the rosetta stone of a hidden RNA language? *Cell*, **146**, 353–358.
- Paci,P., Colombo,T. and Farina,L. (2014) Computational analysis identifies a sponge interaction network between long non-coding RNAs and messenger RNAs in human breast cancer. *BMC Syst. Biol.*, **8**, 83.
- Chiu,H.-S., Llobet-Navas,D., Yang,X., Chung,W.-J., Ambesi-Impiombato,A., Iyer,A., Kim,H.R., Seviour,E.G., Luo,Z., Sehgal,V. et al. (2014) Cupid: simultaneous reconstruction of microRNA-target and ceRNA networks. *Genome Res.*, **25**, 257–267.
- Tay,Y., Kats,L., Salmena,L., Weiss,D., Tan,S.M., Ala,U., Karreth,F., Poliseno,L., Provero,P., Di Cunto,F. et al. (2011) Coding-independent regulation of the tumor suppressor PTEN by competing endogenous mRNAs. *Cell*, **147**, 344–357.
- Peng,W., Si,S., Zhang,Q., Li,C., Zhao,F., Wang,F., Yu,J. and Ma,R. (2015) Long non-coding RNA MEG3 functions as a competing endogenous RNA to regulate gastric cancer progression. *J. Exp. Clin. Cancer Res.*, **34**, 79.
- Yue,B., Sun,B., Liu,C., Zhao,S., Zhang,D., Yu,F. and Yan,D. (2015) Long non-coding RNA FER1L4 suppresses oncogenesis and exhibits prognostic value by associating with miR-106a-5p in colon cancer. *Cancer Sci.*, **106**, 1323–1332.
- Sanchez-Mejias,A. and Tay,Y. (2015) Competing endogenous RNA networks: tying the essential knots for cancer biology and therapeutics. *J. Hematol. Oncol.*, **8**, 30.
- Xu,J., Li,Y., Lu,J., Pan,T., Ding,N., Wang,Z., Shao,T., Zhang,J., Wang,L. and Li,X. (2015) The mRNA related ceRNA–ceRNA landscape and significance across 20 major cancer types. *Nucleic Acids Res.*, **43**, 8169–8182.
- Sumazin,P., Yang,X., Chiu,H.S., Chung,W.J., Iyer,A., Llobet-Navas,D., Rajbhandari,P., Bansal,M., Guarnieri,P., Silva,J. et al. (2011) An extensive microRNA-mediated network of RNA–RNA interactions regulates established oncogenic pathways in glioblastoma. *Cell*, **147**, 370–381.
- Ergun,S. and Oztuzcu,S. (2015) Oncocers: ceRNA-mediated cross-talk by sponging miRNAs in oncogenic pathways. *Tumour Biol.*, **36**, 3129–3136.
- Tay,Y., Rinn,J. and Pandolfi,P.P. (2014) The multilayered complexity of ceRNA crosstalk and competition. *Nature*, **505**, 344–352.
- Karreth,F.A. and Pandolfi,P.P. (2013) ceRNA cross-talk in cancer: when ce-bling rivalries go awry. *Cancer Discov.*, **3**, 1113–1121.
- Hill,C.G., Jabbari,N., Matyunina,L.V. and McDonald,J.F. (2014) Functional and evolutionary significance of human microRNA seed region mutations. *PLoS One*, **9**, e115241.
- Mencia,A., Modamio-Hoybjor,S., Redshaw,N., Morin,M., Mayo-Merino,F., Olavarrieta,L., Aguirre,L.A., del Castillo,I., Steel,K.P., Dalmay,T. et al. (2009) Mutations in the seed region of human miR-96 are responsible for nonsyndromic progressive hearing loss. *Nat. Genet.*, **41**, 609–613.

33. Iliff, B.W., Riazuddin, S.A. and Gottsch, J.D. (2012) A single-base substitution in the seed region of miR-184 causes EDICT syndrome. *Invest. Ophthalmol. Vis. Sci.*, **53**, 348–353.
34. Bhattacharya, A. and Cui, Y. (2015) miR2GO: comparative functional analysis for microRNAs. *Bioinformatics*, **31**, 2403–2405.
35. Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Hausser, J., Berninger, P., Rothballer, A., Ascano, M. Jr, Jungkamp, A.C., Munschauer, M. *et al.* (2010) Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell*, **141**, 129–141.
36. Hafner, M., Lianoglou, S., Tuschl, T. and Betel, D. (2012) Genome-wide identification of miRNA targets by PAR-CLIP. *Methods*, **58**, 94–105.
37. Chi, S.W., Zang, J.B., Mele, A. and Darnell, R.B. (2009) Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature*, **460**, 479–486.
38. Moore, M.J., Zhang, C., Gantman, E.C., Mele, A., Darnell, J.C. and Darnell, R.B. (2014) Mapping Argonaute and conventional RNA-binding protein interactions with RNA at single-nucleotide resolution using HITS-CLIP and CIMS analysis. *Nat. Protoc.*, **9**, 263–293.
39. Darnell, R.B. (2010) HITS-CLIP: panoramic views of protein–RNA regulation in living cells. *Wiley Interdiscip. Rev. RNA*, **1**, 266–286.
40. Kudla, G., Granneman, S., Hahn, D., Beggs, J.D. and Tollervey, D. (2011) Cross-linking, ligation, and sequencing of hybrids reveals RNA-RNA interactions in yeast. *Proc. Natl. Acad. Sci. U.S.A.*, **108**, 10010–10015.
41. Helwak, A., Kudla, G., Dudnakova, T. and Tollervey, D. (2013) Mapping the human miRNA interactome by CLASH reveals frequent noncanonical binding. *Cell*, **153**, 654–665.
42. Helwak, A. and Tollervey, D. (2014) Mapping the miRNA interactome by cross-linking ligation and sequencing of hybrids (CLASH). *Nat. Protoc.*, **9**, 711–728.
43. Hudson, T.J., Anderson, W., Artez, A., Barker, A.D., Bell, C., Bernabe, R.R., Bhan, M.K., Calvo, F., Eerola, I., Gerhard, D.S. *et al.* (2010) International network of cancer genome projects. *Nature*, **464**, 993–998.
44. Downing, J.R., Wilson, R.K., Zhang, J., Mardis, E.R., Pui, C.H., Ding, L., Ley, T.J. and Evans, W.E. (2012) The Pediatric Cancer Genome project. *Nat. Genet.*, **44**, 619–622.
45. Bhattacharya, A. and Cui, Y. (2015) Knowledge-based analysis of functional impacts of mutations in microRNA seed regions. *J. Biosci.*, **40**, 791–798.
46. Hansen, T.B., Kjems, J. and Damgaard, C.K. (2013) Circular RNA and miR-7 in Cancer. *Cancer Res.*, **73**, 5609–5612.
47. Qu, S., Yang, X., Li, X., Wang, J., Gao, Y., Shang, R., Sun, W., Dou, K. and Li, H. (2015) Circular RNA: a new star of noncoding RNAs. *Cancer Lett.*, **365**, 141–148.
48. Spizzo, R., Almeida, M.I., Colombatti, A. and Calin, G.A. (2012) Long non-coding RNAs and cancer: a new frontier of translational research. *Oncogene*, **31**, 4577–4587.
49. Prensner, J.R. and Chinnaiyan, A.M. (2011) The emergence of lncRNAs in cancer biology. *Cancer Discov.*, **1**, 391–407.
50. Wapinski, O. and Chang, H.Y. (2011) Long noncoding RNAs and human disease. *Trends Cell Biol.*, **21**, 354–361.
51. Kozomara, A. and Griffiths-Jones, S. (2011) miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res.*, **39**, D152–D157.
52. Kozomara, A. and Griffiths-Jones, S. (2014) miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res.*, **42**, D68–D73.
53. Garcia, D.M., Baek, D., Shin, C., Bell, G.W., Grimson, A. and Bartel, D.P. (2011) Weak seed-pairing stability and high target-site abundance decrease the proficiency of lsy-6 and other microRNAs. *Nat. Struct. Mol. Biol.*, **18**, 1139–1146.
54. Li, J.H., Liu, S., Zhou, H., Qu, L.H. and Yang, J.H. (2014) starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. *Nucleic Acids Res.*, **42**, D92–D97.
55. Karolchik, D., Hinrichs, A.S., Furey, T.S., Roskin, K.M., Sugnet, C.W., Haussler, D. and Kent, W.J. (2004) The UCSC Table Browser data retrieval tool. *Nucleic Acids Res.*, **32**, D493–D496.
56. Karolchik, D., Barber, G.P., Casper, J., Clawson, H., Cline, M.S., Diekhans, M., Dreszer, T.R., Fujita, P.A., Gurusvadoo, L., Haussler, M. *et al.* (2014) The UCSC Genome Browser database: 2014 update. *Nucleic Acids Res.*, **42**, D764–D770.
57. Forbes, S.A., Beare, D., Gunasekaran, P., Leung, K., Bindal, N., Boutselakis, H., Ding, M., Bamford, S., Cole, C., Ward, S. *et al.* (2015) COSMIC: exploring the world’s knowledge of somatic mutations in human cancer. *Nucleic Acids Res.*, **43**, D805–D811.
58. Volders, P.J., Verheggen, K., Menschaert, G., Vandepoele, K., Martens, L., Vandesompele, J. and Mestdagh, P. (2015) An update on LNCpedia: a database for annotated human lncRNA sequences. *Nucleic Acids Res.*, **43**, D174–D180.
59. Glazar, P., Papavasileiou, P. and Rajewsky, N. (2014) circBase: a database for circular RNAs. *RNA*, **20**, 1666–1670.
60. Ellwanger, D.C., Buttner, F.A., Mewes, H.W. and Stumpflen, V. (2011) The sufficient minimal set of miRNA seed types. *Bioinformatics*, **27**, 1346–1350.
61. Kertesz, M., Iovino, N., Unnerstall, U., Gaul, U. and Segal, E. (2007) The role of site accessibility in microRNA target recognition. *Nat. Genet.*, **39**, 1278–1284.
62. Kanehisa, M., Goto, S., Sato, Y., Kawashima, M., Furumichi, M. and Tanabe, M. (2014) Data, information, knowledge and principle: back to metabolism in KEGG. *Nucleic Acids Res.*, **42**, D199–D205.
63. Hindorf, L.A., Sethupathy, P., Junkins, H.A., Ramos, E.M., Mehta, J.P., Collins, F.S. and Manolio, T.A. (2009) Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proc. Natl. Acad. Sci. U.S.A.*, **106**, 9362–9367.
64. Schully, S.D., Yu, W., McCallum, V., Benedicto, C.B., Dong, L.M., Wulf, A., Clyne, M. and Houry, M.J. (2011) Cancer GAMAdb: database of cancer genetic associations from meta-analyses and genome-wide association studies. *Eur. J. Hum. Genet.*, **19**, 928–930.