



# HHS Public Access

Author manuscript

*Biochim Biophys Acta*. Author manuscript; available in PMC 2016 November 01.

Published in final edited form as:

*Biochim Biophys Acta*. 2015 November ; 1848(11 0 0): 2985–2997. doi:10.1016/j.bbamem.2015.08.008.

## Equilibrium Fluctuation Relations for Voltage Coupling in Membrane Proteins

Ilsoo Kim and Arieh Warshel<sup>1</sup>

Department of Chemistry, University of Southern California, SGM 418, 3620 McClintock Avenue, Los Angeles, CA 900089, USA

### Abstract

A general theoretical framework is developed to account for the effects of an external potential on the energetics of membrane proteins. The framework is based on the free energy relation between two (forward/backward) probability densities, which was recently generalized to non-equilibrium processes, culminating in the work-fluctuation theorem. Starting from the probability densities of the conformational states along the reaction coordinate of “voltage coupling”, we investigate several interconnected free energy relations between these two conformational states, considering voltage activation of ion channels. The free energy difference at zero membrane potential (i.e., between the two “non-equilibrium” conformational states) is shown to be equivalent to the free energy difference between the two “equilibrium” conformational states along the one-dimensional reaction coordinate of voltage coupling. Furthermore, the requirement that the application of linear response approximation to the free energy functions (free energies) of voltage coupling should satisfy the general free energy relations, yields a novel expression for the gating charge in terms of other experimentally measurable quantities. This connection is familiar in statistical mechanics, known as the equilibrium fluctuation-response relation. The theory is illustrated by considering the movement of a unit charge within the membrane under the influence of an external potential, using a coarse-graining (CG) model of membrane proteins, which includes the membrane, the electrolytes and the electrodes. The CG model yields Marcus-type voltage dependent free energy parabolas for the two conformational states, which allow for quantitative estimations of an equilibrium free energy difference, a free energy of barrier, and the voltage dependency of channel activation (Q-V curve) for the unit charge movement. In addition, our analysis offers a quantitative rationale for the correlation between the free energy landscapes (parabolas) and the Q-V curve, upon site-directed mutagenesis or drug binding. Taken together, by introducing the voltage coupling as a reaction coordinate of energy gap, the present theory offers a firm physical foundation from the equilibrium theory of statistical mechanics for the thermodynamic models of voltage activation in voltage-sensitive membrane proteins. This formulation also provides a powerful bridge between the CG model and the conventional macroscopic treatments, offering an intuitive and quantitative framework for a better understating of the structure-function correlations

<sup>1</sup>Correspondence should be addressed to: Arieh Warshel (warshel@usc.edu).

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of voltage gating in ion channels as well as electrogenic phenomena in ion pumps and transporters.

---

## 1. Introduction

The advances in structural elucidation of voltage activated ion channels as well as in biophysical studies (e.g. Refs. [1–7]) have provided major clues about the relationship between the membrane voltage and the gating process. However, despite these great progress we still do not have a clear understanding of the corresponding structure-function correlation. Apparently, although there have been a significant progress in the computational/theoretical modeling of the energetics of ion channels (e.g. Refs. [8–22]) and the control of ion selectivity (e.g. Refs. [23–30]), the quantitative understanding of the voltage activation process is still somewhat limited. In addition to the obvious need for more structural information, the ability to obtain a microscopic description of the energetics of the conformational transition and the coupling to the external voltage is far from satisfactory. Similar problems occur with regards to the molecular understanding of the nature of the gating charge, where despite the enormous insight provided by macroscopic approaches [13, 14, 31], it is hard to be fully comfortable with the corresponding physical picture, which does not include the electrolytes and the electrodes explicitly (see discussion in Refs. [20, 32]).

A promising way for advancing our understanding of ion channels and related systems has been offered by the recent development of our coarse-grained (CG) model of voltage coupling [20, 21, 32, 33], which considers the entire membrane–protein, electrolytes and electrodes explicitly. The power and insight of this model have been illustrated in several works, but it seems that these advances have not been widely recognized, due in part to the use of descriptions that are very different than the familiar macroscopic formulation and the fact that the model does not use the straightforward fully microscopic treatment (that unfortunately does not offers yet converging free energy results [17, 34] nor a clear description of the nature of gating charge [35]). Thus we try to explore in the present work the relationship between our CG model of voltage coupling and the corresponding macroscopic continuum results.

Furthermore, more importantly, we introduce the general theoretical description of voltage coupling in membrane proteins, extending and generalizing the previous thermodynamic models of voltage activation in ion channels [1,19, 36–41]. This is done in the framework of equilibrium fluctuation relations, where linear response approximation to the free energy function of voltage coupling is introduced, resulting in the Marcus-Warshel (MW) type parabola, with the energy gap of voltage coupling as a reaction coordinate. A novel closed expression for the gating charge is then derived in the framework of the equilibrium fluctuation relations and linear response approximation and explored by using the CG model of voltage coupling. The corresponding results are validated against the macroscopic continuum results and applied to determining voltage dependent MW parabola for the movement of a unit charge within the membrane. We then use the MW type free energy parabolas in the framework of equilibrium fluctuation relations to formulate the quantitative

relationship between the free energy landscapes and the Q-V curves, thus providing a new insight on the molecular information content of these curves.

## 2. Theory and methods

### 2.1. Free energy relations

We will start by defining formal free energy relationships that will result in free energy functions of the energy gap reaction coordinate, introduced by Warshel [36] and used extensively in studies of reactions in condensed phase (e.g. [37–41]). This will be done while exploring several variants of free energy relations in the general framework of equilibrium fluctuation relation.

The equilibrium probability densities for the initial (0) and the final (1) states ( $p_\lambda(u)$  { $\lambda = 0, 1$ }) are related to each other [42], given by:

$$\frac{p_0(u)}{p_1(u)} = \exp(u - \Delta G_{conf}), \quad (1)$$

where  $k_B T = 1$  and  $G_{conf}$  is the free energy difference between, for example, two conformational states in the present study. This relationship is referred to “the equilibrium fluctuation relation”, in an analogy with its generalization to non-equilibrium processes, known as the work-fluctuation theorem [43, 44]).

The equilibrium fluctuation relation is easily derived (see SI for a detailed derivation) by considering the system as being perturbed from its initial to final states by  $U$ , with the Hamiltonian of the form  $H_1 = H_0 + U$ :

$$\begin{aligned} p_0(u) &\equiv \langle \delta(u - \Delta U(x)) \rangle_0 \\ &= \frac{\int dx \exp(-H_0(x)) \delta(u - \Delta U(x))}{Z_0} \\ &= \frac{Z_1}{Z_0} \exp(u) \frac{\int dx \exp(-H_1(x)) \delta(u - \Delta U(x))}{Z_1} \quad (2) \\ &= \exp(u - \Delta G_{conf}) p_1(u) \end{aligned}$$

where  $p_\lambda(u)$  { $\lambda = 0, 1$ } are the probability densities of finding a particular value of  $u$  along a reaction coordinate  $U(x)$ , where  $x$  is a point in the  $6N$  – dimensional phase space. In fact, this relation has been known to the community for some time ago as the theory of Bennett overlapping histogram (BOH) [41, 45], which is also here referred to the free energy relation of the first kind. In the following, the free energy difference between the two conformational states ( $G_{conf}$ ) is defined by two different ways, eventually leading to an equivalent relation.

Notice that  $G_{conf}$  is identified as the value ( $u^*$ ) of reaction coordinate  $U$  where the two probability densities intersect [41, 46]:

$$\frac{p_0(u^*)}{p_1(u^*)} = 1 = \exp(u^* - \Delta G_{conf}),$$

leading to

$$\Delta G_{conf} = U^* = \Delta U(x^*) \equiv \Delta U^* \quad (3)$$

Rearranging Eq. (1) yields the free energy relation of the second kind:

$$\Delta f_1(u) - \Delta f_0(u) = u + \ln \frac{p_{1,\max}}{p_{0,\max}} \quad (4)$$

or

$$\Delta g_1(u) - \Delta g_0(u) = u - \Delta G_{conf} + \ln \frac{p_{1,\max}}{p_{0,\max}}, \quad (5)$$

where  $g_\lambda(u)$  and  $f_\lambda(u)$  ( $\lambda = 0, 1$ ) are free energy functions or potentials of mean force (PMF) along the reaction coordinate of energy gap [36, 37, 41], which are related to each other by:

$$\Delta f_1(u) \equiv \Delta g_1(u) + \Delta G_{conf} \equiv -\ln \frac{p_1(u)}{p_{1,\max}} + \Delta G_{conf} \quad (6)$$

$$\Delta f_0(u) \equiv \Delta g_0(u) = -\ln \frac{p_0(u)}{p_{0,\max}} \quad (7)$$

Here,  $p_\lambda^{\max}$  ( $\lambda = 0, 1$ ) are defined as the maximum values of the probability densities, such that  $g_\lambda$  ( $\lambda = 0, 1$ ) have their respective global minimum set to zero [41], i.e.,  $g_\lambda(u_{\lambda,\min}) = -\ln[p_\lambda(u_{\lambda,\min})/p_{\lambda,\max}] = 0$ , which results in the free energy relation of third kind:

$$\Delta f_1(u_{1,\min}) - \Delta f_0(u_{0,\min}) = \Delta G_{conf} \quad (8)$$

This expression states that the free energy difference between the two equilibrium conformational states along a one-dimensional reaction coordinate of energy gap, ( $f_1(u_{1,\min}) - f_0(u_{0,\min})$ ), defined from the second free energy relation, is equal to the free energy difference,  $G_{conf}(= U^*)$ , determined using the equilibrium fluctuation relation or the first free energy relation. In other words, the free energy difference,  $G_{conf}(= U^*)$ , defined in two different ways from Eqs. (3) and (4) or (5), leads to an equivalent relation of Eq. (8), which is thus termed as the conformational free energy along the one-dimensional reaction coordinate of energy gap. A closely related derivation to Eq. (8) along “positional” reaction coordinates is found in Ref. [41] and applied to the thermodynamics of ion binding in a  $K^+$  channel [28], supporting the earlier proposal of the multi-ion mechanism of ion selectivity in  $K^+$  channels [27].

The additive constant on the right hand side in Eq. (4) or (5) vanishes in the case where  $g_\lambda$  ( $\lambda = 0, 1$ ) are functions with equal maximum values (curvatures) of the probability

densities at their respective minimum (i.e.,  $p_1^{\max}=p_0^{\max}$ ). In the present study, the constant will be dropped out without a loss of generality [38, 41] and the approximation introduced here will be discussed in conjunction with the linear response expression for free energy functions (see Section 2.3). The second free energy relation of Eq. (4) was first recognized by Warshel and others [37–39] in formulating microscopic treatment of electron transfer in condensed phases and relating it to the framework of the macroscopic Marcus theory [47, 48]

Note that the free energy functions ( $f_\lambda(u)\{\lambda=0,1\}$ ), whose final state ( $\lambda=1$ ) is shifted by the free energy ( $G_{conf}$ ) from the reference free energy function ( $g_1(u)$ ), intersect at  $u=0$  along the energy gap coordinate [37–39], leading to:

$$\Delta f_1(0)=\Delta f_0(0) \quad (9)$$

or

$$\Delta g_1(0)-\Delta g_0(0)=-\Delta G_{conf} \quad (10)$$

The free energy relations examined so far is illustrated in Fig. 1 with  $U=Q_g V$ , where  $Q_g$  and  $V$  are, respectively, the gating charge and the externally applied potential. This will be further investigated in the following section, with the energy gap of “voltage coupling” in ion channels to derive the fundamental free energy relations (see e.g. Ref.[1]) used to describe the voltage dependency of channel activation in ion channels.

## 2.2 Free energy relations for voltage activation in ion channels

The general free energy relations investigated in the previous section can be applied to account for the kinetics/thermodynamics of voltage activation of ion channels by using the voltage dependent free energy functions. As a background for such considerations, we start by noting that the effect of an external potential on the activation of the voltage sensor domain (VSD) in ion channels is described by the voltage coupling to the gating charge  $Q_g$ . This coupling reflects the response of the system to the application of an external potential but before the ions are allowed to pass through the channel [1]. The gating charge,  $Q_g$ , is defined by determining the fraction of activated (up) and resting (down) channels as a function of the applied potential and asking what is the Boltzmann probability for the voltage induced structural change [1]. This assumption leads to the expression [1]:

$$Q_g \Delta V_{ext} = \Delta G^{down \rightarrow up}, \quad (11)$$

where  $G^{down \rightarrow up}$  is the phenomenological free energy associated with conformational change between two “equilibrium” states (i.e., the conformational free energy of Eq. 8), which can be measured experimentally in Q-V measurements [1]. Basically this is equivalent to the assumption that the free energy needed to move the gating charge,  $Q_g$ , in the membrane electric field is equal to the work of moving the protein charges between the two conformations (activated and resting), under the membrane electric field. Now, Eq. (11) is basically a reasonable formal definition of the gating charge that does not tell us what is

the relationship of this parameter to physical measured observables such as the integral of the current that flows to the electrodes,  $Q_{\text{ext}}$ , although many accept the identity of  $Q_{\text{ext}}$  with  $Q_g$ . The structure-based evaluation of  $Q_g$  is almost always done using reasonable but not necessarily microscopic assumptions. It is also not certain that in all cases that the actual electrode potential  $V_{\text{ext}}$  is equal to the membrane potential,  $V_m$ . These issues will be addressed below but at this point we just note that one of the problem is that the assumption that leads to Eq. (11) implies that the potential is linear across the membrane. This is likely to be a reasonable approximation and it has been used in previous macroscopic studies [13, 49]. Obviously, such a picture would be justified if the potential across the protein/membrane system was obtained by converging microscopic simulations. However, in such cases, it is unlikely that we will have the same potential in different sites with the same  $z$  value that is normal to the membrane surface. Thus such a treatment does not really provide a microscopic description even if a few of the relevant quantities (e.g. the average displacements) are evaluated microscopically [17]. Other related problems are discussed in our previous work [32].

In view of the above discussion we can write

$$\begin{aligned} Q_{\text{ext}} &= c_1 Q_{g,m} \\ V_{\text{ext}} &= c_2 V_m, \end{aligned} \quad (12)$$

where  $Q_{g,m}$  is the charge movement within the membrane and  $V_m$  is the change in the potential across the membrane. However, for simplicity we will assume below that  $c_2$  and  $c_1$  are approximately equal to one, while exploring this assumption by examining the correspondence between the energetics and charge movement, considering both only the membrane-protein system and the entire electrolyte/membrane-protein system. We start by considering only the membrane-protein system, whose Hamiltonian is given by [19]:

$$H_{1,m} = H_{0,m} + \Delta U \equiv H_{0,m} + Q_{g,m} V_m, \quad (13)$$

where the subscripts 0 and 1 represent initial (activated) and final (resting) states, respectively, of, for example, VSD. The gating charge ( $Q_{g,m}$ ) arises from the displacement of charged residues within membrane associated with conformational changes of VSD by  $V_m$ .

By taking an energy gap reaction coordinate of ( $U = Q_{g,m} V_m$ ) (referred to as the ‘‘voltage coupling’’), we obtain from Eq. (3):

$$\Delta G_{\text{conf}} = Q_{g,m} V_m^* = Q_{g,m} V_{m,1/2}, \quad (14)$$

where  $V_{m,1/2}$  is a half voltage at which two conformational states have equal population. This relation along with Eqs. (19) and (20) is basically the same as the phenomenological free energy of conformational change of Eq. (11).

The equilibrium fluctuation relation or the first free energy relation of Eq. (1) for voltage activation is therefore expressed as:

$$\frac{p_0(V)}{p_1(V)} = \exp(Q_{g,m}V_m - Q_{g,m}V_{m,1/2}) \quad (15)$$

In fact, this fundamental relation has been known to the ion channel community to describe the kinetics/thermodynamics of voltage activation, e.g., as empirically derived in Hille's book [1], which goes back to Hodgkin and Huxley for their formulation to quantify membrane currents and thus action potentials (spikes) in a nerve cell [50].

Using Eqs. (14) and (15), the second free energy relation of Eq. (4) or (5) leads to:

$$\Delta f_0(V_m) - \Delta f_1(V_m) = -Q_{g,m}V_m \quad (16)$$

or

$$\Delta g_0(V_m) - \Delta g_1(V_m) = Q_{m,g}V_{m,1/2} - Q_{m,g}V_m, \quad (17)$$

In addition, the free energy functions ( $f_\lambda(V)\{\lambda = 0,1\}$ ) of Eq. (16) intersect at  $V=0$  along the reaction coordinate of voltage coupling, leading to the expression:

$$\Delta f_0(0) = \Delta f_1(0) \quad (18)$$

or from Eq. (17)

$$\Delta g_0(0) - \Delta g_1(0) = Q_{g,m}V_{m,1/2} \quad (19)$$

The expression (Eq. (16) or Eqs. (17)) are the fundamental free energy relation (equivalent to Eq. (15)), used to describe the voltage dependency of channel activation (Q-V curve) in voltage gated ion channels [19, 51]. The free energy difference between the two conformational states at zero membrane potential, ( $g_0(0) - g_1(0)$ ), (i.e., between the two non-equilibrium states) is commonly known as the "chemical" component of the total free energy in ion channels under the external potential [19], which is represented by  $G_c$  below. Combining Eq. (19) and Eq. (14) yields

$$\Delta G_c (\equiv \Delta g_0(0) - \Delta g_1(0)) = Q_{g,m}V_{m,1/2} = \Delta G_{conf} \quad (20)$$

Thus we showed here the chemical component of the voltage dependent total free energy,  $G_c (\equiv g_0(0) - g_1(0))$ , is equal to the conformational free energy of  $G_{conf}$  along a one-dimensional reaction coordinate of voltage coupling.

Finally, Eq. (20) combined with the third free energy relation of Eq. (8) implies that

$$\Delta G_c = (\Delta G_{conf} = Q_g V_{1/2}) = \Delta f_1(V_{1,\min}) - \Delta f_0(V_{0,\min}), \quad (21)$$

which is an explicit expression of the free energy difference between the two conformational states at zero membrane potential,  $G_c \equiv g_0(0) - g_1(0)$ , (i.e., between the two non-equilibrium states) in terms of the free energy difference between the two “equilibrium” conformational states ( $f_1(V_{1,\min}) - f_0(V_{0,\min})$ ) along the one-dimensional reaction coordinate of voltage coupling, via the conformational free energy of  $G_{conf}$ , determined using the equilibrium fluctuation relation or the first free energy relation.

The free energy relations, investigated so far, are illustrated in Fig. 1, using the actual simulation data for the model system in section 3.

### 2.3 Linear response approximation satisfies the free energy relations (or the equilibrium fluctuation relation)

If we consider the effect of the overall electrode potential, i.e., an externally applied voltage,  $V_{ext}$ , on the energetics of the entire electrolyte/membrane-protein system, instead of just the membrane potential on the energetics of the membrane-protein system, the Hamiltonian in Eq. 12 may be rewritten formally as:

$$H_1 = H_0 - Q_{ext} V_{ext} \quad (22)$$

where  $Q_{ext}$  is the charge that flows through the electrodes and assumed to be equal to gating charge (see Eq. 12 and Fig. 2), which will be reviewed shortly below.

The state-dependent free energy of the voltage activation can be expressed by modifying the treatment of [52–54] and writing:

$$\Delta f_\lambda(V_{ext}) = \Delta f_\lambda(0) - \left[ \sum_j q_j \phi_{mp,\lambda}(x_j) \right] V_{ext} + \frac{1}{2} C'_\lambda V_{ext}^2 \quad (23)$$

where  $f_\lambda(0)$  is related to the chemical component of free energy of Eq. (18) or (19).

Here the second term is the continuum expression for  $Q_{\lambda,g,m} V_{ext}$  (assuming that  $V_{ext} = V_m$ ) and the last term is the interaction of the external potential with electrolytes, where  $C'_\lambda$  is the capacitance of the electrolyte/membrane-protein system.

The function  $\phi_{\lambda,mp}(x_j)$ , which represents a fraction of membrane potential that falls on the  $j$ th charged amino acid [52, 53, 55], can be obtained by solving a modified Poisson-Boltzmann equation (PB-V) [54]. For a linearized membrane potential, it leads to the simple expression

$$\Delta \phi_{mp}(x_j) = \frac{\varepsilon_L z_j}{\varepsilon_j L}, \quad (24)$$

which is often termed as the dielectric distance (see [33] and references therein). The expression is a generalization of a simple geometric distance ( $z_j/L$ ) along the axis normal to the membrane surface, where  $\varepsilon_L$  and  $\varepsilon_j$  are, respectively, the dielectric constant of the membrane and the (frequently ill defined) local dielectric constant in the region of the  $j$ th



charged amino acid. Here  $L$  is the width of membrane and  $z_j$  is a coordinate normal to the membrane.

Using Eqs. 23 and 24, in the case of linear change of the potential across the membrane, the state dependent gating charge becomes

$$Q_{g,\lambda} = \sum_l (z_l q_l^p \varepsilon_L / \varepsilon_l)_\lambda / L, \quad (25)$$

Finally, using the Hamiltonian of Eq. 22 and differentiating  $f_\lambda(V)$  with respect to the external potential gives (see the SI):

$$\begin{aligned} \frac{\partial \Delta f_\lambda}{\partial V_{ext}} \Big|_{V_{ext}=0} &= - \frac{\partial}{\partial V_{ext}} \ln Z_\lambda \Big|_{V_{ext}=0} \\ &= - \langle Q_{ext} \rangle_{\lambda, V_{ext}=0} \end{aligned} \quad (26)$$

We also have from Eq. 23

$$\frac{\partial \Delta f_\lambda}{\partial V_{ext}} \Big|_{V_{ext}=0} = \left[ \left( - \sum_j q_j \phi_{\lambda, mp}(x_j) \right) + C' V_{ext} \right] \Big|_{V_{ext}=0} = - Q_{\lambda, g, m} \quad (27)$$

This proves that  $Q_{\lambda, g, m} = \langle Q_{ext} \rangle_{\lambda, V_{ext}=0}$  but it should be noticed that the equality reflects the assumption that  $Q_{g, m}$  is equal to  $Q_{ext}$ , each of which corresponds to charge coupling to the membrane potential and the external potential in their respective Hamiltonians (see Eqs. (13) and (22)).

In fact, the quadratic free energy expression in Eq. (23), which is familiar in the Marcus theory of electron transfer [37], is thought of as a realization of linear response approximation (LRA) [56–58] that is likely to be satisfied by many dimensional systems in their response to charging processes (e.g. by the responses to an external potential in the present study). Here, the LRA expression for the free energy functions ( $f_\lambda(V) \{ \lambda=0,1 \}$ ) is shown to satisfy the second free energy relation (Eq. 16), or equivalently, the equilibrium fluctuation relation (Eq. 15), leading to a novel expression for gating charge in terms of other experimentally measurable quantities.

The Hamiltonian used in the present study may also be expressed as follows:

$$H_1 = H_0 - Q_{ext}(V - V_{0, min}), \quad (28)$$

where the subscript of  $V$  (“ext”) was dropped. The free energy functions of voltage activation ( $f_\lambda(V) \{ \lambda=0,1 \}$ ) can be expanded around their respective minima up-to a second order (i.e., using the equilibrium fluctuation-response relation) and expressed as (see SI for a detailed derivation):

$$\begin{aligned}
\Delta f_0(V) &= \Delta f_0(V_{0,\min}) - \frac{1}{2} \langle \delta Q_{ext}^2 \rangle_{V_{0,\min}} (V - V_{0,\min})^2 \\
&= \Delta f_0(V_{0,\min}) - \frac{1}{2} \langle \delta Q_{ext}^2 \rangle_{V_{0,\min}} V_{0,\min}^2 + \langle \delta Q_{ext}^2 \rangle_{V_{0,\min}} V_{0,\min} V - \frac{1}{2} \langle \delta Q_{ext}^2 \rangle_{V_{0,\min}} V^2, \quad (29) \\
&= \Delta f_0(0) + \langle \delta Q_{ext}^2 \rangle_{V_{0,\min}} V_{0,\min} V - \frac{1}{2} \langle \delta Q_{ext}^2 \rangle_{V_{0,\min}} V^2
\end{aligned}$$

where  $\delta Q_{ext} = Q_{ext} - \langle Q_{ext} \rangle$ . Similarly, for the up state (state (1)) we have,

$$\Delta f_1(V) = \Delta f_1(0) + \langle \delta Q_{ext}^2 \rangle_{V_{1,\min}} V_{1,\min} V - \frac{1}{2} \langle \delta Q_{ext}^2 \rangle_{V_{1,\min}} V^2, \quad (30)$$

Notice that these two expressions could be obtained by a direct application of linear response approximation, i.e., differentiation of the free energy function with respect to the external potential gives:

$$\left. \frac{\partial \Delta f_0}{\partial V} \right|_V = -\langle Q_{ext} \rangle_V = -\langle Q_{ext} \rangle_{V_{0,\min}} - \langle \delta Q_{ext} \rangle_V = -\langle \delta Q_{ext} \rangle_V \quad (31)$$

An explicit application of linear response approximation [58] yields:

$$\begin{aligned}
\left. \frac{\partial \Delta f_0}{\partial V} \right|_V &= -\langle \delta Q_{ext} \rangle_V \\
&\simeq -\langle \delta Q_{ext} \rangle_{V_{0,\min}} (V - V_{0,\min}) \quad (32)
\end{aligned}$$

Integration gives the same result as Eq. (29).

Subtracting Eq. (30) from Eq. (29) yields

$$\begin{aligned}
\Delta f_0(V) - \Delta f_1(V) &= f_0(0) - f_1(0) + \left[ \langle \delta Q_{ex}^2 \rangle_{V=V_{0,\min}} V_{0,\min} - \langle \delta Q_{ext}^2 \rangle_{V=V_{1,\min}} V_{1,\min} \right] V \\
&\quad - \frac{1}{2} \left[ \langle \delta Q_{ex}^2 \rangle_{V=V_{0,\min}} - \langle \delta Q_{ext}^2 \rangle_{V=V_{1,\min}} \right] V^2 \\
&= \left[ \langle \delta Q_{ex}^2 \rangle_{V=V_{0,\min}} V_{0,\min} - \langle \delta Q_{ext}^2 \rangle_{V=V_{1,\min}} V_{1,\min} \right] V - \frac{1}{2} \left[ \langle \delta Q_{ex}^2 \rangle_{V=V_{0,\min}} - \langle \delta Q_{ext}^2 \rangle_{V=V_{1,\min}} \right] V^2 \\
&= C' \Delta V_{\min} V, \quad (33)
\end{aligned}$$

where  $C' \equiv -\langle \delta Q_{ext}^2 \rangle_{V=V_{1,\min}} = -\langle \delta Q_{ext}^2 \rangle_{V=V_{0,\min}}$  is the capacitance of the channel-membrane system and  $V_{\min} = V_{1,\min} - V_{0,\min}$ . The equality in the first line comes directly from Eq. (18) that the free energy functions intersect at  $V=0$ . The approximation of equal curvature for the quadratic free energy functions was used in the second line, leading to

$C' \equiv -\langle \delta Q_{ext}^2 \rangle_{V=V_{1,\min}} = -\langle \delta Q_{ext}^2 \rangle_{V=V_{0,\min}}$ . In fact, this approximation is the same level of approximation ( $p_1^{\max} = p_0^{\max}$ ) used to drop a constant for the second free energy relation of Eq. (16), when it moves from Eq. (5).

A comparison of Eq. (33) with Eq. (16) (i.e., the requirement that the LRA expression for free energy functions satisfy the second free energy relation) yields a novel expression for gating charge of the form:

$$Q_g = -C' \Delta V_{\min}, \quad (34)$$

This gating charge expression is the key relation of our paper, where a measure of voltage sensitivity (gating charge) is expressed in terms of the basic properties of channels, as probably conceived by Hodgkins and Huxley [50]. This relation may offer a simple rationale for the differences in gating charges among members of the voltage gated ion channels [59]. In addition, the LRA expressions of Eqs. (29) and (30) should satisfy the equilibrium fluctuation relation of Eq. (15) with the probability densities of finding the  $\lambda$  state at  $V$ , i.e.,

$$p_\lambda(V) = \frac{\exp(-f_\lambda(V))}{Z_\lambda};$$

$$\begin{aligned} \frac{p_0(V)}{p_1(V)} &= \frac{Z_1}{Z_0} \exp(\Delta f_1(V) - \Delta f_0(V)) \\ &\simeq \frac{Z_1}{Z_0} \exp(-C' \Delta V_{\min} V) \\ &= \exp(-C' \Delta V_{\min} V - \Delta G_{conf}) \end{aligned} \quad (35)$$

The approximation in the second line comes directly from Eq. (33). A comparison with the equilibrium fluctuation relation of Eq. (15) yields the same gating charge expression as Eq. (34).

Last, the state dependent gating charge ( $Q_{g,\lambda}$ ) can be defined from Eqs. (33) and (34) as:

$$Q_{g,\lambda} = \left\langle \delta Q_{ext}^2 \right\rangle_{V=V_{\lambda,\min}} V_{\lambda,\min}, \quad (36)$$

The above expression may be rewritten as

$$\left. \frac{\partial Q_{g,\lambda}}{\partial V} \right|_{V=V_{\lambda,\min}} = \left\langle \delta Q_{ext}^2 \right\rangle_{V=V_{\lambda,\min}}, \quad (37)$$

This relation is nothing but the fluctuation-response relation in voltage gated ion channels, where the response of the gating charge to the external potential ( $Q_{\lambda,g} = \langle Q_{ext} \rangle_\lambda$  as in the equivalence of Eq. (26) with Eq. (27)) is related to its fluctuation, by noticing the equivalent relation:

$$\left. \frac{\partial \langle Q_{ext} \rangle_\lambda}{\partial V} \right|_{V=V_{\lambda,\min}} = \left\langle \delta Q_{ext}^2 \right\rangle_{V=V_{\lambda,\min}}, \quad (38)$$

as implicated in Eq. (29) or (30) (see SI for a detailed derivation).

Considering the observation that we have quadratic Marcus-type free energies of equal curvature (which is reflected in the capacitance), we find out that the reorganization energy ( $\Lambda$ ) for voltage coupling in membrane proteins is given by [60]:

$$\Lambda = \frac{1}{2} C' (\Delta V_{\min})^2 = -\frac{1}{2} Q_g \Delta V_{\min}, \quad (39)$$

which yields the free energy of (deactivation) barrier (from  $\lambda=0$  to  $\lambda=1$ ), given by:

$$\Delta G^\ddagger = \frac{\Lambda}{4} \left(1 + \frac{\Delta G_{conf}}{\Lambda}\right)^2 = -\frac{1}{8} Q_g \Delta V_{\min} \left(1 - \frac{2V_{1/2}}{\Delta V_{\min}}\right)^2 \quad (40)$$

Notice that the gating charge has a negative sign as we have consider the conformational transition from the activated ( $\lambda=0$ ) to resting ( $\lambda=1$ ) states.

## 2.4 Key Features of the Coarse-Graining (CG) Model

The energetics of our CG model is different than most other models as it focuses on reliable treatment of the electrostatic energy ( $G_{elec}$ ), considering the self-energy of ionizable residues and the charge-charge interaction with a realistic dielectric as well as the electrostatic energetics of protein-membrane system [20, 32]. The model also considers the hydrophobic contribution ( $G_{hydro}$ ) to the CG model that has been constantly refined over past years. Most importantly, the influence of an applied voltage was recently incorporated into the CG model ( $G_{Iyte-voltage}(V_{ext})$ ) – CG (semi-microscopic) model of voltage coupling in membrane proteins, referred to as the Kim-Dryga-Warshel (KDW) model – as a part of electrostatic contribution ( $G_{elec}$ ) [61]. The KDW model was successfully applied to evaluating the CG energetics of voltage coupling, as well as the gating currents in ion channels (Kv1.2) [21], and gating charge and voltage changes in Bacterial Reaction Center [33].

The energetics of the CG model is given by:

$$\Delta G^{tot} = \Delta G^{fold} = \Delta G_{main} + \Delta G_{side} \quad (41)$$

where the total free energy is taken relative to the free energy of the unfolded system in water at zero applied potential. The main chain energy is given by backbone and hydrogen bonds contributions with weights ( $w$ ) optimized to yield the observed absolute folding free energies:

$$\Delta G_{main} = w_{back} \Delta G_{back} + w_{HB} \Delta G_{HB},$$

while the side chain contribution is decomposed into four terms:

$$\Delta G_{side} = \Delta G_{elec}^0 + \Delta G_{hydro} + \Delta G_{polar} + \Delta G_{vdw} \quad (42)$$

Finally, in the case of the presence of electrodes and electrolytes,  $\Delta G_{lyte-voltage}^{fold}(V_{ext})$  is added to the total free energy (Eq. 41), resulting in

$$\Delta G^{tot}(V_{ext}) = \Delta G^{fold} + \Delta G_{lyte-voltage}(V_{ext}), \quad (43)$$

where the  $\Delta G_{lyte-voltage}^{fold}(V_{ext})$  is the CG representation of the effect of the external potential (KDW model), and the nature of this term will be elaborated below.

In the following, we consider the electrostatic contributions to the folding energy as a state dependent function of the form, given by:

$$\Delta G_{elec,\lambda}(V_{ext}) = \Delta G_{elec,\lambda}^0 + \Delta G_{lyte-voltage,\lambda}(V_{ext}) \quad (44)$$

The first term represents the (voltage independent) electrostatic free energy of the “ionizable” residues, which is the sum of two contributions: a local-environment and membrane-depth dependent Born-type self-energy and charge-charge interaction energy with a distant dependent dielectric that approaches  $\sim 40$  in an infinite distance:

$$\Delta G_{elec,\lambda}^0 = \Delta G_{self} + 332 \sum_{l,m} \frac{q_l^p q_m^p}{\epsilon_{lm}^{eff}(r_{lm}) r_{lm}}, \quad (45)$$

where we use a distant dependent effective dielectric function,

$\epsilon_{lm}^{eff} = 1 + 60[1 - \exp(-0.5r_{lm})]$ . The  $\Delta G_{lyte-voltage,\lambda}(V_{ext})$  term represents voltage-dependent energetics of membrane proteins coupled to the externally applied potential, which includes the effects of electrolytes on the electrostatic energetics [33]:

$$\Delta G_{lyte-voltage,\lambda}(V_{ext}) = \frac{1}{2} \sum_j V_{ext}^j q_j^g + \frac{1}{2} 332 \sum_{k \neq j} \frac{q_j^h q_k^g}{\epsilon_{kj}^{eff,lyte}(r_{jk}) r_{jk}} + \sum_l V_{ext}^l q_l^p + \frac{1}{2} 332 \sum_{k,m} \frac{q_m^p q_k^g}{\epsilon_{km}^{eff}(r_{km}) r_{km}}, \quad (46)$$

where the units are kcal/mol rather than the esu units in the above macroscopic expression and distant dependent effective dielectric constants of 40 and 60 were used, respectively, for the electrolyte response to the charging of protein residues ( $\epsilon_{km}^{eff}$ ) and external potentials ( $\epsilon_{kj}^{eff,lyte}$ ). The first and second terms represents the contribution that arises from the polarization of electrolyte effective charges ( $q^g$ ) by the external potential and all the other charge of the system. The third term represents voltage coupling to protein charges ( $q^p$ ) and the last term represents the interaction between protein charges and electrolytes that arises when protein ionizable residues are charged.

More specifically, the electrolyte charges in this expression ( $q_j^g$ ) are represented by a grid with a charge distribution of the form:

$$q_j^\pm = \frac{z^\pm N_{box}^\pm e^{\mp\beta\phi_j}}{\sum_{k \in box} e^{\mp\beta\phi_k}}, \quad (47)$$

where ( $q_j^g = q_j^+ + q_j^-$ ) is determined by solving iteratively with the interactions between all the charges in the system, considering the local potential on each grid point [20, 32]:

$$\varphi_j = 332 \sum_m \frac{q_m^P}{\epsilon_{jm}^{eff}(r_{jm})r_{jm}} + 332 \sum_{k \neq j} \frac{q_k^g}{\epsilon_{kj}^{eff}(r_{jk})r_{jk}} + V_j^{ext}, \quad (48)$$

where  $V_j^{ext}$ , is evaluated using the macroscopic formula [20, 32]:

$$V_j^{ext} = \int_{Z_0}^{Z_j} D_z^0 / \bar{\epsilon}(Z) dZ \quad (49)$$

where  $Z_0$  is the Z coordinate at the left electrode. Alternatively we can consider the potential from the electrode charges and using periodic boundary conditions.

The KDW model evaluates the gating charge in a direct manner by computing the cumulative electrolyte charges near the electrode that arises in response to charge movements within the membrane, using an integration through the relation:

$$Q_g = \int_{Z_0}^{Z'} dZ \left( \Delta(\Delta q^{grid}(Z, V) / \Delta Z) \right) = \sum_{i=0}^{i=i'} \sum_{j,k} q_{i,j,k}^{grid} \quad (50)$$

where  $q_{grid}(Z, V)$  is the difference in the accumulated sum of  $q_{grid}$  of the initial and final state, and  $Z'$  is the point to the left of the membrane where the electrolyte charge distribution changes sign. At this point, the integrated charge reaches a plateau and then starts to decrease. The formal integral in the second term of Eq. 50 corresponds to the explicit summation on the grid points (in the third term), where j and k run on all the grid [points in the X and Y directions, whereas i runs from points near the electrode to  $Z'$ .

The KDW model was recently applied to estimating gating charge in voltage gated  $K^+$  ion channel (Kv1.2) [21] and voltage changes in response to proton/electron transfer in bacterial reaction center [33].

With the above formulation we can try to look for the analogy to the macroscopic continuum treatment. The most obvious analogy come with the capacitor model where we note that the first two terms in Eq. (46) correspond to the free energy of charging (the membrane) between two electrolytes by an electric field in the macroscopic continuum approach. Here we can try to look for correspondence between the CG and continuum models and note that in the absence of the protein charge we have:

$$\frac{1}{2} \sum_j V_j^{ext} q_j^g + \frac{1}{2} 332 \sum_{k \neq j} \frac{q_j^g q_k^g}{\epsilon_{jk}^{eff} (r_{jk}) r_{jk}} \leftrightarrow \frac{1}{2} C V_{ext}^2 \quad (51)$$

This correspondence states that the free energy stored in the membrane capacitor is equal to the free energy of polarizing electrolyte solutions. The validity of this relationship will be explored in Section 3.1 (see also Fig. 4B).

Now we investigate the relationship between the gating charge obtained by the macroscopic continuum and the KDW model, by formally rewriting the state dependent total CG energy as the free energy function, introduced before:

$$\Delta G_\lambda^{total}(V_{ext}) = \Delta f_\lambda(V_{ext}) \quad (52)$$

While the CG model of voltage coupling was used to evaluate gating charge in a direct manner by computing the cumulative electrolyte charges on the electrode generated in response to charge movements within the membrane, we can look for the trend in a state dependent gating charge,  $Q_{g,\lambda}$ , using the differentiation of Eq. 52, in an analogy with Eq. 27, which can be expressed as:

$$Q_{g,\lambda} = \partial \Delta f_{\lambda, fold} / \partial V_{ext} |_{V_{ext}=0} \quad (53)$$

This expression can clearly be explored by numerical differentiation. Furthermore, an inspection of our CG energetics (i.e. the third term in Eq. 46) against Eq. (23) yields the gating charge, through:

$$Q_{g,\lambda} = \sum_l (V_m^l q_l^p)_\lambda / V_m \quad (54)$$

This expression should have the same trend as the (linearized) macroscopic expression as Eq. 25. Both Eqs. 53 and 54 will be applied in Section 3 to estimate the gating charge.

## 3. Results

### 3.1 Gating charge

We start by illustrating the application of the free energy relations derived above to the analysis of the thermodynamics of voltage activation in ion channels. This is done by using our CG model to determine the energetics of moving a unit charge between the two sides of a membrane, in the presence of an external (see Fig. 2). The area (A) and width (L) of membrane are  $80 \times 80 \text{ \AA}^2$  and  $60 \text{ \AA}$  respectively, where the electrolyte concentration is taken as 250mM. The unit charge is initially positioned near the membrane on the left side (i.e., state 0). An application of a positive external potential drives the unit charge to the right side (i.e., state 1) of the membrane. Such a “non-equilibrium” movement of the charge within the membrane leads to accumulation of electrolyte charges near the membrane, where opposite

electrolyte charges are accumulated near the electrode (see bottom in Fig. 2). The difference in electrolyte charges near the electrode before and after the charge movement is the actual gating charge measured in experiments. The charge movement can also generate voltage change (unless we keep the voltage constant), and be expressed as a change in the electrode potential.

Such a direct evaluation of the gating charge has been possible through the development of our CG model of membrane proteins that include membrane, electrolytes, and electrodes [20, 32], which is now called the KDW model in the present study. This model focuses on the actual measured quantity, namely the displacement current (e.g. [35]), rather than on its interpretation. Thus, the gating charge is evaluated by considering the fact that the gating current is due to the motion and accumulation of the electrolyte charges.

Figure 3A shows the gating charge determined by such a direct procedure, while using Eq. 50, yielding the gating charge of  $\sim 0.73e$ . It turns out that  $q_g(Z, V)$  is independent of the external potential [33], where the accumulated charge (gating charge) was produced at zero potential (also see inset of Fig 3A that shows the gating charge at an external potential of 100 mV).

The second approach, referred to here as the first “indirect method”, uses the free energy of voltage coupling to the protein charges, whose expression is obtained by treating the energetics of voltage activation from our KDW model (Eq. 53) in analogy to Eq. 27. The numerical differentiation of the CG free energies at zero membrane potential yields a gating charge of  $\sim 0.74e$ . In addition, we can use the second indirect approach (Eq. 54), which is an analogous expression to the corresponding macroscopic continuum expression (Eq. 25). This approach yields a gating charge of  $\sim 0.75e$ , where the linear component of capacitance charge is subtracted, as shown in Fig. 3B.

The third approach uses a new expression of gating charge derived in the present study (Eq. 34, i.e.,  $Q_g = -C' V_{\min}$ ), using a minimum voltage difference between two states and a capacitance of the system. The voltage difference, where the two states have their respective minimum, was obtained from the voltage dependent free energies using our CG model (Fig. 4A), yielding a value of 300mV ( $V_{1,\min} - V_{0,\min}$ ). The capacitance of the model system, i.e., fluctuation of an external charge flowing through a complete circuit, can be estimated as a second derivative of the voltage dependent free energies with respect to an external potential, yielding the capacitance of  $\sim 2.52e/V$ , which is close to the membrane-only (between two electrolytes) capacitance of  $\sim 2.33e/V$  ( $0.58\mu F/cm^2$ ), determined by using the (linearized) macroscopic continuum approach [62]:

$$C = C_m \frac{1}{1 + \frac{2\epsilon_m}{kL\epsilon_w}}, \quad (55)$$

where  $l/k$  Debye length,  $L(=60)$  is the width of membrane,  $\epsilon_m (=4)$  and  $\epsilon_w (=80)$  are dielectric constants of membrane and water respectively, and  $C_m$  is the capacitance of the pure membrane:



$$C_m = \frac{\epsilon_m A}{4\pi L} \quad (56)$$

Notice that our CG energetics of voltage coupling (the KDW model) also yields the free energy parabola that yields a membrane capacitance of  $\sim 2.34\text{e/V}$ , which is excellent agreement with the prediction of the macroscopic capacitor formula of  $\sim 2.33\text{e/V}$  (Fig. 4B), as predicted by the correspondence between our CG model of energetics of voltage coupling and the macroscopic continuum model (Eq. 51). At any rate, the newly developed expression for gating charge (Eq. 34, i.e.,  $Q_g = -C' V_{\min}$ ) yields a similar gating charge ( $Q_g \sim 0.77\text{e}$  with  $C'=2.58\text{ e/V}$  and  $V_{\min}=300\text{mV}$ ) to the charges obtained by the other approaches ( $0.73\text{e}$  from the direct approach and  $0.74\text{e}/0.76\text{e}$  from the indirect approach). Last, for this particular simple system, assuming a linear membrane potential across the membrane, the simple dielectric distant formula with a uniform dielectric constant (Eq. 24 or 25) yields a value of  $40/60\text{e} \sim 0.67\text{e}$ . The fact that we obtained similar results with different approaches is a comforting confirmation of the validity of our CG formulation that can be used in more challenging cases where the macroscopic approximation can be questionable.

### 3.2 Voltage dependent free energies and voltage dependency of activation (Q-V curve)

Figure 4A shows the voltage dependent CG (the KDW model) free energy functions ( $f_\lambda(V)$   $\{\lambda=0,1\}$ ) for moving the unit charge within the membrane as a result of the external potential. These type of free energy parabolas are familiar in the Marcus theory of electron transfer except that they are obtained here along the voltage coupling coordinate and evaluate microscopically. The parabolic dependence is a realization of linear response approximation, which should be satisfied approximately by any system that involves free energies of charging (by an external potential in the present study) along chosen reaction coordinates. The free energy difference between two equilibrium conformational states, i.e.,  $f_1(V_{1,\min}) - f_0(V_{0,\min})$ , is about  $\sim 0.3\text{ kcal/mol}$  (Figs. 4A and 1B), yielding approximately the same result as the “chemical” component of free energy ( $\sim 0.2\text{ kcal/mol}$ ) obtained using the gating charge and half voltage ( $V_{1/2}$ ) (see below and Eq. 20). Notice that  $f_\lambda(V)$   $\{\lambda=0,1\}$  interest at  $V=0$  along the reaction coordinate of voltage coupling, which is a natural consequence of equilibrium fluctuation relation (or the first free energy relation) that any system of two states described by the reaction coordinate of energy gap (voltage coupling in the present study) should satisfy. Last, the free energy of (deactivation) barrier (see Eq. (40)) yields of a value of  $\sim 0.7\text{ kcal/mol}$  (Figs. 4A and 1B).

The determination of voltage dependent free energies using our CG model of voltage coupling (the KDW model) allows for a direct evaluation of Q-V curve. That is, starting with the relation

$$p_\lambda(V) \propto \exp(-\Delta f_\lambda(V)) \quad (57)$$

By normalization (or by an application of Bayes’ theorem with a uniform prior [63]), we obtain:

$$p_0(V) (\equiv Q_g(V)/Q_g^{\max}) = \frac{p_0(V)}{p_0(V)+p_1(V)} = \frac{1}{1+p_1(V)/p_0(V)}, \quad (58)$$

where  $p_\lambda(V)$  ( $\lambda = 0, 1$ ) are the probability densities of finding state  $\lambda$  of voltage sensor domain (VSD) at a given voltage  $V$ . Notice that the 0<sup>th</sup> state is treated as the activated one (see Eq. (13)). Application of equilibrium fluctuation relation (Eq. 15) to Eq. 58 leads to the familiar Fermi-Dirac logistic function, given by

$$p_0(V) (\equiv Q_g(V)/Q_g^{\max}) = \frac{1}{1+\exp(Q_g V_{1/2} - Q_g V)}, \quad (59)$$

This analytic form of Q-V curve is commonly used in the field of ion channels to fit the observed open (activated) probability of voltage activation [19] to extract the thermodynamic information of voltage coupling such as a half voltage (where the initial (0) and final (1) states have equal probability), gating charge (notice, however,  $Q_g$  is equal to  $Q_g^{\max}$  only for the two state model) and thus chemical component of free energy of  $G_c$ . Figure 5 shows the numerical Q-V curve for the unit charge movement using the voltage dependent CG free energies (Eqs. 57 and 58). The least square fit [64] of the voltage dependent 0<sup>th</sup> (activated) CG probability density (Eq. 58) to the analytic form of Q-V curve (Eq. 59) yields the half voltage ( $V_{1/2}$ ) of  $\sim -9.2\text{mV}$  and gating charge of  $\sim -0.78e$ , thereby resulting in the chemical component of free energy of  $\sim 0.2\text{ kcal/mol}$  using

$G_c \equiv g_0(0) - g_1(0) = Q_g V_{1/2}$ . As expected from Eq. 20, the chemical component of free energy ( $G_c$ ) is approximately equal to the conformational free energy ( $G_{conf}$ ) ( $\sim 0.2\text{ kcal/mol}$ ), obtained from the CG voltage dependent free energies using the equilibrium free energy difference ( $f_1(V_{1,min}) - f_0(V_{0,min})$ ) along the reaction coordinate of voltage coupling.

### 3.3 Free energy functions (parabolas) along voltage coupling offers a new insight into the mechanisms of voltage activation in ion channels

Our understanding of voltage activation and gating processes benefits largely from the site-directed mutagenesis studies of selected residues that are critical for channel functions. The effects of such site-directed mutations on the functions of channels that contains voltage sensors modules have been commonly examined through electrophysiological measurements of macroscopic currents and gating currents. In particular, the resulting Q-V curve (and also shifts in the Q-V curve upon site-directed mutations) offers important thermodynamic information of voltage activation of ion channels. For example, specific mutations of key arginine or lysine residues of the S4 helix in the VSD of Shaker channel, which is due to the stabilization of the open-activated (up) state, causes a left shift in the Q-V curve (toward hyperpolarization) [65]. Also a specific set of mutations in the S3b helix of VSD in the same channel causes a right shift in the Q-V by destabilizing the open-activated (up) state of the channel [66]. However, quantitative interpretation of the Q-V curve in terms of the free energy landscapes of channel activation has been hampered due to several reasons; these include the lack of quantitative criteria exists about which conformational state is stabilized or destabilized to account for the observed shifts in the Q-V curve. It also reflects the lack of a quantitative relation between the free energy of stabilization or

destabilization and the observed shifts in the Q-V curve exists (partly due to the absence of a reaction coordinate to describe thermodynamics of voltage activation).

To progress in analyzing the above mutation effects we can exploit the findings of the previous subsection, where we show that the Marcus-type free energy parabolas along the reaction coordinate of voltage coupling allows one to obtain quantitative predictions of the Q-V curve. For example, Figure 6 show such a correlation between the free energies of voltage activation and the Q-V curve. Here, the “hypothetical” Marcus-type free energy parabolas of the 0 and 1 states are related each other by equilibrium fluctuation relation, intersecting each other at  $V=0$  along the reaction coordinate of voltage coupling and yielding the conformational free energy of  $Q_g V_{1/2}$ . This figure shows that the hypothetical Q-V curve (Fig. 6C) could be shifted toward the left (toward hyperpolarization) in either way by the stabilization of the open activated (0) state (red dagger in Fig. 6A) or the destabilization of closed resting (1) state (red dagger in Fig. 6B). In addition, there exists a simple quantitative relation between the free energy of stabilization or destabilization and shift in the Q-V curve, given by  $Q_g V_{1/2} \approx G_{conf}$  from Eq. (20), which can be explored by the CG model of voltage coupling. Thus, the proposed theory and the CG energetics of voltage coupling (the KDW model) offers a clear graphical and quantitative analysis on the correlation between the stabilization/destabilization in the free energies of voltage activation and shifts in Q-V curve by integrating equilibrium fluctuation relations and Marcus type parabolas using the reaction coordinate of voltage coupling. The accurate determination of equilibrium free energy changes of conformational states and corresponding shifts in activation free energy, upon site-directed mutations, have strong implications in designing pharmacophore in drug discovery targeting voltage sensing domains.

#### 4. Discussion

Starting from the equilibrium fluctuation relation that makes a connection between the conformational free energy between the two (forward/backward) probability densities, along the one-dimensional reaction of “voltage coupling”, we have offer a rigorous physical foundation for the fundamental free energy relations in the voltage gated ion channels, used to quantify kinetics/thermodynamics of voltage activation [1,19, 36–41]. We have shown that the chemical component of the voltage dependent free energy, namely, the free energy difference between two non-equilibrium conformational states at zero membrane potential ( $Q_g V_{1/2}$ ), is equivalent to the free energy difference between two “equilibrium” conformational states in a one-dimensional space of voltage coupling.

Application of linear response approximation (LRA) to the free energies of voltage activation in the framework of equilibrium fluctuation relations yields a remarkable relation between gating charge and its fluctuation, i.e., in terms of the basic properties of channels. This relationship, which is known as the fluctuation-response relation in statistical mechanics, is reminiscent of the Nyquist relation for current fluctuation (as a realization of the fluctuation-dissipation theorem in an electric circuit) [67, 68], which was employed to estimate the number of channels and thus the gating charge per channel (see Ref. [69] and references therein). The gating charge expression developed in the present study could thus be considered a time independent version of Nyquist relation (as a realization of fluctuation-

response relation in an analogous neuro-electric circuit). A recent realization of the non-equilibrium (and non-linear) extension of the relation in a quantum coherent conductor [70, 71] may further implicate the roles played by the Nyquist relation in ion channels in the framework of the non-equilibrium fluctuation theorem [72].

The accuracy of our CG model of the energetics of voltage coupling (the KDW model) in membrane proteins, simulated by using semi-“microscopic” grid-based electrolytes and implicit electrodes, was examined considering a well-defined test case (i.e., the membrane between two electrolytes) where we have clear results from macroscopic continuum model [62]. It was found that the free energy of charging (and also capacitance) obtained by using the KDW model, yields an excellent agreement with the macroscopic continuum prediction. Thus, our model offers a new explicit microscopic insight into the nature of the voltage coupling in membrane proteins.

The KDW model for a unit charge movement displaced by an external potential was also applied to simulate gating charge using three different approaches, including the new formula, yielding approximately equal result to each other. The KDW model was then used to illustrate the free energy relations for voltage activation in the simple model of the unit charge movement. The KDW model yields Marcus type parabolas along the reaction coordinate of voltage coupling, as predicted by linear response approximation, allowing for a direct prediction of voltage activation profiles, e.g., gating charge vs. voltage curve (Q-V curve). These free energy parabolas along the one-dimension reaction coordinate of voltage activation have allowed for a clear understanding of the correlation between the changes in the free energies of voltage activation and the shifts in Q-V curve, upon site-directed mutagenesis or drug binding.

Overall, the presented theory and the CG simulation of voltage couplings offers new quantitative tools for understanding the molecular basis of the action of mechanisms of voltage gating in voltage gated ion channels. This was done in the framework of the equilibrium fluctuation relation and LRA, thus laying a firm quantitative foundation for a more unified understanding of voltage effects in ion channels, ion pumps, and transporters in general. Application of the framework developed in the present study to (real) membrane proteins is under way, considering the voltage sensor domain from a voltage sensitive phosphatase [73]. Note that, however, while we expect LRA to be very effective in approximating the energetics of the conformational changes, as it has been in our studies of processes in condensed phases and in proteins (e.g. [74]), the validity is not essential for the new derivation of the gating charge and for the CG treatment of the gating charge/current. That is, we emphasize the CG model of gating charge/charges follows LRA and gives very reasonable capacitor-like results because the electrolytes grid follows nicely LRA. This means that regardless of whether the protein free energy landscapes and conformational transitions follow LRA or not, we do have a reliable description of the gating charges when the structures of the open activated and closed resting state are known. This is indeed the case in our studies of Kv1.2, where we reproduced the observed gating charge/current [21].

In addition, the direct approach of the KDW model provides an effective way for determining the gating charge and the resulting changes in electrode potentials (voltage

generation), that has been used to estimate electrogenicity in membrane proteins (e.g.[75]). This important feature has been illustrated in studies of the electrogenicity in the charge separation in bacterial RC [33], should offer a new insight into the mechanisms of proton/electron transfer in Cytochrome c Oxidase. The work in this direction is also in progress.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

This work was supported by the NIH grant GM40283, the National Science Foundation Grant MCB- 0836400. We acknowledge the University of Southern California's High Performance Computing and Communications Center for computer time.

## References


1. Hille, B. Ion channels of excitable membranes. Vol. 507. Sinauer; Sunderland, MA: 2001.
2. Sakmann, B.; Neher, E. Single-channel recording. Springer Science & Business Media; 2009.
3. Doyle DA, et al. The structure of the potassium channel: molecular basis of K<sup>+</sup> conduction and selectivity. *Science*. 1998; 280(5360):69–77. [PubMed: 9525859]
4. Zhou Y, et al. Chemistry of ion coordination and hydration revealed by a K<sup>+</sup> channel–Fab complex at 2.0 Å resolution. *Nature*. 2001; 414(6859):43–48. [PubMed: 11689936]
5. Long SB, et al. Atomic structure of a voltage-dependent K<sup>+</sup> channel in a lipid membrane-like environment. *Nature*. 2007; 450(7168):376–382. [PubMed: 18004376]
6. Payandeh J, et al. The crystal structure of a voltage-gated sodium channel. *Nature*. 2011; 475(7356):353–358. [PubMed: 21743477]
7. Zhang X, et al. Crystal structure of an orthologue of the NaChBac voltage-gated sodium channel. *Nature*. 2012; 486(7401):130–134. [PubMed: 22678295]
8. Åqvist J, Luzhkov V. Ion permeation mechanism of the potassium channel. *Nature*. 2000; 404(6780):881–884. [PubMed: 10786795]
9. Berneche S, Roux B. Energetics of ion conduction through the K<sup>+</sup> channel. *Nature*. 2001; 414(6859):73–77. [PubMed: 11689945]
10. Burykin A, et al. Simulations of ion current in realistic models of ion channels: the KcsA potassium channel. *Proteins: Structure, Function, and Bioinformatics*. 2002; 47(3):265–280.
11. Garofoli S, Jordan P. Modeling permeation energetics in the KcsA potassium channel. *Biophysical journal*. 2003; 84(5):2814–2830. [PubMed: 12719216]
12. Lecar H, Larsson HP, Grabe M. Electrostatic model of S4 motion in voltage-gated ion channels. *Biophysical journal*. 2003; 85(5):2854–2864. [PubMed: 14581190]
13. Grabe M, et al. A quantitative assessment of models for voltage-dependent gating of ion channels. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101(51):17640–17645. [PubMed: 15591352]
14. Chanda B, et al. Gating charge displacement in voltage-gated ion channels involves limited transmembrane movement. *Nature*. 2005; 436(7052):852–856. [PubMed: 16094369]
15. Sadovsky E, Yifrach O. Principles underlying energetic coupling along an allosteric communication trajectory of a voltage-activated K<sup>+</sup> channel. *Proceedings of the National Academy of Sciences*. 2007; 104(50):19813–19818.
16. Silva JR, et al. A multiscale model linking ion-channel molecular dynamics and electrostatics to the cardiac action potential. *Proceedings of the National Academy of Sciences*. 2009; 106(27):11102–11106.
17. Khalili-Araghi F, et al. Calculation of the gating charge for the Kv1. 2 voltage-activated potassium channel. *Biophysical journal*. 2010; 98(10):2189–2198. [PubMed: 20483327]

18. Schwaiger CS, et al. The free energy barrier for arginine gating charge translation is altered by mutations in the voltage sensor domain. *PloS one*. 2012; 7(10):e45880. [PubMed: 23094020]
19. Chowdhury S, Chanda B. Thermodynamics of electromechanical coupling in voltage-gated ion channels. *The Journal of general physiology*. 2012; 140(6):613–623. [PubMed: 23183697]
20. Dryga A, et al. Realistic simulation of the activation of voltage-gated ion channels. *Proceedings of the National Academy of Sciences*. 2012; 109(9):3335–3340.
21. Kim I, Warshel A. Coarse-grained simulations of the gating current in the voltage-activated Kv1. 2 channel. *Proceedings of the National Academy of Sciences*. 2014; 111(6):2128–2133.
22. Delemotte L, et al. Free-energy landscape of ion-channel voltage-sensor–domain activation. *Proceedings of the National Academy of Sciences*. 2015; 112(1):124–129.
23. Allen TW, Kuyucak S, Chung SH. Molecular dynamics study of the KcsA potassium channel. *Biophysical journal*. 1999; 77(5):2502–2516. [PubMed: 10545352]
24. Luzhkov VB, Åqvist J. K<sup>+</sup>/Na<sup>+</sup> selectivity of the KcsA potassium channel from microscopic free energy perturbation calculations. *Biochimica et Biophysica Acta (BBA)-Protein Structure and Molecular Enzymology*. 2001; 1548(2):194–202. [PubMed: 11513964]
25. Burykin A, Kato M, Warshel A. Exploring the origin of the ion selectivity of the KcsA potassium channel. *Proteins: Structure, Function, and Bioinformatics*. 2003; 52(3):412–426.
26. Noskov SY, Berneche S, Roux B. Control of ion selectivity in potassium channels by electrostatic and dynamic properties of carbonyl ligands. *Nature*. 2004; 431(7010):830–834. [PubMed: 15483608]
27. Thompson AN, et al. Mechanism of potassium-channel selectivity revealed by Na<sup>+</sup> and Li<sup>+</sup> binding sites within the KcsA pore. *Nature structural & molecular biology*. 2009; 16(12):1317–1324.
28. Kim I, Allen TW. On the selective ion binding hypothesis for potassium channels. *Proceedings of the National Academy of Sciences*. 2011; 108(44):17963–17968.
29. Roux B, et al. Ion selectivity in channels and transporters. *The Journal of general physiology*. 2011; 137(5):415–426. [PubMed: 21518830]
30. Nimigean CM, Allen TW. Origins of ion selectivity in potassium channels from the perspective of channel block. *The Journal of general physiology*. 2011; 137(5):405–413. [PubMed: 21518829]
31. Delemotte L, et al. Intermediate states of the Kv1. 2 voltage sensor from atomistic molecular dynamics simulations. *Proceedings of the National Academy of Sciences*. 2011; 108(15):6109–6114.
32. Dryga A, et al. Coarse grained model for exploring voltage dependent ion channels. *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 2012; 1818(2):303–317. [PubMed: 21843502]
33. Kim I, et al. Modeling gating charge and voltage changes in response to charge separation in membrane proteins. *Proceedings of the National Academy of Sciences*. 2014; 111(31):11353–11358.
34. Jensen MØ, et al. Mechanism of voltage gating in potassium channels. *Science*. 2012; 336(6078):229–233. [PubMed: 22499946]
35. Armstrong, CM.; Bezanilla, F. Currents related to movement of the gating particles of the sodium channels. 1973.
36. Warshel A. Dynamics of reactions in polar solvents. Semiclassical trajectory studies of electron-transfer and proton-transfer reactions. *The Journal of Physical Chemistry*. 1982; 86(12):2218–2224.
37. King G, Warshel A. Investigation of the free energy functions for electron transfer reactions. *The Journal of Chemical Physics*. 1990; 93(12):8682–8692.
38. Zhou HX, Szabo A. Microscopic formulation of Marcus' theory of electron transfer. *The Journal of chemical physics*. 1995; 103(9):3481–3494.
39. Warshel A, Parson WW. Dynamics of biochemical and biophysical reactions: insight from computer simulations. *Quarterly reviews of biophysics*. 2001; 34(04):563–679. [PubMed: 11852595]
40. Hummer G, Szabo A. Calculation of free-energy differences from computer simulations of initial and final states. *The Journal of chemical physics*. 1996; 105(5):2004–2010.

41. Kim I, Allen TW. Bennett's acceptance ratio and histogram analysis methods enhanced by umbrella sampling along a reaction coordinate in configurational space. *The Journal of chemical physics*. 2012; 136(16):164103. [PubMed: 22559466]
42. Bennett CH. Efficient estimation of free energy differences from Monte Carlo data. *Journal of Computational Physics*. 1976; 22(2):245–268.
43. Crooks GE. Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences. *Physical Review E*. 1999; 60(3):2721.
44. Jarzynski C. Nonequilibrium equality for free energy differences. *Physical Review Letters*. 1997; 78(14):2690.
45. Frenkel, D.; Smit, B. *Understanding molecular simulation: from algorithms to applications*. Vol. 1. Academic press; 2001.
46. Collin D, et al. Verification of the Crooks fluctuation theorem and recovery of RNA folding free energies. *Nature*. 2005; 437(7056):231–234. [PubMed: 16148928]
47. Marcus RA. On the Theory of Oxidation-Reduction Reactions Involving Electron Transfer I. *J Chem Phys*. 1956; 24:966–978.
48. Marcus RA. Chemical and electrochemical electron transfer theory. *Ann Rev Phys Chem*. 1964; 15:155–196.
49. Pathak MM, et al. Closing in on the resting state of the Shaker K<sup>+</sup> channel. *Neuron*. 2007; 56(1):124–140. [PubMed: 17920020]
50. Hodgkin AL, Huxley AF. A quantitative description of membrane current and its application to conduction and excitation in nerve. *The Journal of physiology*. 1952; 117(4):500–544. [PubMed: 12991237]
51. Roux B. The membrane potential and its representation by a constant electric field in computer simulations. *Biophysical journal*. 2008; 95(9):4205–4216. [PubMed: 18641071]
52. Stevens C. Interactions between intrinsic membrane protein and electric field. An approach to studying nerve excitability. *Biophysical journal*. 1978; 22(2):295. [PubMed: 656546]
53. Sigworth FJ. Voltage gating of ion channels. *Quarterly reviews of biophysics*. 1994; 27(01):1–40. [PubMed: 7520590]
54. Roux B. Influence of the membrane potential on the free energy of an intrinsic protein. *Biophysical Journal*. 1997; 73(6):2980. [PubMed: 9414213]
55. Lauger, P. *Electrogenic ion pumps*. Vol. 5. Sinauer Associates; Sunderland: 1991.
56. Marcus RA. On the theory of electron-transfer reactions. VI. Unified treatment for homogeneous and electrode reactions. *The Journal of Chemical Physics*. 1965; 43(2):679–701.
57. Warshel A, Russell ST. Calculations of electrostatic interactions in biological systems and in solutions. *Quarterly reviews of biophysics*. 1984; 17(03):283–422. [PubMed: 6098916]
58. Simonson T. Gaussian fluctuations and linear response in an electron transfer protein. *Proceedings of the National Academy of Sciences*. 2002; 99(10):6544–6549.
59. Ishida IG, et al. Voltage-dependent gating and gating charge measurements in the Kv1. 2 potassium channel. *The Journal of general physiology*. 2015 p. jgp. 201411300.
60. Rosta E, Warshel A. Origin of Linear Free Energy Relationships: Exploring the Nature of the Off-Diagonal Coupling Elements in SN2 Reactions. *Journal of chemical theory and computation*. 2012; 8(10):3574–3585. [PubMed: 23329895]
61. Vicatos S, et al. An effective coarse-grained model for biological simulations: recent refinements and validations. *Proteins*. 2014; 82(7):1168–85. [PubMed: 25050439]
62. Lauger P, et al. Electrical properties of bimolecular phospholipid membranes. *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 1967; 135(1):20–32. [PubMed: 6031508]
63. Anastasio, TJ. *Tutorial on neural systems modeling*. Sinauer Associates, Incorporated; 2009.
64. Jones, E.; Oliphant, T.; Peterson, P. {SciPy}: Open source scientific tools for {Python}. 2014.
65. Tao X, et al. A gating charge transfer center in voltage sensors. *Science*. 2010; 328(5974):67–73. [PubMed: 20360102]
66. Xu Y, et al. Energetic role of the paddle motif in voltage gating of Shaker K<sup>+</sup> channels. *Nature structural & molecular biology*. 2013; 20(5):574–581.


67. Nyquist H. Thermal agitation of electric charge in conductors. *Physical review*. 1928; 32(1):110–113.
68. Johnson JB. Thermal agitation of electricity in conductors. *Physical review*. 1928; 32(1):97.
69. Islas LD, Sigworth FJ. Voltage sensitivity and gating charge in Shaker and Shab family potassium channels. *The Journal of general physiology*. 1999; 114(5):723–742. [PubMed: 10539976]
70. Nakamura S, et al. Nonequilibrium fluctuation relations in a quantum coherent conductor. *Physical review letters*. 2010; 104(8):080602. [PubMed: 20366923]
71. Pekola JP. Towards quantum thermodynamics in electronic circuits. *Nature Physics*. 2015; 11(2): 118–123.
72. Evans DJ, Cohen E, Morriss G. Probability of second law violations in shearing steady states. *Physical Review Letters*. 1993; 71(15):2401. [PubMed: 10054671]
73. Li Q, et al. Structural mechanism of voltage-dependent gating in an isolated voltage-sensing domain. *Nature structural & molecular biology*. 2014; 21(3):244–252.
74. Muegge I, et al. The reorganization energy of cytochrome c revisited. *The Journal of Physical Chemistry B*. 1997; 101(5):825–836.
75. Belevich I, et al. Exploring the proton pump mechanism of cytochrome c oxidase in real time. *Proc Natl Acad Sci U S A*. 2007; 104(8):2685–90. [PubMed: 17293458]






**Figure 1.**

Illustration of fundamental free energy relations found in the present study with the CG model system of Fig. 2. The calculations were done by moving the positive charge in a straight line between the left and right points (say from the activated to the resting) thus changing both  $f_0$  and  $f_1$ . (A) The probability densities ( $p_\lambda(V)$ )  $\{\lambda = 0, 1\}$  of finding the conformational states  $\lambda$  at an external potential of  $V$  ( $\equiv V_{ext}$  is assumed to be equal to the membrane potential of  $V_m$ ), for the movement of a unit charge displaced by the external potential, where the subscripts 0 and 1 represent initial (activated) and final (resting) states. These probability densities are obtained from the voltage dependent CG free energy functions (parabolas)  $f_\lambda(V)$   $\{\lambda = 0, 1\}$  with  $p_\lambda(V) \propto \exp(-f_\lambda(V))$  (see Fig. 4A).  $G_{conf}$  is identified as the value ( $Q_g V^*$ ) of the reaction coordinate of voltage coupling ( $Q_g V$ ) where two probability densities intersect (i.e., have equal probability), leading to  $Q_g V_{1/2} = G_{conf}$ . A reference free energy function ( $g_1(V)$ ), whose minimum is zero, is shifted by the conformational free energy of  $Q_g V_{1/2} = G_{conf}$ , resulting in  $f_1(V)$ . (B) The corresponding Marcus free energy parabolas along the reaction coordinate of voltage coupling. The parabolas are reconstructed from (A) using the Eqs. (6) and (7) with  $u = Q_g V$ .  $\Lambda$  represents reorganization energy, given by the form of  $\Lambda = -1/2 Q_g V_{min}$  in voltage gated ion channel, leading to the free energy of barrier, given by the form of  $G^\ddagger = \Lambda/4(1 + G_{conf}/\Lambda)^2$  (see Eq. (40) for details). The chemical component of free energy ( $g_0(0) - g_1(0)$ ), the free energy difference between two non-equilibrium conformational states (i.e., at zero membrane potential), is shown to be equal to the free energy difference between two “equilibrium” conformational states along the reaction coordinate of voltage coupling ( $f_1(V_{1,min}) - f_0(V_{0,min})$ ), via the conformational free energy ( $Q_g V_{1/2} = G_{conf}$ ), determined using the equilibrium fluctuation relation of Eq. (15). Free energy function  $f_1(V)$  is shifted by the reference free energy from  $g_1(V)$  (, whose global minimum set to zero) (dashed line), resulting in the intersection with  $f_0(V)$  at  $V = 0$ , i.e.,  $f_0(0) = f_1(0)$ , leading to  $g_0(0) - g_1(0) = Q_g V_{1/2}$ . Free energy functions (for 0 and 1 states) are related to each other by the second free energy relation  $f_0(V) - f_1(V) = -Q_g V$  or  $g_0(V) - g_1(V) = Q_g V_{1/2} - Q_g V$ , which is equivalent to the equilibrium fluctuation relation for voltage activation (i.e., the equilibrium fluctuation relation or the first free energy relation of  $p_0(V)/p_1(V) = \exp(Q_g V - Q_g V_{1/2})$ ).




**Figure 2.**

Description of the model system (i.e., a unit charge embedded in the membrane between electrolytes that incorporates electrodes), used to illustrate the theoretical framework and the CG model of voltage coupling. The electrode potential (as in the Gouy-Chapman model), provided by a virtual battery, is determined by Eqs. (47), (48) and (49) with the appropriate boundary condition (i.e., the denominator in Eq. 47).  $I_g$  (defined as the gating current in the macroscopic continuum approach, of which time integration yields  $Q_g$  (see Eqs. (13), (23), and (27))) is shown to be equal to the current that flows through the electrode (defined as  $I_{ext}$ , of which time integration yields  $Q_{ext}$ ; see Eqs. (22), (26), and (36)). The actual gating current measured in experiment corresponds to  $I_c$  in the external circuit. The area (A) and width (L) of membrane is  $80 \times 80 \text{Å}^2$  and  $40 \text{Å}$  width, respectively, and an electrolyte concentration of 250mM was used. (Top) The unit positive charge is initially positioned near the membrane surface on the left (i.e., state 0). An application of a positive external potential drives the unit charge to the right side (i.e., state 1) of the membrane. (Bottom) Net electrolyte charge distributions [ $e/\text{Å}^3$ ] before (left) and after (right) the unit charge movement at 100mV; the integration of charge distribution along the axis normal to the membrane surface (multiplied by membrane area) yields state dependent gating charge shown in Fig. 3.




**Figure 3.**

Evaluations of the gating charge by several different approaches: (A) The gating charge obtained using the direct CG approach of Eq. (50) at zero potential (and at 100mV in inset). The difference in cumulative electrolyte charges near the electrode that arises in response to charge movements within the membrane gives a gating charge of  $\sim 0.73e$ . (B) The gating charge obtained by the second indirect approach (i.e., using Eq. (54)). In this case a linear least square fits of data points from several different potentials gives approximately zero slope (zero dependence on the external potential) for both the 0 and 1 states, yielding a gating charge of  $\sim 0.76e$  estimated from the gap between the two parallel lines. This implies that the (total) gating charge is independent of external potentials without the consideration of voltage dependent population of each state.

**Figure 4.**


The voltage dependent Marcus-Warshel parabolas, obtained using the CG model of voltage coupling (the KDW model). (A) The free energy of a unit charge movement as a function of external potentials using the CG energetics of voltage coupling, which is approximately quadratic, as predicted by linear response approximation applied to the voltage coupling of charge displacement. Notice that  $f_\lambda(V)$  ( $\lambda=0,1$ ) intersect at  $V=0$  along the reaction coordinate of voltage coupling, which is a natural consequence of the equilibrium fluctuation relation of Eq. (15) that any system of two equilibrium states described by a reaction coordinate of energy gap (voltage coupling in the present study) should satisfy.  $g_1(V)$  (dashed light blue) is obtained by shifting  $f_1(V)$  (blue) by the conformational free energy of  $-G_{conf}$ . (B) The free energy of charging the membrane between two electrolytes (i.e., the model system of (A) but in the absence of any charge within the membrane). The free energy parabola obtained by our KDW model shows an excellent agreement with the capacitor formula (Eq. (55)) obtained from the macroscopic continuum approach by solving a modified Poisson-Boltzmann equation (PB-V), yielding a capacitance of  $\sim 2.34e/V$  (in comparison with  $2.33e/V$  from the capacitor formula of Eq. (55)).



**Figure 5.**

The voltage dependent open (activated) probability (Q-V curve) obtained using the CG model of voltage coupling (see Fig. 4A) with  $p_{\lambda}(V) \propto \exp(-f_{\lambda}(V))$ . The least square fit of the voltage dependent open (final state) probability density of Eq. (58) to the analytic form of Q-V curve of Eq. (59) yields the half voltage ( $V_{1/2}$ ) of  $\sim -9.2$  mV and gating charge of  $\sim -0.78e$ , resulting in the chemical component of free energy of  $\sim -0.2$  kcal/mol using

$G_c \equiv g_0(0) - g_1(0) = Q_g V_{1/2}$ . As expected from Eq. (21),  $G_c$  is approximately equal to the free energy difference ( $\sim -0.2$  kcal/mol) between the two equilibrium conformational states along a reaction coordinate of voltage activation, i.e.,  $f_1(V_{1,\min}) - f_1(V_{0,\min})$ .



**Figure 6.** Schematic illustrations of the quantitative correlation between free energy landscapes (parabolas) and a Q-V curve. (A) The stabilization of an open activated state, for example, upon mutations and drug binding, leads to a left shift (toward hyperpolarization) in the Q-V curve (C). (B) The destabilization of a closed resting state also results in the same shift in the Q-V curve. The shift in the Q-V curve in terms of  $V_{1/2}$  is correlated with the stabilization or destabilization of the conformational states, with a quantitative relation of the form,  $Q_g V_{1/2} \approx \Delta G_{\text{conf}}$  (see Eq. 20).