

# Genitourinary mast cells and survival

Theoharis C. Theoharides<sup>1,2</sup>, Julia M. Stewart<sup>1</sup>

<sup>1</sup>Molecular Immunopharmacology and Drug Discovery Laboratory, Department of Integrative Physiology and Pathobiology, Tufts University School of Medicine, Boston, MA, USA; <sup>2</sup>Department of Internal Medicine, Tufts University School of Medicine, Boston, MA, USA

**Contributions:** (I) Conception and design: TC Theoharides; (II) Administrative support: JM Stewart; (III) Provision of study materials or patients: TC Theoharides; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: TC Theoharides; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

**Correspondence to:** Theoharis C. Theoharides, MS, PhD, MD, FAAAAI. Department of Integrative Physiology and Pathobiology, Tufts University School of Medicine, 136 Harrison Avenue, J304, Boston, MA 02111, USA. Email: theoharis.theoharides@tufts.edu.

**Abstract:** Mast cells (MCs) are ubiquitous in the body, but they have historically been associated with allergies, and most recently with regulation of immunity and inflammation. However, it remains a puzzle why so many MCs are located in the diencephalon, which regulates emotions and in the genitourinary tract, including the bladder, prostate, penis, vagina and uterus that hardly ever get allergic reactions. A number of papers have reported that MCs have estrogen, gonadotropin and corticotropin-releasing hormone (CRH) receptors. Moreover, animal experiments have shown that diencephalic MCs increase in number during courting in doves. We had reported that allergic stimulation of nasal MCs leads to hypothalamic-pituitary adrenal (HPA) activation. Interestingly, anecdotal information indicates that female patients with mastocytosis or mast cell activation syndrome may have increased libido. Preliminary evidence also suggests that MCs may have olfactory receptors. MCs may, therefore, have been retained phylogenetically not only to “smell danger”, but to promote survival and procreation.

**Keywords:** Arousal; cervix; contractions; mast cells (MCs); prostate; uterus

Submitted Sep 16, 2015. Accepted for publication Sep 29, 2015.

doi: 10.3978/j.issn.2223-4683.2015.10.04

**View this article at:** <http://dx.doi.org/10.3978/j.issn.2223-4683.2015.10.04>

## Introduction

Mast cells (MCs) develop from hematopoietic precursors in response to stem cell factor (SCF) the ligand of the CD117 (KIT) tyrosine kinase receptor. Precursors then migrate from the blood into all tissues where they acquire their tissue-specific phenotype, influenced by the local microenvironment (1).

MCs are best known for causing allergic reactions when activated through exposure to an antigen (allergen) that crosslinks allergen-specific immunoglobulin E (IgE), already bound to the high affinity Fc epsilon receptor 1 (FcεRI) (2). MCs can also be activated by anaphylatoxins (C3a, C5a), hormones, physical stimuli (pressure and temperature changes), as well as cytokines and neuropeptides (3) such as corticotropin-releasing hormone (CRH), neurotensin (4), and substance P (SP) (5). MCs express receptors for diverse

ligands (6), including toll-like receptors (7) that can be activated by bacterial and viral products (8). MC stimulation can be enhanced by SCF and IL-33 (9), which together with SP, induce vascular endothelial growth factor (VEGF) release (5) and act as “sensors of cell injury” (10). Each MC contains about 500,000 secretory granules filled with numerous biologically active molecules (11).

MC stimulation leads to secretion of numerous vasoactive, muscle contracting, neurosensitizing and proinflammatory mediators (3,12). In particular, histamine causes muscle contraction, exocrine gland secretion and vasodilation; it also activates the hypothalamic-pituitary adrenal (HPA) axis (13), as does MC-derived IL-6 (14) and CRH (15). In fact, MCs have been implicated in the regulation of HPA axis both in the brain (16), and its equivalent in the skin (17,18).

MCs can secrete the content of individual granules (19)

or individual mediators, such as serotonin, selectively without degranulation (20). MCs can also communicate with neurons by transgranulation (21) or undergo “polarized” exocytosis of proteolytic enzymes at surface sites called “antibody-dependent degranulation synapse” (22). MCs can also secrete nanovesicles (exosomes) (23) containing many different biologically active molecules (24), in a manner that may be guided by antigens embedded in their phospholipid envelope (25). Such exosomes could participate in immune (26,27) and neuropsychiatric diseases (28,29).

The ability of MCs to secrete some mediators selectively (30), permits MCs to participate in diverse processes without causing allergic or inflammatory reactions (12). For instance, we showed that IL-1 can stimulate selective release of IL-6 (31) and so did SCF (32). We also showed that CRH stimulated selective release of VEGF (33) and so did prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) (34), all without any degranulation. Taken together, available data suggest that MCs are capable of releasing a panoply of molecules that may participate in many pathophysiological processes such as innate immunity (1,35), autoimmunity (36), and neuroinflammation (37), but may also have immunomodulatory (38) functions.

### Effect of stress

Emotional stress is the most common trigger of symptoms in patients with systemic mastocytosis, characterized by increased number and degree of activation of MCs (39). In one case, symptoms worsened with stress and there was elevated serum CRH levels, with bone marrow MCs express CRHR-1 (40). MCs (15) and other immune cells (41) can produce CRH (42). Amazingly, even corticosterone has been localized inside MC secretory granules (43).

It is not well understood how animals “smell danger” or are attracted to their mates through odors since the olfactory nerve does not connect directly to the hypothalamus. We and others reported that stimulation of nasal MCs leads to activation of the HPA (44-46) driving the organism into a fight-or-flight mode. Recently olfactory and taste receptors were identified in subpopulations of human circulating leukocytes (47). MCs may turn out to also express such receptors since brain MCs were reported to be influenced by chemosensory cues associated with estrus induction (48).

Surprisingly, MC numbers and reactivity have been reported to undergo daily rhythmic variations (49) and the reactivity of individual MCs was further shown to follow a “circadian clock” (50,51). In this context, it is of interest

that mast cell behavior is affected by the pineal through the expression of melatonin receptors and MCs release melatonin, themselves (52).

Excessive stress can lead to pathological outcomes in various tissues (53). Stress has been reported to induce inflammatory change in rat bladders (54), as well as selective release of VEGF (55), effects that are absent in MC deficient mice (56). Acute stress (57,58) and locally secreted CRH (59) activated MCs (53,59) leading to neurogenic inflammation with subsequent chronic nerve sensitization (60).

MCs (15), immune cells (41), human endometrium, intrauterine pregnancy tissues (61,62), and local nerve endings (42) can produce CRH (42). We reported high levels of CRH and tryptase in products of conception from women with habitual spontaneous abortions (63). Maternal stress has been also linked to preterm delivery (64) and high levels of CRH expression has been reported for placenta, decidua and fetal membranes, where it induces prostaglandin production and promotes labor (65). The decidua of women with high levels of stress have also been reported to have high number of tryptase-positive MC (66), as also reported in aborted deciduas (67). Endometriosis tissue has also been associated with high number of activated MCs (68), which were shown to be increased in response to stress which exacerbated endometriosis in a rat model (69).

### Genitourinary MCs

MCs are present in animal and human bladder (70,71), prostate (72-75), uterus (76-79), penis (80,81), vagina (76,82) and placenta (83). However, their role in these tissues is unknown especially since they are not known to undergo allergic reactions. Uterine MCs are increased during pregnancy and may be important for reproductive processes (84,85). MCs can release muscle contracting and vasodilatory substances that could contribute to clitoral enlargement and uterine contractions (78,86,87). IgE-independent MC activation has been reported to augment contractility of guinea pig (88), mouse (89) and human (86) myometrium. Activation of MCs also leads to angiogenesis in the rat uterine cervix during pregnancy (90). MC degranulation also modulates cervical contractility as shown in the guinea pig (91).

Current evidence from clinical and laboratory studies confirms that MC play a central role in the pathophysiology of bladder pain syndrome/interstitial cystitis (BPS/IC) (92,93) and possibly prostate hyperplasia (74) and sterile prostate inflammation (73). Damaged or dysfunctional

urothelial cells produce cytokines, such as SCF, that can stimulate proliferation and/or activation of MCs (71). In fact, MCs are increased in the detrusor of BPS/IC (71,94-97), and are maximally activated by SCF (98,99) and nerve growth factor (NGF), which is increased in patients with BPS/IC (100). We have shown that CRH activates rat bladder MCs (101) and CRH is involved in signaling in feline bladder urothelial cells (102). In fact, CRH has been considered a mediator of emotional influences on bladder function (103).

### Effect of sex hormones

Human MCs express estrogen receptors (104) activation of which increase MC stimulation (105,106). Estradiol also induced MC migration into the uterus and their degranulation (107). Treatment of mice with leutinizing hormone (LH), follicle stimulating hormone (FSH) or estradiol increased the number and extent of MC degranulation in the ovaries (108). Estrogen receptors are also expressed on bladder (109-111), and lung (112,113) MCs. Human MCs also express progesterone (114,115) and testosterone (116) receptors, but their activation appears to have an inhibitory effect.

One laboratory has reported gonadotropin-releasing hormone-like immunoreactive MCs in the habenula of doves (equivalent to the hypothalamic infundibulum in humans) (117), which increased during courting (118). MCs in ovarian, uterine and brain tissues change their histamine content throughout the rat estrus cycle; moreover, MCs are absent from the thalamus during pro-estrus but are present in the hypothalamus only during the estrus phase (119).

MC-derived mediators, especially histamine, are considered to be important in sexual arousal and coitus (120). Anecdotal information suggests that patients with mastocytosis or MC activation (3) may have increased libido. Uterine MCs were shown to have oxytocin receptors, activation of which prevented serotonin uptake and increased serotonin availability (121) that may positively affect sexual behavior. Circulating levels of oxytocin are known to increase during sexual arousal and orgasm in both men and women (122). It is interesting that intranasal oxytocin was reported to increase libido and related sexual behavior in a male subject (123).

### Conclusions

MCs have been retained throughout the phylogenetic

tree (124). Moreover, MC ability to produce numerous hormonal, immune and neural substances resemble that of the unicellular organism *Tetrahymena* dating from some 500 million years ago (125,126). MCs are present in all mammals and may be necessary for survival of the species by regulating immunity (127), protecting the organism against external triggers (53), supporting pregnancy (128), augmenting delivery and also ensuring optimal conditions for procreation.

### Acknowledgements

None.

### Footnote

*Conflicts of Interest:* The authors have no conflicts of interest to declare.

### References

1. Galli SJ, Borregaard N, Wynn TA. Phenotypic and functional plasticity of cells of innate immunity: macrophages, mast cells and neutrophils. *Nat Immunol* 2011;12:1035-44.
2. Rivera J, Fierro NA, Olivera A, et al. New insights on mast cell activation via the high affinity receptor for IgE. *Adv Immunol* 2008;98:85-120.
3. Theoharides TC, Valent P, Akin C. Mast Cells, Mastocytosis, and Related Disorders. *N Engl J Med* 2015;373:163-72.
4. Donelan J, Boucher W, Papadopoulou N, et al. Corticotropin-releasing hormone induces skin vascular permeability through a neurotensin-dependent process. *Proc Natl Acad Sci U S A* 2006;103:7759-64.
5. Theoharides TC, Zhang B, Kempuraj D, et al. IL-33 augments substance P-induced VEGF secretion from human mast cells and is increased in psoriatic skin. *Proc Natl Acad Sci U S A* 2010;107:4448-53.
6. Migalovich-Sheikhet H, Friedman S, Mankuta D, et al. Novel identified receptors on mast cells. *Front Immunol* 2012;3:238.
7. Sandig H, Bulfone-Paus S. TLR signaling in mast cells: common and unique features. *Front Immunol* 2012;3:185.
8. Abraham SN, St John AL. Mast cell-orchestrated immunity to pathogens. *Nat Rev Immunol* 2010;10:440-52.
9. Fux M, Pecaric-Petkovic T, Odermatt A, et al. IL-33 is a mediator rather than a trigger of the acute allergic

- response in humans. *Allergy* 2014;69:216-22.
10. Enoksson M, Lyberg K, Möller-Westerberg C, et al. Mast cells as sensors of cell injury through IL-33 recognition. *J Immunol* 2011;186:2523-8.
  11. Wernersson S, Pejler G. Mast cell secretory granules: armed for battle. *Nat Rev Immunol* 2014;14:478-94.
  12. Theoharides TC, Alysandratos KD, Angelidou A, et al. Mast cells and inflammation. *Biochim Biophys Acta* 2012;1822:21-33.
  13. Scaccianoce S, Lombardo K, Nicolai R, et al. Studies on the involvement of histamine in the hypothalamic-pituitary-adrenal axis activation induced by nerve growth factor. *Life Sci* 2000;67:3143-52.
  14. Mastorakos G, Chrousos GP, Weber JS. Recombinant interleukin-6 activates the hypothalamic-pituitary-adrenal axis in humans. *J Clin Endocrinol Metab* 1993;77:1690-4.
  15. Kempuraj D, Papadopoulou NG, Lytinas M, et al. Corticotropin-releasing hormone and its structurally related urocortin are synthesized and secreted by human mast cells. *Endocrinology* 2004;145:43-8.
  16. Theoharides TC, Donelan JM, Papadopoulou N, et al. Mast cells as targets of corticotropin-releasing factor and related peptides. *Trends Pharmacol Sci* 2004;25:563-8.
  17. Banerjee A, Romero-Lorenzo E, Sur M. MeCP2: making sense of missense in Rett syndrome. *Cell Res* 2013;23:1244-6.
  18. Arck PC, Slominski A, Theoharides TC, et al. Neuroimmunology of stress: skin takes center stage. *J Invest Dermatol* 2006;126:1697-704.
  19. Theoharides TC, Douglas WW. Secretion in mast cells induced by calcium entrapped within phospholipid vesicles. *Science* 1978;201:1143-5.
  20. Theoharides TC, Bondy PK, Tsakalos ND, et al. Differential release of serotonin and histamine from mast cells. *Nature* 1982;297:229-31.
  21. Wilhelm M, Silver R, Silverman AJ. Central nervous system neurons acquire mast cell products via transgranulation. *Eur J Neurosci* 2005;22:2238-48.
  22. Joulia R, Gaudenzio N, Rodrigues M, et al. Mast cells form antibody-dependent degranulatory synapse for dedicated secretion and defence. *Nat Commun* 2015;6:6174.
  23. Skokos D, Goubran-Botros H, Roa M, et al. Immunoregulatory properties of mast cell-derived exosomes. *Mol Immunol* 2002;38:1359-62.
  24. Shefler I, Salamon P, Hershko AY, et al. Mast cells as sources and targets of membrane vesicles. *Curr Pharm Des* 2011;17:3797-804.
  25. Bryniarski K, Ptak W, Jayakumar A, et al. Antigen-specific, antibody-coated, exosome-like nanovesicles deliver suppressor T-cell microRNA-150 to effector T cells to inhibit contact sensitivity. *J Allergy Clin Immunol* 2013;132:170-81.
  26. Fujita Y, Yoshioka Y, Ito S, et al. Intercellular communication by extracellular vesicles and their microRNAs in asthma. *Clin Ther* 2014;36:873-81.
  27. Benito-Martin A, Di Giannatale A, Ceder S, et al. The new deal: a potential role for secreted vesicles in innate immunity and tumor progression. *Front Immunol* 2015;6:66.
  28. Tsilioni I, Panagiotidou S, Theoharides TC. Exosomes in neurologic and psychiatric disorders. *Clin Ther* 2014;36:882-8.
  29. Kawikova I, Askenase PW. Diagnostic and therapeutic potentials of exosomes in CNS diseases. *Brain Res* 2015;1617:63-71.
  30. Theoharides TC, Kempuraj D, Tagen M, et al. Differential release of mast cell mediators and the pathogenesis of inflammation. *Immunol Rev* 2007;217:65-78.
  31. Kandere-Grzybowska K, Letourneau R, Kempuraj D, et al. IL-1 induces vesicular secretion of IL-6 without degranulation from human mast cells. *J Immunol* 2003;171:4830-6.
  32. Gagari E, Tsai M, Lantz CS, et al. Differential release of mast cell interleukin-6 via c-kit. *Blood* 1997;89:2654-63.
  33. Cao J, Papadopoulou N, Kempuraj D, et al. Human mast cells express corticotropin-releasing hormone (CRH) receptors and CRH leads to selective secretion of vascular endothelial growth factor. *J Immunol* 2005;174:7665-75.
  34. Abdel-Majid RM, Marshall JS. Prostaglandin E2 induces degranulation-independent production of vascular endothelial growth factor by human mast cells. *J Immunol* 2004;172:1227-36.
  35. Kumar V, Sharma A. Mast cells: emerging sentinel innate immune cells with diverse role in immunity. *Mol Immunol* 2010;48:14-25.
  36. Rottem M, Mekori YA. Mast cells and autoimmunity. *Autoimmun Rev* 2005;4:21-7.
  37. Theoharides TC, Stewart JM, Hatziagelaki E, et al. Brain "fog," inflammation and obesity: key aspects of neuropsychiatric disorders improved by luteolin. *Front Neurosci* 2015;9:225.
  38. Galli SJ, Grimbaldston M, Tsai M. Immunomodulatory mast cells: negative, as well as positive, regulators of immunity. *Nat Rev Immunol* 2008;8:478-86.
  39. Jennings S, Russell N, Jennings B, et al. The Mastocytosis Society survey on mast cell disorders: patient experiences

- and perceptions. *J Allergy Clin Immunol Pract* 2014;2:70-6.
40. Theoharides TC, Petra AI, Stewart JM, et al. High serum corticotropin-releasing hormone (CRH) and bone marrow mast cell CRH receptor expression in a mastocytosis patient. *J Allergy Clin Immunol* 2014;134:1197-9.
  41. Karalis K, Muglia LJ, Bae D, et al. CRH and the immune system. *J Neuroimmunol* 1997;72:131-6.
  42. Chrousos GP. The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *N Engl J Med* 1995;332:1351-62.
  43. Csaba G, Oláh I, Kiss J, et al. The localization of H3-corticosterone in mast cell granules by electron microscopic autoradiography. *Experientia* 1967;23:944.
  44. Matsumoto I, Inoue Y, Shimada T, et al. Brain mast cells act as an immune gate to the hypothalamic-pituitary-adrenal axis in dogs. *J Exp Med* 2001;194:71-8.
  45. Bugajski AJ, Chłap Z, Gadek-Michalska A, et al. Degranulation and decrease in histamine levels of thalamic mast cells coincides with corticosterone secretion induced by compound 48/80. *Inflamm Res* 1995;44 Suppl 1:S50-1.
  46. Kalogeromitros D, Syrigou EK, Makris M, et al. Nasal provocation of patients with allergic rhinitis and the hypothalamic-pituitary-adrenal axis. *Ann Allergy Asthma Immunol* 2007;98:269-73.
  47. Malki A, Fiedler J, Fricke K, et al. Class I odorant receptors, TAS1R and TAS2R taste receptors, are markers for subpopulations of circulating leukocytes. *J Leukoc Biol* 2015;97:533-45.
  48. Kriegsfeld LJ, Hotchkiss AK, Demas GE, et al. Brain mast cells are influenced by chemosensory cues associated with estrus induction in female prairie voles (*Microtus ochrogaster*). *Horm Behav* 2003;44:377-84.
  49. Catini C, Legnaioli M. Role of mast cells in health: daily rhythmic variations in their number, exocytotic activity, histamine and serotonin content in the rat thyroid gland. *Eur J Histochem* 1992;36:501-16.
  50. Molyva D, Kalokasidis K, Poullos C, et al. Rupatadine effectively prevents the histamine-induced up regulation of histamine H1R and bradykinin B2R receptor gene expression in the rat paw. *Pharmacol Rep* 2014;66:952-5.
  51. Nakao A, Nakamura Y, Shibata S. The circadian clock functions as a potent regulator of allergic reaction. *Allergy* 2015;70:467-73.
  52. Maldonado MD, Mora-Santos M, Naji L, et al. Evidence of melatonin synthesis and release by mast cells. Possible modulatory role on inflammation. *Pharmacol Res* 2010;62:282-7.
  53. Theoharides TC, Cochrane DE. Critical role of mast cells in inflammatory diseases and the effect of acute stress. *J Neuroimmunol* 2004;146:1-12.
  54. Ercan F, Akici A, Ersoy Y, et al. Inhibition of substance P activity prevents stress-induced bladder damage. *Regul Pept* 2006;133:82-9.
  55. Cao J, Boucher W, Kempuraj D, et al. Acute stress and intravesical corticotropin-releasing hormone induces mast cell dependent vascular endothelial growth factor release from mouse bladder explants. *J Urol* 2006;176:1208-13.
  56. Kandere-Grzybowska K, Gheorghe D, Priller J, et al. Stress-induced dura vascular permeability does not develop in mast cell-deficient and neurokinin-1 receptor knockout mice. *Brain Res* 2003;980:213-20.
  57. Ercan F, San T, Cavdar S. The effects of cold-restraint stress on urinary bladder wall compared with interstitial cystitis morphology. *Urol Res* 1999;27:454-61.
  58. Spanos C, Pang X, Ligris K, et al. Stress-induced bladder mast cell activation: implications for interstitial cystitis. *J Urol* 1997;157:669-72.
  59. Theoharides TC, Singh LK, Boucher W, et al. Corticotropin-releasing hormone induces skin mast cell degranulation and increased vascular permeability, a possible explanation for its proinflammatory effects. *Endocrinology* 1998;139:403-13.
  60. Theoharides TC, Sant GR. Neuroimmune Connections and Regulation of Function in the Urinary Bladder. In: Bienenstock J, Goetzl E, Blennerhassett M, editors. *Autonomic Neuroimmunology*. Lausanne: Hardwood Academic Publishers, 2003:345-69.
  61. Hillhouse EW, Grammatopoulos DK. Role of stress peptides during human pregnancy and labour. *Reproduction* 2002;124:323-9.
  62. Reis FM, Petraglia F. The placenta as a neuroendocrine organ. *Front Horm Res* 2001;27:216-28.
  63. Madhappan B, Kempuraj D, Christodoulou S, et al. High levels of intrauterine corticotropin-releasing hormone, urocortin, tryptase, and interleukin-8 in spontaneous abortions. *Endocrinology* 2003;144:2285-90.
  64. Sun L, Tao F, Hao J, et al. Vaginal bleeding in early pregnancy and associations with physical, psychological and environmental factors among Chinese women: from the C-ABC cohort study. *Gynecol Obstet Invest* 2012;73:330-6.
  65. Lockwood CJ. Stress-associated preterm delivery: the role of corticotropin-releasing hormone. *Am J Obstet Gynecol* 1999;180:S264-6.
  66. Arck PC, Rose M, Hertwig K, et al. Stress and immune mediators in miscarriage. *Hum Reprod* 2001;16:1505-11.

67. Marx L, Arck P, Kieslich C, et al. Decidual mast cells might be involved in the onset of human first-trimester abortion. *Am J Reprod Immunol* 1999;41:34-40.
68. Kempuraj D, Papadopoulou N, Stanford EJ, et al. Increased numbers of activated mast cells in endometriosis lesions positive for corticotropin-releasing hormone and urocortin. *Am J Reprod Immunol* 2004;52:267-75.
69. Cuevas M, Flores I, Thompson KJ, et al. Stress exacerbates endometriosis manifestations and inflammatory parameters in an animal model. *Reprod Sci* 2012;19:851-62.
70. Theoharides TC, Sant GR, el-Mansoury M, et al. Activation of bladder mast cells in interstitial cystitis: a light and electron microscopic study. *J Urol* 1995;153:629-36.
71. Theoharides TC, Kempuraj D, Sant GR. Mast cell involvement in interstitial cystitis: a review of human and experimental evidence. *Urology* 2001;57:47-55.
72. Hofmeister MA, He F, Ratliff TL, et al. Mast cells and nerve fibers in interstitial cystitis (IC): an algorithm for histologic diagnosis via quantitative image analysis and morphometry (QIAM). *Urology* 1997;49:41-7.
73. Theoharides TC, Flaris N, Cronin CT, et al. Mast cell activation in sterile bladder and prostate inflammation. *Int Arch Allergy Appl Immunol* 1990;92:281-6.
74. Papadoukakis S, Kyroudi-Voulgari A, Truss MC, et al. Quantitative study of mast cells in experimentally induced benign prostatic hyperplasia. *Urol Int* 2010;84:100-4.
75. Shafik A, Shafik I, el-Sibai O. Identification of c-kit-positive cells in the human prostate: the interstitial cells of Cajal. *Arch Androl* 2005;51:345-51.
76. Walter J, Klein C, Wehrend A. Distribution of mast cells in vaginal, cervical and uterine tissue of non-pregnant mares: investigations on correlations with ovarian steroids. *Reprod Domest Anim* 2012;47:e29-31.
77. Menzies FM, Higgins CA, Shepherd MC, et al. Mast cells reside in myometrium and cervix, but are dispensable in mice for successful pregnancy and labor. *Immunol Cell Biol* 2012;90:321-9.
78. Garfield RE, Irani AM, Schwartz LB, et al. Structural and functional comparison of mast cells in the pregnant versus nonpregnant human uterus. *Am J Obstet Gynecol* 2006;194:261-7.
79. Sivridis E, Giatromanolaki A, Agnantis N, et al. Mast cell distribution and density in the normal uterus--metachromatic staining using lectins. *Eur J Obstet Gynecol Reprod Biol* 2001;98:109-13.
80. Shafik A. Study of interstitial cells in the penis: human study. *J Sex Med* 2007;4:66-71.
81. Sathananthan AH, Adaikan PG, Lau LC, et al. Fine structure of the human corpus cavernosum. *Arch Androl* 1991;26:107-17.
82. Shafik A, El-Sibai O, Shafik I, et al. Immunohistochemical identification of the pacemaker cajal cells in the normal human vagina. *Arch Gynecol Obstet* 2005;272:13-6.
83. Purcell WM, Hanahoe TH. A novel source of mast cells: the human placenta. *Agents Actions* 1991;33:8-12.
84. Woidacki K, Jensen F, Zenclussen AC. Mast cells as novel mediators of reproductive processes. *Front Immunol* 2013;4:29.
85. Menzies FM, Shepherd MC, Nibbs RJ, et al. The role of mast cells and their mediators in reproduction, pregnancy and labour. *Hum Reprod Update* 2011;17:383-96.
86. Bytautiene E, Vedernikov YP, Saade GR, et al. Degranulation of uterine mast cell modifies contractility of isolated myometrium from pregnant women. *Am J Obstet Gynecol* 2004;191:1705-10.
87. Rudolph MI, Reinicke K, Cruz MA, et al. Distribution of mast cells and the effect of their mediators on contractility in human myometrium. *Br J Obstet Gynaecol* 1993;100:1125-30.
88. Bytautiene E, Vedernikov YP, Saade GR, et al. IgE-independent mast cell activation augments contractility of nonpregnant and pregnant guinea pig myometrium. *Int Arch Allergy Immunol* 2008;147:140-6.
89. Allix S, Reyes-Gomez E, Aubin-Houzelstein G, et al. Uterine contractions depend on KIT-positive interstitial cells in the mouse: genetic and pharmacological evidence. *Biol Reprod* 2008;79:510-7.
90. Varayoud J, Ramos JG, Bosquiaz VL, et al. Mast cells degranulation affects angiogenesis in the rat uterine cervix during pregnancy. *Reproduction* 2004;127:379-87.
91. Bytautiene E, Vedernikov YP, Saade GR, et al. Endogenous mast cell degranulation modulates cervical contractility in the guinea pig. *Am J Obstet Gynecol* 2002;186:438-45.
92. Whitmore KE, Theoharides TC. When to suspect interstitial cystitis. *J Fam Pract* 2011;60:340-8.
93. Sant GR, Hanno PM. Interstitial cystitis: current issues and controversies in diagnosis. *Urology* 2001;57:82-8.
94. Larsen S, Thompson SA, Hald T, et al. Mast cells in interstitial cystitis. *Br J Urol* 1982;54:283-6.
95. Feltis JT, Perez-Marrero R, Emerson LE. Increased mast cells of the bladder in suspected cases of interstitial cystitis: a possible disease marker. *J Urol* 1987;138:42-3.
96. Lynes WL, Flynn SD, Shortliffe LD, et al. Mast cell involvement in interstitial cystitis. *J Urol* 1987;138:746-52.
97. Larsen MS, Mortensen S, Nordling J, et al. Quantifying mast cells in bladder pain syndrome by

- immunohistochemical analysis. *BJU Int* 2008;102:204-7; discussion 207.
98. Pang X, Sant G, Theoharides TC. Altered expression of bladder mast cell growth factor receptor (c-kit) in interstitial cystitis. *Urology* 1998;51:939-44.
  99. Peeker R, Enerbäck L, Fall M, et al. Recruitment, distribution and phenotypes of mast cells in interstitial cystitis. *J Urol* 2000;163:1009-15.
  100. Lowe EM, Anand P, Terenghi G, et al. Increased nerve growth factor levels in the urinary bladder of women with idiopathic sensory urgency and interstitial cystitis. *Br J Urol* 1997;79:572-7.
  101. Boucher W, Kempuraj D, Michaelian M, et al. Corticotropin-releasing hormone-receptor 2 is required for acute stress-induced bladder vascular permeability and release of vascular endothelial growth factor. *BJU Int* 2010;106:1394-9.
  102. Hanna-Mitchell AT, Wolf-Johnston A, Roppolo JR, et al. Corticotropin-releasing factor family peptide signaling in feline bladder urothelial cells. *J Endocrinol* 2014;222:113-21.
  103. Klausner AP, Steers WD. Corticotropin releasing factor: a mediator of emotional influences on bladder function. *J Urol* 2004;172:2570-3.
  104. Zaitsu M, Narita S, Lambert KC, et al. Estradiol activates mast cells via a non-genomic estrogen receptor-alpha and calcium influx. *Mol Immunol* 2007;44:1977-85.
  105. Narita S, Goldblum RM, Watson CS, et al. Environmental estrogens induce mast cell degranulation and enhance IgE-mediated release of allergic mediators. *Environ Health Perspect* 2007;115:48-52.
  106. Alevizos M, Karagkouni A, Kontou-Fili K, et al. A probable case report of stress-induced anaphylaxis. *Ann Allergy Asthma Immunol* 2014;112:383-4.
  107. Jensen F, Woudwyk M, Teles A, et al. Estradiol and progesterone regulate the migration of mast cells from the periphery to the uterus and induce their maturation and degranulation. *PLoS One* 2010;5:e14409.
  108. Jaiswal K, Krishna A. Effects of hormones on the number, distribution and degranulation of mast cells in the ovarian complex of mice. *Acta Physiol Hung* 1996;84:183-90.
  109. Theoharides TC, Pang X, Letourneau R, et al. Interstitial cystitis: a neuroimmunoendocrine disorder. *Ann N Y Acad Sci* 1998;840:619-34.
  110. Spanos C, el-Mansoury M, Letourneau R, et al. Carbachol-induced bladder mast cell activation: augmentation by estradiol and implications for interstitial cystitis. *Urology* 1996;48:809-16.
  111. Pang X, Cotreau-Bibbo MM, Sant GR, et al. Bladder mast cell expression of high affinity oestrogen receptors in patients with interstitial cystitis. *Br J Urol* 1995;75:154-61.
  112. Hox V, Desai A, Bandara G, et al. Estrogen increases the severity of anaphylaxis in female mice through enhanced endothelial nitric oxide synthase expression and nitric oxide production. *J Allergy Clin Immunol* 2015;135:729-36.e5.
  113. Zhao XJ, McKerr G, Dong Z, et al. Expression of oestrogen and progesterone receptors by mast cells alone, but not lymphocytes, macrophages or other immune cells in human upper airways. *Thorax* 2001;56:205-11.
  114. Vasiadi M, Kempuraj D, Boucher W, et al. Progesterone inhibits mast cell secretion. *Int J Immunopathol Pharmacol* 2006;19:787-94.
  115. Vliagoftis H, Dimitriadou V, Theoharides TC. Progesterone triggers selective mast cell secretion of 5-hydroxytryptamine. *Int Arch Allergy Appl Immunol* 1990;93:113-9.
  116. Guhl S, Artuc M, Zuberbier T, et al. Testosterone exerts selective anti-inflammatory effects on human skin mast cells in a cell subset dependent manner. *Exp Dermatol* 2012;21:878-80.
  117. Silverman AJ, Millar RP, King JA, et al. Mast cells with gonadotropin-releasing hormone-like immunoreactivity in the brain of doves. *Proc Natl Acad Sci U S A* 1994;91:3695-9.
  118. Silver R, Ramos CL, Silverman AJ. Sexual behavior triggers the appearance of non-neuronal cells containing gonadotropin-releasing hormone-like immunoreactivity. *J Neuroendocrinol* 1992;4:207-10.
  119. Aydin Y, Tunçel N, Gürer F, et al. Ovarian, uterine and brain mast cells in female rats: cyclic changes and contribution to tissue histamine. *Comp Biochem Physiol A Mol Integr Physiol* 1998;120:255-62.
  120. Devidze N, Lee AW, Zhou J, et al. CNS arousal mechanisms bearing on sex and other biologically regulated behaviors. *Physiol Behav* 2006;88:283-93.
  121. Rudolph MI, Oviedo C, Vega E, et al. Oxytocin inhibits the uptake of serotonin into uterine mast cells. *J Pharmacol Exp Ther* 1998;287:389-94.
  122. Meston CM, Frohlich PF. The neurobiology of sexual function. *Arch Gen Psychiatry* 2000;57:1012-30.
  123. MacDonald K, Feifel D. Dramatic improvement in sexual function induced by intranasal oxytocin. *J Sex Med* 2012;9:1407-10.
  124. Wong GW, Zhuo L, Kimata K, et al. Ancient origin of mast cells. *Biochem Biophys Res Commun* 2014;451:314-8.
  125. Csaba G. The unicellular Tetrahymena as a model cell for

- receptor research. *Int Rev Cytol* 1985;95:327-77.
126. Csaba G. Biogenic amines at a low level of evolution: Production, functions and regulation in the unicellular Tetrahymena. *Acta Microbiol Immunol Hung* 2015;62:93-108.
127. Theoharides TC. The mast cell: a neuroimmunoendocrine master player. *Int J Tissue React* 1996;18:1-21.
128. Rudolph MI, Rojas IG, Penissi AB. Uterine mast cells: a new hypothesis to understand how we are born. *Biocell* 2004;28:1-11.

**Cite this article as:** Theoharides TC, Stewart JM. Genitourinary mast cells and survival. *Transl Androl Urol* 2015;4(5):579-586. doi: 10.3978/j.issn.2223-4683.2015.10.04