



Review:

Impact of DNA mismatch repair system alterations on human fertility and related treatments*

Min-hao HU, Shu-yuan LIU, Ning WANG, Yan WU, Fan JIN^{†‡}

(Key Laboratory of Reproductive Genetics (Zhejiang), Ministry of Education, and Centre of Reproductive Medicine, Women's Hospital, School of Medicine, Zhejiang University, Hangzhou 310006, China)

[†]E-mail: jinfan@zju.edu.cn

Received June 28, 2015; Revision accepted Nov. 7, 2015; Crosschecked Dec. 16, 2015

Abstract: DNA mismatch repair (MMR) is one of the biological pathways, which plays a critical role in DNA homeostasis, primarily by repairing base-pair mismatches and insertion/deletion loops that occur during DNA replication. MMR also takes part in other metabolic pathways and regulates cell cycle arrest. Defects in MMR are associated with genomic instability, predisposition to certain types of cancers and resistance to certain therapeutic drugs. Moreover, genetic and epigenetic alterations in the MMR system demonstrate a significant relationship with human fertility and related treatments, which helps us to understand the etiology and susceptibility of human infertility. Alterations in the MMR system may also influence the health of offspring conceived by assisted reproductive technology in humans. However, further studies are needed to explore the specific mechanisms by which the MMR system may affect human infertility. This review addresses the physiological mechanisms of the MMR system and associations between alterations of the MMR system and human fertility and related treatments, and potential effects on the next generation.

Key words: DNA mismatch repair, Infertility, Assisted reproductive technology
<http://dx.doi.org/10.1631/jzus.B1500162>

CLC number: R711.6

1 Introduction

Cellular DNA damage accumulates as a result of exposure to exogenous agents (biological, physical, or chemical) and endogenous sources including oxidative stress and errors associated with DNA processing. DNA damage, if unrepaired, creates the possibility of mutagenesis in somatic or germline cells, which can alter normal function and result in diseases, even in the next generation. Consequently, there are multiple mechanisms to repair DNA damage and maintain the stability of the DNA sequence in all

tissues and cells. One of them, known as DNA mismatch repair (MMR), mainly repairs base-pair mismatches and insertion/deletion loops (IDLs) (Jascur and Boland, 2006). MMR repairs DNA mismatches arising during replication, thereby preventing deleterious mutations and maintaining genomic stability. The MMR system also plays a critical role in the meiotic process and gametogenesis. Defects in the MMR system are related to predisposition to certain types of cancer, resistance to certain therapeutic drugs, and infertility, collectively known as DNA MMR defective abnormalities (Maduro *et al.*, 2003; Li, 2008).

Infertility is a common and complex condition affecting about 10%–15% of couples of reproductive age (Gnoth *et al.*, 2005), and the causes of a considerable proportion of infertility cases remain unknown. Based on their important physiological functions and a growing amount of evidence, MMR genes and

[‡] Corresponding author

* Project supported by the National Basic Research Programs (973) of China (Nos. 2012CB944901 and 2014CB943302), the National Natural Science Foundation of China (Nos. 81200475, 81370760, and 81571500), and the Zhejiang Provincial Natural Science Foundation of China (No. LZ13H040001)

ORCID: Min-hao HU, <http://orcid.org/0000-0003-0426-0275>

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2016

proteins demonstrate an impact on human infertility, even in the next generation conceived by assisted reproductive technology (ART). The molecular mechanisms of the DNA damage and repair in spermatogenesis have been reviewed recently (Gunes *et al.*, 2015), focusing on the process of spermatogenesis, origin of DNA damage, and five types of DNA repair mechanisms. This review will concentrate on the relationship between DNA MMR alterations and human fertility, fertility-related treatments, and the potential influence on ART offspring.

2 Components and mechanisms of the MMR system

First identified in *Escherichia coli*, MMR is a biological pathway highly conserved throughout evolution to maintain genomic integrity. The bacterial MMR system is the best studied biochemically (Lahue *et al.*, 1989; Modrich, 1991), and yeast and mouse systems have provided valuable insights because of the power of genetic models (Kolodner and Marsischky, 1999). In humans, the MMR system is an excision and re-synthesis system that can be divided into three steps: recognition of the mismatch, excision of the incorrect fragment, and DNA re-synthesis (Table 1). The human MMR protein, MutS, which is a heterodimer composed of MutS homologues MSH2 and MSH6 (MutS α) or MSH2 and MSH3 (MutS β), is an ATPase that plays a critical role in mismatch recognition and initiation of repair. MutS α preferentially recognizes base-pair mismatches and short IDLs, while MutS β recognizes larger IDLs. Then the DNA-MutS complex recruits MutL α , a heterodimer of MutL homologues MLH1 and PMS2. Other potential proteins like exonuclease 1 (EXO1), DNA polymerase δ (Pol δ) and its cofactors proliferation cell nuclear antigen (PCNA), and replication factor C (RFC) are recruited to accomplish the repair activity. Other members of the MMR system are yet to be found or confirmed such as DNA methyltransferase 1 (Dnmt1) (Guo *et al.*, 2004; Kim *et al.*, 2004).

In addition to DNA MMR activity, MMR is also associated with a series of DNA damage signaling pathways. Plenty of articles have reported interactions between MMR and DNA damage regulators including MLH1 and ATM (Brown *et al.*, 2003),

MSH2 and ATR, Bcl-2 (Wang and Qin, 2003; Youn *et al.*, 2005), MLH1, PMS2 and p53, p73 (Shimodaira *et al.*, 2003; Chen and Sadowski, 2005), MLH1, PMS1, and PMS2 in cell cycle arrest (Stojic *et al.*, 2005; Cannavo *et al.*, 2007), all of which are involved in cell signaling/cycle arrest/apoptosis. Thus, the MMR system recognizes and repairs mismatches and eliminates severely damaged cells, preventing mutagenesis in the short term and tumorigenesis in the long term.

Table 1 Human MMR components and functions

MMR component	Function
MutS α (MSH2-MSH6)	DNA mismatch recognition
MutS β (MSH2-MSH3)	
MutL α (MLH1-PMS2)	Molecular matchmaker;
MutL β (MLH1-PMS1)	recruitment of other related
MutL γ (MLH1-MLH3)	proteins
EXO1	DNA mismatch excision
Pol δ	DNA re-synthesis
PCNA	Initiation of DNA re-synthesis
RFC	Loading and unloading PCNA
RPA	Single strand DNA protection;
	termination of DNA excision
DNA ligase I	Nick ligation
Other members unidentified	

MMR proteins also take part in the meiotic process and are involved in gametogenesis. They have been implicated in somatic hypermutation, interstrand-crosslink repair, immunoglobulin class switching, trinucleotide repeat (TNR) expansion, and other processes. With so many proteins and regulators involved in the MMR system, it is not surprising that mistakes may occur and manifest themselves in different phenotypes. Loss of MMR function leads to failure to repair base-pair mismatches and IDLs, including short repetitive sequences known as microsatellites. Shortening or lengthening of microsatellites, referred to as microsatellite instability (MSI), is the hallmark of MMR system deficiency. The MSI status is commonly determined by five microsatellite markers (BAT25, BAT26, D2S123, D5S346, and D17S250).

The contribution of defective MMR to the development of human cancer has been recognized for more than two decades (Peltomaki, 2003). Alterations in MLH1, MSH2, MSH6, and PMS2 lead to the most common form of cancer, hereditary nonpolyposis colorectal cancer (HNPCC) or Lynch syndrome (LS).

Loss of MMR function caused by either genetic or epigenetic variation of MMR genes is associated with various human cancers such as colorectal, endometrial, ovarian, cervical, breast, gastric, urological, skin, and other rare cancers (Watson and Lynch, 1994; Karamurzin *et al.*, 2012). Defects in MMR can also trigger a multidrug resistance phenotype, resulting cellular resistance to certain alkylating, methylating, and platinating agents, antimetabolites, topoisomerase inhibitors, and DNA minor groove binders (Valentini *et al.*, 2006) since the system is involved in DNA repair, cell cycle arrest, and many other metabolic pathways. In addition to its involvement in human cancer and drug resistance, the role of the MMR system in the meiotic process and gametogenesis should never be neglected. MMR alterations could manifest themselves in human infertility and infertility treatment. More research is needed to understand the origin and development of infertility.

3 MMR alterations and infertility

Infertility is a worldwide problem that occurs in 10%–15% of couples. Male factors account for about half of all infertility cases. However, the etiology of male infertility remains unknown in about half of all cases. Such cases are classified as idiopathic. This is particularly relevant to men with non-obstructive azoospermia or severe oligospermia. The human testis is composed mainly of three types of cells: spermatogenic cells to produce spermatids; Sertoli cells to protect, support, and supply spermatogenic cells, making sure that spermatids are well produced; and Leydig cells to secrete androgen. Male infertility could be an outcome of genetic, epigenetic, endocrine, or environmental changes that lead to aberrant sperm production or function. Substantial evidence has indicated that several members of the MMR family participate in the meiotic recombination process and gametogenesis (Kunkel and Erie, 2005; Iyer *et al.*, 2006; Jiricny, 2006). A defective MMR system has been shown to be related to spermatogenic failure and spermatid dysfunction in some infertile individuals and animal models (Mukherjee *et al.*, 2010). Later, we will discuss recent progress in understanding the genetic and epigenetic variation in the MMR system associated with male infertility.

Animal knockout models have shown the importance of MMR genes in both male and female infertility. First discovered in 1995, mice with defective *PMS2* exhibit male infertility due to failure of chromosomal synapsis (Baker *et al.*, 1995). Disruption of other genes, including *MLH1*, *MLH3*, and *EXO1* (Baker *et al.*, 1996; Lipkin *et al.*, 2002; Wei *et al.*, 2003), also leads to phenotypes of infertility (Table 2). However, there has been little progress in understanding links between the MMR system and human male infertility. Maduro *et al.* (2003) proposed that MSI and defective MMR proteins like *MLH1* and *MSH2* are presented in both testicular and peripheral blood cells in some azoospermic men, predominantly in Sertoli cell-only patients. This indicates that defects in MMR may underlie some forms of male infertility such as Sertoli cell-only, hypospermatogenesis type, and maturation arrest. Recently, Sertoli cells have been recognized as having several common metabolic features analogous to those of cancer cells and could be used as a good model for exploring new perspectives of the Warburg effect (Oliveira *et al.*, 2015). Sertoli cells may play a significant role in obesity-induced male infertility (Martins *et al.*, 2015). The above studies signify important linkages between MMR alterations and human infertility and Sertoli cells, which deserve further detailed study.

Table 2 MMR genes involved in mouse infertility models

Gene	Phenotype
<i>MSH4</i>	Infertile: failure of spermatogonial maturation beyond zygonema
<i>MSH5</i>	Infertile: incomplete and non-homologous chromosomal pairing
<i>MLH1</i>	Infertile: failure of crossing over
<i>PMS2</i>	Male infertile: disruption of normal chromosomal synapsis
<i>MLH3</i>	Infertile
<i>EXO1</i>	Infertile

MMR proteins participate in the meiotic recombination process in yeast and mammals. Among these proteins there are two MutS homologues (*MSH4*, *MSH5*) and three MutL homologues (*MLH1*, *MLH3*, *PMS2*) (Surtees *et al.*, 2004; Her *et al.*, 2007). Terribas *et al.* (2010) studied 13 infertile patients each with one of two types of spermatogenic arrest: maturation arrest or hypospermatogenesis. These two types showed significantly decreased MMR expression

values (MLH1, MLH3, MSH4, MSH5) in testicular samples, except for PMS2, and the more serious type, the maturation arrest group, showed a greater reduction. Ferras *et al.* (2007) showed that the testicular *MLH3* gene in 13 infertile patients with spermatogenic arrest contained four missense (T2896C, C2531T) and eight intronic (IVS9+66G/A) variants. Among these 13 patients, the combination of C2531T and IVS9+66G/A variants was identified only in patients with primary spermatogenic arrest, indicating that the presence of two simultaneous *MLH3* variants might be a cause of the arrest. MLH3 is certainly involved in the mammalian meiotic process, but its role in MMR during DNA replication is controversial (Lipkin *et al.*, 2000; Wu *et al.*, 2001; Hienonen *et al.*, 2003; Liu *et al.*, 2003; Korhonen *et al.*, 2007). Cytological studies in mice have shown that MLH3 is found in zygotene, combines with recombination nodules in early pachytene, and associates with MLH1 from mid-pachytene to the early diplonema stages (Lipkin *et al.*, 2002; Santucci-Darmanin *et al.*, 2002; Marcon and Moens, 2003). Also, both male and female *MLH3*^{-/-} mice manifested infertility (Lipkin *et al.*, 2002). Therefore, human *MLH3* variants might interfere in the meiotic process resulting in male infertility.

Ferras *et al.* (2012) found that one patient with two *MLH3* mutations, the combination of C2531T and IVS9+66G/A, showed overexpression of MLH3 and MLH1. Since MLH3 and PMS2 share the same interaction domain on MLH1, it is possible that the quantitative balance of the MLH1-binding partners (MLH3 and PMS2) plays a crucial role in specifying different outcomes during meiosis (Kondo *et al.*, 2001; Korhonen *et al.*, 2007). Accordingly, mutation-related overexpression of MLH3 and MLH1 may result in a predominance of MLH1-MLH3 complexes and a relative reduction in MLH1-PMS2 complexes. Consequently, meiotic failure might occur as MLH1-PMS2 complexes ought to replace MLH1-MLH3 complexes during diplonema. However, due to the relatively small sample sizes in the above studies, further evaluation of the correlations and mechanisms linking MMR defects and male infertility is required.

Recently, numerous sequencing analyses of single nucleotide polymorphisms (SNPs) in candidate genes have helped to clarify the etiology and susceptibility of both male and female infertility (Kang *et al.*,

2014; Ni *et al.*, 2014). Xu *et al.* (2010) identified two SNPs, *MSH5* (C85T, Pro29Ser) and *MLH3* (C2531T, Pro844Leu) associated with male infertility, especially in non-obstructive azoospermia or severe oligozoospermia. The allele carriers *MSH5* Pro29Ser and *MLH3* Pro844Leu demonstrated a 2.89-fold and 2.25-fold increased risk of azoospermia or oligozoospermia, respectively. However, two major problems have limited the value of medical sequencing studies. First, the actual rates of these reported SNPs are quite low and could explain only a small percentage of idiopathic male infertility. Growing evidence suggests that common genetic variations are rarely responsible for the whole disease phenotype (Frazer *et al.*, 2009; Manolio *et al.*, 2009). The “common disease, common variant” model is not suitable for most complex diseases, including male infertility. Further studies have suggested that risk or causative genetic variants for most cases of idiopathic male infertility might be related to the identification of rare polymorphisms and copy number variants (CNVs). Second, most reports of SNPs associated with male infertility have either not been followed up with validation studies, or often failed validation in follow-up studies.

Ji *et al.* (2012) conducted a prospective case-control study of 1292 idiopathic male infertility patients and 480 fertile controls in a Chinese population. The idiopathic infertility patients were divided into two subgroups: 524 patients with non-obstructive azoospermia or oligozoospermia and 768 with a normal sperm count. Twenty-one tagging SNPs in five MMR genes (*MLH1*, *MLH3*, *PMS2*, *MSH4*, and *MSH5*) were examined by the sequence detection system and the SNPstream 12-plex platform. The results showed that the genotype frequencies of three SNPs, *MLH1* (rs4647269), *PMS2* (rs1059060, Ser775Asn), and *MSH5* (rs2075789, Pro29Ser), were significantly increased (by from 6% to 17%) in patients with azoospermia or oligozoospermia. Therefore, the presence of these SNPs seems to be a risk factor for the development of azoospermia or oligozoospermia. Another SNP in *PMS2* rs1059060 appeared to contribute to the risk of male infertility in patients with a normal sperm count. Guerrette *et al.* (1999) localized the MLH1-PMS2 interaction region to amino acids 506–675 of MLH1 and 675–850 of PMS2. This study provided evidence that the *PMS2*

Ser775Asn variant attenuates the binding of MLH1 and PMS2 detected by fluorescence resonance energy transfer (FRET) and co-immunoprecipitation assay, further supporting the potential interaction between MLH1-PMS2 and MLH1-MLH3 discussed above. The *MSH5* Pro29Ser polymorphism is located within the MSH4-MSH5 interacting domain and leads to a weakened formation of the MSH4-MSH5 hetero-complex (Yi *et al.*, 2005). This was supported by the study of Xu *et al.* (2010). Gene knockout of *MSH4* or *MSH5* results in infertility in mice since they are unable to resolve meiotic chromosomal crossovers (de Vries *et al.*, 1999; Edlmann *et al.*, 1999; Kneitz *et al.*, 2000). Consequently, *MSH5* Pro29Ser alteration is related to a significantly increased risk of male infertility. However, the detailed molecular mechanisms are unknown.

Compared with the current understanding of the MMR system in male infertility, its potential involvement in female infertility has received much less attention. A case-control study including 41 women with premature ovarian failure and 39 controls suggested that *MSH5* Pro29Ser polymorphism might be an explanation for premature ovarian failure (Mandon-Pepin *et al.*, 2008), indicating that MMR gene mutation is likely to be related to female infertility. Pashaiefar *et al.* (2013) suggested that *MLH3* C2531T polymorphism can be associated with the risk of unexplained fertility in Iranian women. Perry *et al.* (2014) showed that gene mutation-related low expression of *MSH6* due to one SNP in *MSH6* (rs1800932) is associated with earlier menopause, suggesting that MMR is a key process in determining female reproductive lifespan and could be a crucial therapeutic target for female infertility.

With an increased incidence and a younger age of patients, certain forms of MMR-related genital system cancers which require surgical treatment may contribute to female “structural” infertility. For instance, endometrial cancer has an intimate connection with genetic and epigenetic variation of the MMR system. Studies demonstrated that aberrant methylation of *MLH1* is detected in about 40% of endometrial cancer cases and is considered to be an important process in the early stage of endometrial carcinogenesis (Muraki *et al.*, 2009). Conservative surgical and hormonal therapy is currently practical in clinical treatment. This kind of therapy ensures that a con-

siderable proportion of women with endometrial cancer successfully achieve pregnancies resulting in live births with or without ART (Park *et al.*, 2013). However, some other problems still exist. Although current studies have demonstrated that there is no definitive evidence of a significant association between MMR status and survival in endometrial cancer (Diaz-Padilla *et al.*, 2013), deficient MMR is associated with a higher risk of high-grade endometrial cancer and worse clinical outcomes in women aged 40 years or younger (Garg *et al.*, 2009; Shih *et al.*, 2011). Also, MMR status appears to be linked to body mass index (BMI) and endometrial cancer (Win *et al.*, 2011; Joehlin-Price *et al.*, 2014), indicating a possible interactive system between MMR and some types of metabolic diseases and cancers. Therefore, cautious and comprehensive assessment should be carried out before, during, and after conservative therapy in young women who have a strong desire to preserve fertility, to maintain the safety and health of both the mother and offspring.

Apart from the genetic components involved in human infertility, a growing body of evidence clearly indicates the importance of other factors such as epigenetic variation (Dada *et al.*, 2012; Richardson *et al.*, 2014). Whether epigenetic variation in the MMR system may be etiologic of infertility is unknown, but the fast developing field of epigenetics may be yet another area related to infertility and should be considered together with genetic factors.

4 MMR alterations and ART

With the application of ART including in vitro fertilization (IVF), intracytoplasmic sperm injection (ICSI), and preimplantation genetic diagnosis (PGD), it is now practical to bypass the natural barriers to produce offspring. Interestingly, Terribas *et al.* (2010) considered the expression of testicular *MSH4* could be useful as a surrogate marker for intratesticular elongated spermatids in patients with non-obstructive azoospermia and may be useful in predicting the viability of assisted reproduction. This approach needs further evaluation for sperm quality selection. Zheng *et al.* (2005) showed that in vitro culture could lead to dysregulation of many genes and possibly affect the expression of *MSH2*, which may affect the viability

of Rhesus monkey embryos. This raises the possibility of the same possible biological activity in humans and implies that the MMR system might play a role in the ART process and affect the outcome. ART offspring are associated with various risks such as preterm delivery, low birth weight, congenital abnormality, and genomic imprinting syndromes (Maher *et al.*, 2003; Kurinczuk *et al.*, 2004; de Ligt *et al.*, 2012). However, it is difficult to establish whether these risks are related to the technology itself or the inherited genetic background. Whether the incidence of congenital abnormalities increases following ART is controversial (Halliday *et al.*, 2010; Simon *et al.*, 2010; Woldringh *et al.*, 2010).

ART has been shown to have a definite relationship with an increased number of genetic and epigenetic alterations (Kochanski *et al.*, 2013). ART may create de novo mutations through diverse chromosomal and molecular mechanisms. It enables the transmission of pre-existing mutations which are related to infertility and the natural process of early pregnancy termination. A Y-chromosome microdeletion is found in about 1 in 4000 in the general population and is responsible for male infertility. Feng *et al.* (2008) proposed that ART offspring may show a rising risk of a gene static mutation, azoospermia factor (AZF) microdeletion, even when their fathers have a normal spermatogenesis and genetic background. Bianchi *et al.* (2002) showed that Y chromosome MSI and AZF microdeletion testing in five cases of HNPCC strongly suggested a correlation between heterozygosity for *MLH1* or other MMR gene mutations, Y chromosome instability, and AZF microdeletions. Thus, MMR gene mutations may play a role in the origin of AZF microdeletions. Given the fact of increased vertical transmission of Y-chromosome microdeletions in the offspring via ICSI (Serebrovska *et al.*, 2006), the potential benefits and risks of adopting ART for those specific individuals who carry defective MMR genes should be considered.

Zheng *et al.* (2013) indicated that ART conceived babies display an increased incidence of gene dynamic mutation, also defined as TNR instability, which is related to more than 20 neuromuscular and neurodegenerative disorders including Huntington's disease, myotonic dystrophy, and fragile X syndrome. Due to the specificity of TNR structure, the DNA

damage repair system plays an important role in the formation of TNR alterations, except for its function in repairing de novo mutations (Fig. 1). As MMR protein expression can alter TNR in diverse ways by either enhancing or suppressing different human TNR types (Lin and Wilson, 2009; Halabi *et al.*, 2012; Pluciennik *et al.*, 2013), ART application should be considered more cautiously to prevent severe diseases.




Fig. 1 Proposed function model of MutS and MutL in relation to TNR instability

TNR-related hairpin structure occurring during DNA replication is initially recognized by the MutSβ complex, and then the MutL complex is recruited. Various pathways including canonical MMR and noncanonical MMR (ncMMR) may be engaged, leading to different outcomes of TNR instability: DNA damage repaired in time or otherwise, eventual TNR mutation

As for epigenetic variation, possible risks related to ART may be caused by either the use of incomplete reprogrammed sperm or in vitro embryo culture during a time of epigenetic reprogramming (Lucifero *et al.*, 2004; Niemitz and Feinberg, 2004; Thompson and Williams, 2005). Geneticists have reported an increased incidence of selected epigenetic defects in ART offspring, such as Beckwith-Weidemann syndrome and Angelman syndrome (Maher *et al.*, 2003; de Ligt *et al.*, 2012). Although there is no evidence that the MMR system is involved in the epigenetic stability of ART offspring, we can assume that there might be an interaction since both MMR and epigenetic biological mechanisms are extremely active during gametogenesis and embryonic development.

Our unpublished data show that ART procedures and the infertility background might affect MMR epigenetic modifications like DNA methylation. This needs further investigation.

Studies in mice and humans have reported that some genetic and epigenetic modifications caused by ART can be transmitted to the next or even to the F₂ generation (van Montfoort *et al.*, 2012; Wang *et al.*, 2013). However, there have been no reports of transgenerational effects caused by MMR alterations. In consideration of its function in the parental generation, the inheritance of abnormal MMR gene status might affect aspects of an offspring's health condition including predisposition to certain types of cancer, resistance to certain therapeutic drugs, and infertility. This area needs further study.

Given that the above, augmented genetic mutations seen in infertile individuals may raise the probability of these variants being inherited, conferring increased risk of infertility and then possible somatic diseases in later life in ART offspring. Research has shown that epigenetic mechanisms not only lead to inappropriate expression of the affected gene but may also expose hidden genetic variation, contributing to a predisposition to epigenetic instability in offspring conceived by ART (Sollars *et al.*, 2003). Alternatively, the ART procedures and the genetic background may alter the genetic and epigenetic status of these offspring. More detailed information about infertility-related factors, gene mutations, DNA SNPs, other genetic and epigenetic forms of the MMR system, and correlated biomarkers should be found and ultimately applied clinically to provide better counseling and treatment for infertile couples considering IVF, ICSI, or even PGD, which aims to avoid inheriting defective genes. Also, long term follow-up studies of children born from ART are required to understand the possible effects on these offspring.

5 Conclusions

Apart from its primary role to correct replication-associated base-pair mismatches and IDLs, MMR also plays a crucial role in DNA recombination and is involved in gametogenesis. Infertility is an extremely complex disorder caused by various genetic, epige-

netic, endocrine, and environmental factors. Although the etiology of human infertility is still largely unknown, the impact of genetic and epigenetic MMR system alterations on infertility is beyond debate. However, further studies are required to elucidate the molecular mechanisms of MMR genes and other additive risk factors in infertility. A better understanding of MMR alterations might be helpful in reducing DNA damage, developing gene targeted therapy, and providing diagnostic applications for ART.

The diverse functions of the MMR system in the human body, the applicability to humans of research based on transgenic, knock-out or knock-in mice models, and the great difficulty in obtaining and ethically using human gametes, especially oocytes and embryos, are huge obstacles to improving our understanding of MMR mechanisms in human infertility. More than thirty years of ART application and thirty-six years of observation of the first "in vitro" baby are still insufficient to thoroughly understand the potential changes and risks in later life resulting from infertility treatments. More work is urgently needed to investigate the interaction between MMR alterations and DNA integrity, human infertility, and ART security, to establish a better medical system with improved preventive, diagnostic, and therapeutic treatments of human infertility.

There is no doubt that ART will continue to be a major medical approach for infertile couples. Much work remains to be done to maximize the safety and reduce the risks of ART offspring. Long-term follow-up programs will yield significant results for fertility specialists providing ART, neonatologists working with ART offspring and geneticists counseling infertile parents. The efforts of global scientists, increasing maturity of the ART process, and the development of scientific technologies such as microarrays and single cell level sequencing, will lead to a better understanding of the cellular and molecular mechanisms of the MMR system involved in human infertility and related treatments. This will provide a better perspective to improve diagnostic capability, patient care, and ultimately the quality of the offspring.

Compliance with ethics guidelines

Min-hao HU, Shu-yuan LIU, Ning WANG, Yan WU, and Fan JIN declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Baker, S.M., Bronner, C.E., Zhang, L., et al., 1995. Male mice defective in the DNA mismatch repair gene *PMS2* exhibit abnormal chromosome synapsis in meiosis. *Cell*, **82**(2): 309-319.
[http://dx.doi.org/10.1016/0092-8674\(95\)90318-6](http://dx.doi.org/10.1016/0092-8674(95)90318-6)
- Baker, S.M., Plug, A.W., Prolla, T.A., et al., 1996. Involvement of mouse *Mlh1* in DNA mismatch repair and meiotic crossing over. *Nat. Genet.*, **13**(3):336-342.
<http://dx.doi.org/10.1038/ng0796-336>
- Bianchi, N.O., Richard, S.M., Peltomaki, P., et al., 2002. Mosaic AZF deletions and susceptibility to testicular tumors. *Mutat. Res.*, **503**(1-2):51-62.
[http://dx.doi.org/10.1016/S0027-5107\(02\)00072-6](http://dx.doi.org/10.1016/S0027-5107(02)00072-6)
- Brown, K.D., Rathi, A., Kamath, R., et al., 2003. The mismatch repair system is required for S-phase checkpoint activation. *Nat. Genet.*, **33**(1):80-84.
<http://dx.doi.org/10.1038/ng1052>
- Cannavo, E., Gerrits, B., Marra, G., et al., 2007. Characterization of the interactome of the human MutL homologues MLH1, PMS1, and PMS2. *J. Biol. Chem.*, **282**(5):2976-2986.
<http://dx.doi.org/10.1074/jbc.M609989200>
- Chen, J., Sadowski, I., 2005. Identification of the mismatch repair genes PMS2 and MLH1 as p53 target genes by using serial analysis of binding elements. *PNAS*, **102**(13): 4813-4818.
<http://dx.doi.org/10.1073/pnas.0407069102>
- Dada, R., Kumar, M., Jesudasan, R., et al., 2012. Epigenetics and its role in male infertility. *J. Assist. Reprod. Genet.*, **29**(3):213-223.
<http://dx.doi.org/10.1007/s10815-012-9715-0>
- de Ligt, J., Willemsen, M.H., van Bon, B.W., et al., 2012. Diagnostic exome sequencing in persons with severe intellectual disability. *N. Engl. J. Med.*, **367**(20):1921-1929.
<http://dx.doi.org/10.1056/NEJMoa1206524>
- de Vries, S.S., Baart, E.B., Dekker, M., et al., 1999. Mouse MutS-like protein Msh5 is required for proper chromosome synapsis in male and female meiosis. *Genes. Dev.*, **13**(5):523-531.
<http://dx.doi.org/10.1101/gad.13.5.523>
- Diaz-Padilla, I., Romero, N., Amir, E., et al., 2013. Mismatch repair status and clinical outcome in endometrial cancer: a systematic review and meta-analysis. *Crit. Rev. Oncol. Hematol.*, **88**(1):154-167.
<http://dx.doi.org/10.1016/j.critrevonc.2013.03.002>
- Edelmann, W., Cohen, P.E., Kneitz, B., et al., 1999. Mammalian MutS homologue 5 is required for chromosome pairing in meiosis. *Nat. Genet.*, **21**(1):123-127.
<http://dx.doi.org/10.1038/5075>
- Feng, C., Wang, L., Dong, M., et al., 2008. Assisted reproductive technology may increase clinical mutation detection in male offspring. *Fertil. Steril.*, **90**(1):92-96.
<http://dx.doi.org/10.1016/j.fertnstert.2007.06.004>
- Ferras, C., Zhou, X.L., Sousa, M., et al., 2007. DNA mismatch repair gene *hMLH3* variants in meiotic arrest. *Fertil. Steril.*, **88**(6):1681-1684.
<http://dx.doi.org/10.1016/j.fertnstert.2007.01.063>
- Ferras, C., Fernandes, S., Silva, J., et al., 2012. Expression analysis of MLH3, MLH1, and MSH4 in maturation arrest. *Reprod. Sci.*, **19**(6):587-596.
<http://dx.doi.org/10.1177/1933719111428521>
- Frazer, K.A., Murray, S.S., Schork, N.J., et al., 2009. Human genetic variation and its contribution to complex traits. *Nat. Rev. Genet.*, **10**(4):241-251.
<http://dx.doi.org/10.1038/nrg2554>
- Garg, K., Shih, K., Barakat, R., et al., 2009. Endometrial carcinomas in women aged 40 years and younger: tumors associated with loss of DNA mismatch repair proteins comprise a distinct clinicopathologic subset. *Am. J. Surg. Pathol.*, **33**(12):1869-1877.
<http://dx.doi.org/10.1097/PAS.0b013e3181bc9866>
- Gnoth, C., Godehardt, E., Frank-Herrmann, P., et al., 2005. Definition and prevalence of subfertility and infertility. *Hum. Reprod.*, **20**(5):1144-1147.
<http://dx.doi.org/10.1093/humrep/deh870>
- Guerrette, S., Acharya, S., Fishel, R., 1999. The interaction of the human MutL homologues in hereditary nonpolyposis colon cancer. *J. Biol. Chem.*, **274**(10):6336-6341.
<http://dx.doi.org/10.1074/jbc.274.10.6336>
- Gunes, S., Al-Sadaan, M., Agarwal, A., 2015. Spermatogenesis, DNA damage and DNA repair mechanisms in male infertility. *Reprod. BioMed. Online*, **31**(3):309-319.
<http://dx.doi.org/10.1016/j.rbmo.2015.06.010>
- Guo, G., Wang, W., Bradley, A., 2004. Mismatch repair genes identified using genetic screens in Blm-deficient embryonic stem cells. *Nature*, **429**(6994):891-895.
<http://dx.doi.org/10.1038/nature02653>
- Halabi, A., Ditch, S., Wang, J., et al., 2012. DNA mismatch repair complex MutSβ promotes GAA·TTC repeat expansion in human cells. *J. Biol. Chem.*, **287**(35):29958-29967.
<http://dx.doi.org/10.1074/jbc.M112.356758>
- Halliday, J.L., Ukoumunne, O.C., Baker, H.W., et al., 2010. Increased risk of blastogenesis birth defects, arising in the first 4 weeks of pregnancy, after assisted reproductive technologies. *Hum. Reprod.*, **25**(1):59-65.
<http://dx.doi.org/10.1093/humrep/dep364>
- Her, C., Zhao, N., Wu, X., et al., 2007. MutS homologues hMSH4 and hMSH5: diverse functional implications in humans. *Front. Biosci.*, **12**(1):905-911.
<http://dx.doi.org/10.2741/2112>
- Hienonen, T., Laiho, P., Salovaara, R., et al., 2003. Little evidence for involvement of MLH3 in colorectal cancer predisposition. *Int. J. Cancer.*, **106**(2):292-296.
<http://dx.doi.org/10.1002/ijc.11218>
- Iyer, R.R., Pluciennik, A., Burdett, V., et al., 2006. DNA mismatch repair: functions and mechanisms. *Chem. Rev.*, **106**(2):302-323.
<http://dx.doi.org/10.1021/cr0404794>

- Jascur, T., Boland, C.R., 2006. Structure and function of the components of the human DNA mismatch repair system. *Int. J. Cancer.*, **119**(9):2030-2035. <http://dx.doi.org/10.1002/ijc.22023>
- Ji, G., Long, Y., Zhou, Y., et al., 2012. Common variants in mismatch repair genes associated with increased risk of sperm DNA damage and male infertility. *BMC Med.*, **10**(1):49. <http://dx.doi.org/10.1186/1741-7015-10-49>
- Jiricny, J., 2006. The multifaceted mismatch-repair system. *Nat. Rev. Mol. Cell. Biol.*, **7**(5):335-346. <http://dx.doi.org/10.1038/nrm1907>
- Joehlin-Price, A.S., Perrino, C.M., Stephens, J., et al., 2014. Mismatch repair protein expression in 1049 endometrial carcinomas, associations with body mass index, and other clinicopathologic variables. *Gynecol. Oncol.*, **133**(1):43-47. <http://dx.doi.org/10.1016/j.ygyno.2014.01.017>
- Kang, S., Li, Y., Li, B., et al., 2014. Genetic variation of the E-cadherin gene is associated with primary infertility in patients with ovarian endometriosis. *Fertil. Steril.*, **102**(4):1149-1154. <http://dx.doi.org/10.1016/j.fertnstert.2014.07.005>
- Karamurzin, Y., Zeng, Z., Stadler, Z.K., et al., 2012. Unusual DNA mismatch repair-deficient tumors in Lynch syndrome: a report of new cases and review of the literature. *Hum. Pathol.*, **43**(10):1677-1687. <http://dx.doi.org/10.1016/j.humpath.2011.12.012>
- Kim, M., Trinh, B.N., Long, T.I., et al., 2004. Dnmt1 deficiency leads to enhanced microsatellite instability in mouse embryonic stem cells. *Nucleic Acids Res.*, **32**(19):5742-5749. <http://dx.doi.org/10.1093/nar/gkh912>
- Kneitz, B., Cohen, P.E., Avdievich, E., et al., 2000. MutS homolog 4 localization to meiotic chromosomes is required for chromosome pairing during meiosis in male and female mice. *Genes Dev.*, **14**(9):1085-1097. <http://dx.doi.org/10.1101/gad.14.9.1085>
- Kochanski, A., Merritt, T.A., Gadzinowski, J., et al., 2013. The impact of assisted reproductive technologies on the genome and epigenome of the newborn. *J. Neonatal. Perinatal. Med.*, **6**(2):101-108. <http://dx.doi.org/10.3233/NPM-1366812>
- Kolodner, R.D., Marsischky, G.T., 1999. Eukaryotic DNA mismatch repair. *Curr. Opin. Genet. Dev.*, **9**(1):89-96. [http://dx.doi.org/10.1016/S0959-437X\(99\)80013-6](http://dx.doi.org/10.1016/S0959-437X(99)80013-6)
- Kondo, E., Horii, A., Fukushige, S., 2001. The interacting domains of three MutL heterodimers in man: hMLH1 interacts with 36 homologous amino acid residues within hMLH3, hPMS1 and hPMS2. *Nucleic Acids Res.*, **29**(8):1695-1702. <http://dx.doi.org/10.1093/nar/29.8.1695>
- Korhonen, M.K., Raevaara, T.E., Lohi, H., et al., 2007. Conditional nuclear localization of hMLH3 suggests a minor activity in mismatch repair and supports its role as a low-risk gene in HNPCC. *Oncol. Rep.*, **17**(2):351-354. <http://dx.doi.org/10.3892/or.17.2.351>
- Kunkel, T.A., Erie, D.A., 2005. DNA mismatch repair. *Annu. Rev. Biochem.*, **74**(1):681-710. <http://dx.doi.org/10.1146/annurev.biochem.74.082803.133243>
- Kurinczuk, J.J., Hansen, M., Bower, C., 2004. The risk of birth defects in children born after assisted reproductive technologies. *Curr. Opin. Obstet. Gynecol.*, **16**(3):201-209. <http://dx.doi.org/10.1097/00001703-200406000-00002>
- Lahue, R.S., Au, K.G., Modrich, P., 1989. DNA mismatch correction in a defined system. *Science*, **245**(4914):160-164. <http://dx.doi.org/10.1126/science.2665076>
- Li, G.M., 2008. Mechanisms and functions of DNA mismatch repair. *Cell Res.*, **18**(1):85-98. <http://dx.doi.org/10.1038/cr.2007.115>
- Lin, Y., Wilson, J.H., 2009. Diverse effects of individual mismatch repair components on transcription-induced CAG repeat instability in human cells. *DNA Repair (Amst)*, **8**(8):878-885. <http://dx.doi.org/10.1016/j.dnarep.2009.04.024>
- Lipkin, S.M., Wang, V., Jacoby, R., et al., 2000. MLH3: a DNA mismatch repair gene associated with mammalian microsatellite instability. *Nat. Genet.*, **24**(1):27-35. <http://dx.doi.org/10.1038/71643>
- Lipkin, S.M., Moens, P.B., Wang, V., et al., 2002. Meiotic arrest and aneuploidy in MLH3-deficient mice. *Nat. Genet.*, **31**(4):385-390. <http://dx.doi.org/10.1038/ng931>
- Liu, H.X., Zhou, X.L., Liu, T., et al., 2003. The role of hMLH3 in familial colorectal cancer. *Cancer Res.*, **63**(8):1894-1899.
- Lucifero, D., Mann, M.R., Bartolomei, M.S., et al., 2004. Gene-specific timing and epigenetic memory in oocyte imprinting. *Hum. Mol. Genet.*, **13**(8):839-849. <http://dx.doi.org/10.1093/hmg/ddh104>
- Maduro, M.R., Casella, R., Kim, E., et al., 2003. Microsatellite instability and defects in mismatch repair proteins: a new aetiology for Sertoli cell-only syndrome. *Mol. Hum. Reprod.*, **9**(2):61-68. <http://dx.doi.org/10.1093/molehr/gag013>
- Maher, E.R., Afnan, M., Barratt, C.L., 2003. Epigenetic risks related to assisted reproductive technologies: epigenetics, imprinting, ART and icebergs? *Hum. Reprod.*, **18**(12):2508-2511. <http://dx.doi.org/10.1093/humrep/deg486>
- Mandon-Pepin, B., Touraine, P., Kuttann, F., et al., 2008. Genetic investigation of four meiotic genes in women with premature ovarian failure. *Eur. J. Endocrinol.*, **158**(1):107-115. <http://dx.doi.org/10.1530/EJE-07-0400>
- Manolio, T.A., Collins, F.S., Cox, N.J., et al., 2009. Finding the missing heritability of complex diseases. *Nature*, **461**(7265):747-753. <http://dx.doi.org/10.1038/nature08494>
- Marcon, E., Moens, P., 2003. MLH1p and MLH3p localize to precociously induced chiasmata of okadaic-acid-treated mouse spermatocytes. *Genetics*, **165**(4):2283-2287.

- Martins, A.D., Moreira, A.C., Sa, R., et al., 2015. Leptin modulates human Sertoli cells acetate production and glycolytic profile: a novel mechanism of obesity-induced male infertility? *Biochim. Biophys. Acta*, **1852**(9):1824-1832. <http://dx.doi.org/10.1016/j.bbadis.2015.06.005>
- Modrich, P., 1991. Mechanisms and biological effects of mismatch repair. *Annu. Rev. Genet.*, **25**(1):229-253. <http://dx.doi.org/10.1146/annurev.ge.25.120191.001305>
- Mukherjee, S., Ridgeway, A.D., Lamb, D.J., 2010. DNA mismatch repair and infertility. *Curr. Opin. Urol.*, **20**(6):525-532. <http://dx.doi.org/10.1097/MOU.0b013e32833f1c21>
- Muraki, Y., Banno, K., Yanokura, M., et al., 2009. Epigenetic DNA hypermethylation: clinical applications in endometrial cancer (Review). *Oncol. Rep.*, **22**(5):967-972. <http://dx.doi.org/10.3892/or.00000523>
- Ni, B., Ma, H., Lin, Y., et al., 2014. Genetic variants in Ser-Arg protein-coding genes are associated with the risk of nonobstructive azoospermia in Chinese men. *Fertil. Steril.*, **101**(6):1711-1717. <http://dx.doi.org/10.1016/j.fertnstert.2014.02.033>
- Niemitz, E.L., Feinberg, A.P., 2004. Epigenetics and assisted reproductive technology: a call for investigation. *Am. J. Hum. Genet.*, **74**(4):599-609. <http://dx.doi.org/10.1086/382897>
- Oliveira, P.F., Martins, A.D., Moreira, A.C., et al., 2015. The Warburg effect revisited—lesson from the Sertoli cell. *Med. Res. Rev.*, **35**(1):126-151. <http://dx.doi.org/10.1002/med.21325>
- Park, J.Y., Seong, S.J., Kim, T.J., et al., 2013. Pregnancy outcomes after fertility-sparing management in young women with early endometrial cancer. *Obstet. Gynecol.*, **121**(1):136-142. <http://dx.doi.org/10.1097/AOG.0b013e31827a0643>
- Pashaiefar, H., Sheikha, M.H., Kalantar, S.M., et al., 2013. Analysis of MLH3 C2531T polymorphism in Iranian women with unexplained infertility. *Iran. J. Reprod. Med.*, **11**(1):19-24.
- Peltomaki, P., 2003. Role of DNA mismatch repair defects in the pathogenesis of human cancer. *J. Clin. Oncol.*, **21**(6):1174-1179. <http://dx.doi.org/10.1200/JCO.2003.04.060>
- Perry, J.R., Hsu, Y.H., Chasman, D.I., et al., 2014. DNA mismatch repair gene MSH6 implicated in determining age at natural menopause. *Hum. Mol. Genet.*, **23**(9):2490-2497. <http://dx.doi.org/10.1093/hmg/ddt620>
- Pluciennik, A., Burdett, V., Baitinger, C., et al., 2013. Extrahelical (CAG)/(CTG) triplet repeat elements support proliferating cell nuclear antigen loading and MutLα endonuclease activation. *PNAS*, **110**(30):12277-12282. <http://dx.doi.org/10.1073/pnas.1311325110>
- Richardson, M.E., Bleiziffer, A., Tuttelmann, F., et al., 2014. Epigenetic regulation of the *RHOX* homeobox gene cluster and its association with human male infertility. *Hum. Mol. Genet.*, **23**(1):12-23. <http://dx.doi.org/10.1093/hmg/ddt392>
- Santucci-Darmanin, S., Neyton, S., Lespinasse, F., et al., 2002. The DNA mismatch-repair MLH3 protein interacts with MSH4 in meiotic cells, supporting a role for this MutL homolog in mammalian meiotic recombination. *Hum. Mol. Genet.*, **11**(15):1697-1706. <http://dx.doi.org/10.1093/hmg/11.15.1697>
- Serebrovska, Z.A., Serebrovska, T.V., Pyle, R.L., et al., 2006. Transmission of male infertility and intracytoplasmic sperm injection (mini-review). *Fiziol. Zh.*, **52**(3):110-118.
- Shih, K.K., Garg, K., Levine, D.A., et al., 2011. Clinicopathologic significance of DNA mismatch repair protein defects and endometrial cancer in women 40 years of age and younger. *Gynecol. Oncol.*, **123**(1):88-94. <http://dx.doi.org/10.1016/j.ygyno.2011.06.005>
- Shimodaira, H., Yoshioka-Yamashita, A., Kolodner, R.D., et al., 2003. Interaction of mismatch repair protein PMS2 and the p53-related transcription factor p73 in apoptosis response to cisplatin. *PNAS*, **100**(5):2420-2425. <http://dx.doi.org/10.1073/pnas.0438031100>
- Simon, L., Brunborg, G., Stevenson, M., et al., 2010. Clinical significance of sperm DNA damage in assisted reproduction outcome. *Hum. Reprod.*, **25**(7):1594-1608. <http://dx.doi.org/10.1093/humrep/deq103>
- Sollars, V., Lu, X., Xiao, L., et al., 2003. Evidence for an epigenetic mechanism by which Hsp90 acts as a capacitor for morphological evolution. *Nat. Genet.*, **33**(1):70-74. <http://dx.doi.org/10.1038/ng1067>
- Stojic, L., Cejka, P., Jiricny, J., 2005. High doses of SN1 type methylating agents activate DNA damage signaling cascades that are largely independent of mismatch repair. *Cell Cycle*, **4**(3):473-477. <http://dx.doi.org/10.4161/cc.4.3.1528>
- Surtees, J.A., Argueso, J.L., Alani, E., 2004. Mismatch repair proteins: key regulators of genetic recombination. *Cytogenet. Genome Res.*, **107**(3-4):146-159. <http://dx.doi.org/10.1159/000080593>
- Terribas, E., Bonache, S., Garcia-Arevalo, M., et al., 2010. Changes in the expression profile of the meiosis-involved mismatch repair genes in impaired human spermatogenesis. *J. Androl.*, **31**(4):346-357. <http://dx.doi.org/10.2164/jandrol.109.008805>
- Thompson, J.R., Williams, C.J., 2005. Genomic imprinting and assisted reproductive technology: connections and potential risks. *Semin. Reprod. Med.*, **23**(3):285-295. <http://dx.doi.org/10.1055/s-2005-872457>
- Valentini, A.M., Armentano, R., Pirrelli, M., et al., 2006. Chemotherapeutic agents for colorectal cancer with a defective mismatch repair system: the state of the art. *Cancer Treat. Rev.*, **32**(8):607-618. <http://dx.doi.org/10.1016/j.ctrv.2006.08.001>
- van Montfoort, A.P., Hanssen, L.L., de Sutter, P., et al., 2012. Assisted reproduction treatment and epigenetic inheritance. *Hum. Reprod. Update*, **18**(2):171-197. <http://dx.doi.org/10.1093/humupd/dmr047>

- Wang, L.Y., Wang, N., Le, F., et al., 2013. Persistence and intergenerational transmission of differentially expressed genes in the testes of intracytoplasmic sperm injection conceived mice. *J. Zhejiang Univ.-Sci. B (Biomed. & Biotechnol.)*, **14**(5):372-381.
<http://dx.doi.org/10.1631/jzus.B1200321>
- Wang, Y., Qin, J., 2003. MSH2 and ATR form a signaling module and regulate two branches of the damage response to DNA methylation. *PNAS*, **100**(26):15387-15392.
<http://dx.doi.org/10.1073/pnas.2536810100>
- Watson, P., Lynch, H.T., 1994. The tumor spectrum in HNPCC. *Anticancer Res.*, **14**(4B):1635-1639.
- Wei, K., Clark, A.B., Wong, E., et al., 2003. Inactivation of Exonuclease 1 in mice results in DNA mismatch repair defects, increased cancer susceptibility, and male and female sterility. *Genes Dev.*, **17**(5):603-614.
<http://dx.doi.org/10.1101/gad.1060603>
- Win, A.K., Dowty, J.G., Antill, Y.C., et al., 2011. Body mass index in early adulthood and endometrial cancer risk for mismatch repair gene mutation carriers. *Obstet. Gynecol.*, **117**(4):899-905.
<http://dx.doi.org/10.1097/AOG.0b013e3182110ea3>
- Woldringh, G.H., Besselink, D.E., Tillema, A.H., et al., 2010. Karyotyping, congenital anomalies and follow-up of children after intracytoplasmic sperm injection with non-ejaculated sperm: a systematic review. *Hum. Reprod. Update*, **16**(1):12-19.
<http://dx.doi.org/10.1093/humupd/dmp030>
- Wu, Y., Berends, M.J., Sijmons, R.H., et al., 2001. A role for MLH3 in hereditary nonpolyposis colorectal cancer. *Nat. Genet.*, **29**(2):137-138.
<http://dx.doi.org/10.1038/ng1001-137>
- Xu, K., Lu, T., Zhou, H., et al., 2010. The role of MSH5 C85T and MLH3 C2531T polymorphisms in the risk of male infertility with azoospermia or severe oligozoospermia. *Clin. Chim. Acta*, **411**(1-2):49-52.
<http://dx.doi.org/10.1016/j.cca.2009.09.038>
- Yi, W., Wu, X., Lee, T.H., et al., 2005. Two variants of MutS homolog hMSH5: prevalence in humans and effects on protein interaction. *Biochem. Biophys. Res. Commun.*, **332**(2):524-532.
<http://dx.doi.org/10.1016/j.bbrc.2005.04.154>
- Youn, C.K., Cho, H.J., Kim, S.H., et al., 2005. Bcl-2 expression suppresses mismatch repair activity through inhibition of E2F transcriptional activity. *Nat. Cell Biol.*, **7**(2):137-147.
<http://dx.doi.org/10.1038/ncb1215>
- Zheng, P., Schramm, R.D., Latham, K.E., 2005. Developmental regulation and in vitro culture effects on expression of DNA repair and cell cycle checkpoint control genes in rhesus monkey oocytes and embryos. *Biol. Reprod.*, **72**(6):1359-1369.
<http://dx.doi.org/10.1095/biolreprod.104.039073>
- Zheng, Y.M., Li, L., Zhou, L.M., et al., 2013. Alterations in the frequency of trinucleotide repeat dynamic mutations in offspring conceived through assisted reproductive technology. *Hum. Reprod.*, **28**(9):2570-2580.
<http://dx.doi.org/10.1093/humrep/det294>

中文概要

题目: DNA 错配修复系统的改变对人类生育能力及相关治疗的影响

概要: 简要概括 DNA 损伤修复系统在人体中的作用和机制, 并探讨其改变与人类生殖能力以及通过辅助生殖技术诞生的子代之间的相互影响。希望更多相关工作的进行能够为人类不孕症的预防、诊断和治疗工作建立一个更好的医疗体系。

关键词: DNA 错配修复; 不孕; 辅助生殖技术; 后代