

ERRATUM

Erratum to: Rad51 Expression Is a Useful Predictive Factor for the Efficacy of Neoadjuvant Chemoradiotherapy in Squamous Cell Carcinoma of the Esophagus

Tomonori Nakanoko, MD¹, Hiroshi Saeki, MD, Phd, FACS¹, Masaru Morita, MD, Phd, FACS¹, Yuichiro Nakashima, MD, Phd¹, Koji Ando, MD, Phd¹, Eiji Oki, MD, Phd, FACS¹, Takefumi Ohga, MD, Phd, FACS¹, Yoshihiro Kakeji, MD, Phd, FACS¹, Yasushi Toh, MD, Phd, FACS², and Yoshihiko Maehara, MD, Phd, FACS¹

¹Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan;

²Department of Gastrointestinal Surgery, National Kyushu Cancer Center, Fukuoka, Japan

ERRATUM TO: ANN SURG ONCOL DOI 10.1245/S10434-013-3220-2

On page 3 of the original article, the last sentence in the second paragraph is revised as follows:

In multivariate analysis, Rad51 was not an independent prognostic factor (Table 2, $P = 0.9509$).

Table 2 is revised as follows:

TABLE 2 Multivariate analysis of Rad51 expression in the patients without NACRT

Factor	Object	Control	Odds ratio	95 % CI	<i>P</i> value
Depth of invasion	T3/T4	T1/T2	3.04	1.31–7.48	0.0088
Lymph node metastasis	Positive	Negative	1.88	0.82–4.56	0.1390
Rad51	Positive	Negative	1.02	0.48–2.33	0.9509

NACRT neoadjuvant chemoradiotherapy

The online version of the original article can be found under doi:[10.1245/s10434-013-3220-2](https://doi.org/10.1245/s10434-013-3220-2).

© Society of Surgical Oncology 2013

Published Online: 19 October 2013

H. Saeki, MD, Phd, FACS

e-mail: h-saeki@surg2.med.kyushu-u.ac.jp