

ARTICLE

# Systematic analysis of variants related to familial hypercholesterolemia in families with premature myocardial infarction

Ingrid Brænne<sup>1,2,8</sup>, Mariana Kleinecke<sup>1,2,8</sup>, Benedikt Reiz<sup>1,2</sup>, Elisabeth Graf<sup>3</sup>, Tim Strom<sup>3</sup>, Thomas Wieland<sup>3</sup>, Marcus Fischer<sup>4</sup>, Thorsten Kessler<sup>5</sup>, Christian Hengstenberg<sup>5,6</sup>, Thomas Meitinger<sup>3,6,7</sup>, Jeanette Erdmann<sup>\*,1,2,8</sup> and Heribert Schunkert<sup>5,6,8</sup>

Familial hypercholesterolemia (FH) is an oligogenic disorder characterized by markedly elevated low-density lipoprotein cholesterol (LDLC) levels. Variants in four genes have been reported to cause the classical autosomal-dominant form of the disease. FH is largely under-diagnosed in European countries. As FH increases the risk for coronary artery disease (CAD) and myocardial infarction (MI), it might be specifically overlooked in the large number of such patients. Here, we systematically examined the frequency of potential FH-causing variants by exome sequencing in 255 German patients with premature MI and a positive family history for CAD. We further performed co-segregation analyses in an average of 5.5 family members per MI patient. In total, we identified 11 potential disease-causing variants that co-segregate within the families, that is, 5% of patients with premature MI and positive CAD family history had FH. Eight variants were previously reported as disease-causing and three are novel (LDLR.c.811G>A p.(V271I)), PCSK9.c.610G>A (p.(D204N)) and STAP1.c.139A>G (p.(T47A))). Co-segregation analyses identified multiple additional family members carrying one of these FH variants and the clinical phenotype of either FH ( $n=2$ ) or FH and premature CAD ( $n=15$ ). However, exome sequencing also revealed that some variants in FH genes, which have been reported to cause FH, do not co-segregate with FH. The data reveal that a large proportion of FH patients escape the diagnosis, even when they have premature MI. Hence, systematic molecular-genetic screening for FH in such patients may reveal a substantial number of cases and thereby allow a timely LDLC-lowering in both FH/MI patients as well as their variant-carrying family members.

*European Journal of Human Genetics* (2016) 24, 191–197; doi:10.1038/ejhg.2015.100; published online 3 June 2015

## INTRODUCTION

Familial hypercholesterolemia (FH) is an autosomal-dominant disease that leads to markedly elevated low-density lipoprotein cholesterol (LDLC) levels and increased risk for coronary artery disease (CAD) and myocardial infarction (MI). The prevalence of FH is estimated as high as one in 200–500,<sup>1</sup> with even higher frequencies in populations with founder effects.<sup>2</sup> FH is mainly caused by variants in genes coding for proteins affecting hepatic LDLC uptake including the LDL receptor (LDLR), in which most disease-causing variants are found, as well as apolipoprotein B-100 (APOB) and proprotein convertase subtilisin/kexin type 9 (PCSK9).<sup>1,3–6</sup> More recently, *STAP1* has been proposed as a fourth gene causing FH.<sup>7</sup> Cumulatively, variants in these genes explain around 40% of FH cases.<sup>8</sup>

The phenotype may vary in variant carriers or even copied by clustering of common LDL-modifying variants, each affecting LDLC levels by only a small extent.<sup>9</sup> The large number of patients with apparently monogenic FH but without currently known variants also

suggests that other genes, which have not been identified so far, may cause FH.

Multiple studies document the preventive effect of intensive medical LDL-lowering at young age to prevent cardiovascular events.<sup>10–12</sup> Therefore, it has been suggested that incidental detection of variants leading to FH should be communicated to the affected individual and the family.<sup>13</sup> In fact, owing to the high frequency of FH, several guidelines recommend programs to systematically unravel variants and to facilitate medical treatment already at young age.<sup>14–16</sup> Despite the knowledge of causal genes and the obvious advantages of early therapy only 1–15% of FH cases are diagnosed in most European countries.<sup>17</sup> Notable exceptions are Norway (43%) and the Netherlands (71%), in which national screening programs had been initiated.<sup>1,18,19</sup> There are several reasons why FH is vastly under-diagnosed. First, LDLC levels and other clinical presentations of FH are variable.<sup>20</sup> Second, a small family size may obscure the inherited nature of FH,<sup>21–23</sup> and third, with FH being only one of multiple genetic and exogenous conditions

<sup>1</sup>Institut für Integrative und Experimentelle Genomik, Universität zu Lübeck, Lübeck, Germany; <sup>2</sup>DZHK (German Research Centre for Cardiovascular Research), partner site Hamburg/Lübeck/Kiel, Lübeck, Germany; <sup>3</sup>Institute of Human Genetics, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany; <sup>4</sup>Klinik und Poliklinik für Innere Medizin II, Universitätsklinikum Regensburg, Regensburg, Germany; <sup>5</sup>Deutsches Herzzentrum München, Technische Universität München, München, Germany; <sup>6</sup>DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich, Germany; <sup>7</sup>Institute of Human Genetics, Technische Universität München, Munich, Germany

<sup>8</sup>These authors contributed equally to this work.

\*Correspondence: Dr J Erdmann, Institut für Integrative und Experimentelle Genomik, Universität zu Lübeck, Maria-Goeppert-Str. 1, Lübeck 23562, Germany. Tel: +49 451 500 4857; Fax: +49 451 500 5288; E-mail: jeanette.erdmann@iieg.uni-luebeck.de

Received 17 December 2014; revised 2 April 2015; accepted 17 April 2015; published online 3 June 2015

affecting CAD risk, it might be overlooked in the large number of CAD/MI patients.

It is assumed that FH explains about 20% of premature CAD cases with familial clustering.<sup>24,25</sup> However, no systematic analysis is available to quantitate FH variants by molecular-genetic screening in individuals with premature CAD. In this work, we evaluate the frequency of FH due to variants in the *LDLR*, *APOB*, *PCSK9* and *STAP1* genes in 255 unselected patients with premature MI (clinical manifestation before the age of 60) and a positive family history.

## MATERIALS AND METHODS

### Patients

The ascertainment strategy of MI families is described elsewhere.<sup>26,27</sup> In brief, index patients had suffered from MI before the age of 60 years. If at least one additional living sibling was affected with MI or severe CAD (defined by percutaneous coronary intervention or coronary artery bypass grafting) before the age of 70 years, the entire family (index patient, available parents and siblings) was contacted and invited to participate in the study. For the present study, we chose one affected individual each from 255 such families for whole-exome sequencing. Clinical characteristics are detailed in Table 1.

All subjects analyzed in this study gave written informed consent. The local ethical committee (University of Regensburg, Germany) approved the study.

### Exome sequencing

Exome sequencing was performed as 54-bp (base pair) paired end runs on a GenomeAnalyzer Ix system (Illumina, San Diego, CA, USA) after in-solution enrichment of exonic sequences (SureSelect Human All Exon 38 Mb kit, Agilent, Santa Clara, CA, USA), yielding on average 6.2 giga bases (Gb) of sequence per individual. Read alignment was performed with BWA (v. 0.5.8) using the default parameters. We used the human genome assembly hg19 (GRCh37) as reference. A small percentage of duplicate reads (4–5%) were removed. Single nucleotide variants and small insertions and deletions (indels) were detected using SAMtools (v 0.1.7). For the variant filter part of SAMtools, we used the default parameters with the exception of setting a maximum read depth to 9999. Furthermore, we required putative single nucleotide variants to fulfil the following criteria: (i) median base quality of the variant bases of at least 15; (ii) a minimum of 15% of reads showing the variant base; and (iii) the variant base is indicated by at least 5% of reads coming from different strands.

### Variant validation

We used annovar<sup>28</sup> to annotate the single nucleotide variant. Annotation was based on several databases provided by annovar such as UCSC known gene,<sup>29</sup> dbSNP (<http://www.ncbi.nlm.nih.gov/SNP/>), Exome Sequencing Project (<http://evs.gs.washington.edu/EVS/>) and 1000 Genomes.<sup>30</sup> In addition, we also

**Table 1** Clinical characteristics of the MI patients analyzed in this work

Clinical characteristics	MI patients (n = 255 <sup>a</sup> )
Age at inclusion	53.6
Sex	
Male, n (%)	78.3
Female, n (%)	21.7
BMI	27.5
Age at first MI	42.5
LDL <sup>c</sup> (mg/dl)	175.8
Statin therapy, n (%)	62.9
LDL <sup>c</sup> > 190	42.5
Diabetes, n (%)	9.9
Smoking, n (%)	82.6

<sup>a</sup>The mean values are calculated based on the data available.

<sup>b</sup>Adjusted for statin intake according to the CURVES study.<sup>35</sup>

annotated several functional prediction scores such as SIFT,<sup>31</sup> CADD,<sup>32</sup> PolyPhen2<sup>33</sup> and Mutation Taster.<sup>34</sup> Variant validation was performed using PCR and Sanger sequencing. For co-segregation analysis, validated variants were screened in affected and unaffected family members.

Primers that were used for validation are listed in Supplementary Table 1. PCR was carried out in 20  $\mu$ l volume containing 50 ng genomic DNA, 1  $\mu$ l of each primer and either 8  $\mu$ l of Mastermix (5PRIME, Hamburg, Germany) or 0.1  $\mu$ l Taq-Polymerase and 4  $\mu$ l Taq-buffer mix (Bioline Pharmaceutical AG, Baar, Switzerland). Samples were processed in a Sensoquest labcycler with a standard touchdown PCR program (annealing temperature from 59 to 65 °C).

The variants identified in this work were submitted to the publicly funded database LOVD (<http://databases.lovd.nl/shared/variants/>) with the LOVD individual IDs 00033731, 00033768, 00033771–00033807.

## RESULTS AND DISCUSSION

We studied 255 unselected MI/CAD patients from families with strong familial clustering of MI/CAD. The average age at disease manifestation was 42.5 years. The families of the index patients had an average size of 5.5 individuals with an average number of 2.3 affected family members.

Whole-exome sequencing yielded on average 6.2 Gb of sequence per individual. The average read depth was 78 with between 84.5 and 85.6% of the target regions covered at least 20 $\times$ . In total, we identified 259 single nucleotide variants in the *LDLR*, *APOB*, *PCSK9* and *STAP1* genes.

We considered individuals with LDLC levels of > 160 mg/dl after adjusting for statin intake for further analysis.<sup>35</sup> To identify potential disease-causing variants, we filtered these variants based on three assumptions. First, we expect a strong functional impact and therefore, we removed synonymous and intronic variants outside splice-site regions. Second, we expect the variant to be rare in the general population, as FH affects 1 in 200 individuals at the most. Hence, we filtered based on a frequency of 1% in public databases. Third, we filtered variants in segmental duplications due to the high false-positive rate in these regions.

Additionally, a decision tree (Supplementary Figure 1) was developed to further filter the remaining 54 variants. This decision tree includes several considerations, such as the predicted deleterious/damaging effect of a variant on the protein function or whether the variant is known as potentially disease-causing in HGMD.<sup>36</sup> Figure 1 shows the pedigrees of families where a potential disease-causing variant was identified.


### Variant spectrum in *LDLR*, *APOB*, *PCSK9* and *STAP1* genes

In total, we identified 13 rare variants with potentially functional effect in the *LDLR*, 9 variants in *APOB*, 8 variants in *PCSK9* and 1 variant in *STAP1*. Twenty-three variants are missense variants, three are deletions, two are nonsense variants, two are splice-site variants and one is an insertion. Of these 31 variants, 24 were confirmed using Sanger sequencing (see Table 2). Fourteen have been previously reported to cause FH and listed in HGMD (access date March 2014).

### *LDLR* gene variants

*Co-segregating new variant: c.811G>A (p.(V271I))*. We identified only one variant in the *LDLR* gene that has not been previously reported (c.811G>A (p.(V271I))). Both CAD/MI-affected members of family 7421 available for the genetic study carry this variant. Both individuals show elevated LDLC levels of 286 and 261 mg/dl after adjustment for statin treatment. The variant is also found in an family member unaffected by CAD but with markedly elevated LDLC levels (220 mg/dl).

Variant c.811G>A (p.(V271I)) lies in the domain that interacts with *APOB*<sup>37</sup> and is in close proximity with several disease-causing amino acid changes listed in HGMD, for instance p.N272T, p.C270S,



**Figure 1** Pedigrees of families where a potential disease-causing variant was identified. Squares are males and circles are females. Individuals affected with MI/CAD are shown as dark symbols. The first row of numbers below the symbols are the individual IDs, the second line show the genotype for the identified variant and the third row the corrected LDLC level. (LDLC levels marked with \* are not corrected for statin treatment.) Detailed information is shown in Table 2. The potential disease-causing variant per family are: Family ID 4349: LDLR. c.131G>A (p.(W44\*)); Family ID7520: LDLR. c.1285G>A (p.(V429M)); Family ID8450: LDLR. c.G1775 (p.(G592E)); Family ID 9242: LDLR. c.1444G>A (p.(D482N)); Family ID 7500: PCSK9. c.610G>A (p.(D204N)); Family ID 8797: STAP1. c.139A>G (p.(T47A)); Family ID 6548 LDLR. c.2231G>A (p.(R744Q)); Family ID 7421: LDLR. c.811G>A (p.(V271I)); Family IDs 8400, 8615, 9192: APOB. c.10580G>A (p.(R3527Q)) //APOB. c.7696G>A (p.(E2566K)); Family ID6502: LDLR.c.1359-1G>A; Family ID 6565: LDLR. c.757C>T (p.(R253W)).

p.C270R, p.C276R, p.G269D or p.C270Y.<sup>38–43</sup> On the basis of the position of the identified variant with respect to previously reported variants, it is very likely that c.811G>A (p.(V271I)) is responsible for the increase of LDLC in the variant carriers.

*Co-segregating known variants:* c.1285G>A (p.(V429M)), c.1444G>A (p.(D482N)), c.G1775 (p.(G592E)), c.2231G>A (p.(R744Q)), c.757C>T (p.(R253W)), c.131G>A (p.(W44\*)), c.798T>A (p.(D266E)) and c.828C>A (p.(C276\*)). The missense variants

**Table 2 Validated variants identified by exome sequencing of 255 MI-affected individuals**

Family ID	Gene	Position and change	HGVs (nucleotide and protein)	Reference sequence	Co-segregation with elevated LDLC	Index ID	Mean corrected LDLC in variant carriers (mg/dl)	Mean corrected LDLC in non-carriers (mg/dl)	Frequency internal controls (n = 1,4xx)	HGMD
4349	LDLR	g.11210962G>A	c.131G>A(p.(W44*))	NM_000527.4	+	501	483.5	107	0	+
4318	LDLR	g.11213464T>C	c.313+2T>C	NM_000527.4	-	501	225*	247	0	+
6565	LDLR	g.11217303C>T	c.757C>T(p.(R253W))	NM_000527.4	+	502	185	143	0	+
7080	LDLR	g.11217344T>A	c.798T>A(p.(D266E))	NM_000527.4	-	502	310	187	2	+
7421	LDLR	g.11217357G>A	c.811G>A(p.(V271I))	NM_000527.4	+	501	265.7	-	0	+
8985	LDLR	g.11218078C>A	c.828C>A(p.(C276*))	NM_000527.4	+	501	204	205	0	+
7520	LDLR	g.11224052G>A	c.1285G>A(p.(V429M))	NM_000527.4	+	502	341	-	0	+
6502	LDLR	g.11224210G>A	c.1359-1G>A	NM_000527.4	+	501	366.4	-	0	+
9242	LDLR	g.11224296G>A	c.1444G>A(p.(D482N))	NM_000527.4	+	501	388.5	130.5	0	+
8450	LDLR	g.11227604G>A	c.1775G>A(p.(G592E))	NM_000527.4	+	501	446	-	0	+
8700	LDLR	g.11233886C>T	c.2177C>T(p.(T726I))	NM_000527.4	- (2/6)	503	114*	160	0	+
7442						501	162.5	139	25	+
6035						501	195	238.7		
7421						502	265.7			
8887						501	114*	162		
4373						502	154			
6548	LDLR	g.11233940G>A	c.2231G>A(p.(R744Q))	NM_000527.4	+	501	237	-	1	+
8615	APOB	g.21229160C>T	c.10580G>A (p.(R3527Q))	NM_000384.2	+	503	275	160	6	+
9192						501	246	212		
8400						501	338.5	-		
9603	APOB	g.21229703C>A	c.10037G>T (p.(S3346I))	NM_000384.2	-	501	171.2	167	0	-
9076	APOB	g.21232044C>T	c.7696G>A (p.(E2566K))	NM_000384.2	-	502	175	191	21	+
8910						502	326	223		
4016						501	188.5	-		
8708						501	92.5	148.5		
8615						503	275	160		
9192						501	246	212		
8400						501	338.5	-		
9082	APOB	g.21232606G>T	c.7134C>A (p.(S2378R))	NM_000384.2	-	501	372	118	0	-
6491	APOB	g.21233102_21233104del	c.6636_6638del (p.(2212_2213del))	NM_000384.2	-	501	161	-	47	-
8209						505	155	157		
6431						502	184	-		
4211						501	215	169.5		
4368						501	172	132		
8531	APOB	g.21234086T>C	c.5654A>G (p.(Y1885C))	NM_000384.2	-	501	173.5	-	0	-
4318						501	187	247		
9082	APOB	g.21234674C>T	c.5066G>A (p.(R1689H))	NM_000384.2	-	501	372	118	27	+
6688						503	162.5	81		
6553						501	203	140.5		
9403	APOB	g.21246416A>G	c.2585T>C(p.(V862A))	NM_000384.2	-	501	224	179.5	0	-
4335	APOB	g.21260868G>A	c.4990>T(p.(P167S))	NM_000384.2	-	501	158.5	137.7	0	-
7438	APOB	g.55512245_55512246del	c.449_450del (p.(L150_150del))	NM_174936.3	-	501	168.8	-	0	-
8610	PCSK9	g.55512245_55512246del	c.449_450del (p.(L150_150del))	NM_174936.3	-	501	198.5	163.5	0	-
7500	PCSK9	g.55518037G>A	c.610G>A(p.(D204N))	NM_174936.3	+	502	314	104	1	-
8797	STAP1	g.68436820A>G	c.139A>G(p.(T47A))	NM_012108.2	+	501	247.5	-	0	-

Abbreviations: AA, amino acid; HGMD, Human Gene Mutation Database; LDLC, LDL cholesterol. LDLC levels marked with \* are not corrected for statin treatment. The reference sequence used is hg19. LDLR gene is located on chromosome 19, APOB on chromosome 2, PCSK9 on chromosome 1 and STAP1 on chromosome 4.

c.1285G>A (p.(V429M)) (first found in the African population<sup>44</sup>), c.1444G>A (p.(D482N)), c.G1775 (p.(G592E)), c.2231G>A (p.(R744Q)), c.757C>T (p.(R253W)) and c.131G>A (p.(W44\*)) show co-segregation with elevated LDLC levels (mean 344 mg/dl after adjustment for statin treatment) and MI in the families under study. Hence, on the basis of our own results and evidence in previously reported studies,<sup>38,44–46</sup> these variants probably cause the disease in these patients/families.

The splice site variant c.1359-1G>A has also been reported to cause FH.<sup>47</sup> In addition, we have previously also identified this variant as probably disease-causing in one of our extended MI/FH families.<sup>48</sup> Co-segregation analysis showed that all five family members carry the variant. Four family members are affected with MI/CAD and all have elevated LDLC levels even after statin therapy (mean LDLC of 366 mg/dl). Hence, our data confirm previous findings.

The missense variant (c.798T>A (p.(D266E))) and the nonsense variant (c.828C>A (p.(C276\*))) were also found in individuals with markedly elevated LDLC levels and MI. Both variants have been reported as disease-causing in HGMD.<sup>38,49</sup> Here, we only identified both variants in the index patients of family 7080 and 8985 and not in the two other affected family members. Both index patients have markedly elevated LDLC levels of 310 and 204 mg/dl, respectively. Interestingly, the two non-carriers within these families also have elevated LDLC levels of 187 and 205 mg/dl.

Whereas c.828C>A (p.(C276\*)) is not found in our internal non CAD/FH patients, c.798T>A (p.(D266E)) is found twice. However, as FH is a common disease, we also expect to find a small number of FH cases in population-based controls.

**Non-co-segregating known variant: rs45508991.** One LDL receptor variant, c.2177C>T (p.(T726I),rs45508991), is found in 6 of the 255 patients. This variant is reported to cause FH and is labelled as likely disease-causing in HGMD but the variant may be only pathogenic in combination with another variant in the *LDLR* gene.<sup>50,51</sup> In our data, we checked for co-segregation with LDLC and observed a positive co-segregation only in one family (family 7421) that also carries an additional co-segregating LDLR variant (c.811G>A (p.(V271I))). The other five families show poor or no evidence that the variant causes FH. In fact, in carriers of this variant, LDLC levels range from 147 to 268 mg/dl, and in non-carriers from 86 to 257 mg/dl.

Despite incomplete co-segregation with elevated LDLC levels, c.2177C>T (p.(T726I)) is only found in one MI/CAD-affected family member (Supplementary Figure 2). In our data set, we do not find variant carriers with neither MI/CAD nor FH. To further check whether c.2177C>T (p.(T726I)) is associated with an increased risk of CAD or FH, we compared the number of variant carriers in our sample within the general population. As c.2177C>T (p.(T726I)) is mainly found in individuals of European ancestry,<sup>50</sup> we compared the frequency only in European samples. With a frequency of 0.007 in 1000 Genomes (1kG) data set (European samples, phase 3 version 5) *versus* 0.023 (6/255) in our sample set, there is evidence that the variant indeed increases the risk of CAD/FH. However, we also found c.2177C>T (p.(T726I)) in 25 of 1462 (0.017) internal European non-CAD patients (*P*-value=0.477). Hence, as the variant is equally common in affected *versus* unaffected individuals, we expect the accumulation of c.2177C>T (p.(T726I)) rather to be a founder effect than related to an increased risk of CAD/FH.

**Non-co-segregating known variant: c.313+2T>C.** LDLR variant c.313+2T>C does neither co-segregate with elevated LDLC levels nor with MI/CAD. This splice-site variant is only found in the index patient and

not in the two relatives with elevated LDLC levels (mean LDLC 217 mg/dl) of which one is also MI/CAD-affected. Variant c.313+2T>C is reported to cause FH<sup>52</sup> and is labelled as disease-causing in HGMD. However, on the basis of the results of co-segregation analysis in family 4318, we do not expect the variant to be disease-causing, at least not to be the primary cause of MI/CAD in this family.

**Double variant: LDLR. c.131G>A (p.(W44\*)) and PCSK9.c.137G>T (p.(R46L)).** For all validated LDLR-variant carriers, we checked for a second variant in *PCSK9*, *APOB* and *STAP1* gene. We found one compound heterozygote patient with a variant in the *LDLR* gene (c.131G>A (p.(W44\*))) and *PCSK9* gene (c.137G>T (p.(R46L))) (family 4349, patient 501). The *PCSK9* variant, c.137G>T (p.(R46L)), is reported to be associated with a significant reduction of LDLC.<sup>53</sup> Co-segregation analysis revealed that one of the three c.131G>A (p.(W44\*)) carriers also carry c.137G>T (p.(R46L)) (patient 501), in addition to one family member without the *LDLR* variant (patient 504).

Reduced *PCSK9* activity leads to increased density of LDLR<sup>54,55</sup> at the cell membrane. As we have heterozygous variant carriers expressing one healthy *LDLR* allele, we would not expect to see a strong effect, but expect to see reduced levels of LDLC for the double variant carriers. However, the LDLC level is markedly elevated for all three c.131G>A (p.(W44\*)) variant carriers (mean LDLC 322 mg/dl). We do not identify a higher level of LDLC in the affected relative with the protective *PCSK9* variant (LDLC level of 432 mg/dl). On the contrary, this patient has the highest LDLC level in the family. This could, however, imply that the protective effect of the *PCSK9* variant is negligible compared with the LDLC level increase caused by the *LDLR* variant. The unaffected relative carrying only the reported protective variant has similar LDLC levels as the second unaffected relative lacking both variants. In summary, we do not see a reduction in LDLC levels in c.137G>T (p.(R46L)) carriers and would not expect the variant to markedly reduce LDLC levels.

#### **APOB gene variants**

Of the nine *APOB* variants, three were found in HGMD: c.7696G>A (p.(E2566K)) and c.5066G>A (p.(R1689H)) are reported to be associated with high triglyceride and c.10580G>A (p.(R3527Q)) with high LDLC levels. Of the nine variants, only one co-segregates with FH and MI/LDLC levels, and is previously reported. In the following, we will only discuss the three HGMD variants.

**c.10580G>A (p.(R3527Q)) and c.7696G>A (p.(E2566K)).** Interestingly, c.10580G>A (p.(R3527Q)) is always found in combination with the c.7696G>A (p.(E2566K)) variant. Vice versa, c.7696G>A (p.(E2566K)) is also found without the c.10580G>A (p.(R3527Q)) variant. As c.10580G>A (p.(R3527Q)) appears to be on the same allele as c.7696G>A (p.(E2566K)), c.10580G>A (p.(R3527Q)) seems to be the more recent variant.

The c.10580G>A (p.(R3527Q)) variant was found in three families (8615, 9192, 8400). Variant c.10580G>A (p.(R3527Q)) has been reported to cause defective binding of ApoB to the *LDLR*.<sup>56</sup> All five c.10580G>A (p.(R3527Q)) carriers show elevated LDLC levels (mean LDLC 300 mg/dl). However, two siblings with elevated LDLC (mean 206 mg/dl), do not carry the variant. These non-carriers could, however, have another cause of disease. Hence, there is evidence that c.10580G>A (p.(R3527Q)) may cause elevated LDLC levels. Therefore, our results support previous findings.

Variant c.7696G>A (p.(E2566K)) was identified in seven families. This variant does neither co-segregate with elevated LDLC levels nor

with MI. In the seven families, we find three affected individuals (mean 214 mg/dl) without the variant as well as four unaffected variant carriers (mean 133 mg/dl).

*Non-co-segregating known variant: c.5066G>A (p.(R1689H)).* The c.5066G>A (p.(R1689H)) variant was found in three of the 255 exome-sequenced MI-patients. Two non-carriers have LDLC levels above 190 mg/dl while four variant carriers do not show elevated cholesterol levels (mean 156 mg/dl). Hence, we do not expect this variant to be the cause of disease in these families nor increase the risk of FH/MI in general.

#### PCSK9 gene variant

We identified two variants in the *PCSK9* gene. Both variants have not been reported earlier. On the basis of co-segregation analysis with LDLC levels, the c.610G>A (p.(D204N)) variant might cause FH. The affected variant carrier has LDLC levels of 314 mg/dl and the MI-affected brother has low LDLC levels (104 mg/dl). Hence, the cause of MI may differ in this individual. The variant c.449\_450del (p.(150\_150del)) does not co-segregate with FH in the family.

#### STAP1 gene variant

Variant c.139A>G (p.(T47A)) is the only variant we found in the *STAP1* gene and has not been reported previously. Both family members available for genetic studies are MI-affected and carry the amino acid substitution. In addition, both show elevated LDLC levels (mean 248 mg/dl). Hence, we might have identified the causal variant in this family but functional studies are necessary to further evaluate its functional implication.

#### Potential polygenic cause of FH

Of the analyzed CAD patients, 48% have LDLC levels above 190 mg/dl after correction for statin intake<sup>35</sup> (LDLC levels were available for 212 CAD patients). We find rare potential disease-causing variants in 12.7% of these patients CAD patients with high LDLC levels. It has recently been shown that the clinical phenotype of FH can also be caused by the accumulation of common variants with small effects.<sup>9</sup> Indeed, it has been reported that a score of six SNPs enables to discriminate FH patients from healthy controls.<sup>57</sup> We calculated the score as described by Futema *et al* for the patients with high LDLC levels but without a rare disease-causing variant and compared the score with the controls of the German MI Family Study II<sup>58</sup> ( $n=1298$ ). Not all SNPs were covered by exome sequencing, so we calculated the score based on GeneChip Human Mapping 500 K Array Set (Affymetrix) available for 234 of the 255 CAD patients. The mean score in controls (0.63) are significantly lower than the mean score (0.69) in patients with high LDLC levels ( $P$ -value = 0.025). Our values are in range with the scores reported by Futema *et al* (0.63 in the control cohort and 0.71 in the mutation negative FH patients). Hence, the mutation-negative patients with elevated LDLC levels might have a polygenic cause of disease.

#### CONCLUSION

Here, we screened 255 patients with premature MI/CAD for variants in genes known to cause FH. If we only account for variants in the *LDLR* gene, 3.1% of the patients carry potential FH-causing variants. If we also account for variants in *APOB*, *PCSK9* and *STAP1*, we have a cumulative frequency of FH-causing variants of 5.1%. Indeed, the frequency of potential disease-causing variants in our sample is probably underestimated. Large rearrangements are reported to account for 11% of the *LDLR* variants.<sup>59</sup> Unfortunately, the coverage

profile of our data was not uniform enough to allow a screening for such variants and hence we might have missed some of these.

In summary, the high frequency of potential FH-causing variants in these unselected MI/CAD patients supports the hypothesis that FH is overseen in a substantial number of patients with MI/CAD and demonstrates that genetic screening also of MI/CAD patients can improve diagnosis of FH.

Additionally, we also screened family members of the index patients for the identified variants. This revealed that 17 family members also carry the potential FH-causing variant. Hence, our findings underline the need for a systematic molecular-genetic screening to enable an early diagnosis of FH and to allow timely preventive treatment.

A further interesting finding is that the functional effect of several variants reported as disease-causing, for instance in HGMD, is questionable. We observed that five reported causal variants show none or minor functional impact in our analyzed families. Consequently, given the far-reaching implications of the diagnosis of FH, each variant has to be carefully evaluated. In fact, a co-segregation analysis is advisable to determine whether a variant truly is disease-causing.

In summary, our work demonstrates that exome sequencing can be used for FH-variant screening. In addition, the quality of the exome sequencing has improved over the last years, allowing identification not only of small nucleotide variants but also large rearrangements.<sup>60</sup> Also, as the sequencing costs have decreased dramatically, exome sequencing might become the method of choice for molecular genetic screening of, for instance, FH.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### ACKNOWLEDGEMENTS

We thank all the family members who participated in this research. Without the continuous support of these patients over more than 15 years, the present work would not have been possible. We like to thank Sandra Wrobel for technical assistance. We also would like to thank Drs Björn Mayer, Ute Hubauer and Anika Großhennig for help with the German Myocardial Infarction Study. The study is supported by the German Federal Ministry of Education and Research (BMBF) in the context of the e:Med program (e:AtheroSysMed and sysINFLAME) and the FP7 European Union project CVgenes@target (261123). Further grants were received by the local focus program 'Medizinische Genetik' of the Universität zu Lübeck, as well as by the Fondation Leducq (CADgenomics: Understanding Coronary Artery Disease Genes, 12CVD02). This study was also supported through the Deutsche Forschungsgemeinschaft (DFG) cluster of excellence 'Inflammation at Interfaces'.

#### AUTHOR CONTRIBUTIONS

JE, IB and HS designed the study; IB, MK and BR performed the bioinformatic analysis; EG, TS, TM and TW performed the NGS; MK, MF and CH collected the family data; IB, MK, JE and HS wrote the manuscript. All authors read and approved the final manuscript.

- 1 Nordestgaard BG, Chapman MJ, Humphries SE *et al*: Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: guidance for clinicians to prevent coronary heart disease: consensus statement of the European Atherosclerosis Society. *Eur Heart J* 2013; **34**: 3478–3490a.
- 2 Foody JM: Familial hypercholesterolemia: an under-recognized but significant concern in cardiology practice. *Clin Cardiol* 2014; **37**: 119–125.
- 3 De Castro-Oros I, Pocovi M, Civeira F: The genetic basis of familial hypercholesterolemia: inheritance, linkage, and mutations. *App Clin Genet* 2010; **3**: 53–64.

- 4 Austin MA, Hutter CM, Zimmern RL, Humphries SE: Genetic causes of monogenic heterozygous familial hypercholesterolemia: a HuGE prevalence review. *Am J Epidemiol* 2004; **160**: 407–420.
- 5 Umans-Eckenhausen MA, Defesche JC, Sijbrands EJ, Scheerder RL, Kastelein JJ: Review of first 5 years of screening for familial hypercholesterolemia in the Netherlands. *Lancet* 2001; **357**: 165–168.
- 6 Leren TP, Manshaus T, Skovholt U *et al*: Application of molecular genetics for diagnosing familial hypercholesterolemia in Norway: results from a family-based screening program. *Semin Vasc Med* 2004; **4**: 75–85.
- 7 Fouchier SW, Dallinga-Thie GM, Meijers JC *et al*: Mutations in STAP1 Are Associated With Autosomal Dominant Hypercholesterolemia. *Circ Res* 2014; **115**: 552–555.
- 8 Santos RD, Maranhao RC: What is new in familial hypercholesterolemia? *Curr Opin Lipidol* 2014; **25**: 183–188.
- 9 Talmud PJ, Shah S, Whittall R *et al*: Use of low-density lipoprotein cholesterol gene score to distinguish patients with polygenic and monogenic familial hypercholesterolemia: a case-control study. *Lancet* 2013; **381**: 1293–1301.
- 10 Vermissen J, Oosterveer DM, Yazdanpanah M *et al*: Efficacy of statins in familial hypercholesterolemia: a long term cohort study. *BMJ* 2008; **337**: a2423.
- 11 Wiegman A, de Groot E, Hutten BA *et al*: Arterial intima-media thickness in children heterozygous for familial hypercholesterolemia. *Lancet* 2004; **363**: 369–370.
- 12 Marks D, Thorogood M, Neil HA, Humphries SE: A review on the diagnosis, natural history, and treatment of familial hypercholesterolemia. *Atherosclerosis* 2003; **168**: 1–14.
- 13 Green RC, Berg JS, Grody WW *et al*: ACMG recommendations for reporting of incidental findings in clinical exome and genome sequencing. *Genet Med* 2013; **15**: 565–574.
- 14 Reiner Z, Catapano AL, De Backer G *et al*: [ESC/EAS Guidelines for the management of dyslipidaemias]. *Revista espanola de cardiologia* 2011; **64**: 1168.e1–1168.e60.
- 15 Minhas R, Humphries SE, Qureshi N, Neil HA, Group NGD: Controversies in familial hypercholesterolemia: recommendations of the NICE Guideline Development Group for the identification and management of familial hypercholesterolemia. *Heart* 2009; **95**: 584–587, discussion 587–591.
- 16 National-Institute-of-Health. Third Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) 2004.
- 17 Brice P, Burton H, Edwards CW, Humphries SE, Aitman TJ: Familial hypercholesterolemia: a pressing issue for European health care. *Atherosclerosis* 2013; **231**: 223–226.
- 18 Goldberg AC, Hopkins PN, Toth PP *et al*: Familial hypercholesterolemia: screening, diagnosis and management of pediatric and adult patients: clinical guidance from the National Lipid Association Expert Panel on Familial Hypercholesterolemia. *J Clin Lipidol* 2011; **5**: S1–S8.
- 19 Benn M, Watts GF, Tybjaerg-Hansen A, Nordestgaard BG: Familial hypercholesterolemia in the danish general population: prevalence, coronary artery disease, and cholesterol-lowering medication. *J Clin Endocrinol Metab* 2012; **97**: 3956–3964.
- 20 Nauck MS, Koster W, Dorfer K *et al*: Identification of recurrent and novel mutations in the LDL receptor gene in German patients with familial hypercholesterolemia. *Hum Mutat* 2001; **18**: 165–166.
- 21 Pötzsch O, Weinmann J, Haustein T: Geburtstrends und familiensituation in deutschland. *Statistisches Bundesamt, Wiesbaden* 2013, 2013.
- 22 Office-for-National-Statistics. Family size in 2012, 2013.
- 23 Testa MR: Family sizes in Europe: evidence from the 2011 Eurobarometer Survey. *European Demographic Research Papers* 2. Vienna: Vienna Inst of Demography, 2012.
- 24 Goldstein JL, Schrott HG, Hazzard WR, Bierman EL, Motulsky AG: Hyperlipidemia in coronary heart disease. II. Genetic analysis of lipid levels in 176 families and delineation of a new inherited disorder, combined hyperlipidemia. *J Clin Invest* 1973; **52**: 1544–1568.
- 25 Neeffjes LA, Ten Kate GJ, Alexia R *et al*: Accelerated subclinical coronary atherosclerosis in patients with familial hypercholesterolemia. *Atherosclerosis* 2011; **219**: 721–727.
- 26 Broeckel U, Hengstenberg C, Mayer B *et al*: A comprehensive linkage analysis for myocardial infarction and its related risk factors. *Nat Genet* 2002; **30**: 210–214.
- 27 Fischer M, Broeckel U, Holmer S *et al*: Distinct heritable patterns of angiographic coronary artery disease in families with myocardial infarction. *Circulation* 2005; **111**: 855–862.
- 28 Wang K, Li M, Hakonarson H: ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* 2010; **38**: e164.
- 29 Karolchik D, Barber GP, Casper J *et al*: The UCSC Genome Browser database: 2014 update. *Nucleic Acids Res* 2014; **42**: D764–D770.
- 30 Abecasis GR, Altshuler D, Auton A *et al*: 1000 Genomes Project Consortium A map of human genome variation from population-scale sequencing. *Nature* 2010; **467**: 1061–1073.
- 31 Ng PC, Henikoff S: Predicting deleterious amino acid substitutions. *Genome Res* 2001; **11**: 863–874.
- 32 Kircher M, Witten DM, Jain P, O’Roak BJ, Cooper GM, Shendure J: A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet* 2014; **46**: 310–315.
- 33 Adzhubei IA, Schmidt S, Peshkin L *et al*: A method and server for predicting damaging missense mutations. *Nat Methods* 2010; **7**: 248–249.
- 34 Schwarz JM, Rodelsperger C, Schuelke M, Seelow D: MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods* 2010; **7**: 575–576.
- 35 Pincus J: Comparative dose efficacy study of atorvastatin versus simvastatin, pravastatin, lovastatin, and fluvastatin in patients with hypercholesterolemia (the CURVES study). *Am J Cardiol* 1998; **82**: 406–407.
- 36 Loux N, Saint-Jore B, Collod G *et al*: Screening for new mutations in the LDL receptor gene in seven French familial hypercholesterolemia families by the single strand conformation polymorphism method. *Hum Mutat* 1992; **1**: 325–332.
- 37 Russell DW, Brown MS, Goldstein JL: Different combinations of cysteine-rich repeats mediate binding of low density lipoprotein receptor to two different proteins. *J Biol Chem* 1989; **264**: 21682–21688.
- 38 Hobbs HH, Brown MS, Goldstein JL: Molecular genetics of the LDL receptor gene in familial hypercholesterolemia. *Hum Mutat* 1992; **1**: 445–466.
- 39 Duskova L, Kopeczkova L, Jansova E *et al*: An APEX-based genotyping microarray for the screening of 168 mutations associated with familial hypercholesterolemia. *Atherosclerosis* 2011; **216**: 139–145.
- 40 Fouchier SW, Kastelein JJ, Defesche JC: Update of the molecular basis of familial hypercholesterolemia in The Netherlands. *Hum Mutat* 2005; **26**: 550–556.
- 41 Kuhrova V, Francova H, Zapletalova P *et al*: Spectrum of low density lipoprotein receptor mutations in Czech hypercholesterolemic patients. *Hum Mutat* 2001; **18**: 253.
- 42 Bertolini S, Cantafora A, Aversa M *et al*: Clinical expression of familial hypercholesterolemia in clusters of mutations of the LDL receptor gene that cause a receptor-defective or receptor-negative phenotype. *Arterioscler Thromb Vasc Biol* 2000; **20**: E41–E52.
- 43 Mozas P, Castillo S, Tejedor D *et al*: Molecular characterization of familial hypercholesterolemia in Spain: identification of 39 novel and 77 recurrent mutations in LDLR. *Hum Mutat* 2004; **24**: 187.
- 44 Leitersdorf E, Van der Westhuyzen DR, Coetzee GA, Hobbs HH: Two common low density lipoprotein receptor gene mutations cause familial hypercholesterolemia in Afrikaners. *J Clin Invest* 1989; **84**: 954–961.
- 45 Ward AJ, O’Kane M, Young I, Nicholls DP, Nevin NC, Graham CA: Three novel mutations in the EGF precursor homology domain of the low-density lipoprotein receptor gene in Northern Irish patients with familial hypercholesterolemia. *Hum Mutat* 1995; **6**: 254–256.
- 46 Sun XM, Patel DD, Knight BL, Soutar AK: Influence of genotype at the low density lipoprotein (LDL) receptor gene locus on the clinical phenotype and response to lipid-lowering drug therapy in heterozygous familial hypercholesterolemia. The Familial Hypercholesterolemia Regression Study Group. *Atherosclerosis* 1998; **136**: 175–185.
- 47 Peeters AV, Van Gaal LF, du Plessis L, Lombardi MP, Havekes LM, Kotze MJ: Mutational and genetic origin of LDL receptor gene mutations detected in both Belgian and Dutch familial hypercholesterolemic. *Hum Genet* 1997; **100**: 266–270.
- 48 Brønne I, Reiz B, Medack A *et al*: Whole-exome sequencing in an extended family with myocardial infarction unmasks familial hypercholesterolemia. *BMC Cardiovasc Disord* 2014; **14**: 108.
- 49 Mozas P, Cenarro A, Civeira F, Castillo S, Ros E, Pocovi M: Mutation analysis in 36 unrelated Spanish subjects with familial hypercholesterolemia: identification of 3 novel mutations in the LDL receptor gene. *Hum Mutat* 2000; **15**: 483–484.
- 50 Tejedor MT, Cenarro A, Tejedor D *et al*: Haplotype analyses, mechanism and evolution of common double mutants in the human LDL receptor gene. *Mol Genet Genomics* 2010; **283**: 565–574.
- 51 Alharbi KK, Aldahmesh MA, Spanakis E *et al*: Mutation scanning by meltMADGE: validations using BRCA1 and LDLR, and demonstration of the potential to identify severe, moderate, rare, and paucimorphic mutations in the general population. *Genome Res* 2005; **15**: 967–977.
- 52 Lombardi P, Sijbrands EJ, van de Giessen K *et al*: Mutations in the low density lipoprotein receptor gene of familial hypercholesterolemic patients detected by denaturing gradient gel electrophoresis and direct sequencing. *J Lipid Res* 1995; **36**: 860–867.
- 53 Cohen JC, Boerwinkle E, Mosley TH Jr., Hobbs HH: Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med* 2006; **354**: 1264–1272.
- 54 Youngblom E, Knowles JW: Familial hypercholesterolemia In: Pagon RA, Adam MP, Bird TD (eds): *GeneReviews*. Seattle (WA): University of Washington. 1993.
- 55 Wu NQ, Li JJ: PCSK9 gene mutations and low-density lipoprotein cholesterol. *Clin Chim Acta* 2014; **431C**: 148–153.
- 56 Soria LF, Ludwig EH, Clarke HR, Vega GL, Grundy SM, McCarthy BJ: Association between a specific apolipoprotein B mutation and familial defective apolipoprotein B-100. *Proc Natl Acad Sci USA* 1989; **86**: 587–591.
- 57 Futema M, Shah S, Cooper JA *et al*: Refinement of variant selection for the LDL cholesterol genetic risk score in the diagnosis of the polygenic form of clinical familial hypercholesterolemia and replication in samples from 6 countries. *Clin Chem* 2015; **61**: 231–238.
- 58 Erdmann J, Großhennig A, Braund PS *et al*: New susceptibility locus for coronary artery disease on chromosome 3q22.3. *Nat Genet* 2009; **41**: 280–282.
- 59 Goldmann R, Tichy L, Freiberger T *et al*: Genomic characterization of large rearrangements of the LDLR gene in Czech patients with familial hypercholesterolemia. *BMC Med Genet* 2010; **11**: 115.
- 60 Futema M, Plagnol V, Whittall RA *et al*: Use of targeted exome sequencing as a diagnostic tool for Familial Hypercholesterolemia. *J Med Genet* 2012; **49**: 644–649.