

Evaluation of Toxicity Effects of Asafetida on Biochemical, Hematological, and Histological Parameters in Male Wistar Rats

Seyyed Majid Bagheri, Maryam Yadegari¹, Aghdas Mirjalily¹, Mohammad Ebrahim Rezvani²

Department of Physiology, Herbal Medicine Research Center, Shahid Sadoghi University of Medical Sciences, ¹Department of Anatomy, Shahid Sadoghi University of Medical Sciences, ²Neurobiomedical Research Center, Shahid Sadoghi University of Medical Sciences, Yazd, Iran

ABSTRACT

Objectives: Asafetida is traditionally used in folklore medicine for the treatment of various ailments. To validate its use in traditional medicine, it is important to evaluate its toxicity in the animal system. Therefore, this study aimed to evaluate the toxicological effects of asafetida in Wistar albino rats. **Materials and Methods:** Acute toxicity tests were conducted by the oral administration of 250, 500, and 1,000 mg/kg body weight of the animal. In chronic study, animals were administered with various doses of asafetida (25, 50, 100, and 200 mg/kg body weight) for a period of 6 weeks. At end of experiment, the effects of asafetida on hematological, renal, and hepatic markers and histological parameters were analyzed. **Results:** In acute toxicity study, no mortality was seen up to 72 h of the administration of asafetida. No signs of neurological and behavioral changes were noticed within 24 h. In the chronic study, the asafetida intake has changed the hematological parameters such as red blood cell (RBC), white blood cell (WBC), hematocrit (HCT), and platelets. Aspartate aminotransferase (AST) and lactate dehydrogenase (LDH) were significantly increased in treated animals. The plasma level of urea and creatinine were not altered by the administration of asafetida throughout the study. Histopathology study indicates hepatotoxicity, but no signs of prominent pathological changes in kidney. **Conclusions:** Asafetida did not show any acute toxicity, but chronic administration could have undesirable effects on hepatocytes and hematological factors.

Key words: Asafetida, hematology, kidney, liver, toxicity

INTRODUCTION

Asafetida, an oleo-gum-resin, is extracted by incisions in the stem and roots of *Ferula assafoetida* L. and some others *Ferula* species.^[1] The resin is an important pharmacological and industrial agent and is being used in traditional

medicines for treatment of various disorders like stomach ache, indigestion, bronchitis, asthma, and whooping cough;^[2] and for treatment of intestinal parasites.^[3] Nepalian people use asafetida 50–200 mg twice a week mainly as sedative, carminative, antispasmodic, diuretic, antihelmintic, emmenagogue, and aphrodisiac agent.^[4] In Ayurveda, asafetida is introduced as a valuable remedy for flatulence, hysteria, gastrointestinal disorders, inflammation, stomach ache, spasmodic, and helminthic.^[1] New studies

Address for correspondence: Mr. Seyyed Majid Bagheri, Department of Physiology, Shahid Sadoughi University of Medical Sciences, Prof. Hesabi Bulvd. Shohadaye Gomnam Bulvd, Yazd-8915173149, Iran.
E-mail: boss_bagheri@yahoo.com

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Bagheri SM, Yadegari M, Mirjalily A, Rezvani ME. Evaluation of toxicity effects of asafetida on biochemical, hematological, and histological parameters in male wistar rats. *Toxicol Int* 2015;22:61-5.

Access this article online

Quick Response Code:



Website:

www.toxicologyinternational.com

DOI:

10.4103/0971-6580.172258

showed that this oleo-gum-resin has antiviral, antifungal, cancer chemopreventive, antidiabetic, and cytotoxicity effect.^[1] It also has antispasmodic, anticonvulsant, and antinociceptive effects.^[5-7] Although there are different studies about pharmacological properties of asafoetida, to our knowledge, there is no comprehensive toxicological study of asafoetida on animal models. There is only one case report that methemoglobinemia has been observed after administration of asafoetida in a 5-week-old black male infant.^[8] It is also recommended that asafoetida should not be used during pregnancy as it may increase the risk of abortion.^[9]

The toxicity effects of asafoetida have been studied mainly on parasites and some protozoan animals suggesting it antiparasitic,^[10-12] antifungal,^[13] and antibacterial.^[14] Kumar and Singh reported that different root extracts of *Ferula assafoetida* have anti molluscicidal activity against the snail *Lymnaea acuminata*.^[10] Bagheri et al., showed that asafoetida has cytotoxicity effect on the brine shrimp.^[15] Some of old studies showed that asafoetida has a weak exchange-inducing on sister chromatid in spermatogonia^[16] and clastogenicity in mouse spermatocytes.^[17] The study is aimed to study the toxic effects of asafoetida on blood parameters, kidney and liver factors, and histopathology of kidney and liver in male Wistar rats.

MATERIALS AND METHODS

Animals

Male Wistar rats weighing between 150 and 180g and 6–8 weeks old were bred and maintained in the animal house unit of the faculty of medicine under controlled temperature at $21 \pm 1^\circ\text{C}$ in 12 h light: 12 h darkness schedule. Animals were housed in plastic cages and food and water was made available *ad libitum*. The study was approved by Institutional Animal Ethical Committee of Shahid Sadoughi University of Medical Science.

Acute toxicity studies

For acute toxicity studies, rats were divided into four groups ($n = 5$; each) and were administered orally with three doses of asafoetida corresponding to 250, 500, and 1,000 mg/kg body weight. The control group received 1 ml of physiological saline. They were all placed under observation immediately for 24 h for any behavioral, neurological changes and then mortality for the next 72 h.^[18]

Chronic toxicity studies

The rats were divided into five groups ($n = 4$; each) and were administered four different doses of asafoetida corresponding to 25, 50, 100, and 200 mg/kg orally every

day for 6 weeks' duration. Another group of five rats that received 1 ml of physiological saline served as control.

Plant oleo-gum resin

Ferula assafoetida oleo-gum-resin was collected from Tabas region (Yazd province, Iran) during the summer and the plant species was botanically identified by Dr. Abbas Zarezadeh in Yazd Agricultural Research Center. The dried powder of asafoetida was soaked in distilled water overnight at room temperature and the yielded suspension was used orally. Concentrations and dosages of the extract were expressed as crude amount of the dried oleo-gum-resin used in preparing the stock solution.

Biochemical analysis

Blood was collected from orbital sinus of rats. Serum was prepared by centrifugation (3,000 rpm, 20 min) and stored frozen until biochemical assay. The urea, creatinine, lactate dehydrogenase (LDH), and aspartate aminotransferase (AST) were determined using the suitable kits and according to the manufacturer's instructions.

Hematological tests

The blood with ethylenediaminetetraacetic acid (EDTA) was used for the counting of RBC and total count of WBC and platelet by standard procedures. Hematocrit percentage (HCT %) was also determined.^[19]

Histopathology evaluation

For histopathological studies, liver and kidney tissue were collected immediately in 10% neutral buffered formalin. The tissues were processed by routine histological methods and embedded in paraffin blocks. The sections were cut by a rotary microtome and subsequently stained with Ehrlich's hematoxylin and eosin (H and E). The histological sections were studied under Olympus light microscopy (Olympus, Japan, magnification $\times 40$) to evaluate histopathological changes.

Statistical analysis

Statistical data were assessed with one-way analysis of variance (ANOVA), followed by *post hoc* Tukey's test using Graph pad prism version 5. Results were expressed as mean \pm standard error of the mean (SEM). A value of $P < 0.05$ was considered significant.

RESULTS

Acute toxicity studies

After the administration of different doses of asafoetida, the rats were monitored for the signs of toxicity. The animals

showed normal activity and no mortality were observed till the end of the study.

Biochemical factors

The result of serum kidney and liver parameters analysis in chronic study was shown in Table 1. There is no significant difference between the urea and creatinine concentration in control and treated animals with asafoetida. Our results also showed that AST and LDH significantly increased in all treated animals with asafoetida compared to control group ($P < 0.05$).

Hematological changes

Table 2 reveals that the asafoetida significantly decreased the concentrations of WBC, RBC, platelets, and HCT % after 6 weeks of treatment with asafoetida. The HCT % in asafoetida 25 group did not show any significant difference.

Histopathological findings of liver

The liver sections of rats in control group showed normal hepatic architecture and normal hepatocytes with prominent nucleus, central vein, and portal areas with no sign of inflammation or necrosis [Figure 1a]. Histopathological examination of extract treated rats showed little degenerative changes in some hepatic cells [Figure 1b]. Findings demonstrated that with increase in the extract doses (50, 100, and 200 mg/kg respectively), hepatocytes became larger in size with prominent nucleus compared to control group. Hypotrophied Kupffer cells

were also prominent with increase in extract doses and blood vessels expansion and dilated hepatic sinusoids were noticed [Figure 1c-e]. However, the general structures of lobules of liver are normal and clear.

Histopathological findings of kidney

Histological examination of the control group showed normal appearance and there were no changes in the renal tubular and glomerulus of kidneys [Figure 2a]. Sections of kidneys from extract groups (25, 50, 100, and 200 mg/kg) showed no signs of prominent pathological changes except occasional renal tubular necrosis [Figure 2b-d]. In some sections of extract groups, mild infiltration of inflammatory cells around the blood vessels and in the interstitial spaces was found. In extract group (200 mg/kg), some of the glomeruli appeared to be slightly expanded and there were little signs of tubular degenerative changes [Figure 2e].

DISCUSSION

The herbal drugs are playing an important role in healthcare programs throughout the world with a general belief that herbal drugs are always safe as these are natural. However,

Table 1: Effect of asafoetida on kidney and liver biochemical parameters in male rats

Groups	Urea	Creatinine	AST	LDH
Control	40.5±4.4	0.5±0.1	60.2±8.9	880.7±8.3
Asafoetida 25	46.4±6.2	0.4±0.1	95.4*±10.2	924.2*±8.4
Asafoetida 50	45.0±5.9	0.6±0.1	122.2*±12.3	1900.4*±17.5
Asafoetida 100	44.5±6.1	0.6±0.1	111.5*±10.8	1881.3*±16.7
Asafoetida 200	47.3±7.3	0.5±0.1	117.5*±12.3	1694.3*±14.9

*Represents significant increases and decreases, respectively, at $P < 0.05$ when compared to control values. Values are means±SEM. SEM = Standard error of the mean, AST = Aspartate transaminase, LDH = Lactate dehydrogenase

Table 2: Effect of oral consumption of asafoetida on hematological parameters in male rats

Groups	WBC ($\times 10^9/L$)	RBC ($\times 10^6/\mu L$)	HCT (%)	PLT ($\times 10^3/\mu L$)
Control	6.3±1.9	7.8±0.4	45.1±2.7	901±336
Asafoetida 25	4.8±1.9*	7.1±0.5*	41.3±3.1	622±178.8*
Asafoetida 50	4.5±1.1*	6.4±0.5*	38.2±2.9*	725.7±109.2*
Asafoetida 100	5.5±1.5*	6.6±0.2*	38.8±1.5*	893.6±85.9*
Asafoetida 200	4±1.3*	6.8±0.3*	39.8±1.2*	787.3±107.6*

*Represent significant increases and decreases, respectively, at $P < 0.05$ when compared to control values. Values are means±SEM. SEM = Standard error of the mean, WBC = White blood cell, RBC = Red blood cell, HCT = Hematocrit, PLT = Platelet




Figure 1: Histological architecture of the liver in different experimental groups. (a) No histopathological changes were seen in control group and the structures of lobules and hepatocyte of liver are normal. (b) Little degenerative changes of some hepatic cells (#) were seen. (c-e) In treatment groups (50, 100, and 200 mg/kg respectively) an increase in size of hepatocytes (arrows) were noticed. Hypotrophied Kupffer's cells (head arrow), dilated hepatic sinusoids (+) were also seen with increase in extract doses (H and E, $\times 400$). H and E = hematoxylin and eosin




Figure 2: (a) Paraffin section of kidney from control group showing normal histology. (b-d) Treatment groups (25, 50, and 100 mg/kg, respectively) showing no prominent histopathological changes except little renal tubular necrosis (#) and mild infiltrations of inflammatory cells in the interstitial spaces (arrow). (e) Showing mild degeneration of the cells lining the kidney tubules (+) and some of the slightly expanded glomeruli (head arrow) in treatment group at 200 mg/kg (H and E, $\times 400$)

adverse effects do occur which are usually mild and affect a small number of people. Asafoetida is used in different countries not only as a culinary spice but also traditionally used to treat various diseases. Although there are no experimental data about toxicity effects of asafoetida, in Iranian traditional, it was emphasized that intake of larger dosages of asafoetida can lead to swelling of the lips, digestive complaints such as flatulence and diarrhea, discomfort, and headache.^[3] The nontoxic nature of the asafoetida and its oral safety at selected dose level was confirmed as the administration of the drug does not cause mortality in rats. According to Clarke and Clarke,^[20] any compound or drug with the oral LD50 estimate greater than 1,000 mg/kg could be considered of low toxicity. This supports that asafoetida was found to be safe up to 1,000 mg/kg body weight in terms of mortality. In the present study, acute toxicity of asafoetida caused a significant increase in AST and LDH level in treated animals as compared to control group [Table 1]. There are many enzymes found in the serum that did not originally originate from the serum. During tissue damage, some of these enzymes find their way into the serum, probably by leakage.^[21] Serum enzyme measurements are therefore a valuable tool in clinical diagnosis, providing information on the effect and nature of pathological damage to any tissue. The increase in serum LDH and AST activity may indicate hepatic damage

probably by the altered cell membrane permeability leading to the leakage of the enzymes from the tissues to the serum.^[22,23] LDH and AST are considered to be sensitive indicators of hepatocellular damage and within limit can provide a quantitative evaluation of the degree of damage to the liver.^[24] In the present study, there was not any significant difference in urea and creatinine concentrations following treatment with asafoetida. Destruction of glomeruli causes significant decrease in the glomerular filtration rate (GFR) and increases the blood urea and creatinine resulting chronic renal failure.^[25] Since the urea and creatinine are markers of kidney function,^[26] the data suggest that asafoetida is not nephrotoxic. Hematological data obtained in our work indicated a significant decline in RBC and WBC and platelet count [Table 2]. Altered RBC, WBC, and platelet counts reflect that asafoetida would have some mild effect on the hematological parameter. Since blood cells are originated from bone marrow, the inertness of the extract on this organ is evident and its use in pharmacological evaluation is again certified. Histopathological observations revealed that treatment with asafoetida resulted little degenerative changes of some hepatic cells. Sections of kidneys from extract groups (25, 50, 100, and 200 mg/kg) showed no signs of prominent pathological changes in the cortex and the medulla, but little renal tubular necrosis was seen. Our finding also demonstrated that these changes were dose-dependent, respectively, and these histopathological evidences were in agreement with biochemical results.

CONCLUSION

The data showed that asafoetida did not have any acute toxicity, but the chronic consumption of this oleo-gum-resin caused reverse effects on liver and blood parameters. This study suggested that asafoetida may be used in minimum possible doses.

Acknowledgment

This research was supported by the Foundation of Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran.

Financial support and sponsorship

This research was supported by a grant from Shahid Sadoughi University of Medical Sciences Research Council.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Iranshahy M, Iranshahi M. Traditional uses, phytochemistry and pharmacology of asafoetida (*Ferula assa-foetida* oleo-gum-resin) -a review. *J Ethnopharmacol* 2011;134:1-10.

2. Zargari A. Medicinal Plants. Tehran: Tehran University Publications; 1996. p. 592-602.
3. Emami A, Fasihi S, Mehregan I. Medicinal Plants. Tehran: Andisheh Avar Publications; 2010. 1:24-8.
4. Eigner D, Scholz D. The magic book of Gyani Dolma. Pharm Unserer Zeit 1990;19:141-52.
5. Bagheri S, Hejazian Sh, Dashti-R M. The Relaxant Effect of Seed's Essential Oil and Oleo-gum-resin of *Ferula assa-foetida* on Isolated Rat's Ileum. Ann Med Health Sci Res 2014;4:238-41.
6. Bagheri SM, Rezvani ME, Vahidi AR, Esmaili M. Anticonvulsant effect of *Ferula assa-foetida* oleo gum resin on chemical and amygdala-kindled rats. N Am J Med Sci 2014;6:408-12.
7. Bagheri SM, Dashti-R MH, Morshedi A. Antinociceptive effect of *Ferula assa-foetida* oleo-gum-resin in mice. Res Pharm Sci 2014;9:207-12.
8. Kelly KJ, Nue J, Camitta BM, Honig GR. Methemoglobinemia in an infant treated with the folk remedy glycerited asafoetida. Pediatrics 1984;73:717-9.
9. Eigner D, Scholz D. *Ferula asa-foetida* and *Curcuma longa* in traditional medical treatment and diet in Nepal. J Ethnopharmacol 1999;67:1-6.
10. Kumar P, Singh DK. Molluscicidal activity of *Ferula asafoetida*, *Syzygium aromaticum* and *Carum carvi* and their active components against the snail *Lymnaea acuminata*. Chemosphere 2006;63:1568-74.
11. Ramadan NI, Abdel-Aaty HE, Abdel-Hameed DM, El Deeb HK, Samir NA, Mansy SS, et al. Effect of *Ferula assafoetida* on experimental murine *Schistosoma mansoni* infection. J Egypt Soc Parasitol 2004;34:1077-94.
12. Ramadan NI, Al Khadrawy FM. The *in vitro* effect of Assafoetida on *Trichomonas vaginalis*. J Egypt Soc Parasitol 2003;33:615-30.
13. Sitara U, Niaz I, Naseem J, Sultana N. Antifungal effect of essential oils on *in vitro* growth of pathogenic fungi. Pak J Bot 2008;40:409-14.
14. Garg DK, Banerjea AC, Verma J. The role of intestinal Clostridia and the effect of asafoetida (Hing) and alcohol in flatulence. Indian J Microbiol 1980;20:194-7.
15. Bagheri SM, Sahebkar A, Gohari AR, Saeidnia S, Malmir M, Iranshahi M. Evaluation of cytotoxicity and anticonvulsant activity of some Iranian medicinal *Ferula* species. Pharm Biol 2010;48:242-6.
16. Abraham SK, Kesavan PC. Genotoxicity of garlic, turmeric and asafetida in mice. Mutat Res 1984;136:85-8.
17. Walia K. Effect of asafoetida (7-hydroxycoumarin) on mouse spermatocytes. Cytologia (Tokyo) 1973;38:719-24.
18. Devaki K, Beulah U, Akila G, Gopalakrishnan VK. Effect of Aqueous Extract of *Passiflora edulis* on Biochemical and Hematological Parameters of Wistar Albino Rats. Toxicol Int 2012;19:63-7.
19. Sanderson JH. Philips: An atlas of laboratory animal haematology. New York: Oxford University Press; 1981; 1:150-55.
20. Clarke EG, Clarke ML. Landers veterinary toxicology in London. London: Bailliere Tindall; 1997; 2:456-61.
21. Will's DE. Biochemical basis of medicine. Bristol: John Wright and Sons Ltd; 1985;1:45-66.
22. Appidi JR, Yakubu MT, Grierson DS, Afolayan AJ. Toxicological evaluation of aqueous extracts of *Hermannia incana* Cav. Leaves in male wistar rats. Afr J Biotechnol 2009;8:2016-20.
23. Akanji MA, Yakubu MT. Alpha tocopherol protects against metabisulphite- induced tissue damage in rats. J Biochem Mol Biol 2000;15:179-83.
24. Al-Habori M, Al-Aghbari A, Al-Mamary M, Baker M. Toxicological evaluation of *Catha edulis* leaves: A long term feeding experiment in animals. J Ethnopharmacol 2002;83:209-17.
25. Ramakrishnan S, Swami R. Text Book of Clinical Biochemistry and Immunology. Chennai: T.R. Publications; 1995;3:245-75.
26. Odoula T, Adeniyi FA, Bello IS, Subair HG. Toxicity studies on an unripe *Carica papaya* aqueous extract: Biochemical and haematological effects in Wistar albino rats. J Med Plants Res 2007;1:1-4.