



Partial denture metal framework may harbor potentially pathogenic bacteria

Cristiane Machado Mengatto^{1*}, Leonardo Marchini², Luciano Angelo de Souza Bernardes³,
Sabrina Carvalho Gomes¹, Alecsandro Moura Silva⁴, Célia Marisa Rizzatti-Barbosa⁵

¹Department of Conservative Dentistry, Federal University of Rio Grande do Sul, School of Dentistry, Porto Alegre, Rio Grande do Sul, Brazil

²Department of Preventive and Community Dentistry, University of Iowa, College of Dentistry, Iowa City, Iowa, USA

³Department of Science and Technology, State University of Santa Cruz, School of Computer Science, Ilheus, Bahia, Brazil

⁴Department of Dental Materials and Prosthodontics, State University Julio de Mesquita Filho, School of Dentistry, Sao Jose dos Campos, Sao Paulo, Brazil

⁵Department of Prosthodontics and Periodontology, State University of Campinas, Piracicaba Dental School, Piracicaba, Sao Paulo, Brazil

PURPOSE. The aim of this study was to characterize and compare bacterial diversity on the removable partial denture (RPD) framework over time. **MATERIALS AND METHODS.** This descriptive pilot study included five women who were rehabilitated with free-end mandibular RPD. The biofilm on T-bar clasps were collected 1 week (t_1) and 4 months (t_2) after the RPD was inserted (t_0). Bacterial 16S rDNA was extracted and PCR amplified. Amplicons were cloned; clones were submitted to cycle sequencing, and sequences were compared with GenBank (98% similarity). **RESULTS.** A total of 180 sequences with more than 499 bp were obtained. Two phylogenetic trees with 84 (t_1) and 96 (t_2) clones represented the bacteria biofilm at the RPD. About 93% of the obtained phylotypes fell into 25 known species for t_1 and 17 for t_2 , which were grouped in 5 phyla: Firmicutes ($t_1=82\%$; $t_2=60\%$), Actinobacteria ($t_1=5\%$; $t_2=10\%$), Bacteroidetes ($t_1=2\%$; $t_2=6\%$), Proteobacteria ($t_1=10\%$; $t_2=15\%$) and Fusobacteria ($t_1=1\%$; $t_2=8\%$). The libraries also include 3 novel phylotypes for t_1 and 11 for t_2 . Library t_2 differs from t_1 ($P=.004$); t_1 is a subset of the t_2 ($P=.052$). Periodontal pathogens, such as *F. nucleatum*, were more prevalent in t_2 . **CONCLUSION.** The biofilm composition of the RPD metal clasps changed along time after RPD wearing. The RPD framework may act as a reservoir for potentially pathogenic bacteria and the RPD wearers may benefit from regular follow-up visits and strategies on prosthesis-related oral health instructions.

[J Adv Prosthodont 2015;7:468-74]

KEY WORDS: Framework; Denture plaque; Microorganisms; 16S rDNA gene; Removable partial denture

INTRODUCTION

There has been a longstanding controversy over the influ-

ence of wearing removable partial denture (RPD) on oral health. While many studies report that the RPD alters neither caries prevalence nor periodontal status in the long-term¹⁻⁴ others argue that there is a detrimental effect of RPD on abutment teeth.⁵⁻⁷ Furthermore, the literature agrees that dentures, both partial and complete, are considered reservoirs for various microorganism species, particularly opportunistic pathogens associated with systemic and oral diseases, such as caries, dentoalveolar abscesses, noma lesions, endocarditis, aphthous ulcers, denture stomatitis, and periodontal disease.⁸⁻¹⁸ For example, caries-associated bacteria, such as *Streptococcus mutans*^{19,20} and *Bifidobacteria spp.*¹⁴, and periodontal pathogens^{11,17,21} were identified in the complete denture plaque and oral cavity of patients even after the extraction of all teeth. However, no study has ever investigated the microorganisms adhering to the surface of the RPD framework. Thus, the microbial species adhering to

Corresponding author:
Cristiane Machado Mengatto
Department of Conservative Dentistry, School of Dentistry, Federal University of Rio Grande do Sul, Rua Ramiro Barcelos #2492, Porto Alegre, Rio Grande do Sul, Zipcode 90035-003, Brazil
Tel. 55 51 9991 4176; e-mail, cristiane.mengatto@ufrgs.br
Received June 5, 2015 / Last Revision September 15, 2015 / Accepted October 20, 2015

© 2015 The Korean Academy of Prosthodontics
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

This research was supported by the The State of Sao Paulo Research Foundation, Brazil (Grant# 01058-3).

the inner surface of the cobalt-chromium alloy clasp remain unknown.

Culture-dependent efforts were made to analyze changes in oral bacterial composition of denture wearers.^{5,6,20-23} However, these studies have been limited by the difficulties of *in vitro* growth techniques. Advances in molecular techniques have allowed the identification of many species of oral microorganisms, including those that cannot be cultivated.^{8,10,15,16,24,25} A recent *in vitro* study showed that potentially pathogenic microorganisms from human saliva adhere more largely at framework alloys than at acrylic resin surface and that cobalt-chromium alloy has a unique pattern of microorganism biofilm formation compared to other investigated surfaces.²⁶ Other authors evaluated the microbial diversity of supra and subgingival plaque of RPD wearers, using denaturing gradient gel electrophoresis (DGGE), a culture-independent method.⁷ They observed that health-associated genera tended to decrease, whereas the disease-associated species including *Streptococcus mutans* tended to increase. Thus, it is possible that the oral environment may be altered by RPD treatments.

To the best of our knowledge, no study determined bacterial diversity on RPD framework *in vivo* through culture-independent methods. Therefore, the objectives of this preliminary descriptive study are 1) to characterize bacterial community adhered to the RPD clasps and 2) identify changes in bacterial diversity at two time points: 1 week and 4 months after RPD placement. Our hypothesis is that the bacterial diversity on the RPD framework would change over time.

MATERIALS AND METHODS

The dental records of 130 partially dentate patients referred to the Piracicaba School of Dentistry, State University of Campinas, Sao Paulo, Brazil, were assessed. Individuals were eligible if they were completely edentulous at the maxilla and used previous complete denture and partially edentate at the mandible with no previous RPD (n=54). Twenty eight individuals did not respond to the telephone contact. The remaining (n=26) were clinically examined and

included if they were systemically healthy, without periodontitis (clinical attachment loss no more than 3 mm with no bleeding on probing), no smokers, with no presence of restorations, active caries, and crowns, no need of pre-prosthetic surgery, no prescription of systemic antibiotics or chemical plaque control in the last six months before the study commencement. Only patients without mandibular posterior teeth and at least 6 mandibular teeth were to be included in order to standardize RPD framework design.

A total of five women (60 to 74 years, mean age 67.6) composed the sample. The study was approved by the institutional ethical review board and the subjects signed informed consent.

During a pre-experimental period (approximately 45 days before RPD delivery) supragingival scaling and polishing were performed and oral hygiene instructions (OHI) were given. The OHI included how to properly floss and how to brush the RPD and the teeth with soft toothbrush and specific toothpaste (Colgate Triple Action, Colgate Palmolive Industry Com., Sao Paulo, Brazil). Subjects were instructed to remove the dentures for cleaning after each meal and before sleeping. The Fig. 1 shows the fluxogram of the study.

The clinical procedures to fabricate maxillary complete dentures and mandibular RPDs were performed by a single professional and followed a strict protocol for the production and placement of the dentures.²⁷ The cobalt-chromium RPD design was standardized and fabricated by the same dental technician. The clasps used were T-bar (Roach) retainers for direct abutment teeth (canines). The clasps were about 1 mm distant from the gingival edge. The denture bases and/or retainers were connected by a lingual plate that was seated on cingulum rests.

At t_0 (day 0, baseline), the new maxillary complete denture and mandibular RPD were delivered to patients. Oral hygiene was checked and instructions were repeated at the delivery appointment. Patients were oriented to abstain from any form of chemical control during the study. They also received reinstructions on how to perform tooth brushing as well as denture cleaning.

Biofilm was sampled after 1 week (t_1) and 4 months (t_2)




Fig. 1. Timeline of the study after enrolment process, interventions, and assessments performed on participants. (RPD) removable partial denture; (OHI) oral hygiene instructions.

of RPD delivery. All subjects consumed no food or beverages 30 minutes prior to the biofilm collection. First, the RPD was removed from the patient's mouth, washed with saline solution for 10 seconds, and left dry on paper towel for 1 minute. Then, the biofilm adhered on the inner surface of the T-bar clasps was collected with sterile curettes, pooled in a test tube containing 100 μ L of DNA extraction solution, and stored at -20°C. rDNA extraction and ethanol precipitation steps followed a previously described protocol.¹⁵ Concentrated rDNA was suspended in 20 μ L 10 mM Tris-HCl pH 7.5, 1 mM EDTA solution and PCR-amplified by using universally conserved primers D88 and E94.16 Briefly, 1 μ L of each sample was added to 49 μ L of reaction mixture containing 5 μ L 10 \times PCR Buffer (Promega Corporation, Madison, WI, USA), 1.25 unit Taq DNA Polymerase (Promega Corporation), 20 pmol of each primer, and 0.2 mM of each deoxyribonucleotides. PCR amplification was performed in a DNA thermal cycler (PTC-100, MJ Research, Watertown, MA, USA), as reported previously.¹⁵ The PCR products were purified by using spin columns (Wizard Clean-up System, Promega Corporation).

A total of 3 μ L of the purified PCR products were cloned by using the plasmid vector (TOPO TA Cloning[®] Kit for Sequencing, Invitrogen Life Technologies, Carlsbad, CA). Transformation used chemically competent cells (One Shot[®] TOP10 Chemically Competent *E. coli*, Invitrogen Life Technologies). The next steps for cloning and plasmid DNA purification followed previously described procedures.¹⁵

Purified plasmid DNA was sequenced using primer D88.¹⁶ Sequencing reactions were performed in a DNA thermal cycler (MJ Research PTC-100, MJ Research, Inc., Waltham, MA, USA) using DYEnamic ET Terminator Cycle Sequencing Kit (Amersham Biosciences Corp., Piscataway, NJ, USA). Once chain termination was complete, DNA sequencing was carried out in an ABI Prism 3100 Genetic Analyzer (Life Technologies, Carlsbad CA, USA). The sequencing protocol was 95°C, for 30 seconds, followed by 40 cycles of 95°C, 20 seconds; 50°C, 15 seconds, and 60°C, 2 minutes; with a final hold step at 4°C. The resultant 250 sequences presenting 499 or more nucleotides of good quality, which was ascertained by inspecting the chromatograms, were pooled and grouped into two 16S rDNA clone libraries, named t_1 and t_2 . Chimeric sequences were identified using Bellerophon (<http://foo.maths.uq.edu.au/huber/bellerophon.pl>) and discarded. A total of 180 reminiscent sequences, including 84 from t_1 and 96 from t_2 , were compared with sequences of known phylotypes from GenBank (<http://www.ncbi.nlm.nih.gov>) using the BLASTn algorithm. Sequence similarity of 98% or greater was used as the definition of a species-level cluster. Clones that were less than 98% similar to the closest known organisms were supposed to represent novel phylotypes.¹⁶ Contiguous sequences were assembled with the Phred/Phrap/Consed software package (<http://www.phrap.org>), aligned with the CLUSTAL W software. Phylogenetic trees were constructed with MEGA 5 software, according to the calculation of a Jukes and Cantor distance matrix using the neighbor-join-

ing method. Bootstrap confidence values for branching nodes were inferred by generating 1,000 resampling trees. The percentage of coverage was calculated by Good's method with the formula $[1 - (n / N)] \times 100$, where n is the number of phylotypes in a sample represented by one clone (singletons) and N is the total number of sequences in that sample. The computer program J-LIBSHUFF (<http://whitman.myweb.uga.edu/libshuff.html>) was used to statistically compare the clone libraries between time points ($P \leq .05$). Chao²⁸ and ACE²⁹ (Abundance-based Coverage Estimator) statistics were calculated using MOTHUR (www.mothur.org). The Chao and ACE estimators are statistical approaches used to estimate the species diversity and richness of microbial communities. Chao calculation considers the number of observed species, the number of singletons (species captured once), and the number of doubletons (species captured twice).²⁸ The observed species (Sobs) represent the microorganisms collected and identified. The ACE calculation considers the data from all species with fewer than 10 individuals, rather than just singletons and doubletons.²⁹ Bacteria diversity was considered the dependent variable and time was considered independent variable. The sequences representing novel phylotypes were deposited at GenBank under the following accession numbers KF715649 to KF715662.

RESULTS

Phylogenetic trees are shown in Fig. 2 and Fig. 3 and reveal 25 known phylotypes for t_1 and 17 known phylotypes for t_2 . The library t_1 also included 3 novel phylotypes while t_2 had 11 novel phylotypes. The phylotypes accounted for 86% of t_1 and 80% of t_2 cloned sequences, according to Good's population coverage.

The J-Libshuff analysis indicated that library t_1 is contained in t_2 ($P = .05$), but t_2 differed from t_1 ($P = .004$). So, community structure of the first library can be considered a subset of the second. When OTU (operational taxonomic unit) definition of 0.02 is considered, the t_2 shared approximately 40% of its membership (11 phylotypes) with those of t_1 , while 17 phylotypes were exclusive of t_1 and 17 were unique to t_2 .

The sequences were assigned to five phyla: Firmicutes (82% of t_1 and 60% of t_2), Actinobacteria (5% of t_1 ; 10% of t_2), Bacteroidetes (2% of t_1 ; 6% of t_2), Proteobacteria (10% of t_1 ; 15% of t_2) and Fusobacteria (1% of t_1 ; 8% of t_2) (Fig. 2 and Fig. 3). The major phyla were different between the libraries: the Firmicutes were decreased, while the Proteobacteria as well as other minor phyla were increased in t_2 . Sequences with more than 98% homology to Genbank represented 13 genera in t_1 and 9 genera in t_2 . Seven genera were shared between t_1 and t_2 libraries (Veillonella, Streptococcus, Capnocytophaga, Fusobacterium, Haemophilus, Neisseria, and Lautropia). The Collector's curves, estimated by Chao and ACE, represent that the species richness of the two libraries differs; it was highest in t_1 but lower in t_2 (Fig. 4).




Fig. 2. Phylogenetic tree with the highest-scored BLAST search results from 84 clones of library t_1 . The right side of the Fig. shows grayscale bars that represent the distribution of phylotypes among 5 phyla. The scale bar represents evolutionary distance (5% nucleotide sequence divergence). The organism names in bold letters mark the new phylotypes identified in the project. Final codes correspond to GenBank accession numbers.




Fig. 3. Phylogenetic tree with the highest-scored BLAST search results from 96 clones of library t_2 . The right side of the Fig. shows grayscale bars that represent the distribution of phylotypes among 5 phyla. The scale bar represents evolutionary distance (5% nucleotide sequence divergence). The organism names in bold letters mark the new phylotypes identified in the project. Final codes correspond to GenBank accession numbers.




Fig. 4. Collector's curves of observed (Sobs) and estimated (Chao and ACE) phylotype richness as a function of the number of clones recovered from libraries t_1 and t_2 . The number of unseen phylotypes is represented by the gap between the observed and estimated phylotypes. After the sampling of about 55 clones for t_1 and about 70 clones for t_2 , the gap between the observed and estimated phylotype richness was relatively constant, indicating repeated sampling of same phylotypes within samples. The increased sampling effort for t_1 would have yielded more phylotypes. In contrast, the horizontal shape of the curves in t_2 library indicates a trend of diminishing likelihood of finding new phylotypes as sampling continues.

DISCUSSION

The present study revealed that the RPD may act as a reservoir of bacteria species, many of them putative pathogens mainly associated with periodontal disease and caries.

The literature has posed that RPD may be associated with increased risk of dental caries and periodontal disease.^{5-7,19,20} However, the bacterial pathogens of such association are not clearly described. Even though more than 700 bacterial species have been identified in distinct oral habitats such as saliva, supra and subgingival biofilms, tongue, hard and soft palate in healthy or diseased conditions,^{8,10,15,16,24,25} few studies have aimed at evaluating the bacterial composition and shift related to RPD use. To the best of our knowledge, this pilot study is the first to attend this subject by means of culture-independent molecular methods. The advantages of such methods are highlighted at the literature and the present results illustrate them. Additionally, this is the first clinical study which described the diversity of biofilms adhered to the Co-Cr framework in contact to the enamel tooth surface. Most previous studies about the accumulation of biofilm in prosthesis wearers have based their observations on the acrylic denture base biofilm presence and/or composition.^{7,14,20,30} However, it is considered that the bacterial species adhered to the Co-Cr surfaces may differ from that of acrylic resin surfaces.²⁶ It was shown not only several species colonizing the RPD clasps but also an expressive shift of microbial species over time. The libraries population contains a small number of dominant species (*Streptococcus sanguinis* and *Veillonella parvula*) but presents a large number of unique phylotypes, i.e., a high degree of richness. Streptococcus was the most abundant genus in both libraries, comprising about a half of the clones ($t_1 =$

56%; $t_2 = 49\%$). Other studies showed that 63 to 86% of the initial colonizing bacteria in dental plaque were Streptococcus along with some Veillonella and Actinomyces.³¹ The present study did not find Actinomyces genus in t_1 , but it was present in t_2 .

Most Streptococcus of the t_1 fell in 3 species (*S. sanguinis*, *S. mitis*, and *S. oralis*). The second library (t_2) had more varied species of Streptococcus (*S. sanguinis*, *S. oralis*, *S. mitis*, *S. gordonii*, *S. infantis*, and *S. pneumonia*) some of which may be considered opportunistic pathogens. Previous studies reported that certain Streptococcus species, including *S. sanguinis*, *S. mitis*, *S. oralis*, and *S. gordonii* act as primary colonizers in bacterial adhesion to tooth surfaces.^{8,14} Initial adhesion involving Streptococcus and surfaces with similar hydrophobic properties would be facilitated on metal alloys, particularly for small coccus (1 μm), like *S. oralis*.²² Actinomyces, which presents larger cells (2 to 3 μm) compared to Streptococcus, need greater surface defects for retention or depend on Streptococcus first colonizers when the metal surface is polished.³² No sequence representative of *S. mutans* was found. The present results agree with previous literature, reporting that *S. mutans*, shown to be less hydrophobic than *S. sanguinis* and *S. oralis*, is less likely to adhere to alloy surfaces.²²

Rarefaction curves show that the t_1 library represents a small part of the overall population and t_2 library seems to be a development of t_1 community. The different genus found on the present study indicates a complex microbial community on the metal clasp surface, formed by bacteria of Actinomyces, Yellow, Green, Purple, and Orange periodontal complexes.³³ Moreover, the reduction of cocci over time (56% vs. 46%) may be indicative of a more mature plaque in t_2 . In t_2 , bacterial species diversity is also increased

compared to t_1 ($P = .004$). Fusobacteria filum was represented more in t_2 . Although *Fusobacterium spp.* have been identified in supragingival plaque of healthy individuals,²⁴ they have been typically associated with periodontal diseases.¹⁶ Denture plaque status was associated with the increased detection rate of *F. nucleatum* in edentulous patients wearing dentures.²¹ The present results demonstrated that Firmicutes were reduced, while Fusobacteria were increased in t_2 compared to t_1 , which suggests that the RPD clasps is a reservoir of bacteria that shift from non-maleficent to potentially-disease-related bacteria over time, as reported previously.⁷ Whether oral and abutment tooth conditions change from health to periodontal disease will require further investigation.

Many clones were considered novel phylotypes. In t_1 , two sequences exhibited low similarity to *S. sanguinis* (93.3% and 96.8%) and one clone had *Capnocytophaga ochracea* as closest relative (96.8%). In t_2 , 4 sequences displayed very low identity with *Leptotrichia goodfellowii* (89.6%, 90.2%, 92.1%, and 91.1%), commonly isolated from healthy oral cavities.²⁴ Two novel phylotypes had 91.0% and 91.6% identity with *Corynebacterium casei*. Although *Corynebacterium* can be typically isolated from dental plaque, it is not associated with oral diseases. Other novel phylotypes in t_2 exhibited 96.3% similarity to *Eikenella corrodens* and less than 91.5% identity to *Catonella morbi*. *C. morbi* resides in the oral cavity and is associated with primary endodontic infections and endocarditis.³⁴ One clone of t_2 had 96.4% identity to *Selenomonas sputigena*, which is typically found in the oral cavity in congregation to *F. nucleatum* and may be associated with aggressive periodontitis.³⁵ Other novel phylotype had 95.7% identity to its closest relative *Actynomices odontolyticus*, which has been isolated from deep dental caries and is an opportunistic pathogen of immunocompromised patients, involved in bacteremia and pneumonia.³⁶ Obviously, the physiological and clinical relevance of the novel phylotypes detected in this study remain unclear. At one extreme, some of the phylotypes may be transient residents. On the other extreme, these organisms may be critical for maintenance of ecosystem stability and oral health or may be occasional pathogens.

Although 16S rDNA sequencing represents a broad-range analysis, data should be interpreted with caution. A different definition for OTU could provide different results. Additionally, it cannot be inferred that bacteria that have not been detected are not present on the clasp surface. Note that some species may be present but below the limit of detection, and thus may be poorly represented in the clone libraries.

In the present study oral hygiene instructions (OHI) were given up to day zero. In this sense, it is conceivable that the absence of OHI between t_1 and t_2 might have resulted not only in biofilm growth and maturation *per se*, observed herein, but also in an inflammatory response, even subclinical, that, in turn, also can influence the biofilm composition. This retro-filling process is described at the literature.^{37,38} Bearing this two-way influence in mind, the

results herein show that RPD wearers retain mature biofilm at clasps favoring the maintenance of an environment adequate to the maturation of the biofilm as a whole and the long-term impact to the gingival status. On the other hand, it should be noted that the present study established careful oral hygiene instructions and training for subjects up to t_1 . This situation simulates what usually occurs in the dental office, where patients receive oral hygiene instructions during RPD fabrication and adjustment appointments, but the next follow-up appointment takes a long time to be scheduled. The result of this delay is that dentures are used for longer time than they are supposed to be, and the patients lose the ability to perform plaque control.⁶ Previous studies reported that a maintenance interval longer than 6 months was significant predictor for positive red complex bacteria in RPD wearers.²³ The results of the present study demonstrate that, after 4 months, a complex and mature biofilm is formed on RPD clasp surfaces. The establishment of more frequent follow-up appointments for RPD wearers may be a valuable strategy to reduce potential pathogen accumulation on RPD clasps, possibly favoring the gingival condition along time.

CONCLUSION

This study concluded that the RPD clasps harbors a highly diverse bacterial population and the bacterial community on the RPD metal clasp develops into a complex mature community, including putative periodontal and opportunistic pathogens.

ORCID

Cristiane Machado Mengatto <http://orcid.org/0000-0002-1059-8419>

Leonardo Marchini <http://orcid.org/0000-0003-1291-6684>

Célia Marisa Rizzatti-Barbosa <http://orcid.org/0000-0002-8747-0034>

ACKNOWLEDGEMENTS

The authors thank Professor Francisco G. Nobrega for advice and access to laboratory facilities and supplies. The authors are also grateful to the Center for Computational Biotechnology and Biotechnological Information Management (NBCGIB) of the State University of Santa Cruz (UESC) for computational support.

REFERENCES

1. Bergman B, Hugoson A, Olsson CO. A 25 year longitudinal study of patients treated with removable partial dentures. *J Oral Rehabil* 1995;22:595-9.
2. Mullally BH, Linden GJ. Periodontal status of regular dental attenders with and without removable partial dentures. *Eur J Prosthodont Restor Dent* 1994;2:161-3.
3. Schwalm CA, Smith DE, Erickson JD. A clinical study of pa-

- tients 1 to 2 years after placement of removable partial dentures. *J Prosthet Dent* 1977;38:380-91.
4. Zlatarić DK, Celebić A, Valentić-Peruzović M. The effect of removable partial dentures on periodontal health of abutment and non-abutment teeth. *J Periodontol* 2002;73:137-44.
 5. do Amaral BA, Barreto AO, Gomes Seabra E, Roncalli AG, da Fonte Porto Carreiro A, de Almeida EO. A clinical follow-up study of the periodontal conditions of RPD abutment and non-abutment teeth. *J Oral Rehabil* 2010;37:545-52.
 6. Vanzeveren C, D'Hoore W, Bercy P. Influence of removable partial denture on periodontal indices and microbiological status. *J Oral Rehabil* 2002;29:232-9.
 7. Zhu X, Wang S, Gu Y, Li X, Yan H, Yan H, Miyoshi S, Shi L. Possible variation of the human oral bacterial community after wearing removable partial dentures by DGGE. *World J Microbiol Biotechnol* 2012;28:2229-36.
 8. Campos MS, Marchini L, Bernardes LA, Paulino LC, Nobrega FG. Biofilm microbial communities of denture stomatitis. *Oral Microbiol Immunol* 2008;23:419-24.
 9. Danser MM, van Winkelhoff AJ, de Graaff J, van der Velden U. Putative periodontal pathogens colonizing oral mucous membranes in denture-wearing subjects with a past history of periodontitis. *J Clin Periodontol* 1995;22:854-9.
 10. Diaz PI, Chalmers NI, Rickard AH, Kong C, Milburn CL, Palmer RJ Jr, Kolenbrander PE. Molecular characterization of subject-specific oral microflora during initial colonization of enamel. *Appl Environ Microbiol* 2006;72:2837-48.
 11. Fernandes CB, Aquino DR, Franco GC, Cortelli SC, Costa FO, Cortelli JR. Do elderly edentulous patients with a history of periodontitis harbor periodontal pathogens? *Tex Dent J* 2012;129:751-61.
 12. Hutter G, Schlagenhauf U, Valenza G, Horn M, Burgemeister S, Claus H, Vogel U. Molecular analysis of bacteria in periodontitis: evaluation of clone libraries, novel phylotypes and putative pathogens. *Microbiology* 2003;149:67-75.
 13. Ledder RG, Gilbert P, Huws SA, Aarons L, Ashley MP, Hull PS, McBain AJ. Molecular analysis of the subgingival microbiota in health and disease. *Appl Environ Microbiol* 2007;73:516-23.
 14. Mantzourani M, Gilbert SC, Fenlon M, Beighton D. Non-oral bifidobacteria and the aciduric microbiota of the denture plaque biofilm. *Mol Oral Microbiol* 2010;25:190-9.
 15. Marchini L, Campos MS, Silva AM, Paulino LC, Nobrega FG. Bacterial diversity in aphthous ulcers. *Oral Microbiol Immunol* 2007;22:225-31.
 16. Paster BJ, Boches SK, Galvin JL, Ericson RE, Lau CN, Levanos VA, Sahasrabudhe A, Dewhirst FE. Bacterial diversity in human subgingival plaque. *J Bacteriol* 2001;183:3770-83.
 17. Sachdeo A, Haffajee AD, Socransky SS. Biofilms in the edentulous oral cavity. *J Prosthodont* 2008;17:348-56.
 18. Wade WG, Spratt DA, Dymock D, Weightman AJ. Molecular detection of novel anaerobic species in dentoalveolar abscesses. *Clin Infect Dis* 1997;25:S235-6.
 19. Mihalow DM, Tinanoff N. The influence of removable partial dentures on the level of *Streptococcus mutans* in saliva. *J Prosthet Dent* 1988;59:49-51.
 20. Rocha EP, Francisco SB, Del Bel Cury AA, Cury JA. Longitudinal study of the influence of removable partial denture and chemical control on the levels of *Streptococcus mutans* in saliva. *J Oral Rehabil* 2003;30:131-8.
 21. Yasui M, Ryu M, Sakurai K, Ishihara K. Colonisation of the oral cavity by periodontopathic bacteria in complete denture wearers. *Gerodontology* 2012;29:e494-502.
 22. Grivet M, Morrier JJ, Benay G, Barsotti O. Effect of hydrophobicity on in vitro streptococcal adhesion to dental alloys. *J Mater Sci Mater Med* 2000;11:637-42.
 23. Mine K, Fueki K, Igarashi Y. Microbiological risk for periodontitis of abutment teeth in patients with removable partial dentures. *J Oral Rehabil* 2009;36:696-702.
 24. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol* 2005;43:5721-32.
 25. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, Lakshmanan A, Wade WG. The human oral microbiome. *J Bacteriol* 2010;192:5002-17.
 26. Urushibara Y, Ohshima T, Sato M, Hayashi Y, Hayakawa T, Maeda N, Ohkubo C. An analysis of the biofilms adhered to framework alloys using in vitro denture plaque models. *Dent Mater J* 2014;33:402-14.
 27. Carr AB, McGivney GP, Brown DT, McCracken WL. *McCracken's Removable Partial Prosthodontics*. 12 ed. St. Louis: Mosby; 2010.
 28. Chao A. Non-parametric estimation of the number of classes in a population. *Scand J Stat* 1984;11:265-70.
 29. Chao A, Lee SM. Estimating the number of classes via sample coverage. *J Am Stat Assoc* 1992;87:210-7.
 30. Marsh PD, Percival RS, Challacombe SJ. The influence of denture-wearing and age on the oral microflora. *J Dent Res* 1992;71:1374-81.
 31. Nyvad B, Kilian M. Microbiology of the early colonization of human enamel and root surfaces in vivo. *Scand J Dent Res* 1987;95:369-80.
 32. Taylor R, Maryan C, Verran J. Retention of oral microorganisms on cobalt-chromium alloy and dental acrylic resin with different surface finishes. *J Prosthet Dent* 1998;80:592-7.
 33. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL Jr. Microbial complexes in subgingival plaque. *J Clin Periodontol* 1998;25:134-44.
 34. Siqueira JF Jr, Rôças IN. *Catonella morbi* and *Granulicatella adiacens*: new species in endodontic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102:259-64.
 35. Gonçalves LF, Fermiano D, Feres M, Figueiredo LC, Teles FR, Mayer MP, Faveri M. Levels of *Selenomonas* species in generalized aggressive periodontitis. *J Periodontol* 2012;47:711-8.
 36. Cone LA, Leung MM, Hirschberg J. *Actinomyces odontolyticus* bacteremia. *Emerg Infect Dis* 2003;9:1629-32.
 37. Haffajee AD, Teles RP, Patel MR, Song X, Veiga N, Socransky SS. Factors affecting human supragingival biofilm composition. I. Plaque mass. *J Periodontol* 2009;44:511-9.
 38. Rüdiger SG, Carlén A, Meurman JH, Kari K, Olsson J. Dental biofilms at healthy and inflamed gingival margins. *J Clin Periodontol* 2002;29:524-30.