

Activation of integrin $\alpha 5$ mediated by flow requires its translocation to membrane lipid rafts in vascular endothelial cells

Xiaoli Sun^{a,b}, Yi Fu^a, Mingxia Gu^a, Lu Zhang^a, Dan Li^a, Hongliang Li^c, Shu Chien^{d,1}, John Y.-J. Shyy^{e,1}, and Yi Zhu^{a,f,1}

^aDepartment of Physiology and Pathophysiology, Peking University Health Science Center, Beijing 100191, China; ^bDepartment of Medicine, University of California, San Diego, La Jolla, CA 92093; ^cDepartment of Cardiology, Renmin Hospital of Wuhan University, Wuhan 430060, China; ^dDepartment of Bioengineering, University of California, San Diego, La Jolla, CA 92093; ^eCardiovascular Research Center, Xi'an Jiaotong University School of Medicine, Xi'an 710061, China; and ^fDepartment of Physiology and Pathophysiology, Tianjin Medical University, Tianjin 300070, China

Contributed by Shu Chien, December 14, 2015 (sent for review March 26, 2015; reviewed by Peter F. Davies and Hanjoong Jo)

Local flow patterns determine the uneven distribution of atherosclerotic lesions. Membrane lipid rafts and integrins are crucial for shear stress-regulated endothelial function. In this study, we investigate the role of lipid rafts and integrin $\alpha 5$ in regulating the inflammatory response in endothelial cells (ECs) under atheroprone versus atheroprotective flow. Lipid raft proteins were isolated from ECs exposed to oscillatory shear stress (OS) or pulsatile shear stress, and then analyzed by quantitative proteomics. Among 396 proteins redistributed in lipid rafts, integrin $\alpha 5$ was the most significantly elevated in lipid rafts under OS. In addition, OS increased the level of activated integrin $\alpha 5$ in lipid rafts through the regulation of membrane cholesterol and fluidity. Disruption of F-actin-based cytoskeleton and knockdown of caveolin-1 prevented the OS-induced integrin $\alpha 5$ translocation and activation. In vivo, integrin $\alpha 5$ activation and EC dysfunction were observed in the atheroprone areas of low-density lipoprotein receptor-deficient (*Ldlr*^{-/-}) mice, and knockdown of integrin $\alpha 5$ markedly attenuated EC dysfunction in partially ligated carotid arteries. Consistent with these findings, mice with haploinsufficiency of integrin $\alpha 5$ exhibited a reduction of atherosclerotic lesions in the regions under atheroprone flow. The present study has revealed an integrin- and membrane lipid raft-dependent mechanotransduction mechanism by which atheroprone flow causes endothelial dysfunction.

integrin | lipid rafts | shear stress | proteomics | endothelial dysfunction

Shear stress imposed on vascular endothelial cells (ECs) influences vascular phenotype and function. Atherosclerosis preferentially develops at branches and curvatures in the arterial tree where flow is disturbed. In contrast, pulsatile shear stress (PS) in the straight parts of the arteries is atheroprotective (1). At the cellular and molecular levels, disturbed flow pattern increases, while PS inhibits, the inflammatory response in ECs, including the expression of intercellular adhesion molecule 1 (ICAM-1), vascular adhesion molecule 1 (VCAM-1), and interleukin 1 β (IL-1 β) (2). We found that integrins in ECs are shear stress-sensitive (3). Many subsequent studies demonstrated that integrin activation is essential for transmitting mechanical stimuli to intracellular biochemical pathways (4–6). Additionally, mounting evidence indicates that lipid raft microdomains, membrane receptors, cytoskeletal proteins, and extracellular matrices are linked to integrin activation in the context of mechanotransduction (7–11). However, the key mechanism underlying the differential effects of atheroprone versus atheroprotective flows in activating integrins that in turn induces inflammatory response in ECs remains to be elucidated.

Lipid rafts are membrane microdomains that are enriched in cholesterol, sphingolipids, and a variety of signaling molecules, which function as cellular signaling platforms. These microdomains are more ordered and tightly packed than the surrounding membrane. Removal of membrane cholesterol leads to the dissociation of molecules from lipid rafts and hence deactivation of some of

these molecules (12). The F-actin-based cytoskeleton associates with lipid rafts, and many of the structural and functional properties of rafts require an intact F-actin-based cytoskeleton. As many of the lipid raft-associated proteins in ECs are shear stress-sensitive, changes in the cholesterol content of the membrane or disruption of its F-actin-based cytoskeleton abolish mechanosensitivity of the cell.

Both oxidized low-density lipoprotein (OxLDL) and atheroprone flow have been implicated in inducing nuclear factor- κ B (NF- κ B)-mediated inflammation in ECs. Integrin $\alpha 5\beta 1$ is mainly expressed in ECs (13) and is involved in flow activation of NF- κ B (14). With respect to lipid rafts, clustering of ganglioside-abundant lipid rafts has been shown to regulate the activity of integrin $\beta 1$ (15). Furthermore, a recent study by Yurdagul et al. shows that integrin $\alpha 5\beta 1$ is involved in OxLDL-induced activation of NF- κ B in ECs (16). Although it is firmly established that activation of $\alpha 5\beta 1$ mediates the NF- κ B inflammatory axis in ECs, it is unknown whether a common molecular basis exists between the ability of atheroprone flow and OxLDL to activate integrin $\alpha 5\beta 1$ and the subsequent molecular mechanisms by which $\alpha 5\beta 1$ increases the inflammatory response, leading to atherosclerosis susceptibility.

The current study was initiated from findings from a quantitative proteomics analysis of lipid raft-associated proteins in ECs regulated by flow. Surprisingly, this high-throughput screening revealed that oscillatory shear stress (OS) drastically increased integrin $\alpha 5$ in lipid rafts of ECs. Because OxLDL treatment of ECs increases their membrane cholesterol content leading to

Significance

This study reveals that atheroprone flow induces integrin $\alpha 5$ translocation into lipid rafts and hence activation to cause endothelial dysfunction in vitro and in vivo. Consequently, this mechanotransduction event leads to the activation of NLRP3 inflammasome. Knockdown of integrin $\alpha 5$ ameliorated endothelial cell (EC) dysfunction in partially ligated carotid arteries of *Ldlr*^{-/-} mice. Furthermore, haploinsufficiency or functional inhibition of integrin $\alpha 5$ in mice improved EC function in the atheroprone area. These findings reveal a novel mechanism by which atheroprone flow causes endothelial dysfunction that leads to vascular impairments such as atherosclerosis.

Author contributions: X.S., S.C., J.Y.-J.S., and Y.Z. designed research; X.S., Y.F., M.G., L.Z., and D.L. performed research; H.L. contributed new reagents/analytic tools; X.S. and Y.Z. analyzed data; and X.S., S.C., J.Y.-J.S., and Y.Z. wrote the paper.

Reviewers: P.F.D., University of Pennsylvania; and H.J., Emory University and Georgia Institute of Technology.

The authors declare no conflict of interest.

¹To whom correspondence may be addressed. Email: shuchien@ucsd.edu, john.shyy@mail.xjtu.edu.cn, or zhuyi@tmu.edu.cn.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1524523113/-DCSupplemental.

decreased membrane fluidity (17, 18), we hypothesized that alterations in cholesterol composition or lipid packaging in lipid rafts in response to OxLDL or atheroprone flow may alter integrin $\alpha 5$ distribution and/or activity. Furthermore, given that activation of NF- κ B in turn activates Nod-like receptor protein 3 (NLRP3) inflammasome (19), integrin $\alpha 5$ would be upstream of the atheroprone flow-induced NLRP3 inflammasome (20).

In this study, we investigate the roles of lipid rafts and integrin $\alpha 5$ in OS-induced EC dysfunction, including NLRP3 inflammasome and their causative effect on atherogenesis. Our in vitro and in vivo experiments demonstrate that the lipid raft translocation and activation of integrin $\alpha 5$, due to changes in membrane fluidity, is a previously unidentified pathway mediating endothelial dysfunction in response to atheroprone flow.

Results

Differential Effects of OS and PS on Lipid Raft Translocation and Activation of Integrin $\alpha 5$. We isolated total proteins from human umbilical vein ECs (HUVECs) that had been exposed to OS (0.5 ± 4 dyn/cm²) or PS (12 ± 4 dyn/cm²) for 2 h. The cell lysates were fractionated using sucrose density gradient ultracentrifugation to separate the lipid raft-associated proteins. Positive blotting of lipid raft markers including flotillin-2 (Flot2), caveolin-1 (Cav1), and cavin1 in fractions 4–5 (Fig. S1A, Top) validated that these fractions contained the lipid raft-associated proteins. Quantitative proteomic analysis was then performed, and 396 proteins were identified in fractions 4–5 (Datasets S1 and S2). We selected proteins that were significantly upregulated or down-regulated in an OS- or PS-dependent manner. Using the Panther Classification System (www.pantherdb.org) (Dataset S3), we categorized these proteins into different functional groups (binding, enzyme regulator, ion channel, motor, receptor, structural molecular, transcription regulator, translation regulator, and transporter). Of the proteins considered mechanosensitive and significantly differentially regulated, the level of integrin $\alpha 5$ was increased by OS, but decreased by PS, in lipid rafts (1.8- vs. 0.6-fold) (Dataset S3).

To confirm the results from proteomic analysis, ECs were exposed to OS or PS for increasing time points, and the distribution of integrin $\alpha 5$ in lipid rafts was examined. Under OS, the proportion of integrin $\alpha 5$ in lipid rafts increased in a time-dependent manner, up to threefold at 2 h. In contrast, PS had an opposite effect (Fig. S1A, Bottom). In support of this observation, we used equal amounts of protein purified from lipid rafts and nonlipid rafts under OS vs. PS to detect integrin $\alpha 5$ translocation. As expected, although OS induced integrin $\alpha 5$ translocation into lipid rafts, which became significant at 2 h, PS promoted the exit of integrin $\alpha 5$ from lipid rafts (Fig. 1A and B).

We next investigated whether OS regulates the activity of integrin $\alpha 5$. Using an antibody recognizing the activated integrin $\alpha 5$, we found that OS, but not PS, activated integrin $\alpha 5$ (Fig. 1C and D). This differential effect between OS and PS in activating integrin $\alpha 5$ persisted up to 12 h following flow application (Fig. S1B and C). In line with integrin $\alpha 5$ activation, OS enhanced EC adhesion on fibronectin, a specific ligand for integrin $\alpha 5$. By contrast, PS decreased such adhesion (Fig. S1D and E). To validate the in vitro observations, we investigated integrin $\alpha 5$ activation in different regions of the aorta of low-density lipoprotein receptor-deficient (*Ldlr*^{-/-}) mice on normal diet. *En face* immunostaining with the use of an antibody recognizing the activated integrin $\alpha 5$ showed enhanced staining in atheroprone areas [i.e., aortic arch (AA)] in comparison with the atheroprotective area [i.e., thoracic aorta (TA)] (Fig. 1E, Top). Noticeably, there was no difference of total integrin $\alpha 5$ expression between TA and AA (Fig. 1E, Bottom).

Lipid Raft Translocation of Integrin $\alpha 5$ Is Associated with Its Activation. Western blot analysis shown in Fig. 2A indeed verified that OS activated integrin $\alpha 5$ and that the activated integrin




Fig. 1. OS elevates the level of total and activated integrin $\alpha 5$ in lipid rafts in vitro and in vivo. HUVECs were exposed to OS (0.5 ± 4 dyn/cm²) (A and C) or PS (12 ± 4 dyn/cm²) (B and D) for time durations as indicated. Static cells (0 min) were used as a control. (A and B) Representative Western blots of integrin $\alpha 5$ in lipid rafts (Rafts; fractions 4–5), nonlipid rafts (Nonrafts; fractions 6–12), and whole cellular proteins (Total) in static cells or those exposed to OS or PS. (C and D) Lipid raft proteins were isolated for Western blot analysis of activated integrin $\alpha 5$ (Act- $\alpha 5$) and Flot2, a marker for rafts fractions. Bar graphs show the relative ratio of total or activated $\alpha 5$ to Flot2 or β -actin. ANOVA followed by the Bonferroni post hoc test was used for statistical analysis. Data are mean \pm SEM averaged from at least three independent experiments. * $P < 0.05$, rafts vs. static; # $P < 0.05$, nonrafts vs. static. (E) The AA and TA were isolated from *Ldlr*^{-/-} mice (8 wk old; $n = 10$) for *en face* immunostaining with Act- $\alpha 5$ (red), Total $\alpha 5$ ($\alpha 5$, red), Flot2 (green), and nuclear (blue).

$\alpha 5$ was mainly located in the lipid raft fractions. We then examined the causative effect of integrin $\alpha 5$ translocation and activation, namely, whether $\alpha 5$ translocation is necessary for its activation or vice versa. When we knocked down Cav1 to disrupt the structure of lipid rafts to prevent $\alpha 5$ translocation, OS could not activate $\alpha 5$ nor could it enhance EC adhesion to fibronectin (Fig. 2B and C), which indicates that $\alpha 5$ translocation instigates its activation by OS. However, when ECs were treated with an $\alpha 5$ neutralizing antibody ($\alpha 5$ Ab) or ATN161 (an $\alpha 5$ -specific inhibitory peptide) to prevent $\alpha 5$ activation, OS was still able to induce the $\alpha 5$ translocation into lipid rafts (Fig. 2D and E).




Fig. 2. The OS-activated integrin $\alpha 5$ is associated with lipid rafts translocation. (A) HUVECs were exposed to OS or kept static for 2 h, and proteins were isolated for Western blot analysis of activated $\alpha 5$ (Act- $\alpha 5$) in Rafts and Nonrafts. (B) HUVECs pretreated with scramble siRNA (siCav1, -) or Cav1 siRNA (siCav1, +) for 48 h. Lipid raft and EC proteins were isolated for Western blot analysis of Act- $\alpha 5$, Flot2, Cav1, and β -actin. Bar graphs show the relative ratio of Act- $\alpha 5$ to Flot2 or β -actin. (C) HUVECs pretreated with or without siCav1 were subjected to OS for 2 h. ECs adhesion to fibronectin was measured. (D and E) HUVECs were pretreated with integrin $\alpha 5$ neutralizing antibody (10 μ g/mL) for 1 h (D) or with ATN161 or control peptide (10 μ mol/L) for 48 h (E), and then exposed to OS for 2 h. Western blot was performed with antibodies as indicated. Bar graphs show the relative ratio of Raft $\alpha 5$ to Flot2, or Nonraft $\alpha 5$ to β -actin. ANOVA followed by the Bonferroni post hoc test was used for statistical analysis. Data are mean \pm SEM averaged from at least three independent experiments. * P < 0.05, vs. static.

Collectively, results in Fig. 2 suggest that the Cav1-dependent integrin $\alpha 5$ translocation into lipid rafts is necessary for its activation by OS but not vice versa.

Integrin $\alpha 5$ Activity Is Regulated by Membrane Fluidity and Cholesterol Content. As membrane fluidity affects the mobility of membrane proteins, and because we previously reported that shear stress regulates cholesterol content of the EC membrane, in turn altering membrane fluidity (21), we next investigated the involvement of membrane fluidity in flow regulation of integrin $\alpha 5$ activity. We found that OS increased 1,6-diphenyl-1,3,5-hexatriene (DPH) anisotropy, which is indicative of a decrease in membrane fluidity (Fig. 3A). Such change was consistent with the increase in cholesterol content in the membrane (Fig. 3C). These OS-imposed effects were not seen in ECs pretreated with methyl- β -cyclodextrin (β -CD) to deplete cholesterol (Fig. 3A and C). In contrast, PS increased membrane fluidity and attenuated cholesterol content, which were abolished if EC cultures were supplemented with cholesterol (Fig. 3B and D). Importantly, cholesterol depletion with β -CD prevented $\alpha 5$ activation by OS (Fig. 3E), whereas cholesterol supplementation increased the level of activated $\alpha 5$ under PS (Fig. 3F). These findings suggest that membrane fluidity affected by cholesterol content is crucial for OS activation of integrin $\alpha 5$.

Integrin $\alpha 5$ Translocation Is Mediated by F-Actin-Based Cytoskeleton Rearrangements. Given the important role of the F-actin-based cytoskeleton rearrangements in protein translocation from the cytoplasm to lipid rafts (i.e., caveolae) (21, 22), we further investigated whether the OS-induced integrin $\alpha 5$ translocation is dependent on F-actin-based cytoskeleton. Disruption of the F-actin-based cytoskeleton by cytochalasin B (Cyt-B) markedly attenuated the OS-induced integrin $\alpha 5$ translocation and activation (Fig. S2).

OS-Induced EC Dysfunction Is Integrin $\alpha 5$ -Dependent. Integrin $\alpha 5$ activates the NLRP3 inflammasome in macrophages (23), and OS has been shown to induce NLRP3 inflammasome in ECs (20). Because OxLDL-induced NF- κ B is $\alpha 5$ -dependent (16) and NF- κ B activation primes the induction of NLRP3 inflammasome (19), we reasoned that $\alpha 5$ activation is necessary for OS-induced inflammasome activation. Indeed, the OS-induced cleavage of procaspase1 and pro-IL-1 β , hallmarks of NLRP3 inflammasome induction, was reduced by $\alpha 5$ neutralizing antibody in a dose-dependent manner (Fig. 4A). A similar inhibitory effect was seen in ECs in which Cav1 was knocked down (Fig. 4B). Moreover, there was a decrease in OS activation of NLRP3 inflammasome in mouse embryonic fibroblasts (MEFs) isolated from $\alpha 5$ -null embryos (Fig. 4C). Downstream of the OS-induced NLRP3 inflammasome is the enhanced expression of the chemoattractant and adhesion molecules such as VCAM-1 and ICAM-1. The OS-augmented VCAM-1 and ICAM-1 expression was attenuated by either $\alpha 5$ neutralizing antibody or β -CD treatment (Fig. 4D and E). Additionally, the enhanced binding of monocytes to EC




Fig. 3. OS promotes integrin $\alpha 5$ translocation to lipid rafts through alteration of membrane fluidity. HUVECs were treated with β -CD (5 mmol/L) for 2 h (A, C, and E) or Chl (30 μ g/mL) for 2 h (B, D, and F) before being exposed to OS or PS or kept under static condition for 2 h. Cell membrane was then isolated for fluorescence anisotropy (A and B) or cholesterol content (C and D) measurements. (E and F) Lipid raft proteins were isolated for Western blot analysis of Act- $\alpha 5$ and Flot2. ANOVA followed by the Bonferroni post hoc test was used for statistical analysis. Data are mean \pm SEM averaged from at least three independent experiments. * P < 0.05, vs. static controls.




Fig. 4. OS-activated integrin $\alpha 5$ leads to endothelial dysfunction. ECs were pretreated with integrin $\alpha 5$ neutralizing antibody (5 μ g/mL or 10 μ g/mL) for 1 h (A, D, and F), scramble siRNA or Cav1 siRNA (40 nmol/L) for 48 h (B), β -CD (5 mmol/L) for 2 h (E), or caspase-1 inhibitor Z-YVAD-FMK (2 μ mol/L) for 24 h (G); $\alpha 5^{-/-}$ MEFs or littermate wild-type MEFs ($\alpha 5^{+/+}$) were used in C. All cell groups were kept under static condition or exposed to PS or OS for 12 h. In A–E, Western blots were performed for activated integrin $\alpha 5$ (Act- $\alpha 5$), pro-caspase1 (pro-Casp1), caspase1 p20 (Casp1 p20), pro-IL-1 β , IL-1 β p17, ICAM-1, and VCAM-1. Lipid raft fractions (Rafts) were used to detect Act- $\alpha 5$, whereas total cell lysates were used for all other proteins. Bar graphs showed the relative ratio of casp1 p20 and IL-1 β p17 to β -actin. Data are mean \pm SEM normalized to the static control from at least three independent experiments. ANOVA followed by the Bonferroni post hoc test was used. * P < 0.05, vs. static or PS. (F and G). The BCECF-AM-labeled THP1 cells were added to EC monolayer and incubated for 30 min. Fluorescence microscopy was used to assess the attached THP1 cells. The ratio of adherent THP1 cells to ECs was measured. ANOVA followed by the Bonferroni post hoc test was used for statistical analysis. Data are mean \pm SEM averaged from at least three independent experiments. * P < 0.05, vs. static controls.

under OS was blocked by both the $\alpha 5$ neutralizing antibody (Fig. 4F) and Z-YVAD-FMK, a caspase-1 inhibitor (Fig. 4G). Thus, OS activation of integrin $\alpha 5$ is functionally linked to endothelial dysfunction.

EC Dysfunction in Atheroprone Regions in Vivo Is Integrin $\alpha 5$ -Dependent. To demonstrate the relevance of OS-activated integrin $\alpha 5$ in proinflammatory response in ECs in vivo, we used *Ldlr*^{-/-} mice subjected to partial ligation of the carotid artery, which induces an acute disturbed flow pattern (24). Compared with the sham-ligated contralateral artery in the same animal, inflammation was aggravated in the partially ligated artery, as evidenced by the activation of integrin $\alpha 5$ (Fig. 5A) and the upregulation of VCAM-1 and ICAM-1 (Fig. 5B). We used adenovirus (Ad)-mediated integrin $\alpha 5$ shRNA to infect partially ligated carotid arteries in *Ldlr*^{-/-} mice and confirmed the knockdown of integrin $\alpha 5$ (Fig. 5C, first four panels). Compared with control experiments (using Ad-GFP), Ad-integrin $\alpha 5$ shRNA decreased the OS-induced expressions of VCAM-1 and ICAM-1 (Fig. 5C, fifth and sixth panels). Because homozygous deletion of $\alpha 5$ is embryonic lethal, we created $\alpha 5^{+/-}$ mice and their $\alpha 5^{+/+}$ wild-type littermates to further validate the deleterious role of integrin $\alpha 5$ in inducing EC dysfunction in relation




Fig. 5. EC dysfunction in atheroprone region of mouse aorta is integrin $\alpha 5$ -dependent. (A and B) Male *Ldlr*^{-/-} mice (8 wk-old, $n = 10$) that had undergone partial ligation of the carotid artery were fed a WD for 1 wk. The left (partially ligated) and right (sham-operated) carotid arteries (LCA and RCA, respectively) were isolated for *en face* immunostaining. Whereas A shows representative images of Act- $\alpha 5$ (red), Flot2 (green), and nuclear (blue) from RCA and LCA, images in B are VCAM-1 (red), ICAM-1 (green), and nuclear (blue). In C, the experimental conditions and number of animals were the same as those in A and B except that carotid arteries were infused with Ad- $\alpha 5$ shRNA or Ad-GFP during vessel ligation. The first four panels show *en face* immunostaining of integrin $\alpha 5$ ($\alpha 5$, red), GFP (green), nuclear (blue), and merged images, and fifth and sixth panels show VCAM-1 (red, fifth panel) and ICAM-1 (red, sixth panel). (D) *En face* immunostaining of VCAM-1 (red), ICAM-1 (red), $\alpha 5$ (red), and nuclear (blue) in AA and TA from integrin $\alpha 5^{+/-}$ mice and the wild-type littermates ($\alpha 5^{+/+}$) (8 wk-old, $n = 10$ in each group). (E) The expressions of integrin $\alpha 5$ in $\alpha 5^{+/-}$ mice and the wild-type littermates were verified by *en face* immunostaining.




Fig. 6. Atherosclerosis in atheroprone region of mouse aorta is integrin $\alpha 5$ -dependent. (A) The 8-wk-old $Ldlr^{-/-} \alpha 5^{+/+}$ and $Ldlr^{-/-} \alpha 5^{+/-}$ mice were fed a WD for 4 wk. (B) $Ldlr^{-/-}$ mice received i.p. injection of ATN-161 or scramble peptide 1 wk before first receiving WD, which was given every third day until the termination of the experiment. (A and B) Representative of Oil Red O staining of aortas from the various groups of animals. (C and D) Quantification of percent lesion areas in the AA, TA, and total aorta from groups A and B. ANOVA followed by the Bonferroni post hoc test was used for statistical analysis. Data are mean \pm SEM normalized to control. * $P < 0.05$, vs. control.

to atherosclerosis. The reduced expression of integrin $\alpha 5$ in $\alpha 5^{+/-}$ mice was verified by *en face* immunostaining (Fig. 5E). Haploinsufficiency of integrin $\alpha 5$ markedly blocked the OS-induced expression of VCAM-1 and ICAM-1 in the inner curvature of AA (atheroprone area) of integrin $\alpha 5^{+/-}$ mice, compared with the corresponding area in $\alpha 5^{+/+}$ littermates (Fig. 5D).

Atherogenesis in $Ldlr^{-/-}$ Mice Is Integrin $\alpha 5$ -Dependent. We used two different animal models to demonstrate that the OS-regulated $\alpha 5$ is relevant to atherogenesis *in vivo*. First, 8 wk-old $Ldlr^{-/-}$ and $Ldlr^{-/-} \alpha 5^{+/-}$ mice were fed a western diet (WD) for 4 wk. The extent of atherosclerosis in the aorta was determined accordingly. There was a $\sim 50\%$ decrease in total lesion area as well as that in the AA in the $Ldlr^{-/-} \alpha 5^{+/-}$ mice (Fig. 6A and C). In a second model, $Ldlr^{-/-}$ mice were fed with WD and injected with ATN-161, which is an inhibitory peptide of integrin $\alpha 5$. Compared with $Ldlr^{-/-}$ mice injected with a scrambled peptide, inhibition of integrin $\alpha 5$ by ATN-161 significantly decreased the total atherosclerosis area. The lesion areas in both AA and TA were decreased by ATN-161 treatment (Fig. 6B and D). These data demonstrate the importance of integrin $\alpha 5$ in atherogenesis in the $Ldlr^{-/-}$ mice, similar to that in Apolipoprotein E-deficient ($Apoe^{-/-}$) mice, reported by Yurdagul et al. (16).

Discussion

The present study has revealed a mechanotransduction mechanism by which atheroprone flow causes endothelial dysfunction through integrin $\alpha 5$ translocation into lipid rafts and subsequent activation. Consequently, NLRP3 inflammasome is induced by the activated integrin $\alpha 5$. The results of our proteomics analysis of proteins suggested that OS leads to $\alpha 5$ translocation into lipid rafts. In ensuing studies, we confirmed that OS increased integrin $\alpha 5$ translocation into lipid rafts, and that this was a prerequisite for integrin $\alpha 5$ activation. In contrast, PS caused integrin $\alpha 5$ to reside largely outside lipid rafts, and therefore the activity of $\alpha 5$ showed little changes. We also demonstrated a link between integrin $\alpha 5$ activation and NLRP3 inflammasome induction in ECs under OS. Thus, the novelties of our study are twofold. First, OS activates integrin $\alpha 5$ exclusively in the lipid raft. Second, integrin $\alpha 5$ activation in turn mediates NLRP3 inflammasome induction by OS. Thus, the OS-induced activation of integrin $\alpha 5$, in the context of atheroprone flow and hypercholesterolemia, appears to be an essential step in the atherogenic process.

Shear stress may modulate EC functions in a rapid manner through its action on the plasma membrane. We have previously reported that shear stress changes membrane fluidity temporally

and spatially (25). Specifically, OS increases the cholesterol content in EC membranes, which results in decreased membrane fluidity (21). Such alterations of biophysical properties would affect the structure and function of membrane-associated proteins, especially those found in lipid rafts, such as integrin $\alpha 5$. Significantly, OS, but not PS, induced the translocation of integrin $\alpha 5$ into lipid rafts (Fig. 1). Because OS increases the cholesterol content of EC membranes, the cholesterol-enriched lipid rafts may thus sequester integrin $\alpha 5$ to allow for its sustained activation. This notion is supported by our previous observation that Cav1 and the F-actin-based cytoskeleton are involved in protein translocation into lipid rafts (21, 22). Indeed, results in Fig. 2 and Fig. S2 show that Cav1 and F-actin-based cytoskeleton are required for OS-induced $\alpha 5$ translocation.

Atheroprone flow activates NF- κ B and increases the expression of chemoattractants (e.g., monocyte chemoattractant protein-1) and adhesion molecules (e.g., ICAM-1, VCAM-1) *in vitro* and *in vivo* (26–30). Recent studies suggest that atheroprone flow regulates gene expression at the genome level via epigenetic regulation, in particular, DNA methylation (31–33). At the signaling level, atheroprone flow can induce NLRP3 inflammasome in ECs, which contributes to atherosclerosis susceptibility (20). The results in Fig. 4 show that functional blocking or genetic ablation of $\alpha 5$ inhibited the OS-induced NLRP3 inflammasome. Because inflammasome induction leads to the increase in the level of IL-1 β and IL-18, the OS-induced inflammation in ECs is at least in part mediated by integrin $\alpha 5$ activation of NLRP3 inflammasome. The mechanosensitive integrin $\alpha 5$ -inflammation axis *in vivo* was validated by the loss-of-function experiments with the use of integrin $\alpha 5$ shRNA, and haploinsufficiency of integrin $\alpha 5$ in mice. All these experiments showed decreased EC inflammation under atheroprone flow (Fig. 5C and D). Importantly, we demonstrated that the *in vivo* relevance of integrin $\alpha 5$ activation to the atherogenic process using two different murine models that have either haploinsufficiency of $\alpha 5$ or inhibition of $\alpha 5$ activation by an inhibitory peptide led to marked protection against atherosclerosis even in the presence of sustained hypercholesterolemia. In our studies, we used the WD-fed $Ldlr^{-/-}$ model to test the role of integrin $\alpha 5$ in atherogenesis. The study by Yurdagul et al. (16) also demonstrated the important role of integrin $\alpha 5$ activation for the atherogenic process using the $Apoe^{-/-}$ model; they found that $\alpha 5$ activation could also occur via a different mechanism that involved OxLDL (16). The fact that integrin $\alpha 5$ was demonstrated to play an obligatory role in atherogenesis in two different murine models (i.e., $Ldlr^{-/-}$ and $Apoe^{-/-}$), and can be activated by two different mechanisms

(i.e., atheroprone flow and OxLDL), adds to the confidence that this pathway is highly relevant in vivo.

It is intriguing that both OS and OxLDL can activate integrin $\alpha 5$ in vitro. Given the reduced atherosclerosis found in the atheroprone areas in the *Ldlr^{-/-}/ $\alpha 5^{+/-}$* mice, we propose that OS and OxLDL could share similar mechanism in activating integrin $\alpha 5$. Integrin $\alpha 5$ activation would occur at regions of profound atheroprone flow, where atherosclerosis is most prevalent. Seemingly, OxLDL is not present in the circulation, but it is believed that OxLDL is generated in the intima at sites of lesion formation. How this in turn would lead to EC $\alpha 5$ activation is less clear. Whatever the mechanism is, one could envision the regulation of $\alpha 5$ activation by both mechanical and biochemical actions to play a key role in EC dysfunction and atherosclerosis. This thesis is in line with that proposed by Davies and colleagues that the synergism between flow characteristics and hypercholesterolemia deserves “further evaluation of the epigenetic and epigenomic regulation of endothelial phenotype adaptation during early pathological change” (31, 34).

In summary, our study reveals a novel mechanism by which atheroprone flow increases endothelial inflammation that involves

an integrated response of lipid rafts, F-actin-based cytoskeleton, and integrin $\alpha 5$. Our results provide a novel experimental basis for the use of integrin $\alpha 5$ -specific antagonists for preventing dysfunctional endothelium and alleviating atherosclerosis.

Materials and Methods

All animal protocols were approved by Peking University Health Science Center Animal Care and Use Committee. The sources of antibodies and reagents, and detailed methods for cell culture, shear stress, purification of membrane and lipid raft protein, fluorescence anisotropy measurements, iTRAQ labeling and proteomics analysis, *en face* immunostaining, RNA interference, monocyte adhesion, fibronectin adhesion, carotid partial ligation, generation of $\alpha 5^{+/-}$ mice, adenovirus construction, lesion area assessment, and statistical analysis are described in [SI Materials and Methods](#).

ACKNOWLEDGMENTS. The authors thank Dr. Joseph Witztum for his valuable comments. This work was supported by the National Natural Science Foundation of China (81130002 and 30821001 to Y.Z.; 81270349 to J.Y.-J.S.), the Major National Basic Research Grant of China (2010CB517504 to Y.Z.), and National Institutes of Health Grant HL108735 (to J.Y.-J.S. and S.C.).

- Davies PF (2009) Hemodynamic shear stress and the endothelium in cardiovascular pathophysiology. *Nat Clin Pract Cardiovasc Med* 6(1):16–26.
- Humphrey JD, Dufresne ER, Schwartz MA (2014) Mechanotransduction and extracellular matrix homeostasis. *Nat Rev Mol Cell Biol* 15(12):802–812.
- Tzima E, del Pozo MA, Shattil SJ, Chien S, Schwartz MA (2001) Activation of integrins in endothelial cells by fluid shear stress mediates Rho-dependent cytoskeletal alignment. *EMBO J* 20(17):4639–4647.
- Goncalves I, Nesbitt WS, Yuan Y, Jackson SP (2005) Importance of temporal flow gradients and integrin $\alpha_{IIb}\beta_3$ mechanotransduction for shear activation of platelets. *J Biol Chem* 280(15):15430–15437.
- Kapur S, Mohan S, Baylink DJ, Lau KH (2005) Fluid shear stress synergizes with insulin-like growth factor-I (IGF-I) on osteoblast proliferation through integrin-dependent activation of IGF-I mitogenic signaling pathway. *J Biol Chem* 280(20):20163–20170.
- Orr AW, Ginsberg MH, Shattil SJ, Deckmyn H, Schwartz MA (2006) Matrix-specific suppression of integrin activation in shear stress signaling. *Mol Biol Cell* 17(11):4686–4697.
- Chiu JJ, Chien S (2011) Effects of disturbed flow on vascular endothelium: Pathophysiological basis and clinical perspectives. *Physiol Rev* 91(1):327–387.
- Radel C, Carlile-Klusacek M, Rizzo V (2007) Participation of caveolae in $\beta 1$ integrin-mediated mechanotransduction. *Biochem Biophys Res Commun* 358(2):626–631.
- Wang Y, et al. (2002) Interplay between integrins and FLK-1 in shear stress-induced signaling. *Am J Physiol Cell Physiol* 283(5):C1540–C1547.
- Brown WS, Khalil JS, Rodriguez-Cruz TG, Lizee G, McIntyre BW (2014) B-Raf regulation of integrin $\alpha 4\beta 1$ -mediated resistance to shear stress through changes in cell spreading and cytoskeletal association in T cells. *J Biol Chem* 289(33):23141–23153.
- Jalali S, et al. (2001) Integrin-mediated mechanotransduction requires its dynamic interaction with specific extracellular matrix (ECM) ligands. *Proc Natl Acad Sci USA* 98(3):1042–1046.
- Simons K, Ehehalt R (2002) Cholesterol, lipid rafts, and disease. *J Clin Invest* 110(5):597–603.
- Cai WJ, et al. (2009) Activation of the integrins $\alpha 5\beta 1$ and $\alpha 5\beta 3$ and focal adhesion kinase (FAK) during arteriogenesis. *Mol Cell Biochem* 322(1–2):161–169.
- Funk SD, et al. (2010) Matrix-specific protein kinase A signaling regulates p21-activated kinase activation by flow in endothelial cells. *Circ Res* 106(8):1394–1403.
- Mitchell JS, Brown WS, Woodside DG, Vanderslice P, McIntyre BW (2009) Clustering T-cell GM1 lipid rafts increases cellular resistance to shear on fibronectin through changes in integrin affinity and cytoskeletal dynamics. *Immunol Cell Biol* 87(4):324–336.
- Yurdagul A, Jr, et al. (2014) $\alpha 5\beta 1$ integrin signaling mediates oxidized low-density lipoprotein-induced inflammation and early atherosclerosis. *Arterioscler Thromb Vasc Biol* 34(7):1362–1373.
- Thorin E, Hamilton C, Dominiczak AF, Dominiczak MH, Reid JL (1995) Oxidized-LDL induced changes in membrane physico-chemical properties and $[Ca^{2+}]_i$ of bovine aortic endothelial cells. Influence of vitamin E. *Atherosclerosis* 114(2):185–195.
- Lénárt N, et al. (2015) Cultured cells of the blood-brain barrier from apolipoprotein B-100 transgenic mice: Effects of oxidized low-density lipoprotein treatment. *Fluids Barriers CNS* 12(1):17.
- Franchi L, Muñoz-Planillo R, Núñez G (2012) Sensing and reacting to microbes through the inflammasomes. *Nat Immunol* 13(4):325–332.
- Xiao H, et al. (2013) Sterol regulatory element binding protein 2 activation of NLRP3 inflammasome in endothelium mediates hemodynamic-induced atherosclerosis susceptibility. *Circulation* 128(6):632–642.
- Fu Y, et al. (2011) A novel mechanism of $\gamma \delta$ T-lymphocyte and endothelial activation by shear stress: The role of ecto-ATP synthase β chain. *Circ Res* 108(4):410–417.
- Wang T, Chen Z, Wang X, Shyy JY, Zhu Y (2006) Cholesterol loading increases the translocation of ATP synthase beta chain into membrane caveolae in vascular endothelial cells. *Biochim Biophys Acta* 1761(10):1182–1190.
- Jun HK, Lee SH, Lee HR, Choi BK (2012) Integrin $\alpha 5\beta 1$ activates the NLRP3 inflammasome by direct interaction with a bacterial surface protein. *Immunity* 36(5):755–768.
- Nam D, et al. (2009) Partial carotid ligation is a model of acutely induced disturbed flow, leading to rapid endothelial dysfunction and atherosclerosis. *Am J Physiol Heart Circ Physiol* 297(4):H1535–H1543.
- Butler PJ, Norwich G, Weinbaum S, Chien S (2001) Shear stress induces a time- and position-dependent increase in endothelial cell membrane fluidity. *Am J Physiol Cell Physiol* 280(4):C962–C969.
- Davies PF, Civelek M, Fang Y, Fleming I (2013) The atherosusceptible endothelium: Endothelial phenotypes in complex haemodynamic shear stress regions in vivo. *Cardiovasc Res* 99(2):315–327.
- Cheng C, et al. (2005) Shear stress affects the intracellular distribution of eNOS: Direct demonstration by a novel in vivo technique. *Blood* 106(12):3691–3698.
- Hajra L, et al. (2000) The NF- κ B signal transduction pathway in aortic endothelial cells is primed for activation in regions predisposed to atherosclerotic lesion formation. *Proc Natl Acad Sci USA* 97(16):9052–9057.
- Nagel T, Resnick N, Dewey CF, Jr, Gimbrone MA, Jr (1999) Vascular endothelial cells respond to spatial gradients in fluid shear stress by enhanced activation of transcription factors. *Arterioscler Thromb Vasc Biol* 19(8):1825–1834.
- Walpolo PL, Gotlieb AI, Cybulsky MI, Langille BL (1995) Expression of ICAM-1 and VCAM-1 and monocyte adherence in arteries exposed to altered shear stress. *Arterioscler Thromb Vasc Biol* 15(1):2–10.
- Jiang YZ, et al. (2014) Hemodynamic disturbed flow induces differential DNA methylation of endothelial Kruppel-Like Factor 4 promoter in vitro and in vivo. *Circ Res* 115(1):32–43.
- Dunn J, et al. (2014) Flow-dependent epigenetic DNA methylation regulates endothelial gene expression and atherosclerosis. *J Clin Invest* 124(7):3187–3199.
- Zhou J, Li YS, Wang KC, Chien S (2014) Epigenetic mechanism in regulation of endothelial function by disturbed flow: Induction of DNA hypermethylation by DNMT1. *Cell Mol Bioeng* 7(2):218–224.
- Jiang YZ, Manduchi E, Jiménez JM, Davies PF (2015) Endothelial epigenetics in biomechanical stress: Disturbed flow-mediated epigenomic plasticity in vivo and in vitro. *Arterioscler Thromb Vasc Biol* 35(6):1317–1326.
- Zhu Y, et al. (1998) LDL induces transcription factor activator protein-1 in human endothelial cells. *Arterioscler Thromb Vasc Biol* 18(3):473–480.
- Laufs U, Liao JK (1998) Post-transcriptional regulation of endothelial nitric oxide synthase mRNA stability by Rho GTPase. *J Biol Chem* 273(37):24266–24271.
- Zhu Y, et al. (2000) Lipoprotein promotes caveolin-1 and Ras translocation to caveolae: Role of cholesterol in endothelial signaling. *Arterioscler Thromb Vasc Biol* 20(11):2465–2470.
- Wojciak-Stothard B, et al. (2014) Aberrant chloride intracellular channel 4 expression contributes to endothelial dysfunction in pulmonary arterial hypertension. *Circulation* 129(17):1770–1780.
- Gu MX, et al. (2012) Proteomic analysis of endothelial lipid rafts reveals a novel role of statins in antioxidant. *J Proteome Res* 11(4):2365–2373.
- Grönberg M, et al. (2006) Biomarker discovery from pancreatic cancer secretome using a differential proteomic approach. *Mol Cell Proteomics* 5(1):157–171.
- Boukli NM, et al. (2012) Unique and differential protein signatures within the mononuclear cells of HIV-1 and HCV mono-infected and co-infected patients. *Clin Proteomics* 9(1):11.
- Stoeltz O, et al. (2003) Inhibition of integrin $\alpha 5\beta 1$ function with a small peptide (ATN-161) plus continuous 5-FU infusion reduces colorectal liver metastases and improves survival in mice. *Int J Cancer* 104(4):496–503.