

RESEARCH ARTICLE

Is Long-Term Low-Dose Aspirin Therapy Associated with Renal Dysfunction in Patients with Type 2 Diabetes? JPAD2 Cohort Study

Sadanori Okada^{1,2}, Takeshi Morimoto³, Hisao Ogawa⁴, Mio Sakuma³, Hirofumi Soejima⁴, Masafumi Nakayama⁴, Hideaki Jinnouchi⁵, Masako Waki⁶, Yasuhiro Akai¹, Hitoshi Ishii¹, Yoshihiko Saito^{2,7*}, investigators for the Japanese Primary Prevention of Atherosclerosis with Aspirin for Diabetes (JPAD) trial[¶]

1 Department of Diabetology, Nara Medical University, Kashihara, Nara, Japan, **2** First Department of Internal Medicine, Nara Medical University, Kashihara, Nara, Japan, **3** Department of Clinical Epidemiology, Hyogo College of Medicine, Nishinomiya, Hyogo, Japan, **4** Department of Cardiovascular Medicine, Graduate School of Medical Science, Kumamoto University, Kumamoto, Japan, **5** Diabetes Care Center, Jinnouchi Hospital, Kumamoto, Japan, **6** Division of Endocrinology and Metabolism, Department of Internal Medicine, Shizuoka City Hospital, Shizuoka, Japan, **7** Department of Regulatory Medicine of Blood Pressure, Nara Medical University, Kashihara, Nara, Japan

¶ Membership of the JPAD trial is listed in the Acknowledgments.

* yssaito@naramed-u.ac.jp



CrossMark
click for updates

OPEN ACCESS

Citation: Okada S, Morimoto T, Ogawa H, Sakuma M, Soejima H, Nakayama M, et al. (2016) Is Long-Term Low-Dose Aspirin Therapy Associated with Renal Dysfunction in Patients with Type 2 Diabetes? JPAD2 Cohort Study. PLoS ONE 11(1): e0147635. doi:10.1371/journal.pone.0147635

Editor: Paolo Cravedi, Ichan School of Medicine at Mount Sinai, UNITED STATES

Received: September 14, 2015

Accepted: January 4, 2016

Published: January 25, 2016

Copyright: © 2016 Okada et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This study was supported by the Ministry of Health, Labour and Welfare of Japan (HO). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Dr. TM reports receiving research grants from Bayer HealthCare and lecturer's fees from Daiichi-Sankyo, Eisai, Kowa Pharmaceutical, Kyorin Pharmaceutical, and Pfizer

Abstract

Background

Low-dose aspirin is widely recommended for patients at high risk for cardiovascular disease (CVD); however, it remains uncertain whether long-term treatment adversely affects renal function in patients with diabetes.

We investigated whether long-term low-dose aspirin affects renal dysfunction in patients with diabetes.

Methods

We conducted a randomized controlled trial (RCT), the Japanese Primary Prevention of Atherosclerosis with Aspirin for Diabetes (JPAD) trial, to evaluate low-dose aspirin as primary prevention for CVD in patients with type 2 diabetes. We followed the patients with negative urine dipstick albumin of the JPAD trial in a cohort study after the RCT period was completed. Patients were randomly allocated to receive aspirin (81 mg or 100 mg daily, aspirin group) or no aspirin (no aspirin group). After the RCT, the treating physician decided whether to administer aspirin. We evaluated the incidence of positive urine dipstick albumin and annual changes in estimated glomerular filtration rate (eGFR).

Results

Positive urine dipstick albumin developed in 297 patients in the aspirin group (n = 1,075) and 270 patients in the no aspirin group (n = 1,098) during follow-up (median, 8.5 years). Intention-to-treat analysis showed low-dose aspirin did not increase the incidence of

Japan for the past three years. Dr. HO reports receiving research grants from AstraZeneca Pharmaceutical, Astellas Pharma, Bayer HealthCare, Boehringer Ingelheim Pharmaceutical, Bristol-Myers, Chugai Pharmaceutical, Daiichi Sankyo, Sumitomo Dainippon Pharma, Kowa Pharmaceutical, MSD, Novartis, Otsuka Pharmaceutical, Pfizer Japan, Sanofi, and Takeda Pharmaceutical and lecturer's fees from AstraZeneca Pharmaceutical, Bayer HealthCare, Daiichi-Sankyo, MSD, Pfizer Japan, Sanofi, and Taisho Pharmaceutical for the past three years. Dr. HI reports receiving research grant support from Novo Nordisk and Mitsubishi Tanabe Pharma and lecturer's fees from AstraZeneca Pharmaceutical, Eli Lilly Japan, Sanofi, Daiichi Sankyo, Sumitomo Dainippon Pharma, Takeda Pharmaceutical, Mitsubishi Tanabe Pharma, Novartis Pharma, Novo Nordisk, Boehringer Ingelheim Pharmaceutical, and MSD. Dr. YS reports receiving research grant support from Ono Pharmaceutical, MSD, Mitsubishi Tanabe Pharma, Daiichi Sankyo, Takeda Pharmaceutical, Novartis Pharma, Shionogi, Astellas Pharma, Otsuka Pharmaceutical, St. Jude Medical Japan, and Kyowa Hakkō Kirin; lecturer's fees from MSD, Mitsubishi Tanabe Pharma, Takeda Pharmaceutical, Daiichi Sankyo, Otsuka Pharmaceutical, and Pfizer Japan; and consulting fees from Ono Pharmaceutical for the past three years. The rest of the authors have declared that no competing interests exist. Our competing interests do not alter our adherence to PLOS ONE policies on sharing data and materials

positive urine dipstick albumin (hazard ratio [HR], 1.17; 95% confidence interval [CI], 0.995–1.38). On-treatment analysis yielded similar results (HR, 1.08; 95% CI, 0.92–1.28). Multivariable analysis showed the incidence of positive urine dipstick albumin was higher among the elderly and those with elevated serum creatinine, high hemoglobin A1c, or high blood pressure; however, low-dose aspirin did not increase the risk of positive urine dipstick albumin. There were no significant differences in annual changes in eGFR between the groups (aspirin, -0.8 ± 2.9 ; no aspirin, -0.9 ± 2.5 ml/min/1.73m²/year).

Conclusion

Long-term low-dose aspirin does not affect eGFR and positive urine dipstick albumin in patients with type 2 diabetes.

Introduction

Low-dose aspirin therapy is widely recommended for preventing cardiovascular disease (CVD) [1]. Individuals who are at high risk for CVD may take low-dose aspirin for life. However, it remains uncertain whether long-term treatment is safe with respect to renal function because aspirin belongs to a class of cyclooxygenase inhibitors. The inhibition of cyclooxygenase decreases production of prostaglandin in the kidney, and reduces renal blood flow and glomerular filtration rate. In patients with reduced renal function, this may result in retention of water, hypertension and, in some case, to renal failure [2]. Previous clinical studies have yielded conflicting results about aspirin and the risk of chronic kidney disease (CKD) or end-stage renal disease (ESRD) [3–13]. There have been very few studies on the relationship between aspirin and CKD in patients with diabetes.

Low-dose aspirin therapy is also recommended for CVD prevention in high-risk diabetic patients with a history of CVD [14] or CVD risk factors [15]. Patients with diabetes are at high risk not only for CVD, but also CKD secondary to diabetic nephropathy [16, 17]. Albuminuria or proteinuria occur early in the course of diabetic nephropathy, before an appreciable decline in the estimated glomerular filtration rate (eGFR) [18]. However, renal function in previous studies was evaluated based on serum creatinine levels, creatinine clearance, or eGFR, not the presence of albuminuria or proteinuria. Therefore, we thought it was necessary to assess the long-term risk of low-dose aspirin therapy based on the incidence of albuminuria or proteinuria in patients with diabetes.

The Japanese Primary Prevention of Atherosclerosis with Aspirin for Diabetes (JPAD) trial was a randomized controlled trial (RCT) evaluating whether low-dose aspirin prevents CVD in patients with type 2 diabetes and no history of CVD [19]. We followed the patients of the JPAD trial in a cohort study (JPAD2 cohort study) after the RCT was completed. In the JPAD2 cohort study, we analyzed whether long-term low-dose aspirin therapy affects eGFR and the incidence of positive urine dipstick albumin in patients with type 2 diabetes.

Methods

The original JPAD trial was a multicenter, prospective, randomized, open-label, blinded end-point trial conducted at 163 institutions throughout Japan. This trial was performed according to the Declaration of Helsinki and was approved by the ethics committee of each participating hospital (Nara Medical University Ethics Committee and Graduate School of Medical Science, Kumamoto University Ethics Committee). Written informed consent was obtained from each

participant before participation in the original JPAD trial. In the JPAD2 cohort study, the revised study protocol was approved by the ethics committees (Nara Medical University Ethics Committee and Graduate School of Medical Science, Kumamoto University Ethics Committee). Verbal informed consent was obtained at the start of the JPAD2 cohort study, according to the approval by the ethics committees. The study protocol of the JPAD trial was registered at clinicaltrials.gov with the identifier NCT00110448.

Details on the design of the JPAD trial have been previously described [19]. In brief, patient enrollment started in December 2002 and was completed in May 2005. We enrolled 2,536 Japanese patients with type 2 diabetes between 30 and 85 years of age without a history of CVD, who were randomly allocated to receive aspirin (81 mg or 100 mg daily, aspirin group) or no aspirin (no aspirin group). All patients were allowed to undergo all concurrent treatments. After the JPAD trial was completed in May 2008, we started the JPAD2 cohort study. In the JPAD2 cohort study, the decision on whether to administer low-dose aspirin depended on the treating physician's clinical judgment (Fig 1).

In the present analysis, we defined newly developed positive urine dipstick albumin as the primary endpoint. The positive urine dipstick albumin was defined as '1+' and over. We assessed the annual incidence of positive urine dipstick albumin to July 2013. We followed patients with negative urine dipstick albumin at baseline of the original JPAD trial, and assessed the effect of low-dose aspirin on the incidence of positive urine dipstick albumin.


We also evaluated annual changes in eGFR in each group. eGFR (ml/min/1.73 m² of body surface area) was calculated using the new three-variable Japanese equation for estimating glomerular filtration rate: $eGFR = 194 \times \text{serum creatinine}^{-1.094} \times \text{age}^{-0.287} \times 0.739$ (if female) [20]. Annual changes in eGFR were calculated as eGFR divided by time between baseline and follow-up (years). We analyzed differences in the annual change in eGFR between the aspirin and no aspirin groups. Patients whose follow-up period was less than 1 year were excluded from the analysis.

Statistical analyses

Categorical variables were expressed as numbers and percentages. Continuous variables were expressed as means \pm SD or medians (interquartile range [IQR]). Based on their distribution, continuous variables were compared using the Student's *t*-test or Wilcoxon rank sum test as appropriate. Efficacy comparisons were made on the basis of time to the first event according to the intention-to-treat principle. Patients lost to follow-up were censored on the day of their last visit. We used Cox proportional hazards models to estimate hazard ratios (HRs) and 95% confidence intervals (CIs). The cumulative incidence of each endpoint was estimated using the Kaplan-Meier method in each group, and differences between groups were assessed with the log-rank test. Next, we performed an on-treatment analysis, based on whether patients were actually taking aspirin during the follow-up period, using Cox proportional hazards modeling and the Kaplan-Meier method.

We developed multivariable Cox proportional hazards models to assess the effect of aspirin on the incidence of positive urine dipstick albumin in both intention-to-treat and on-treatment analyses. We employed the following variables in the multivariable models: age (≥ 65 years at baseline), sex, baseline serum creatinine level (continuous variable), hemoglobin A1c (HbA1c) (binary variable divided by a median value), blood pressure (systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg), and treatment with angiotensin-converting enzyme (ACE) inhibitors or angiotensin II type 1 receptor blockers (ARBs). The mean HbA1c and blood pressure values at baseline and each annual follow-up were used. Treatment with ACE inhibitors or ARBs was found to be associated with renal function in previous studies [21, 22].

A



B




Fig 1. Timeline of the JPAD2 cohort study. A, The timeline of the JPAD trial and the JPAD2 cohort study. RCT indicates randomized controlled trial. B, The flow chart of the JPAD trial and the JPAD2 cohort study.

doi:10.1371/journal.pone.0147635.g001

Statistical analyses were conducted by an independent statistician (T.M.) with the use of JMP 8.0 (SAS Institute, Cary, NC) and SAS 9.4 (SAS Institute) software. *P* values less than 0.05 were considered statistically significant.

Results

Patient characteristics at baseline


We included 2,173 patients with negative urine dipstick albumin at baseline of the original JPAD trial (Table 1). There were 1,075 patients in the aspirin group and 1,098 patients in the no aspirin group. The mean age ± SD was 65 ± 10 years in the aspirin group and 64 ± 10 years in the no aspirin group (*P* = 0.01). The aspirin group had a slightly higher mean serum creatinine level than the no aspirin group (aspirin group, 68 ± 19 μmol/L; no aspirin group, 66 ± 18 μmol/L; *P* = 0.04), but there were no significant differences in eGFR at baseline (aspirin group, 74.0 ± 19.3 ml/min/1.73m²; no aspirin group, 75.6 ± 20.0 ml/min/1.73m²; *P* = 0.06).

Table 1. Patients' characteristics at baseline of the original JPAD trial.

N	Aspirin	No aspirin	P value
	1075	1098	
Age, y	65 ± 10	64 ± 10	0.01
Male	592 (55)	573 (52)	0.2
BMI, kg/m ²	24 ± 4	24 ± 4	0.4
Duration of diabetes, y	7.2 (2.8–12.2)	6.5 (2.9–12.1)	0.4
Hypertension	604 (56)	605 (55)	0.6
Dyslipidemia	578 (54)	568 (52)	0.3
History of smoking	234 (22)	210 (19)	0.1
Systolic BP, mmHg	135 ± 15	134 ± 14	0.06
Diastolic BP, mmHg	77 ± 9	76 ± 9	0.03
FPG, mmol/L	8.1 ± 2.7	8.0 ± 2.6	0.7
HbA1c, %	7.5 ± 1.4	7.4 ± 1.2	0.09
Serum creatinine, μmol/L	0.77 ± 0.22	0.75 ± 0.20	0.04
eGFR, ml/min/1.73m ²	74.0 ± 19.3	75.6 ± 20.0	0.06
Total cholesterol, mmol/L	5.22 ± 0.88	5.17 ± 0.88	0.1
Fasting triglycerides, mmol/L	1.28 (0.89–1.78)	1.25 (0.89–1.81)	0.5
HDL cholesterol, mmol/L	1.45 ± 0.41	1.45 ± 0.39	0.9
Medications			
Sulfonylurea	619 (58)	604 (55)	0.2
α-glycosidase inhibitor	353 (33)	363 (33)	0.9
Biguanide	134 (12)	156 (14)	0.2
Thiazolidinedione	51 (5)	49 (4)	0.8
Insulin	138 (13)	130 (12)	0.5
Calcium channel blocker	344 (32)	358 (33)	0.8
ACE inhibitor	145 (13)	159 (14)	0.5
ARB	209 (19)	210 (19)	0.9
β-blocker	61 (6)	76 (7)	0.2
Statin	266 (25)	292 (27)	0.3

Duration of diabetes and fasting triglyceride levels are expressed as medians (interquartile range). BMI indicates body mass index; BP, blood pressure; FPG, fasting plasma glucose; eGFR, estimated glomerular filtration rate; HDL, high-density lipoprotein; ACE, angiotensin-converting enzyme; and ARB, angiotensin II type 1 receptor blocker.

doi:10.1371/journal.pone.0147635.t001



No. at risk											
Aspirin	1075	983	869	759	675	591	532	451	393	297	79
No aspirin	1098	1013	922	841	761	662	611	533	462	337	101
Incidence											
Aspirin	0.02	0.09	0.16	0.21	0.24	0.26	0.29	0.31	0.34	0.39	
No aspirin	0.02	0.07	0.11	0.15	0.20	0.22	0.25	0.27	0.29	0.34	

Fig 2. Incidence of positive urine dipstick albumin in patients on long-term low-dose aspirin therapy in the intention-to-treat analysis. Positive urine dipstick albumin developed in 297 patients in the aspirin group and 270 patients in the no aspirin group. The intention-to-treat analysis showed that low-dose aspirin did not increase the incidence of positive urine dipstick albumin (HR, 1.17; 95% CI, 0.995 to 1.38; log-rank $P = 0.057$)


doi:10.1371/journal.pone.0147635.g002

Diastolic blood pressure was significantly higher by 1 mmHg in the aspirin group (aspirin group, 77 ± 9 mmHg; no aspirin group, 76 ± 9 ; $P = 0.03$). There were no differences in systolic blood pressure, prevalence of hypertension, and use of antihypertensives.

Effect of low-dose aspirin on the incidence of positive urine dipstick albumin

The median duration of follow-up was 8.5 years (95% CI, 8.3 to 8.7 years), which includes the RCT period of the JPAD trial. During the follow-up period, positive urine dipstick albumin developed in 297 patients in the aspirin group and 270 patients in the no aspirin group. Kaplan-Meier curves based on the intention-to-treat principle showed that low-dose aspirin does not increase the incidence of positive urine dipstick albumin (HR, 1.17; 95% CI, 0.995 to 1.38; log-rank $P = 0.057$; Fig 2).

Since the JPAD2 cohort study was a cohort study, the use of low-dose aspirin was based on each treating physician’s clinical judgment after the RCT period of the JPAD trial. In July 2013 survey, 881 (82%) patients in the aspirin group and 123 (11%) in the no aspirin group were taking low-dose aspirin (Fig 1B). When we took into account the actual use of low-dose aspirin



No. at risk											
Aspirin	1004	920	816	719	636	559	506	436	375	291	72
No aspirin	1169	1076	975	881	800	694	637	548	480	343	108
Incidence											
Aspirin		0.02	0.09	0.15	0.20	0.23	0.25	0.27	0.29	0.32	0.39
No aspirin		0.02	0.07	0.13	0.16	0.21	0.24	0.26	0.29	0.31	0.34

Fig 3. Incidence of positive urine dipstick albumin in patients on long-term low-dose aspirin therapy in the on-treatment analysis. In the on-treatment analysis, low-dose aspirin had no effect on the incidence of positive urine dipstick albumin (HR, 1.08; 95% CI, 0.92 to 1.28; log-rank $P = 0.32$).

doi:10.1371/journal.pone.0147635.g003

(on-treatment analysis), the incidence of positive urine dipstick albumin was not associated with the use of low-dose aspirin (HR, 1.08; 95% CI, 0.92 to 1.28; log-rank $P = 0.32$; Fig 3).

Multivariable analysis of the incidence of positive urine dipstick albumin

The multivariable Cox proportional hazards models were performed in 2,159 patients, since there were 14 missing data in serum creatinine at baseline. The multivariable analyses showed that low-dose aspirin was not associated with the incidence of positive urine dipstick albumin in the intention-to-treat analysis (HR, 1.12; 95% CI, 0.95 to 1.32) and the on-treatment analysis (HR, 1.08; 95% CI, 0.92 to 1.27) (Table 2). The incidence of positive urine dipstick albumin was higher among the elderly (≥ 65 years old) and those with elevated serum creatinine, high HbA1c ($\geq 7.2\%$), or high blood pressure in both analyses.

Annual changes in eGFR

The analysis of annual changes in eGFR included 1,574 patients (aspirin group, 766; no aspirin group, 808), because there were 599 missing data in eGFR at follow-up. The annual change in

Table 2. Multivariable analysis of the incidence of positive urine dipstick albumin.

	HR	95% CI		
Intention-to-treat analysis				
Aspirin use	1.12	0.95	to	1.32
Age ≥65 years	1.23	1.04	to	1.46
Male sex	0.95	0.79	to	1.15
Serum creatinine	2.59	1.71	to	3.93
HbA1c ≥7.2%	1.38	1.16	to	1.63
Systolic BP ≥140 mmHg or diastolic BP ≥90 mmHg	1.78	1.48	to	2.13
ACE inhibitor or ARB use at baseline	1.11	0.91	to	1.35
ACE inhibitor / ARB use at the time of the follow-up survey in 2009	0.98	0.81	to	1.18
On-treatment analysis				
Aspirin use	1.08	0.92	to	1.27
Age ≥65 years	1.24	1.04	to	1.46
Male sex	0.95	0.79	to	1.15
Serum creatinine	2.61	1.72	to	3.96
HbA1c ≥7.2%	1.38	1.16	to	1.63
Systolic BP ≥140 mmHg or diastolic BP ≥90 mmHg	1.80	1.50	to	2.15
ACE inhibitor or ARB use at baseline	1.10	0.91	to	1.33
ACE inhibitor / ARB use at the time of the follow-up survey in 2009	0.99	0.82	to	1.19

BP indicates blood pressure; ACE, angiotensin-converting enzyme; and ARB, angiotensin II type 1 receptor blocker.

doi:10.1371/journal.pone.0147635.t002

eGFR was -0.8 ± 2.9 ml/min/1.73m²/year in the aspirin group, and -0.9 ± 2.5 ml/min/1.73m²/year in the no aspirin group. There was no significant difference in the both group ($P = 0.2$).

Discussion

Low-dose aspirin therapy is recommended for preventing CVD in high-risk patients with diabetes [14, 15]. However, it remained uncertain whether long-term low-dose aspirin use affects renal function in patients with diabetes. Also in people without diabetes, the association between aspirin and the risk of CKD was controversial.

Previous case-control studies that estimated the cumulative aspirin dose in patients with CKD and those without CKD have produced conflicting results. Some case-control studies have found a positive association between aspirin use and CKD prevalence [4, 5], while other studies have found no association [6, 7].

Several large cohort studies were conducted to examine whether long-term aspirin use affects renal function in healthy people and patients with CKD. One cohort study in healthy people was based on the Physicians' Health Study. In this study, a total of 11,032 initially healthy men were followed for 14 years. The association between renal function based on serum creatinine and creatinine clearance and cumulative aspirin dose was not observed [8]. Another cohort study analyzed the association between changes in eGFR over 11 years and cumulative aspirin dose among 1,697 initially healthy women from the Nurses' Health Study. The study concluded that long-term aspirin use was not associated with renal dysfunction [10]. On the other hand, a cohort study of 19,163 patients with newly diagnosed CKD in Taiwan found that aspirin use was associated with an increased risk of ESRD [11]. The effect of aspirin on renal dysfunction in healthy individuals and patients with CKD may differ.

Most clinical studies evaluated renal function using serum creatinine, creatinine clearance, or eGFR, not albuminuria nor proteinuria. Recently, a nationwide cross-sectional study based on data from the National Health and Nutrition Examination Survey analyzed the association between habitual analgesic use and albuminuria prevalence in 8,057 US adults [13]. It showed that the prevalence of albuminuria (urinary albumin-to-creatinine ratio of 30 mg/g or greater) was similar in people with habitual aspirin use and those without habitual use of any analgesics. Since albuminuria and proteinuria occur before eGFR declines in some conditions such as diabetes [17], this study was notable in using albuminuria as a marker of renal dysfunction. However, this study was a cross-sectional study. Prospective studies were needed to assess the effect of low-dose aspirin on albuminuria or proteinuria.

A previous prospective study reported 'high-dose' aspirin reduced proteinuria in patients with diabetes. This study showed that 6 weeks of aspirin (990 mg/day) and dipyridamole (225 mg/day) reduced urinary protein excretion without changes in blood pressure and plasma glucose levels in 16 patients with type 1 diabetes [23]. Although the sample size was small, the results suggested that short-term high-dose aspirin had beneficial effects on proteinuria in patients with diabetes. Subsequently, more prospective studies were conducted to examine whether 'low-dose' aspirin affects albuminuria in patients with diabetes. These studies recruited patients with diabetes and microalbuminuria (urinary albumin excretion of 30 to 300 mg/day), who were then treated with low-dose aspirin (150 mg/day) for 4 weeks. There were no changes in urinary albumin excretion and eGFR [24, 25], suggesting that low-dose aspirin therapy does not affect albuminuria in patients with diabetes. However, these studies had relatively short follow-up periods. Long-term studies were required to evaluate the effect of low-dose aspirin on albuminuria or proteinuria in patients with diabetes.

The present study assessed for the first time whether long-term low-dose aspirin affects the incidence of positive urine dipstick albumin in patients with type 2 diabetes. Both intention-to-treat analysis and on-treatment analysis showed that low-dose aspirin did not increase the incidence of positive urine dipstick albumin for a median follow-up of 8.5 years. The multivariable Cox proportional hazards models also showed that low-dose aspirin did not increase the risk of positive urine dipstick albumin, adjusting age, sex, serum creatinine, hemoglobin A1c, blood pressure, and treatment with ACE inhibitors and ARBs.

Our study has several limitations. First, we did not have data about habitual analgesic use. Habitual analgesic use might affect the relationship between low-dose aspirin and renal function. Second, we did not quantify the degree of albuminuria. It is possible that we underestimated the effect of aspirin on renal function, because albuminuria is a very early change in diabetic nephropathy. Third, we measured urine dipstick albumin, not by evaluating the urinary albumin-creatinine ratio. Dipstick-positivity may be affected by various conditions, such as sample volume and urine specific gravity. Finally, the JPAD2 cohort study was a cohort study after the RCT period. However, over 80% of patients remained in the original allocation for aspirin as of the 2013 follow-up. The on-treatment analysis and multivariable analysis also showed similar results.

We concluded that low-dose aspirin therapy does not affect eGFR and the incidence of positive urine dipstick albumin during a median follow-up of 8.5 years. Although aspirin is a type of cyclooxygenase inhibitor, our findings suggest that long-term low-dose aspirin therapy is safe with respect to renal function for patients with type 2 diabetes.

Acknowledgments

We thank I. Masuda (Takeda Hospital, Kyoto, Japan) for his commitment to the Committee of Validation of the Data and Events; and M. Ohtorii (Institute for Clinical Effectiveness, Kyoto,

Japan), E. Miyake (Institute for Clinical Effectiveness, Kyoto, Japan), A. Mizutani (Institute for Clinical Effectiveness, Kyoto, Japan), and M. Sakai (Institute for Clinical Effectiveness, Kyoto, Japan) for their roles in data management and analysis. We also thank M. Nagahiro (Kumamoto University, Kumamoto, Japan), Y. Wada (Nara Medical University, Kashihara, Japan), Y. Kamada (Nara Medical University, Kashihara, Japan), and M. Miyagwa (Nara Medical University, Kashihara, Japan) for their secretarial work.

Membership of the JPAD trial / the JPAD2 cohort study

Principal Investigator: Ogawa, Hisao; **Steering Committee:** Saito, Yoshihiko (Chair); Morimoto, Takeshi; Soejima, Hirofumi; Sakuma, Mio; Okada, Sadanori; **Committee on Validation of Data and Events:** Masuda, Izuru (Chair); **Safety Monitoring Board:** Ueshima, Hirotsugu (Chair); Imura, Hiroo; Kimura, Kazuo; **Study Statistician:** Morimoto, Takeshi; **Independent Data Management Center:** Ohtorii, Makiko; Miyake, Eri; Mizutani, Ai; Sakai, Mika.

Independent Statistical Analysis: Dr Morimoto is a statistician at Department of Clinical Epidemiology, Hyogo College of Medicine

JPAD Investigators: Jinnouchi, Hideaki; Hanaoka, Yoichi; Waki, Masako; Kawamura, Kyousuke; Shimabukuro, Michio; Nishiura, Kimiaki; Kawano, Takahiro; Kyoda, Yusuke; Hashiguchi, Jun; Kagoshima, Tadashi; Hanatani, Masakazu; Matsumura, Norihiko; Doi, Naofumi; Nakai, Kenji; Kobayashi, Yoshiyuki; Suzuki, Megumi; Soeda, Tsunenari; Morikawa, Yoshinobu; Horimoto, Masashi; Hasegawa, Atsushi; Yamano, Shigeru; Matsuo, Syuichi; Sakamoto, Yasuhiro; Masuda, Izuru; Yasuno, Akiko; Fujinaga, Yuriko; Horii, Kazuko; Koga, Takeshi; Ogawa, Hiroshi; Ozaki, Ken; Ikemura, Makoto; Hayashi, Motomu; Yabuta, Ikuo; Sugihara, Kiyotaka; Yazaki, Akihiro; Masuda, Joji; Nishitani, Yoshiharu; Naito, Masaki; Ote, Shigenobu; Yamada, Kazuhiko; Wakabayashi, Chikashi; Fukuoka, Yoshiaki; Mahara, Keiji; Kan, Hirofumi; Oshima, Eiji; Sutani, Toshio; Hoda, Koichi; Sawai, Koryo; Yamaga, Kenichi; Nakamura, Tomoki; Okamoto, Shinya; Horie, Hiroaki; Ashihara, Kenichi; Miki, Hiroshi; Makino, Hisaharu; Odo, Takafumi; Iseri, Yoshihisa (retired in 2007); Tanaka, Hiroyuki; Marutsuka, Kousuke; Nakatani, Akira; Murakami, Hironori; Shioya, Yoshiko; Horio, Yutaka; Ikeda, Tsuneo; Machii, Kazuo; Kamura, Masanori; Ban, Keiichiro; Fujii, Yoshihiro; Nishimoto, Kazuo; Misugi, Susumu; Munakata, Tetsuo; Yoshimura, Katsutoshi; Minami, Shigetoshi; Nakashima, Takao; Ogata, Hirofumi; Hifumi, Atuko; Sakurai, Nobuko; Tsurusaki, Ryuichiro; Yamanaka, Yoshito; Yokota, Hiromitsu; Ichihara, Seishi; Yoshinari, Motoki; Sawada, Yoko; Kawashima, Eiji; Goto, Kazuo; Kinoshita, Yoshimi; Kikukawa, Masao; Yamada, Hiroharu; Tanaka, Yuya; Kiyota, Mayumi; Kimura, Yoshihiro; Morikami, Yasuhiro; Fukuda, Masahiro; Takami, Takeshi; Nakatani, Fumihiko; Naomi, Shojiro; Nasu, Toshiaki (retired in 2006); Sawada, Tomohiro; Minagawa, Fuyuki; Haraguchi, Osamu; Kondo, Norifumi; Shono, Hiruyuki; Sugiyama, Hiromichi; Matsuo, Takeshi; Takaoka, Minoru; Nakajima, Tamio; Toihata, Masamitsu; Matsuyama, Kozaburo; Komori, Kenichi; Tsubokura, Toshio; Taguchi, Madoka; Hiramori, Yuko; Okubo, Hiroto; Iemura, Akihiro; Doi, Osamu; Ogihara, Masayuki; Misumi, Kenji; Seo, Koji; Iwai, Ken; Naito, Masatoshi; Ozawa, Seiji; Minoda, Kotaro; Fujii, Hiromi; Miwa, Kimiaki; Egusa, Genshi; Yasuda, Isao; Ueda, Michiaki; Miyata, Junichi; Yoshimura, Midori; Uemyama, Masakuni; Watanabe, Katumi; Haraguchi, Yoshikuni; Tanazawa, Satoshi; Osamura, Yoshiaki; Shibata, Junji; Ono, Takashi; Kamijikkoku, Syuichi; Yoshimoto, Kazumi; Kinuwaki, Etsuo; Kozuma, Kazuo; Onoue, Kenji; Nakano, Yukitaka; Abe, Nanami; Araki, Haruo; Takaoka, Kyoji; Imamoto, Chieko; Suefuji, Hisakazu; Sugimoto, Keisuke; Matsunaga, Terufumi; Oya, Akiko; Onishi, Yoko; Kajiwara, Keizo; Ikuno, Tetsuo; Doi, Michiaki; Igaki, Toshiro; Bando, Hiroshi; Ogura, Tateo; Doijiri, Kenichi; Iwaoka, Taisuke; Akaoshi, Kazunobu; Obata, Kenji; Shimono, Hisashi; Tsuda, Kaoru; Yumoto, Shinya Oka, Keishiro;

Hasegawa, Hironori; Fujimoto, Hisao; Atsumi, Toshiya; Matsutani, Akira; Katsuyama, Yohiyuki; Fukami, Ryo; Iseri, Yoshihisa; Ishibashi, Yutaka; Kudou, Kiyotaka; Kuwahara, Tet-suo; Maeda, Kazutaka; Maki, Akira; Manda, Naiki; Yasue, Hirofumi; Mizuno, Yuji; Momosaki, Sueo; Tokube, Koji; Tomita, Fumishi; Tsuchiya, Tatsuaki; Yabuta, Matahiro; Yamada, Hidei.

Author Contributions

Conceived and designed the experiments: SO TM HO MS HS MN YS. Analyzed the data: SO TM YS. Contributed reagents/materials/analysis tools: SO HO HS HJ MW YA HI YS. Wrote the paper: SO TM YS.

References

1. Antithrombotic Trialists C, Baigent C, Blackwell L, Collins R, Emberson J, Godwin J, et al. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet*. 2009; 373(9678):1849–60. doi: [10.1016/S0140-6736\(09\)60503-1](https://doi.org/10.1016/S0140-6736(09)60503-1) PMID: [19482214](https://pubmed.ncbi.nlm.nih.gov/19482214/)
2. Suleyman H, Demircan B, Karagoz Y. Anti-inflammatory and side effects of cyclooxygenase inhibitors. *Pharmacol Rep*. 2007; 59(3):247–58. PMID: [17652824](https://pubmed.ncbi.nlm.nih.gov/17652824/)
3. Morlans M, Laporte JR, Vidal X, Cabeza D, Stolley PD. End-stage renal disease and non-narcotic analgesics: a case-control study. *British journal of clinical pharmacology*. 1990; 30(5):717–23. PMID: [2271370](https://pubmed.ncbi.nlm.nih.gov/2271370/)
4. Forel CM, Ejerblad E, Lindblad P, Fryzek JP, Dickman PW, Signorello LB, et al. Acetaminophen, aspirin, and chronic renal failure. *The New England journal of medicine*. 2001; 345(25):1801–8. PMID: [11752356](https://pubmed.ncbi.nlm.nih.gov/11752356/)
5. Ibanez L, Morlans M, Vidal X, Martinez MJ, Laporte JR. Case-control study of regular analgesic and nonsteroidal anti-inflammatory use and end-stage renal disease. *Kidney international*. 2005; 67(6):2393–8. PMID: [15882284](https://pubmed.ncbi.nlm.nih.gov/15882284/)
6. Sandler DP, Smith JC, Weinberg CR, Buckalew VM Jr, Dennis VW, Blythe WB, et al. Analgesic use and chronic renal disease. *The New England journal of medicine*. 1989; 320(19):1238–43. PMID: [2651928](https://pubmed.ncbi.nlm.nih.gov/2651928/)
7. Perneger TV, Whelton PK, Klag MJ. Risk of kidney failure associated with the use of acetaminophen, aspirin, and nonsteroidal antiinflammatory drugs. *The New England journal of medicine*. 1994; 331(25):1675–9. PMID: [7969358](https://pubmed.ncbi.nlm.nih.gov/7969358/)
8. Rexrode KM, Buring JE, Glynn RJ, Stampfer MJ, Youngman LD, Gaziano JM. Analgesic use and renal function in men. *JAMA*. 2001; 286(3):315–21. PMID: [11466097](https://pubmed.ncbi.nlm.nih.gov/11466097/)
9. Kurth T, Glynn RJ, Walker AM, Rexrode KM, Buring JE, Stampfer MJ, et al. Analgesic use and change in kidney function in apparently healthy men. *American journal of kidney diseases: the official journal of the National Kidney Foundation*. 2003; 42(2):234–44.
10. Curhan GC, Knight EL, Rosner B, Hankinson SE, Stampfer MJ. Lifetime nonnarcotic analgesic use and decline in renal function in women. *Archives of internal medicine*. 2004; 164(14):1519–24. PMID: [15277282](https://pubmed.ncbi.nlm.nih.gov/15277282/)
11. Kuo HW, Tsai SS, Tiao MM, Liu YC, Lee IM, Yang CY. Analgesic use and the risk for progression of chronic kidney disease. *Pharmacoepidemiology and drug safety*. 2010; 19(7):745–51. doi: [10.1002/pds.1962](https://doi.org/10.1002/pds.1962) PMID: [20582905](https://pubmed.ncbi.nlm.nih.gov/20582905/)
12. Evans M, Forel CM, Bellocco R, Fitzmaurice G, Fryzek JP, McLaughlin JK, et al. Acetaminophen, aspirin and progression of advanced chronic kidney disease. *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association—European Renal Association*. 2009; 24(6):1908–18.
13. Agodoa LY, Francis ME, Eggers PW. Association of analgesic use with prevalence of albuminuria and reduced GFR in US adults. *American journal of kidney diseases: the official journal of the National Kidney Foundation*. 2008; 51(4):573–83.
14. Smith SC Jr., Benjamin EJ, Bonow RO, Braun LT, Creager MA, Franklin BA, et al. AHA/ACC Secondary Prevention and Risk Reduction Therapy for Patients with Coronary and other Atherosclerotic Vascular Disease: 2011 update: a guideline from the American Heart Association and American College of Cardiology Foundation. *Circulation*. 2011; 124(22):2458–73. doi: [10.1161/CIR.0b013e318235eb4d](https://doi.org/10.1161/CIR.0b013e318235eb4d) PMID: [22052934](https://pubmed.ncbi.nlm.nih.gov/22052934/)
15. Pignone M, Alberts MJ, Colwell JA, Cushman M, Inzucchi SE, Mukherjee D, et al. Aspirin for primary prevention of cardiovascular events in people with diabetes: a position statement of the American

Diabetes Association, a scientific statement of the American Heart Association, and an expert consensus document of the American College of Cardiology Foundation. *Circulation*. 2010; 121(24):2694–701. doi: [10.1161/CIR.0b013e3181e3b133](https://doi.org/10.1161/CIR.0b013e3181e3b133) PMID: [20508178](https://pubmed.ncbi.nlm.nih.gov/20508178/)

16. Adler AI, Stevens RJ, Manley SE, Bilous RW, Cull CA, Holman RR, et al. Development and progression of nephropathy in type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS 64). *Kidney international*. 2003; 63(1):225–32. PMID: [12472787](https://pubmed.ncbi.nlm.nih.gov/12472787/)
17. American Diabetes A. Standards of medical care in diabetes—2014. *Diabetes Care*. 2014; 37 Suppl 1: S14–80. doi: [10.2337/dc14-S014](https://doi.org/10.2337/dc14-S014) PMID: [24357209](https://pubmed.ncbi.nlm.nih.gov/24357209/)
18. Mogensen CE, Christensen CK. Predicting diabetic nephropathy in insulin-dependent patients. *The New England journal of medicine*. 1984; 311(2):89–93. PMID: [6738599](https://pubmed.ncbi.nlm.nih.gov/6738599/)
19. Ogawa H, Nakayama M, Morimoto T, Uemura S, Kanauchi M, Doi N, et al. Low-dose aspirin for primary prevention of atherosclerotic events in patients with type 2 diabetes: a randomized controlled trial. *JAMA*. 2008; 300(18):2134–41. doi: [10.1001/jama.2008.623](https://doi.org/10.1001/jama.2008.623) PMID: [18997198](https://pubmed.ncbi.nlm.nih.gov/18997198/)
20. Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations for estimated GFR from serum creatinine in Japan. *American journal of kidney diseases: the official journal of the National Kidney Foundation*. 2009; 53(6):982–92.
21. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. Heart Outcomes Prevention Evaluation Study Investigators. *Lancet*. 2000; 355(9200):253–9. PMID: [10675071](https://pubmed.ncbi.nlm.nih.gov/10675071/)
22. Parving HH, Lehnert H, Brochner-Mortensen J, Gomis R, Andersen S, Arner P, et al. The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *The New England journal of medicine*. 2001; 345(12):870–8. PMID: [11565519](https://pubmed.ncbi.nlm.nih.gov/11565519/)
23. Hopper AH, Tindall H, Davies JA. Administration of aspirin-dipyridamole reduces proteinuria in diabetic nephropathy. *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association—European Renal Association*. 1989; 4(2):140–3.
24. Hansen HP, Gaede PH, Jensen BR, Parving HH. Lack of impact of low-dose acetylsalicylic acid on kidney function in type 1 diabetic patients with microalbuminuria. *Diabetes Care*. 2000; 23(12):1742–5. PMID: [11128344](https://pubmed.ncbi.nlm.nih.gov/11128344/)
25. Gaede P, Hansen HP, Parving HH, Pedersen O. Impact of low-dose acetylsalicylic acid on kidney function in type 2 diabetic patients with elevated urinary albumin excretion rate. *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association—European Renal Association*. 2003; 18(3):539–42.