



# HHS Public Access

Author manuscript

*Acad Pediatr.* Author manuscript; available in PMC 2016 January 26.

Published in final edited form as:

*Acad Pediatr.* 2012 ; 12(4): 269–282. doi:10.1016/j.acap.2012.02.004.

## Interventions to Improve Screening and Follow-Up in Primary Care: A Systematic Review of the Evidence

Jeanne Van Cleave, MD<sup>1</sup>, Karen A. Kuhlthau, PhD<sup>1</sup>, Sheila Bloom, MS<sup>1</sup>, Paul W. Newacheck, DrPH<sup>2</sup>, Alixandra A. Knapp, MS<sup>1</sup>, Charles J. Homer, MD, MPH<sup>3,4</sup>, and James M. Perrin, MD<sup>1</sup>

<sup>1</sup>The Center for Child and Adolescent Health Policy, MassGeneral Hospital for Children, Boston, MA

<sup>2</sup>Philip R. Lee Institute for Health Policy Studies, University of California San Francisco

<sup>3</sup>National Initiative for Children's Healthcare Quality

<sup>4</sup>Department of Pediatrics, Harvard Medical School

### Abstract

**Background**—The American Academy of Pediatrics and other organizations recommend several screening tests as part of preventive care. The proportion of children who are appropriately screened and who receive follow-up care is low.

**Objective**—To conduct a systematic review of the evidence for practice-based interventions to increase the proportion of patients receiving recommended screening and follow-up services in pediatric primary care.

**Data source**—Medline database of journal citations.

**Study eligibility criteria, participants, and interventions**—We developed a strategy to search Medline to identify relevant articles. We selected search terms to capture categories of conditions (e.g., developmental disabilities, obesity), screening tests, specific interventions (e.g., quality improvement initiatives, electronic records enhancements), and primary care. We searched references of selected articles and reviewed articles suggested by experts. We included all studies with a distinct, primary care-based intervention and post-intervention screening data, and studies that focused on children and young adults (< 21 years of age). We excluded studies of newborn screening.

**Study appraisal and synthesis methods**—Abstracts were screened by 2 reviewers and articles with relevant abstracts received full text review and evaluated for inclusion criteria. A structured tool was used to abstract data from selected articles. Because of heterogeneous interventions and outcomes, we did not attempt a meta-analysis.

---

No conflicts of interest

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Results**—From 2547 returned titles and abstracts, 23 articles were reviewed. Nine were pre-post comparisons, 5 were randomized trials, 3 were post-intervention comparisons with a control group, 3 were post-intervention cross-sectional analyses only, and 3 reported time series data. Of 14 articles with pre-intervention or control group data and significance testing, 12 reported increases in the proportion of patients appropriately screened. Interventions were heterogeneous and often multifaceted, and several types of interventions, such as provider/staff training, electronic medical record templates/prompts, and learning collaboratives, appeared effective in improving screening quality. Few articles described interventions to track screening results or referral completion for those with abnormal tests. Data were often limited by single-site, non-randomized design.

**Conclusions**—Several feasible, practice- and provider-level interventions appear to increase the quality of screening in pediatric primary care. Evidence for interventions to improve follow-up of screening tests is scant. Future research should focus on which specific interventions are most effective, whether effects are sustained over time, and what interventions improve follow-up of abnormal screening tests.

### MeSH key words

Mass Screening; Preventive Health Services; Physician's Practice Patterns; Quality of Health Care

---

## Introduction

Prevention of mortality and morbidity secondary to many conditions depends on effective screening and referral procedures in pediatric primary care.<sup>1</sup> For many conditions, such as iron-deficiency anemia, autistic spectrum disorder, and vision and hearing problems, early detection from broad-based, primary screening with timely follow-up care enables children with these conditions to receive treatment that affects long-term health outcomes. The American Academy of Pediatrics, *Bright Futures*, and other organizations recommend screening procedures for several specific conditions.<sup>2,3</sup>

Although many children receive some screening via public health or school-based mechanisms, most screening beyond the newborn period occurs within the context of the primary care office at well-child visits. Even with clear, readily-accessible recommendations, quality of screening in primary care is sub-optimal,<sup>4</sup> leaving children at risk when conditions are not identified. Reasons for this quality gap include lack of knowledge of recommendations,<sup>5,6</sup> presumed patient refusal,<sup>5</sup> lack of time,<sup>6</sup> lack of office staff support,<sup>6</sup> inadequate reimbursement,<sup>7</sup> and inadequate referral resources for those found to have a problem detected through screening.<sup>7</sup>

Several interventions have potential to improve screening in primary care settings<sup>8</sup> and have been studied to some extent in adults.<sup>9</sup> However, which practice-level interventions are most effective for improving screening in pediatric primary care is not known. Interventions in pediatrics may have a different impact compared to adult populations, for several reasons.<sup>10</sup> First, children generally seek health care and make decisions through a proxy, usually a parent. Second, children undergo more rapid developmental changes, and screening recommendations change with each well-child visit. Third, most conditions for which

children are screened are not thought of as potentially life-threatening, in contrast to cancer screening in adults, which may affect the importance providers and parents place on screening in children. Examining interventions that improve receipt of recommended screening in pediatrics may help physicians and policymakers identify changes most likely to benefit a broader population and may inform a research agenda to address questions about how to improve the quality of screening in pediatric practices.

We undertook this systematic review as part of a larger project to examine evidence regarding six core objectives of the Maternal and Child Health Bureau<sup>11</sup> for care for children with special health care needs. Previously, we reviewed the evidence regarding receipt of family-centered care<sup>12</sup> and services to transition to adult providers,<sup>13</sup> having a medical home;<sup>14</sup> and having adequate health insurance coverage.<sup>15</sup> We now review evidence for the objective that all children are screened early and continuously for special health care needs. Because high-quality screening in primary care is necessary for objective, we focused our review on office-based interventions to increase the proportion of children receiving recommended screening. Our specific research question was, what is the evidence for interventions to improve such screening in primary care settings? As a secondary objective, we also examined interventions to improve follow-up or referral completion, once screening tests identified concerns.

## Methods

To guide our search strategy (Table 1), we constructed a logic model<sup>16</sup> (Figure 1) that depicts the health conditions for which screening tests are recommended, interventions, and outcomes of interest. In developing and refining the model, we held a conference with relevant experts, including policymakers, family advocates, and researchers in the field of improving care for children with special health care needs. The purpose of this panel was to guide the systematic reviews around the MCHB core objectives, and the panel discussed and made recommendations for our logic model and search strategy.

### Screening tests

To select the screening tests and corresponding specific conditions for inclusion in our search, we reviewed recommendations for preventive care screening from Bright Futures/<sup>17</sup> American Academy of Pediatrics, the US Preventive Services Task Force, and the Centers for Disease Control. We selected screening tests for conditions such as developmental delay, mental health conditions, vision problems, hearing problems, lead poisoning, anemia, hypertension, sexually transmitted infections, and obesity. We did not include conditions detected by newborn screening or prenatal screening, since testing procedures and much of the follow-up occurs not in primary care but in hospitals and in conjunction with state public health authorities.

### Interventions

We chose search terms to capture primary care interventions designed to improve receipt of recommended screening and follow up. Specific activities were derived from a review of the

literature of interventions to improve quality of other functions of primary care practices (e.g., vaccination) and recommendations from our expert panel.

Interventions included practice-level initiatives such as provider/staff education sessions and materials, quality improvement initiatives, and improvements in office workflow. Our search included interventions to improve patient identification for screening, particularly changes that led to automated identification, such as chart flagging, electronic medical record (EMR) reminders, and patient registries. We also searched for interventions that involved pay-for-performance initiatives targeted toward screening.

## Outcomes

Our primary outcomes were the proportion of children appropriately screened, and proportion of children with abnormal screening results who received follow-up care. Appropriateness of screening was determined by the individual studies. Because follow-up care can vary among patients due to family preferences and available referral options, we broadly defined follow up care as any action by the provider that would advance a plan for additional screening, evaluation or treatment prompted by an abnormal result. This definition included discussing abnormal results with parents and patients, retesting patients, and referring to specialists or community resources for further treatment or evaluation. We also included search terms to capture secondary outcomes derived from the Institute of Medicine domains of healthcare quality.<sup>17</sup>

## Database search

We conducted a systematic search of Medline (Jan 1961–Aug 2010) for titles and abstracts relevant to our research question. We queried for articles containing MeSH terms in each of the columns in Table 1, i.e., containing terms that represented a condition, a setting, and an outcome/intervention. We also reviewed bibliographies of selected articles, as well as bibliographies of review articles related to our search. For the bibliography reviews, when we found a potentially relevant title that was missed during the previous search, we obtained the article's Medical Subject Heading (MeSH) terms from the Medline citation to determine why the article was missed. We then refined the search to include omitted MeSH terms, reran the search and reviewed the additional abstracts. We limited our search to English-language articles studying children and youth aged 0–18 years.

## Selection of articles

Two reviewers (JV and AAK) screened titles and abstracts for inclusion in the group of articles for full-text review. Abstracts were selected if the study examined a recommended screening practice and the study was performed in a primary care setting in the United States. Some returned studies included both adults and adolescents, and we included articles if >50% of participants were under age 21 years. Abstracts that lacked detail to make this determination also underwent full-text review. If the abstract was not appropriate for inclusion in the review but possibly referenced relevant articles, the full-text version was obtained and the bibliography scanned. The reviewers met to resolve discrepancies by discussion and mutual agreement. Each reviewer then abstracted a subset of articles using a structured form to report interventions, populations, settings, and outcomes. After

abstraction, reviewers finalized the list of articles to be included in the review through discussion and agreement. Reviewers overlapped on a random selection of approximately 20% of abstracted articles. Abstractions were qualitatively reviewed to assess for agreement, and abstracted screening rates and descriptions of the interventions were verified through a second review of the full text articles. We did not contact authors of the studies for further details. No formal assessment of study quality was done using standardized tools, but we grouped studies using a hierarchy of study design quality (e.g., RCTs, designs with control groups, and uncontrolled studies) and reported elements of potential bias in our description of the studies.

### Specific categories of excluded studies

We excluded studies to validate screening tools and studies that documented poor-quality screening or follow-up without interventions. We also excluded studies that assessed only feasibility of screening in primary care practices without specific attention to long-term, generalizable changes within the practice (e.g., studies where the intervention was limited to research assistants performing screening procedures). We excluded articles that lacked explicit outcomes related screening or follow-up care.

## Results

The final search strategy identified 2547 titles (Figure 2). After reviewing titles and abstracts, 105 articles underwent full-text review. Eight articles that underwent full-text review were initially identified from bibliographies of selected articles. Reviewers completed data abstraction for 29 of the 105 full-text articles. Of these 29 articles, 23 met criteria for inclusion in the final review (Table 2). Common reasons for exclusion were because no intervention was tested, proportion of patients screened was not measured, or the patient population was primarily adult-aged. The included 23 articles were 5 randomized controlled trials and 18 observational studies. Among the randomized trials, the practice was usually the unit of randomization. Among the observational studies, 9 used pre-post designs, 3 were post-intervention comparisons with a concurrent control group, 3 reported findings using time-series design where the outcome was measured at regular intervals after the intervention was initiated, and 3 were post-intervention, cross-sectional analyses with no comparison group. The diversity of interventions and outcomes prevented any meta-analysis.

### Types of interventions

The studies described several different types of interventions. The most common interventions were 1) changes to office systems, usually part of a formal quality improvement program such as a learning collaborative, 2) physician and staff education, sometimes facilitated by a “physician champion” of a specific screening test, 3) electronic medical record enhancements (e.g., prompts), and 4) distribution of additional tools for physicians to use when screening or counseling patients. Many studies combined intervention types. In some studies where several practices were enrolled in a quality improvement initiative, specific changes were chosen by each practice. In several studies,

quality of preventive care screening was measured along with other preventive care outcomes (e.g., immunizations, preventive care visit attendance, etc).

Twelve articles from ten separate studies<sup>18–29</sup> used interventions based largely on learning collaborative methods, including plan-do-study-act cycles and facilitated contact with other intervention practices. Typically, small teams of practitioners and staff from intervention practices addressed barriers related to office system design, provider and staff knowledge gaps, and workflow. Specific changes included chart flagging or routine chart review by non-physician staff to identify patients behind in testing. For some studies, multiple practices participated, multiple screening tests and other preventive care elements were targeted for improvement, and practices were at liberty to choose from several recommended changes those they deemed most likely to work in their practice. Thus, the specific changes associated with the global intervention varied among individual practices. Post-intervention screening ranged from 39–94% of patients screened appropriately. Improvement from baseline varied widely, from 0–80%. Improvement tended to be greater if pre-intervention screening was low or non-existent and if the focus of the intervention was narrowed to specific screening tests or a specific area, such as the study reported by King et al. from a learning collaborative on developmental screening and services.<sup>24</sup>

Five articles<sup>30–34</sup> described interventions to implement screening using provider training and/or tools for facilitating conversations with parents, such as provider sheets to prompt screening questions or patient questionnaires. These interventions focused on screening for obesity, developmental or mental health problems, or adolescent risky behaviors. Post-intervention screening ranged from 28% (for BMI calculations)<sup>32</sup> to 94% (vision screening).<sup>34</sup>

Two articles<sup>35, 36</sup> examined associations between implementing the Healthy Steps program and screening. Healthy Steps is designed for first-time parents and provides co-located developmental specialists to enhance well-child visits.<sup>35</sup> Parents also receive home visits, telephone access for developmental questions, written materials, and linkages to community resources. Screening of patients enrolled in Healthy Steps was compared to screening of same-aged patients not enrolled in Healthy Steps (e.g., second-born children) after implementation. Screening for lead poisoning and anemia did not markedly change, but developmental screening doubled, from 41–43% to 82–84%.

Three studies<sup>27, 37, 38</sup> examined the effect of EMR enhancements, such as EMR templates and reminders, with varying results. With EMR templates to prompt providers to elicit developmental concerns, screening improved to 65–73% of patients for various areas of development, significant increases from baseline.<sup>37</sup> EMR reminders enabled near universal screening (99%) of patients if providers were able to obtain lead levels at the visit, but only 41% for patients required by insurance to have levels drawn off-site.<sup>38</sup> For Chlamydia screening, reminders had no effect compared to patient charts without reminders.<sup>27</sup>

In two studies,<sup>39, 40</sup> a nurse and a nurse practitioner were employed to identify and track patients in need of screening. Both interventions involved protocols for identifying and tracking which patients were due for testing or follow up of abnormal tests. Hull et al. found

that a nurse-driven protocol to identify and screen patients was highly effective and achieved essentially universal screening in one practice.<sup>39</sup> Block et al. found that a similar intervention achieved improved documentation of a follow up plan for elevated lead levels, but smaller improvements for follow-up testing and parent education.<sup>40</sup>

### **Interventions to increase follow up of abnormal screening results**

We found little evidence about interventions to improve post-visit follow-up or referral completion, once screening tests identified concerns. As mentioned, Block et al.<sup>40</sup> examined the effect of a nurse-driven protocol to increase retesting and parent education for abnormal lead levels. Retesting increased to 65% of those with abnormal levels, and 32% of families with persistently high levels received education. Two other studies<sup>31, 33</sup> examined discussion with patients and parents following screening tests for behavior problems or risky behaviors. Both studies found that patient/provider handouts facilitated discussion of problems detected using formal assessment tools. Schonwald et al.<sup>30</sup> demonstrated that referrals for developmental evaluation remained the same, despite increases in use of formal screening tools.

### **Discussion**

Three key findings emerged from this review of interventions to improve the quality of preventive care screening in pediatric primary care settings. First, most studies reported improved quality of screening post-intervention, usually a modest improvement, although differences were variable across and within studies. Second, because of variable findings, heterogeneous interventions, and relatively few studies with control groups, we could not discern whether a particular type or form of intervention is superior for improving screening. However, we saw patterns where successful interventions tended to emphasize collaborative learning, office-systems changes, and tracking progress over time. Third, we found few interventions that aimed to improve follow-up of abnormal screening results, which offers opportunities for further investigation.

From the articles reviewed, we found screening in pediatric offices generally improved after interventions were implemented. In studies where pre- and post-intervention outcomes with statistical testing were reported, over 80% of interventions demonstrated improvement in at least one area of screening. However, results varied, ranging from no change to an 8-fold increase in the proportion of children screened, and many studies could not control for secular trend with their study designs. The magnitude of the impact of interventions seemed greater when pre-intervention screening was low, and multi-faceted interventions implemented through a learning collaborative structure appeared to be, of all intervention types, more robustly studied and relatively effective. Otherwise, this review identified little regarding the patterns of variable effects or reasons for them, including type of screening or type of intervention. In addition, results varied among practices implementing similar interventions; even when an intervention was introduced in multiple practices as a single study, effects typically varied from practice to practice. No study objectively measured contextual factors (e.g., practice's motivation to change, staff capacity for the intervention),

although some studies included qualitative discussion on contextual reasons for variability in findings across practices (e.g., physician champion left the practice).

With the exception of four studies, fewer than 85% of patients were appropriately screened post-intervention, with most studies reporting post-intervention screening between 50–75%. This finding, which mirrors findings in adult studies,<sup>41</sup> suggests that some patients miss screening despite often intensive office-based improvements. Studies in our review that examined characteristics of patients who were not screened found various associations with less screening, including non-English speaking parents, parents who did not have time to complete the screening tool before seeing the physician, and having to go off-site to complete screening tests.<sup>30, 37, 38</sup> Furthermore, this finding suggests a “ceiling effect” similar to that found with interventions to increase rates of vaccine coverage and well-child visit attendance.<sup>42, 43</sup>

The quality of the studies varied, with many using non-randomized study designs, a limited number of practice sites, and with little account for context of the practices receiving intervention. However, five articles reported on randomized trials with consistent positive effects. Most studies were pre-post designs without randomization, and some lacked comparison groups, making it difficult to assess the effect of natural trends over time. Most studies involved multiple practices, but seven studies used only one practice site, limiting the ability to draw conclusions about how broader-based improvement efforts would increase the quality of screening. Because office staff motivation and technological savvy can play a large role in the success of interventions,<sup>44</sup> practices differing in these contextual factors would likely have different results.

Most interventions were multifaceted, involving several alterations in office workflow, physician and staff education, and changes in staff time allocation. While multifaceted interventions generally had more success, as did interventions tailored to best fit specific practices, no systematic approach examined which elements provide the greatest benefit, or why the same intervention performed better in some practices than others. Findings from such a systematic approach could be used to design more efficient interventions and advance the field of quality improvement research.

Few studies examined the quality of follow-up care, and few interventions contained elements specifically targeting follow-up of abnormal tests. However, the few studies that did have follow-up as an outcome found 35–65% of patients did not receive follow up care after an abnormal screening result. This finding indicates the need to include outcomes related to follow-up in studies of screening, and that measuring screening alone may overestimate changes in identification and treatment of conditions.

We found no studies testing the effects of performance incentives or physician feedback. This strategy has been studied more in adult settings for screening<sup>9, 45</sup> and in pediatrics for immunizations, attendance at well-child visits, and management of chronic conditions.<sup>46</sup> Another review of adult cancer screening interventions focused on motivating patients and reducing barriers to care.<sup>47</sup> These reviews found variable effects among similar interventions, with most interventions associated with some increase in screening.



The review has several limitations. Many quality improvement interventions do not reach publication, which could have limited identification of informative studies. The search terms used may not have captured all relevant studies, particularly studies examining quality of follow-up care, for which search terms were difficult to define. Many studies tested heterogeneous interventions that were modified for each practice; some interventions were multifaceted so that practices could choose specific elements to implement. This “cafeteria” approach makes comparing interventions in separate studies difficult and may limit reliability and generalizability. However, tailoring the intervention to the context of the practice likely increased the chance of the desired effect, and is more representative of how it would be applied in actual practice.

## Conclusion

Although the quality of studies varied, we found a moderate level of evidence that interventions are effective in improving screening in pediatric practices. This review also reveals several avenues for future study that will guide policy makers and practitioners in what specific interventions provide the most value.

Interventions reviewed here appeared to have ceiling effects, which invites the question, given the broad aims of pediatric primary care, what should be the goals for screening, and is there a point of diminishing return where a practice’s extra efforts exceed the value of the gain? Policies around reimbursement based on screening performance should match the right amount of effort to achieve the right rate. Also, improving screening rates from a high baseline will likely require different interventions; near-perfect screening may not be achievable without a large degree of automation and standardization and multiple layers of double-checks performed by non-clinicians or through electronic mechanisms. Lastly, when aiming for high proportions of children appropriately screened, defining the right denominator becomes increasingly important and worth measuring accurately and thoughtfully. A denominator measured by well child visits, versus empanelled patients, might drive different interventions with ultimately different outcomes.

No single type of intervention arose as consistently more effective in increasing screening quality, and few studies addressed the critical issue of assuring adequate follow-up. This review did not identify specific interventions that work better than others, however multifaceted, practice-tailored interventions with ongoing outcome assessment seemed to be effective, and most comprehensively evaluated. Policies supporting such interventions broadly will likely lead to earlier detection and more effective treatment for a large population of children. Quality improvement activities are now required for maintenance of board certification, and many local health systems and payers ask or require practices to participate. Medical societies, such as the American Academy of Pediatrics, can help provide infrastructure to encourage efforts by individual practices.

This review leaves several additional questions: Which components of interventions add to effectiveness, and which are ineffective? What interventions improve follow-up care? How sustainable are the effects of these interventions? Are different interventions more effective for different types of screening procedures (e.g., questionnaires versus blood draws)? How is

practice context best measured, and how is it associated with the success of interventions? Such future avenues for research will help refine interventions to move toward effective, efficient screening in primary care pediatrics.

## Acknowledgments

Funding: Maternal and Child Health Bureau, cooperative agreement # 5 U53MC04473-03-00

We are grateful to the Maternal and Child Health Bureau, who provided funding for this study (Cooperative agreement # 5 U53MC04473-03-00).

## Abbreviations

<b>MCHB</b>	Maternal and Child Health Bureau
<b>HS</b>	Healthy Steps
<b>LC</b>	Learning collaborative
<b>BMI</b>	Body mass index
<b>BP</b>	Blood pressure
<b>QI</b>	Quality improvement
<b>HMO</b>	Health maintenance organization
<b>PEDS</b>	Parents' evaluation of developmental status
<b>EMR</b>	Electronic medical record
<b>EPSDT</b>	Early periodic screening, diagnosis and treatment
<b>ASQ</b>	Ages and stages questionnaire
<b>AAP</b>	American Academy of Pediatrics
<b>RCT</b>	Randomized controlled trial

## References

1. Young PC. Prevention: a new focus for the country but old stuff for pediatricians. *Acad Pediatr*. Nov-Dec;2010 10(6):367–368. [PubMed: 21075314]
2. Committee on Practice and Ambulatory Medicine and Bright Futures Steering Committee. Recommendations for Preventive Pediatric Health Care. *Pediatrics*. 2007; 120:1376.
3. AHRQ Publication No 10-05145. Agency for Healthcare Research and Quality, Rockville, MD: Sep. 2010 Guide to Clinical Preventive Services, 2010–2011. <http://www.ahrq.gov/clinic/pocketgd.htm> [Accessed October 26, 2011]
4. Mangione-Smith R. Bridging the quality chasm for children: need for valid, comprehensive measurement tools. *Arch Pediatr Adolesc Med*. Sep; 2007 161(9):909–910. [PubMed: 17768293]
5. Minniar TD, Gilmore B, Arnold SR, Flynn PM, Knapp KM, Gaur AH. Implementation of and barriers to routine HIV screening for adolescents. *Pediatrics*. Oct; 2009 124(4):1076–1084. [PubMed: 19752084]
6. Honigfeld L, McKay K. Barriers to enhancing practice-based developmental services. *J Dev Behav Pediatr*. Feb; 2006 27(1 Suppl):S30–33. discussion S34–37, S50-32. [PubMed: 16715789]

7. Ayres CG, Griffith HM. Perceived barriers to and facilitators of the implementation of priority clinical preventive services guidelines. *Am J Manag Care*. Mar; 2007 13(3):150–155. [PubMed: 17335358]
8. Klabunde CN, Lanier D, Breslau ES, et al. Improving colorectal cancer screening in primary care practice: innovative strategies and future directions. *J Gen Intern Med*. Aug; 2007 22(8):1195–1205. [PubMed: 17534688]
9. Hulscher ME, Wensing M, Grol RP, van der Weijden T, van Weel C. Interventions to improve the delivery of preventive services in primary care. *American Journal of Public Health*. May; 1999 89(5):737–746. [PubMed: 10224987]
10. Forrest CB, Simpson L, Clancy C. Child health services research. Challenges and opportunities. *JAMA*. Jun 11; 1997 277(22):1787–1793. [PubMed: 9178792]
11. Maternal and Child Health Bureau. [Accessed March 1, 2011] Achieving and measuring success: A national agenda for children with special health care needs. <http://mchb.hrsa.gov/programs/specialneeds/measuresuccess.htm>
12. Kuhlthau KA, Bloom S, Van Cleave J, et al. Evidence for family-centered care for children with special health care needs: a systematic review. *Acad Pediatr*. Mar-Apr;11(2):136–143. e138. [PubMed: 21396616]
13. Bloom S, Kuhlthau K, Van Cleave J, Knapp A, Newacheck P, Perrin J. Health Care Transition for Youth with Special Health Care Needs. *Journal of Adolescent Health*. In press.
14. Homer CJ, Klatka K, Romm D, et al. A review of the evidence for the medical home for children with special health care needs. *Pediatrics*. Oct; 2008 122(4):e922–937. [PubMed: 18829788]
15. Newacheck PW, Houtrow AJ, Romm DL, et al. The future of health insurance for children with special health care needs. *Pediatrics*. May; 2009 123(5):e940–947. [PubMed: 19403486]
16. W.K. Kellogg Foundation. [Accessed April 21, 2010] Logic Model Development Guide. Jan. 2004 <http://www.wkcf.org/knowledge-center/resources/2010/Logic-Model-Development-Guide.aspx>
17. Committee on Quality of Health Care in America, Institute of Medicine. . *Crossing the Quality Chasm: A New Health System for the 21st Century*. National Academies Press; 2001.
18. Young PC, Glade GB, Stoddard GJ, Norlin C. Evaluation of a learning collaborative to improve the delivery of preventive services by pediatric practices. *Pediatrics*. May; 2006 117(5):1469–1476. [PubMed: 16651299]
19. Shaw JS, Wasserman RC, Barry S, et al. Statewide quality improvement outreach improves preventive services for young children. *Pediatrics*. Oct; 2006 118(4):e1039–1047. [PubMed: 16940162]
20. Margolis PA, Lannon CM, Stuart JM, Fried BJ, Keyes-Elstein L, Moore DE Jr. Practice based education to improve delivery systems for prevention in primary care: randomised trial. *BMJ*. Feb 14.2004 328(7436):388. [PubMed: 14766718]
21. Lannon CM, Flower K, Duncan P, Moore KS, Stuart J, Bassewitz J. The Bright Futures Training Intervention Project: implementing systems to support preventive and developmental services in practice. *Pediatrics*. Jul; 2008 122(1):e163–171. [PubMed: 18595961]
22. Bordley WC, Margolis PA, Stuart J, Lannon C, Keyes L. Improving preventive service delivery through office systems. *Pediatrics*. Sep.2001 108(3):E41. [PubMed: 11533359]
23. Shafer MA, Tebb KP, Pantell RH, et al. Effect of a clinical practice improvement intervention on Chlamydial screening among adolescent girls. *JAMA*. Dec 11; 2002 288(22):2846–2852. [PubMed: 12472326]
24. King TM, Tandon SD, Macias MM, et al. Implementing developmental screening and referrals: lessons learned from a national project. *Pediatrics*. 2010 Feb; 125(2):350–360. [PubMed: 20100754]
25. Polacsek M, Orr J, Letourneau L, et al. Impact of a primary care intervention on physician practice and patient and family behavior: keep ME Healthy---the Maine Youth Overweight Collaborative. *Pediatrics*. Jun; 2009 123( Suppl 5):S258–266. [PubMed: 19470601]
26. Pomietto M, Docter AD, Van Borkulo N, Alfonsi L, Krieger J, Liu LL. Small steps to health: building sustainable partnerships in pediatric obesity care. *Pediatrics*. 2009 Jun; 123( Suppl 5):S308–316. [PubMed: 19470608]

27. Scholes D, Grothaus L, McClure J, et al. A randomized trial of strategies to increase chlamydia screening in young women. *Prev Med.* 2006 Oct; 43(4):343–350. [PubMed: 16782182]
28. Earls MF, Hay SS. Setting the stage for success: implementation of developmental and behavioral screening and surveillance in primary care practice--the North Carolina Assuring Better Child Health and Development (ABCD) Project. *Pediatrics.* Jul; 2006 118(1):e183–188. [PubMed: 16818532]
29. Tebb KP, Wibbelsman C, Neuhaus JM, Shafer MA. Screening for asymptomatic Chlamydia infections among sexually active adolescent girls during pediatric urgent care. *Arch Pediatr Adolesc Med.* Jun; 2009 163(6):559–564. [PubMed: 19487613]
30. Schonwald A, Huntington N, Chan E, Risko W, Bridgemohan C. Routine developmental screening implemented in urban primary care settings: more evidence of feasibility and effectiveness. *Pediatrics.* Feb; 2009 123(2):660–668. [PubMed: 19171635]
31. Ozer EM, Adams SH, Lustig JL, et al. Increasing the screening and counseling of adolescents for risky health behaviors: a primary care intervention. *Pediatrics.* Apr; 2005 115(4):960–968. [PubMed: 15805371]
32. Dunlop AL, Leroy Z, Trowbridge FL, Kibbe DL. Improving providers' assessment and management of childhood overweight: results of an intervention. *Ambul Pediatr.* Nov-Dec;2007 7(6):453–457. [PubMed: 17996840]
33. Applegate H, Kelley ML, Applegate BW, Jayasinghe IK, Venters CL. Clinical case study: pediatric residents' discussions of and interventions for children's behavioral and emotional problems. *J Pediatr Psychol.* Jul-Aug;2003 28(5):315–321. [PubMed: 12808008]
34. Hartmann EE, Bradford GE, Chaplin PK, et al. Project Universal Preschool Vision Screening: a demonstration project. *Pediatrics.* Feb; 2006 117(2):e226–237. [PubMed: 16452332]
35. Minkovitz CS, Hughart N, Strobino D, et al. A practice-based intervention to enhance quality of care in the first 3 years of life: the Healthy Steps for Young Children Program. *JAMA.* Dec 17; 2003 290(23):3081–3091. [PubMed: 14679271]
36. Niederman LG, Schwartz A, Connell KJ, Silverman K. Healthy Steps for Young Children program in pediatric residency training: impact on primary care outcomes. *Pediatrics.* Sep; 2007 120(3):e596–603. [PubMed: 17766500]
37. Adams WG, Mann AM, Bauchner H. Use of an electronic medical record improves the quality of urban pediatric primary care. *Pediatrics.* Mar; 2003 111(3):626–632. [PubMed: 12612247]
38. Gioia PC. Quality improvement in pediatric well care with an electronic record. *Proc AMIA Symp.* 2001:209–213. [PubMed: 11825182]
39. Hull PC, Husaini BA, Tropez-Sims S, Reece M, Emerson J, Levine R. EPSDT preventive services in a low-income pediatric population: impact of a nursing protocol. *Clin Pediatr (Phila).* Mar; 2008 47(2):137–142. [PubMed: 17873239]
40. Block B, Szekely K, Escobar M. Difficulties in evaluating abnormal lead screening results in children. *Journal of the American Board of Family Practice.* Nov-Dec;1996 9(6):405–410. [PubMed: 8923397]
41. Holden DJ, Jonas DE, Porterfield DS, Reuland D, Harris R. Systematic review: enhancing the use and quality of colorectal cancer screening. *Ann Intern Med.* May 18; 152(10):668–676. [PubMed: 20388703]
42. Hambidge SJ, Davidson AJ, Phibbs SL, et al. Strategies to improve immunization rates and well-child care in a disadvantaged population: a cluster randomized controlled trial. *Arch Pediatr Adolesc Med.* Feb; 2004 158(2):162–169. [PubMed: 14757608]
43. Baker AM, McCarthy B, Gurley VF, Yood MU. Influenza immunization in a managed care organization. *J Gen Intern Med.* Jul; 1998 13(7):469–475. [PubMed: 9686713]
44. Feifer C, Nemeth L, Nietert PJ, et al. Different paths to high-quality care: three archetypes of top-performing practice sites. *Ann Fam Med.* May-Jun;2007 5(3):233–241. [PubMed: 17548851]
45. Sabatino SA, Habarta N, Baron RC, et al. Interventions to increase recommendation and delivery of screening for breast, cervical, and colorectal cancers by healthcare providers systematic reviews of provider assessment and feedback and provider incentives. *Am J Prev Med.* Jul; 2008 35(1 Suppl):S67–74. [PubMed: 18541190]


46. Chien AT, Conti RM, Pollack HA. A pediatric-focused review of the performance incentive literature. *Current Opinion in Pediatrics*. Dec; 2007 19(6):719–725. [PubMed: 18025943]
47. Baron RC, Rimer BK, Breslow RA, et al. Client-directed interventions to increase community demand for breast, cervical, and colorectal cancer screening a systematic review. *Am J Prev Med*. Jul; 2008 35(1 Suppl):S34–55. [PubMed: 18541187]

Author Manuscript


Author Manuscript

Author Manuscript

Author Manuscript



**Figure 1.**  
 Logic Model for Core Objective: Practice-based interventions to improve screening



**Figure 2.**  
Flow of titles, abstract and articles included in review

**Table 1**

Specific search terms to identify articles testing practice-based interventions to increase the quality of screening in pediatric practices\*

Screening/specific disorders	Setting	Interventions/outcomes
Mass screening Population surveillance Preventive health services Child development Developmental disabilities Language disorders Child behavior disorders Cerebral palsy Autistic disorder Mental retardation Vision disorders Hearing loss Lead poisoning Anemia Iron deficiency Hypertension Obesity Depression Tuberculosis Sexually transmitted infections	Primary health care Community health centers Managed care programs Group practice	Physician’s Practice Patterns Child Health Services Medical Records Systems, Computerized Decision Support Systems, Clinical Information Systems Education, Medical Education, Medical, Continuing Insurance, Health, Reimbursement Total Quality Management Quality Assurance, Health Care Referral and Consultation Primary Prevention Healthcare Disparities Health Care Costs Quality of Health Care Outcome Assessment Process Assessment

\* In PubMed, language was limited to “English” and population was limited to “All child: 0–18 years”

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 2**

Interventions to improve screening and follow-up of abnormal screening tests in pediatric primary care, by type of study design

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screening (% of patients screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
<b>Randomized Controlled Trials</b>					
1. Margolis PA, et al. (2004) RCT <sup>20</sup>	<i>Lead poisoning, anemia, and tuberculosis:</i> Serum lead level: Intervention Control	23% 18%	68% 30%	<0.05	<p><i>Intervention:</i> Process improvement methods (aka “knowledge translation”) to improve office systems around preventive care services.</p> <ul style="list-style-type: none"> <li>• Formation of practice-based improvement teams</li> <li>• Ongoing academic detailing by project staff</li> <li>• Plan-do-study-act cycles with goal setting, workflow mapping, audit/feedback.</li> </ul> <p><i>Setting/population:</i> 44 practices in North Carolina were randomized to intervention vs. usual care; n=660 each for post-intervention control and experimental groups; children aged 24–30 months.</p> <p><i>Other comments:</i> Data were collected pre- and post-intervention for both control and experimental group practices. Tuberculosis screening was PPD, Mantoux test, or risk assessment</p>
	Hematocrit: Intervention Control	65% 64%	79% 71%	<0.05	
	Tuberculosis screening: Intervention Control	34% 30%	54% 32%	<0.05	
2. Minkovitz CS, et al. (2003) RCT <sup>35</sup>	<i>Developmental problems:</i> Parent-reported developmental assessment	41–43%	82–84%	<0.001	<p><i>Intervention:</i> Healthy Steps (HS) program</p> <p>a. Co-located developmental specialists to enhance well-child visits; also conducted home visits, provided telephone information line for parents about development, written materials, parent groups, linkages to community resources</p> <p><i>Setting/population:</i> 15 practices randomized in 14 states; experimental n=2021 patients, control n=1716 patients; post-intervention data were collected for children aged 30–33 months.</p> <p><i>Other comments:</i> Parents reported any developmental screening questions (not specifically whether a formal tool was used)</p>
3. Scholes D, et al. (2006) RCT <sup>27</sup>	<i>Chlamydia infection:</i> Urine Chlamydia screening	EMR reminder: 40.8%	42.6%	0.27	<p><i>Intervention:</i> Practice and patient-level interventions</p> <ul style="list-style-type: none"> <li>• Practice-level intervention—Use of peer opinion leader teams; 1 day training session around implementing screening guidelines; quarterly feedback reports on screening quality</li> </ul>

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screening (% of patients screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
4. Shafer MA, et al. (2002) RCT <sup>23</sup>	<i>Chlamydia infection</i> : Urine Chlamydia screening	EMR reminder: 40.8%  21%	42.6%  65%	0.27  <0.001	<ul style="list-style-type: none"> <li>• Patient level intervention—EMR point-of-care reminder to screen sexually-active adolescent females</li> </ul> <p><i>Setting/population</i>: 23 practices in Washington state; experimental n=3511 patients, control n=3649 patients; females aged 14–20 years.</p> <p><i>Intervention</i>: Quality improvement initiative within managed care network</p> <ul style="list-style-type: none"> <li>• Practices formed improvement teams; monthly meetings to strategize about solutions to self-identified barriers to screening, using Plan-Do-Study-Act cycles; performance monitoring</li> <li>• Intervention targeted preventive care visits</li> </ul> <p><i>Setting/population</i>: 10 pediatric practices in California; experimental n=1017 patients, control n=1194 patients; sexually active adolescent females.</p>
5. Tebb KP, et al. (2009) RCT <sup>29</sup>	<i>Chlamydia infection</i> : Urine Chlamydia screening Intervention Control	26% 32%	42% 30%	<0.001	<p><i>Intervention</i>: Quality improvement initiative within managed care network</p> <ul style="list-style-type: none"> <li>• Practices formed improvement teams; monthly meetings with focus on workflow, performance monitoring using Plan-Do-Study-Act cycles</li> <li>• Intervention targeted urgent care visits</li> </ul> <p><i>Setting/population</i>: 10 pediatric practices in California; n was not reported; sexually active adolescent females <i>Other comments</i>: Data were collected pre- and post-intervention for both control and experimental group practices.</p>
<b>Pre-post intervention design</b>					
6. Adams WG et al. (2003) Pre-post <sup>37</sup>	<i>Developmental problems, anemia, lead poisoning, hearing and vision problems</i> : Language development	65.1%	70.0%	Relative risk (95% confidence interval): 1.07 (0.97–1.09)	<p><i>Intervention</i>: EMR template with prompts to improve preventive care services</p> <ul style="list-style-type: none"> <li>• Prompts included were age-specific milestones regarding development in social, fine/gross motor, and language skills, with checkboxes and normal ranges.</li> </ul>
	Behavior/social development	26.4%	65.7%	1.16 (1.04–1.28)	

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
7. Applegate H, et al. (2003) Pre-post <sup>33</sup>	Motor development	63.8%	73.9%	2.49 (2.00–3.10)	<ul style="list-style-type: none"> <li>Other prompts were for anticipatory guidance and screening for psychosocial problems.</li> </ul> <p><i>Setting/population:</i> One practice in Massachusetts with &gt;28,000 visits/year; pre-intervention n=235 patients; post-intervention n=986 patients; children aged 0–5 years</p> <p><i>Other comments:</i> Pre-intervention group had paper charts with well-child visit templates; sample for specific tests varied because some tests are recommended only for a subset based on age.</p> <p><i>Intervention:</i> Provider education and support tools to implement Pediatric Symptom Checklist (PSC); intervention was 2 stages</p> <ul style="list-style-type: none"> <li>Stage 1: Provider training session about screening tool, importance of screening, screener placed on medical chart</li> <li>Stage 2: Implementation of provider and patient handouts that followed the structure of the PSC and were designed to address specific subgroups of symptoms.</li> </ul> <p><i>Setting/population:</i> One academic pediatric practice; pre-intervention n=16 patients; post-intervention n=38 patients; children aged 6–16 years.</p> <p><i>Other comments:</i> No significance testing reported</p>
	Hematocrit	82.5%	85.3%	1.03 (0.91–1.17)	
	Serum lead level	66.7%	79.1%	1.19 (0.99–1.43)	
	Vision	42.9%	50.0%	1.17 (0.80–1.70)	
	Hearing	33.3%	48.3%	1.45 (0.92–2.28)	
8. Block B, et al. (1996) Pre-post <sup>40</sup>	<i>Behavior, developmental and emotional problems:</i> Discussion about behavior, developmental or emotional problems (# items discussed per visit)	1.6 items	10.4 items per visit after Stage 1; 9.9 items per visit after Stage 2		
	Intervention for behavior and emotional problems (# of interventions per visit)	0 interventions	0.125 interventions per visit after Stage 1; 1.9 interventions per visit after Stage 2		
	<i>Follow up of elevated lead levels:</i> Follow up plan in chart Follow up serum lead level done Parent education about reducing exposure, if persistently high levels	32% 9% Not measured	100% 65% 28%		

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
	Follow up serum lead level done	9%	65%		<i>Other comments:</i> No significance testing reported
9. Bordley WC, et al. (2001) Pre-post <sup>22</sup>	Parent education about <i>Atenich lead poisoning</i> , <i>tuberculosis</i> , <i>Hematochit</i> , <i>persistently high levels</i>	Not measured	28%		<i>Intervention:</i> Quality improvement intervention to improve preventive care: <ul style="list-style-type: none"> <li>Practice improvement teams</li> <li>Specific changes to workflow were individualized by practices and included:                             <ul style="list-style-type: none"> <li>Sending patient reminder cards</li> <li>Chart screening prior to patient being seen</li> <li>Chart flagging</li> <li>Using flowsheets and medical record templates</li> </ul> </li> </ul> <i>Setting/population:</i> 8 practices in North Carolina, pre-intervention n=339 patients; post-intervention n=300; children aged 24–30 months <i>Other comments:</i> Lead and tuberculosis screening was risk assessment and laboratory/skin testing, if indicated
	Lead screening	45%	67%	0.001	
	Tuberculosis screening	12%	48%	0.001	
10. Dunlop AL, et al. (2007) Pre-post <sup>32</sup>	<i>Obesity:</i> BMI percentile documented in chart	12%	15% after Stage 1 28% after Stage 2	NS <0.05	<i>Intervention:</i> Provider training and support tools for obesity, 2 staged intervention: <ul style="list-style-type: none"> <li>Stage 1: 2-hour provider training explaining guidelines for assessing and managing overweight and counseling framework (AIM-- Advise, Identify, Motivate); training on using BMI calculator and growth charts</li> <li>Stage 2: 3 month supply of tools--parent screening tool/counseling guide, BMI charts, "prescription pad" for nutrition/physical activity</li> </ul> <i>Population/setting:</i> 6 academic family medicine and pediatric practices in Georgia; pre-intervention n=466; Stage 1 n=538, Stage 2 n=344; children aged 2–17 years
	Nutrition and activity history	50%	56% after Stage 1 81% after Stage 2	NS <0.05	
	Nutrition and activity counseling	33%	35% after Stage 1 47% after Stage 2	NS <0.05	
11. Lannon CM, et al. (2008) Pre-post <sup>21</sup>	<i>Developmental problems</i> PEDS or ASQ	30% (received any developmental screening)	45% (using structured tool (e.g., ASQ))	NS	<i>Intervention:</i> Bright Futures Training Intervention Project: learning collaborative/quality improvement initiative to improve preventive care services

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
12. Polacsek M, et al. (2009) Pre-post <sup>25</sup>	<i>Obesity:</i> BMI documented in chart Screening with previsit, self-administered tool to assess patient's behavior around nutrition and physical activity	38% Not measured	94% 82%	<0.001 <0.001	<ul style="list-style-type: none"> <li>Key practice-level changes included:                             <ul style="list-style-type: none"> <li>Structured developmental screening (PEDS or ASQ)</li> <li>Chart prompts</li> <li>Patient recall/reminder</li> <li>Linkages with community agencies</li> </ul> </li> <li>Used practice improvement teams and plan-do-study-act cycles</li> </ul> <p><i>Population/setting:</i> 15 practices in 9 states; experimental n=305 patients, control n=171 patients; children aged 0–5 years Other comments: No participating practices used formal developmental screening tools pre-intervention.</p> <p><i>Intervention:</i> Learning collaborative</p> <ul style="list-style-type: none"> <li>Teams of physician, nurse and administrator from each practice; 3 1.5 day learning sessions for teams; practices set goals around nutrition and physical activity screening and counseling.</li> <li>Patient screening instruments and provider decision support tools for obesity management</li> </ul> <p><i>Population/setting:</i> 12 practices in Maine; n=600 patients with visits during both pre and post intervention periods; children aged 5–18 years.</p> <p><i>Interventions:</i> State-wide learning collaborative with 4 1-day learning sessions</p> <ul style="list-style-type: none"> <li>Practices formed teams (physician, nurse, administrator) and chose preventive care outcomes to address through practice improvements.</li> <li>Included periodic statewide gatherings for QI training, collaborative telephone calls, audit/feedback to practices</li> </ul>
13. Shaw JS, et al. (2006) Pre-post <sup>19</sup>	<i>Lead poisoning, anemia, tuberculosis, hypertension:</i> Lead screening	72%	85%	0.001	<p><i>Interventions:</i> State-wide learning collaborative with 4 1-day learning sessions</p> <ul style="list-style-type: none"> <li>Practices formed teams (physician, nurse, administrator) and chose preventive care outcomes to address through practice improvements.</li> <li>Included periodic statewide gatherings for QI training, collaborative telephone calls, audit/feedback to practices</li> </ul>
	Hematocrit	70%	74%	NS	
	Vision screening	62%	75%	0.013	
	Tuberculosis screening	18%	39%	0.001	
	Blood pressure	85%	82%	NS	

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
	Hematocrit	70%	74%	NS	<i>Population/setting:</i> 31 practices in Vermont; pre- and post-intervention n= each approx 930 patients; c
	Vision screening	62%	75%	0.013	31 practices in Vermont; pre- and post-intervention n= each approx 930 patients; c
	Tuberculosis screening	18%	39%	0.001	<i>Other comments:</i> Tuberculosis and lead screening were risk assessment and laboratory/skin testing, if indicated.
14. Young PC, et al. (2006) Pre-post <sup>18</sup>	Blood pressure	85%	82%	NS	<i>Intervention:</i> Learning collaborative <ul style="list-style-type: none"> <li>Practices chose aspects of preventive care to focus improvement efforts. Included QI methodology training, conference calls with participating practices, and chart audit/feedback</li> </ul> <i>Population/setting:</i> 14 practices in Utah; pre-intervention n=544 patients; post-intervention n=517 patients; children aged 2–4 years
	Anemia, vision problems, hypertension, obesity; Hematocrit	49%	57%	0.36	
	Vision screening	46%	75%	0.007	
	BP screening	59%	74%	0.010	
	BMI recorded	32%	45%	0.078	
<b>Post intervention with and without a control group</b>					
15. Gioia PC. (2001) Post intervention without control group <sup>38</sup>	<i>Lead poisoning:</i> Serum lead level	Not measured	81%		<i>Intervention:</i> EMR with point-of-care reminders displayed on screen <i>Population/setting:</i> Single practice in New York, n=208 patients; children born in 1998
	<i>Vision disorders: monocular visual acuity and stereopsis</i> 3 year olds	Not measured	70–85%		<i>Intervention:</i> Vision screening with specific tools for assessing monocular visual acuity and stereopsis. <ul style="list-style-type: none"> <li>Provided written guidelines for referral, follow-up based on screening results.</li> <li>Physician and staff training, either in group sessions or one-on-one training Initiative included both Head Start and primary care practices</li> </ul> <i>Population/setting:</i> 28 practices in Ohio and Tennessee; n=627 patients; children aged 3–4 years.
16. Hartmann EE, et al. (2006) Post-intervention without control group <sup>34</sup>	4 year olds	Not measured	93–94%		
	<i>Lead poisoning, anemia, hearing, vision:</i>	74%	100%	<0.001	<i>Intervention:</i> Nurse-led protocol
17. Hull PC, et al. (2008) Post-intervention					

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
with concurrent control group <sup>59</sup>	"Laboratory testing" (serum lead level and hematocrit)				<ul style="list-style-type: none"> <li>EPSDT screening, carried out by a nurse with a specific preventive care role, using protocol attached to medical record.</li> </ul> <p><i>Population/setting:</i> One academic practice received intervention; control group was sample of children from other practices. Intervention group n=514, control n=115 patients; children aged 0–17 years</p>
	Hearing	12%	100%	<0.001	
	Vision	23%	100%	<0.001	
18. Niederman LG, et al. (2007) Post-intervention with concurrent control group <sup>56</sup>	<i>Anemia and lead poisoning:</i> Hematocrit	77%	73%	NS	<p><i>Intervention:</i> Healthy Steps (HS) program implemented in a resident continuity clinic.</p> <p><i>Population/setting:</i> One academic practice in Illinois; experimental n=71, control n=192 patients; children aged at least 18 months</p> <p><i>Other comments:</i> Control group were patients in the practice but not enrolled in HS</p>
	Serum lead level	64%	67%	NS	
19. Ozer EM, et al. (2005) Post-intervention with concurrent control group <sup>51</sup>	<i>Adolescent health risk behaviors:</i> Adolescent health screening questionnaire	Not measured	80%	NA	<p><i>Intervention:</i> Provider training, patient questionnaire, and prompts to facilitate communication about adolescent risk behaviors 2 stage intervention:</p> <ul style="list-style-type: none"> <li>Stage 1: 8-hour provider training workshop around knowledge and skills regarding adolescent preventive care</li> <li>Stage 2: Introduction of patient questionnaire and provider form to screen for and document discussion and counseling regarding risky behaviors.</li> </ul> <p><i>Population/setting:</i> 4 practices in California (2 practices received the intervention); experimental n=1717, control n=911 patients; adolescents aged 14–17 years</p> <p><i>Other comments:</i> Control practices' screening did not differ over study period</p>
	Provider asked about alcohol use during visit	67%	82% after Stage 1 83% after Stage 2	<0.01 <0.001	
	Provider counseled on alcohol use during visit	59%	77% after Stage 1 81% after Stage 2	<0.01 <0.001	
20. Schonwald A, et al. (2009) Post intervention without concurrent control group <sup>30</sup>	<i>Behavior and development problems:</i> PEDS	Not measured	61%		<p><i>Intervention:</i> Implementation of developmental screening using PEDS</p> <ul style="list-style-type: none"> <li>1-hour provider and staff training; physician champion who was available to answer questions from providers and staff.</li> <li>Offered as option for referral a second-stage screening service at the practice staffed by an educational specialist</li> </ul>

Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
<b>Time Series</b>					
21. Earls M, et al. (2006) Time series <sup>28</sup>	Developmental problems: ASQ	24%	62% at year 2; 76% at year 5		<p><i>Population/setting:</i> 1 practice in Massachusetts; pre-intervention n=338 patients, post-intervention n=338 patients, post-intervention n=338 patients, post-intervention n=338 patients.</p> <p><i>Other comments:</i> Use of structured developmental assessments was not routine pre-intervention; authors reported an increase in developmental concerns identified post-intervention (21% vs. 26%, p=0.05); proportion of children referred for developmental concerns did not change post intervention (10% vs 11%).</p>
22. King TM, et al. (2010) Time series <sup>24</sup>	Development problems: PEDS or ASQ	Not measured	67% at 1 month; 85% at 9 months		<p><i>Intervention:</i> Quality improvement initiative to improve child development services:</p> <ul style="list-style-type: none"> <li>Practices completed Plan-Do-Study-Act cycles</li> <li>Emphasized physician champion, workflow map, staff involvement, and periodic data review</li> <li>Part of a larger state-wide initiative that involved state-level policy changes around child developmental services</li> </ul> <p><i>Population/setting:</i> Several practices in North Carolina; sample size was not reported</p> <p><i>Other comments:</i> No significance testing reported</p> <p><i>Intervention:</i> Provider and staff education, physician champion identification</p> <ul style="list-style-type: none"> <li>One-day workshop for practice teams. Practices teams were a group of three key stakeholders within each practice (physician champion, staff member, and another person).</li> <li>AAP-sponsored national pilot project to implement guideline-adherent developmental screening</li> </ul> <p><i>Population/setting:</i> 17 practices from 15 states; pre- and post-intervention n≈1020 children total; children aged 8–36 months</p> <p><i>Other comments:</i> Post-intervention screening varied among practices (33–100%); no significance testing reported</p>



Author, year, design	Condition(s) being screened and screening test(s)	Pre-Intervention or control group screened, unless otherwise specified)	Post-Intervention or experimental group screening (% of patients screened, unless otherwise specified)	Significance testing (p-value unless otherwise specified)	Nature of the intervention, setting/population, and other comments about the study
23. Pomietto M, et al. (2009) Time series <sup>26</sup>	<i>Obesity:</i> BMI and weight classification documented in chart	Not measured	49% at 1 month; 94% at 9 months		<p><i>Intervention:</i> Learning collaborative, combined with community and policy-level interventions.</p> <ul style="list-style-type: none"> <li>Practices participated in 3 8-hour training sessions, monthly phone calls, and practice-based coaching in QI, which included on-site visits to practices</li> <li>Coincided with community-level efforts to better manage chronic conditions, including obesity</li> </ul> <p><i>Population/setting:</i> 8 practices in Washington state. Chart audits of 20 pediatric patients per month per practice were tracked for 9 months. Age range of patients was not reported.</p> <p><i>Other comments:</i> No significance testing reported</p>

Abbreviations:

- HS – Healthy Steps
- LC – Learning collaborative
- BMI – Body mass index
- BP – Blood pressure
- QI – Quality improvement
- HMO – Health maintenance organization
- PEDS – Parents’ evaluation of developmental status
- EMR – Electronic medical record
- EPSDT – Early periodic screening, diagnosis and treatment
- ASQ – Ages and stages questionnaire
- AAP – American Academy of Pediatrics
- RCT – Randomized controlled trial