

The primary structure of a fungal chitin deacetylase reveals the function for two bacterial gene products

(*Mucor rouxii*/chitosan/NodB/peptidoglycan deacetylase)

DIMITRIS KAFETZOPOULOS*[†], GEORGE THIREOS*, JOHN N. VOURNAKIS[‡], AND VASSILIS BOURIOTIS*^{†§}

*Institute of Molecular Biology and Biotechnology, P.O. Box 1515, Heraklion 711 10, Crete, Greece; [†]Department of Biology, Division of Applied Biology and Biotechnology, University of Crete, P.O. Box 1470, Heraklion 711 10, Crete, Greece; and [‡]Department of Biology, Dartmouth College, Hanover, NH 03755

Communicated by Fotis C. Kafatos, May 4, 1993 (received for review December 12, 1992)

ABSTRACT Chitin deacetylase (EC 3.5.1.41) hydrolyzes the *N*-acetamido groups of *N*-acetyl-D-glucosamine residues in chitin. A cDNA to the *Mucor rouxii* mRNA encoding chitin deacetylase was isolated, characterized, and sequenced. Protein sequence comparisons revealed significant similarities of the fungal chitin deacetylase to rhizobial nodB proteins and to an uncharacterized protein encoded by a *Bacillus stearothersophilus* open reading frame. These data suggest the functional homology of these evolutionarily distant proteins. NodB is a chitooligosaccharide deacetylase essential for the biosynthesis of the bacterial nodulation signals, termed Nod factors. The observed similarity of chitin deacetylase to the *B. stearothersophilus* gene product suggests that this gene encodes a polysaccharide deacetylase.

Chitin exists in the cell wall of several Zygomycetes species in its deacetylated form, referred to as chitosan (1). A chitosan layer has also been identified in the spore wall of *Saccharomyces cerevisiae* (2). The biosynthesis of chitosan in fungi proceeds by the coordinated action of chitin synthase (EC 2.4.1.16) and chitin deacetylase (EC 3.5.1.41). The former enzyme synthesizes chitin by polymerization of *N*-acetylglucosaminyl residues from UDP-*N*-acetylglucosamine, whereas the latter hydrolyzes the *N*-acetamido groups on the nascent chitin chains (3, 4).

Chitosan is a biopolymer with unique properties favorable for a broad variety of industrial and biomedical applications (5–7). Presently, chitosan is produced by the thermochemical deacetylation of crab chitin. To develop an alternative enzymatic process for chitosan production, we have initiated a study of fungal chitin deacetylases.

Chitin deacetylase from the fungus *Mucor rouxii* has been purified to homogeneity (8). The enzyme is an acidic glycoprotein of ≈ 75 kDa with $\approx 30\%$ (wt/wt) carbohydrate content. Further biochemical characterization revealed that the enzyme has a very stringent specificity for β 1-4-linked *N*-acetylglucosamine homopolymers, requires at least four residues (chitotetraose) for catalysis, and can achieve extensive deacetylation on chitin polymers.

We report here the cloning of a cDNA that encodes *M. rouxii* chitin deacetylase, [¶] and we present the striking protein sequence similarities between chitin deacetylase, the rhizobial nodB proteins, and a *Bacillus stearothersophilus* open reading frame located next to a gene encoding class I Fe-S fumarase. These similarities suggest the functional homology of these proteins.

MATERIALS AND METHODS

Materials. *M. rouxii* (ATCC 24905) was obtained from the American Type Culture Collection. The cDNA synthesis kit,

the λ ZAP vector, and the packaging extracts were bought from Stratagene. The Sequenase 2.0 kit was purchased from United States Biochemical. Immobilon-P and nitrocellulose membranes were purchased from Millipore. Restriction enzymes were obtained from Minotech. Synthetic oligonucleotides were obtained from the Institute of Molecular Biology and Biotechnology Microchemistry Facility.

Growth of *M. rouxii* Mycelia. *M. rouxii* was grown with vigorous shaking at 28°C in a medium containing 0.3% yeast extract, 1% peptone, and 2% glucose; the pH was adjusted to 4.5 with H₂SO₄ (1). The medium was inoculated with 10⁸ spores per liter, and the mycelia were harvested at early growth phase by filtration.

Enzyme Purification. Chitin deacetylase was purified to homogeneity from mycelial extracts of the fungus *M. rouxii* as described (8).

Protein Sequencing. The amino-terminal sequence of chitin deacetylase was determined after electroblotting a purified enzyme preparation onto a poly(vinylidene difluoride) membrane. Sequence analysis was performed by the Harvard Microchemistry Facility using methods and instrumentation described (9).

Isolation of poly(A)⁺ RNA. *M. rouxii* mycelia were extracted by grinding using liquid nitrogen, and total RNA was subsequently purified by the guanidinium thiocyanate method of Chirwin *et al.* (10). Further purification by ultracentrifugation was performed as described by Glisin *et al.* (11). Poly(A)⁺ RNA was isolated by three passes through an oligo(dT)-cellulose column according to the method of Aviv and Leder (12).

Construction of the cDNA Library. The oligo(dT)-primed *M. rouxii* cDNA library was constructed using Stratagene's cDNA synthesis and cloning kit. Approximately 5 μ g of poly(A)⁺ RNA was used in the cDNA synthesis, and the double-stranded cDNA was directionally cloned into the *Xho*I-*Eco*RI sites of the λ ZAP vector (13). The constructed cDNA library consisted of $\approx 2.5 \times 10^5$ recombinant phages.

Isolation and Sequencing of Chitin Deacetylase cDNA Clones. Based on the amino-terminal protein sequence analysis data, four degenerate oligonucleotides were designed: 1A, GA(C/T)ACITCIGCIAA(C/T)TA(C/T)TGGC; 1B, GA(C/T)ACIAG(C/T)GCIAA(C/T)TA(C/T)TGGC; 2A, TT(C/T)ACITCICA(A/G)ATIAA(C/T)CC; 2B, TT(C/T)ACIAG(C/T)CA(A/G)ATIAA(C/T)CC. Oligonucleotides 1A and 2A, which gave positive signals when tested by Southern blot analysis of the *Bam*HI-digested library DNA, were initially combined in the cDNA library screening and subsequently alternated in the rescreening. The hybridization buffer used in Southern blot analysis and library screening was as described (14) with the addition of sonicated and

Abbreviation: bv., biovar.

[§]To whom reprint requests should be sent at the * address.

[¶]The nucleotide sequence data reported in this paper has been deposited in the GenBank data base (accession no. Z19109).

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

					5'	TCTACATCCAAAAAGAAATTGAAACG		25
	ATGCAAAATCAAGACATTGCGCCCTTTCAGCT	GCAATTGCACAAGTTGCTACTCTTGCTTTA	GCCGACACCTCCGCAAATTACTGGCAATCA	TTTACTTCTCAAATTAATCCCAAGAACATC				145
-21	MetGlnIleLysThrPheAlaLeuSerAla	AlaIleAlaGlnValAlaThrLeuAlaLeu	AlaAspThrSerAlaAsnTyrTrpGlnSer	PheThrSerGlnIleAsnProLysAsnIle				
	TCCATTCCCTCTATTGAGCAAACCTTACATCC	ATTGACCCCACTCAAGAATGTGCTTACTAC	ACTCCTGATGCTTCATTGTTCACATTCAAC	GCTTCCGAATGGCCCTCTATCTGGGAAGTC				265
20	SerIleProSerIleGluGlnThrSerSer	IleAspProThrGlnGluCysAlaTyrTyr	ThrProAspAlaSerLeuPheThrPheAsn	AlaSerGluTrpProSerIleTrpGluVal				
	GCTACTACCAATGGTATGAATGAGTCTGCC	GAGTTCCTCAGTGTCTACAATTTCTATTGAC	TGGACCAAGGCCCAATATTTCTGTGCGT	ACCCTTGACGCTAACGGCAACTTGGATACC				385
60	AlaThrThrAsnGlyMetAsnGluSerAla	GluPheLeuSerValTyrAsnSerIleAsp	TrpThrLysAlaProAsnIleSerValArg	ThrLeuAspAlaAsnGlyAsnLeuAspThr				
	ACTGGTTACAATACTGCTACTGACCCCTGAT	TGTTGGTGGACAGCTACCACATGTACCTCT	CCCAAGATTTCTGATATCAATGACGATATC	TCCAAGTGTCTGAACCCGAGACTTGGGGT				505
100	ThrGlyTyrAsnThrAlaThrAspProAsp	CysTrpTrpThrAlaThrThrCysThrSer	ProLysIleSerAspIleAsnAspAspIle	SerLysCysProGluProGluThrTrpGly				
	TTGACTTACGATGATGGACCTAAGTCTCT	CACAACGCTTCTGACTACCTTCAAGAG	CAAAAGTTGAAGCCCTCCATGTTTATATT	GTTTCCAATGTTGTTGACTGGCCATACGGT				625
140	LeuThrTyrAspAspGlyProAsnCysSer	HisAsnAlaPheTyrAspTyrLeuGlnGlu	GlnLysLeuLysAlaSerMetPheTyrIle	GlySerAsnValValAspTrpProTyrGly				
	GCTATCGCTGGTGTGTGATGGCCATCAC	ATTGCATCCACACATGGTCTCACCCCTCAA	ATGACAACCAAGACCAATCAAGAGGTCTT	GCTGAATTCATTATCTACTCAAAGGCCATC				745
180	AlaMetArgGlyValValAspGlyHisHis	IleAlaSerHisThrTrpSerHisProGln	MetThrThrLysThrAsnGlnGluValLeu	AlaGluPheTyrTyrThrGlnLysAlaIle				
	AAGCTCGTACTGGTTGACCCTCGTTAC	TGGCGTCTCCTTATGGTGATATCGATGAT	CGTGTCTGGTGGATTGCCTCTCAATTAGGT	TTAACTGCTGTTATTGGAACCTCGATACT				865
220	LysLeuAlaThrGlyLeuThrProArgTyr	TrpArgProProTyrGlyAspIleAspAsp	ArgValArgTrpIleAlaSerGlnLeuGly	LeuThrAlaValIleTrpAsnLeuAspThr				
	GATGATTGGTCTGCTGGTGTCACTACTACT	GTCGAAGCTGTGAGCAAAGTTATTCGGAT	TATATTGCTATGGGTACCAATGGTACTTTT	GCCAACAGTGGTAACATTGTATTGACCCAT				985
260	AspAspTrpSerAlaGlyValThrThrThr	ValGluAlaValGluGlnSerTyrSerAsp	TyrIleAlaMetGlyThrAsnGlyThrPhe	AlaAsnSerGlyAsnIleValLeuThrHis				
	GAAATCAACACAATATGAGTCTCGCTGTC	GAGAACTTGCCCAAGATCATTCTGCCTAT	AAACAAGTCATCGATGTCCTACCTGTGTTAC	AACATTCTCACCTTACTTTGAAGACTAC				1105
300	GluIleAsnThrThrMetSerLeuAlaVal	GluAsnLeuProLysIleIleSerAlaTyr	LysGlnValIleAspValAlaThrCysTyr	AsnIleSerHisProTyrPheGluAspTyr				
	GAATGGACCAATGCTTGAACGGCACAAAA	TCTTCTGCTACCGCCAGTGTATCTGCTACT	TCTGCTAGTGTCTTGGAGGGCTACTACC	GCTGCCGCTCATCCAAGCTTCTACTAGC				1225
340	GluTrpThrAsnValLeuAsnGlyThrLys	SerSerAlaThrAlaSerGlySerAlaThr	SerAlaSerAlaSerGlyGlyAlaThrThr	AlaAlaAlaHisIleGlnAlaSerThrSer				
	GGCGCCATGTCTGCTCTCCCAACCTCGCC	TTGATCTCTGCCTTATTGCTACCCCTGTTG	TTTTAGTCAAACATCGTCATATATCACCTT	CCTGTCATAATTTATAATAGTAAAACATCA				1345
380	GlyAlaMetSerValLeuProAsnLeuAla	LeuIleSerAlaPheIleAlaThrLeuLeu	PheEnd					
	TATTAGATTTTCTACATCTTAAAAAAA	3'						1375

FIG. 1. Nucleotide and deduced amino acid sequence of the *M. rouxii* chitin deacetylase cDNA clone. Amino acid residues are numbered on the left from the first amino acid (indicated by a double underline) of the mature protein. Nucleotide positions are numbered on the right. Underlined asparagine residues indicate the predicted N-glycosylation sites.

denatured calf thymus DNA at a concentration of 100 µg/ml. Hybridizations were performed at 37–42°C for 16–24 hr. The isolated positive clones were subsequently used for the *in vivo* excision of the recombinant pBluescript (Stratagene), performed according to manufacturer's instructions. Double-stranded DNA sequencing was carried out by the dideoxy chain-termination method using specific primers.

Protein Sequence Comparisons. All protein sequence comparisons and the evaluation of the observed homologies were done using the Genetics Computer Group software (15) and the Schartz and Dayhoff data matrix for amino acid comparisons (16). The deduced amino acid sequences of chitin deacetylase and the *B. stearothermophilus* gene (GenBank accession no. L05611) were used in multiple sequence alignments (PILEUP algorithm) with all available nodB protein sequences: *Rhizobium leguminosarum* biovar (bv.) *trifolii*

(17), *R. leguminosarum* bv. *viciae* (18), *Rhizobium fredii* (19), *Rhizobium meliloti* (20), *R. leguminosarum* bv. *phaseoli* (21), *Bradyrhizobium* sp. (*Parasponia*) (22), and *Azorhizobium caulinodans* (23).

RESULTS AND DISCUSSION

We have recently reported the purification and characterization of chitin deacetylase from the fungus *M. rouxii* (6). In this report we present the isolation of a cDNA clone encoding the enzyme, its sequencing, and derived amino acid sequence comparisons with known genes.

Two different enzyme preparations were used for the determination of the amino-terminal sequence of the protein. This analysis provided with high confidence a sequence of 17 amino acids: Asp-Thr-Ser-Ala-Asn-Tyr-Trp-Gln-Ser-Phe-

Table 1. Matrix of the pairwise identity and similarity scores between deacetylases

	<i>M.r.</i> CDA	<i>B.st.</i> PDA	<i>A.c.</i> nodB	<i>R.l.p.</i> nodB	<i>B.sp.</i> nodB	<i>R.m.</i> nodB	<i>R.f.</i> nodB	<i>R.l.v.</i> nodB	<i>R.l.t.</i> nodB
<i>M.r.</i> CDA	100	22 (35)	21 (40)	23 (38)	25 (37)	24 (35)	25 (38)	29 (42)	27 (42)
<i>B.st.</i> PDA		100	22 (39)	29 (46)	33 (47)	31 (48)	29 (48)	31 (47)	32 (47)
<i>A.c.</i> nodB			100	36 (53)	39 (56)	40 (55)	41 (58)	41 (59)	40 (57)
<i>R.l.p.</i> nodB				100	58 (70)	59 (69)	63 (74)	60 (73)	54 (68)
<i>B.sp.</i> nodB					100	58 (69)	66 (76)	65 (75)	56 (68)
<i>R.m.</i> nodB						100	64 (76)	66 (74)	61 (70)
<i>R.f.</i> nodB							100	66 (78)	60 (74)
<i>R.l.v.</i> nodB								100	71 (80)
<i>R.l.t.</i> nodB									100

The numbers show the percentage of identity and similarity (in parentheses) between deacetylases from various organisms. The calculation of the percentage of identity and similarity in the pairwise comparison of the sequences was performed with the DISTANCES algorithm of the Genetics Computer Group software package (15). Similarities were defined according to Dayhoff's table (16), setting the threshold for a match to 0.6. *M.r.*, *M. rouxii*; *B.st.*, *B. stearothermophilus*; *A.c.*, *Azorhizobium caulinodans*; *R.l.p.*, *R. leguminosarum* bv. *phaseoli*; *B.sp.*, *B. sp.* (*Parasponia*); *R.m.*, *R. meliloti*; *R.f.*, *R. fredii*; *R.l.v.*, *R. leguminosarum* bv. *viciae*; *R.l.t.*, *R. leguminosarum* biovar *trifolii*; CDA, chitin deacetylase; PDA, assumed polysaccharide deacetylase.

<i>Mr</i> CDA	116	TCTSPKISDI	NDDISKCEPEP	ETWGLTYDDG	PN-CSHNAFY	DYLQEQKLKA	SMFYIGSNVV
		::	:	:	:	:	:
<i>Rlt</i> nodB	1	MKRRAYISEV	PFDEAGSDD-	<u>RNIYLTEDDG</u>	<u>RNP</u> HCTGQIL	DVLAEHRVPA	TEFVLGGHVK
		:	:	:	:	:	:
<i>Bst</i> PDA	47	KEWNELLNKY	GAFYLGDP	SK	KEIYLTFDNG	YENGYTSKIL	DVLKKHDVHA
		:	:	:	:	:	:
<i>Mr</i> CDA	175	DWPGAMRGV	VDGHHI-ASH	TWHPQMTK	TNQEVLAEFY	YTQKAIKLAT	G-LTPRYWRP
		:	:	:	:	:	:
<i>Rlt</i> nodB	60	DHPDLVRRVA	AEG- <u>HLVANH</u>	<u>TMTHEDLTAC</u>	DSEAIEREIK	ETNEAIVSAC	PQVAVQHIR-
		:	:	:	:	:	:
<i>Bst</i> PDA	107	TAPDLVKRMV	KEG-HIVGNH	SWSHPDMTTI	SADKIKKELD	AVSDKVKELT	QOEGTVVVRP
		:	:	:	:	:	:
<i>Mr</i> CDA	233	PYGDIDRRVR	WIASQLGLTA	VIWNLDTTDD	S-AGVTT---	TVEAVEQSYS	DYIAMGTNGT
		:	:	:	:	:	:
<i>Rlt</i> nodB	118	<u>RYGAW</u> NADVLL	SRSMNAGLRP	VHMSIDPRDW	SRPGVDS---	IVDAVLAA--	-----
		:	:	:	:	:	:
<i>Bst</i> PDA	166	PRGIFSERL	ALSEKYGYRN	IFWSLAFVDW	KVNEQKQWRY	AYDNIINQ--	-----
		:	:	:	:	:	:
<i>Mr</i> CDA	289	FANSGNIVLT	HEINTTMSLA	VENLPKI---	-ISAYKQVID	VATCYNISHP	YFED
		:	:	:	:	:	:
<i>Rlt</i> nodB	163	- <u>ARRGAI</u> VLL	<u>HDGCP</u> EDETG	NCKLTGLRDO	TLSALLAIIE	ALHSRGFSLR	SLPQ
		:	:	:	:	:	:
<i>Bst</i> PDA	214	-IHPGAIILL	H-----	-----TVSKDN	A-DALDEAIK	TLKQGYTFK	SLDD
		:	:	:	:	:	:

FIG. 2. Comparison of chitin deacetylase from *M. rouxii* (*Mr* CDA), nodB protein from *R. leguminosarum* bv. *trifolii* (*Rlt* nodB), and the deduced amino acid sequence of the presumed polysaccharide deacetylase from *B. stearothermophilus* (*Bst* PDA). Invariant residues among all nodB proteins examined are indicated by underlining. Starting amino acid numbers for each protein sequence are given on the left. Vertical lines indicate identities, and colons indicate conservative substitutions, according to the following amino acid similarity groups: V, I, L, M; D, E, N, Q; F, Y, W, H; A, S, T; K, R; P, G.

Thr-Ser-Gln-Ile-Asn-Pro-Lys. On the basis of this information, synthetic oligonucleotide probes (described in the *Materials and Methods*) were designed and used for the screening of a *M. rouxii* cDNA library. Among the positive clones, the one with the longest insert was subsequently used for the determination of the DNA primary structure (Fig. 1).

A search of the European Molecular Biology Laboratory data bank, using the deduced amino acid sequence of chitin deacetylase, revealed striking sequence similarities with the various rhizobial nodB proteins and a *B. stearothermophilus* protein encoded by an uncharacterized open reading frame located next to a fumarase gene (*fumA*). To highlight the most significant similarities, we performed a multiple sequence alignment using the deduced amino acid sequences of these genes. Furthermore, we calculated the pairwise identity and similarity scores between them, as an indication of their relative distances (Table 1).

The nodB protein sequences (≈215 amino acids long) are well conserved among all rhizobial species examined. The *B. stearothermophilus* sequence is somewhat longer (265 amino acids) and exhibits similarities extending from position number 53 toward its carboxyl terminus. Chitin deacetylase is a much longer polypeptide (400 amino acids), and the region that displays homology to the other proteins is located in the central part of the molecule. The amino-terminal (amino acids 1–121) and the carboxyl-terminal (approximately amino acids 325–400) regions do not display any significant similarity to any other known protein sequence. The sequence comparison between the fungal chitin deacetylase and the bacterial proteins is shown in Fig. 2. Invariant residues in all nodB sequences available in the data base are indicated by underlining. It is noteworthy that these invariant nodB residues are also extensively conserved in both chitin deacetylase and the *Bacillus* sequence. In contrast, six of nine predicted N-glycosylation sites in the chitin deacetylase sequence (Fig. 1) are found in the nonconserved regions of the protein, suggesting that these glycosylated domains may be important for the specialized function of the enzyme in cell wall biosynthesis.

The biochemical pathway for Nod factor synthesis is presently under intense investigation. Nod factors from different rhizobial species share a common basic structure: they are all *N*-acetylglucosamine oligomers, with an *N*-acyl substitution at the nonreducing end residue (24–28). This *N*-acyl substitution is required for the biological activity of all Nod factors, since *N*-acetylglucosamine oligomers fail to elicit any

nodulation-specific responses in host plants (25). Furthermore, the similarity of nodC protein to chitin synthase is indicative that *N*-acetylglucosamine oligomers may serve as precursors in Nod factor synthesis (29–31). Given the observed similarity of chitin deacetylase to nodB proteins and the above information, it is reasonable to assume that *nodB* encodes a specific deacetylase for the nonreducing terminal residue of the precursors. This specific deacetylation may in turn provide the necessary free amino group for a subsequent *N*-acylation. While this paper was under review, the above suggested role of nodB protein in Nod factor biosynthesis was demonstrated biochemically by John *et al.* (32).

Some lysozyme-resistant strains of *Bacillus cereus* have been found to contain glucosamine residues in their cell wall peptidoglycan (33), and a specific enzyme activity that catalyzes the deacetylation of *N*-acetylglucosamine residues in peptidoglycan has been identified (34). The carbohydrate part of the peptidoglycan consists of alternating residues of *N*-acetylglucosamine and its lactic acid ether, *N*-acetylmuramic acid, linked by β1-4 glycosidic bonds. On the other hand, *Bacillus* species have been found to produce chitinases and chitosanases degrading chitinous substrates (35). In this process a specific deacetylation of chitin chains may be involved. The observed similarity of chitin deacetylase to the *B. stearothermophilus* gene product, together with the structural similarity of peptidoglycan to chitin, suggests that this gene encodes a peptidoglycan deacetylase, if not a chitin deacetylase.

The fungal chitin deacetylase, the rhizobial nodB proteins, and the deacetylase from *B. stearothermophilus* define a group of proteins with structural and functional homology, although evolutionarily distant. The fact that *N*-acetylglucosamine-6-phosphate deacetylase does not exhibit any significant sequence similarity to this group suggests that the protein conservation relates to both the hydrolysis of the acetamido groups on *N*-acetylglucosamine residues and the polymer structure (i.e., the type and configuration of the glycosidic bond).

The similarities that we have described highlight the catalytic domain in chitin deacetylase sequence and can direct the design of an enzyme with improved efficiency in the deacetylation of chitinous substrates.

We thank W. S. Lane for protein sequencing, A. Athanasiadis and Y. Papanikolaou for their help in sequence comparisons, C. Louis

and C. Savakis for many helpful suggestions, and F. C. Kafatos for critical reading of the manuscript. The chitin deacetylase technology is protected by patents issued to the Institute of Molecular Biology and Biotechnology.

1. Bartnicki-Garcia, S. & Nickerson, W. J. (1962) *Biochim. Biophys. Acta* **58**, 102–119.
2. Briza, P., Ellinger, A., Winkler, G. & Breitenbach, M. (1988) *J. Biol. Chem.* **263**, 11569–11574.
3. Araki, Y. & Ito, E. (1975) *Eur. J. Biochem.* **55**, 71–78.
4. Davis, L. L. & Bartnicki-Garcia, S. (1984) *Biochemistry* **23**, 1065–1073.
5. Rha, C. K., Rodriguez-Sanchez, D. & Kienzle-Sterzer, C. (1984) in *Biotechnology of Marine Polysaccharides*, eds. Colwell, R. R., Pariser, E. R. & Sinskey, A. J. (Hemisphere, Washington, DC), pp. 283–311.
6. Knorr, D. (1984) in *Biotechnology of Marine Polysaccharides*, eds. Colwell, R. R., Pariser, E. R. & Sinskey, A. J. (Hemisphere, Washington, DC), pp. 313–332.
7. Hirano, S. (1989) in *Chitin and Chitosan*, eds. Skjak-Braek, G., Anthonsen, T. & Sandford, P. (Elsevier, Essex, UK), pp. 37–43.
8. Kafetzopoulos, D., Martinou, A. & Bouriotis, V. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 2564–2568.
9. Lane, W. S., Galat, A., Harding, M. W. & Scheiber, S. L. (1991) *J. Protein Chem.* **10**, 151–160.
10. Chirwin, J. M., Przybyla, A. E., MacDonald, R. J. & Rutter, W. J. (1979) *Biochemistry* **18**, 5294–5299.
11. Glisin, V., Crkvenjakov, R. & Byus, C. (1974) *Biochemistry* **13**, 2633–2637.
12. Aviv, H. & Leder, P. (1972) *Proc. Natl. Acad. Sci. USA* **69**, 1408–1412.
13. Short, J. M., Fernandez, J. M., Sorge, J. A. & Huse, W. D. (1988) *Nucleic Acids Res.* **16**, 7583–7600.
14. Church, G. M. & Gilbert, W. (1984) *Proc. Natl. Acad. Sci. USA* **81**, 1991–1995.
15. Devereux, J., Haeberli, P. & Smithies, O. (1984) *Nucleic Acids Res.* **12**, 9509–9524.
16. Scharz, R. M. & Dayhoff, M. O. (1979) in *Atlas of Protein Sequence and Structure*, ed. Dayhoff, M. O. (Natl. Biomed. Res. Fnd., Washington, DC), pp. 353–358.
17. Schofield, P. R. & Watson, J. M. (1986) *Nucleic Acids Res.* **14**, 2891–2903.
18. Rossen, L., Jonhston, A. W. B. & Downie, J. C. (1984) *Nucleic Acids Res.* **12**, 387–395.
19. Krishnan, H. B. & Pueppke, S. G. (1991) *Mol. Pl.-Microbe Interact.* **4**, 512–520.
20. Toeroek, I., Kondorosi, E., Stepkowski, T., Postfai, J. & Kondorosi, A. (1984) *Nucleic Acids Res.* **12**, 9509–9524.
21. Velazquez, M. V., Davalos, A., Las Penas, A., Sanchez, F. & Quinto, C. (1991) *J. Bacteriol.* **173**, 1250–1258.
22. Scott, K. F. (1986) *Nucleic Acids Res.* **14**, 2905–2919.
23. Goethals, K., Gao, M., Toemekpe, K., Van Mondagu, M. & Holsters, M. (1989) *Mol. Gen. Genet.* **219**, 289–298.
24. Lerouge, P., Roch, P., Faucher, C., Maillet, F., Truchet, G., Promé, J. C. & Dénarié, J. (1990) *Nature (London)* **344**, 781–784.
25. Spaink, H. C., Sheeley, J. M., VanBrussel, A. A., Glushken, J., York, W. S., Tak, T., Geiger, O., Kennedy, E. P., Reinhold, V. N. & Lugtenberg, B. J. J. (1991) *Nature (London)* **354**, 125–130.
26. Roche, P., Debelé, F., Maillet, F., Lerouge, P., Faucher, C., Truchet, G., Dénarié, J. & Promé, J. C. (1991) *Cell* **67**, 1131–1143.
27. Truchet, G., Roch, P., Lerouge, P., Vasse, J., Camut, S., de Billy, J., Promé, J. C. & Dénarié, J. (1991) *Nature (London)* **351**, 670–673.
28. Schulze, M., Quiclet-Sire, B., Kondorosi, E., Virelizier, H., Glushka, J. N., Endre, G., Géro, S. & Kondorosi, A. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 192–196.
29. Bulawa, C. E. (1992) *Mol. Cell. Biol.* **12**, 1764–1776.
30. Atkinson, E. M. & Long, S. R. (1992) *Mol. Pl.-Microbe Interact.* **5**, 439–442.
31. Debellé, F., Rosenberg, C. & Dénarié, J. (1992) *Mol. Pl.-Microbe Interact.* **5**, 443–446.
32. John, M., Röhrig, H., Schmidt, J., Wieneke, U. & Shell, J. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 625–629.
33. Araki, Y., Nakatani, T., Nakayama, K. & Ito, E. (1972) *J. Biol. Chem.* **247**, 6312–6322.
34. Araki, Y., Fukuoka, S., Oba, S. & Ito, E. (1971) *Biochem. Biophys. Res. Commun.* **45**, 751–758.
35. Shimahara, K., Takiguchi, Y. & Nagano, Y. (1992) in *Advances in Chitin and Chitosan*, eds. Brine, C. J., Sandford, P. A. & Zikakis, J. P. (Elsevier, Essex, UK), pp. 364–371.