



Published in final edited form as:

Synlett. 2015 ; 26(3): 345–349. doi:10.1055/s-0034-1379319.

Copper-Catalyzed Aerobic C–H Trifluoromethylation of Phenanthrolines

Cheng-Liang Zhu¹, Yong-Qiang Zhang¹, Yong-An Yuan, and Hao Xu

Department of Chemistry, Georgia State University, 100 Piedmont Avenue SE, Atlanta, GA 30303, USA

Hao Xu: hxu@gsu.edu

Abstract

Direct C–H trifluoromethylation of heterocycles is a valuable transformation. In particular, nonprecious metal-catalyzed C–H trifluoromethylation processes, which do not proceed through CF₃ radical species, have been less developed. In this cluster report, a new copper-catalyzed aerobic C–H trifluoromethylation of phenanthrolines is described. This transformation affords trifluoromethylated phenanthrolines that have not been synthesized and preliminary mechanistic studies suggest that the CF₃ group transfer may occur through cooperative activation.

Keywords

copper; fluorine; oxygen; oxidation; heterocycles

Trifluoromethylated arenes and heterocycles are important building blocks for organic synthesis and pharmaceutical research because of their unique electronic and metabolic properties applicable to drug discovery.² Among various catalytic trifluoromethylation methods,³ the direct C–H trifluoromethylation is a straightforward means to afford trifluoromethylated heterocycles. For example, methods for arene C–H trifluoromethylation initiated by CF₃ radical species has received much attention: MacMillan, Baran, and Sanford have each independently discovered radical-based methods for heterocycle trifluoromethylation.⁴ In parallel, nonradical-based methods for C–H trifluoromethylation offer complementary synthetic utilities. Yu, Sanford, and Liu have reported the palladium-catalyzed C–H trifluoromethylation methods of heterocycles.⁵ However, nonprecious metal-catalyzed direct C–H trifluoromethylation which does not proceed through CF₃ radical species has been less explored.

Recently, Qing disclosed a copper-catalyzed oxidative trifluoromethylation of heteroarenes under strongly basic conditions (Scheme 1,A).⁶ In this reaction, a reductive elimination step of a CF₃ group from the high-valent copper center has been proposed as the key step for the C–H trifluoromethylation. Since phenanthrolines are widely used ligands in synthetic

Correspondence to: Hao Xu, hxu@gsu.edu.

¹These two authors contributed equally.

Supporting Information

Supporting information for this article is available online at <http://dx.doi.org/10.1055/s-0034-1379319>.

chemistry and to the best of our knowledge, the synthesis of trifluoromethylated phenanthrolines has not been reported,⁷ we herein describe a Cu(OAc)₂-catalyzed method for aerobic C–H trifluoromethylation of phenanthrolines under nearly neutral conditions (Scheme 1,B). Our preliminary studies reveal that the acetate counterion is crucial for this unique reactivity and that the CF₃-group transfer may occur via Lewis acid–Lewis base cooperative activation.

We initiated catalyst discovery with 1,10-phenanthroline (**1**) as a model substrate for synthetic and mechanistic considerations (Table 1).⁸ When KF was applied to activate TMSCF₃ under O₂ (1.013 bar), we observed that **1** was mostly recovered in the absence of copper catalyst (Table 1, entry 1). CuI and CuBr₂ were subsequently determined ineffective to promote the desired reaction (Table 1, entries 2 and 3). Interestingly, CuOAc catalyzes the *ortho* trifluoromethylation of **1** at 40 °C (Table 1, entry 4, full conversion, 72% yield), and Cu(OAc)₂ is equally active for this reaction (Table 1, entry 5, 75% yield). After exploring the counterion effect with a variety of copper(II) salts, we determined that Cu(OAc)₂ is superior to Cu(TFA)₂ and that Cu(OTf)₂ and CuSO₄ are unreactive (Table 1, entries 6–8). Surprisingly, we discovered that, in the absence of KF, Cu(OAc)₂ catalyzes this reaction, albeit with a lower yield (Table 1, entry 9, 58% yield). Extensive optimization reveals that the combination of Cu(OAc)₂–KOAc or Cu(OAc)₂–KOAc–AcOH promotes the C–H oxidative trifluoromethylation with excellent yield (Table 1, entries 10 and 11). It is interesting to note that no bistrifluoromethylation products were isolated even when a large excess amount of TMSCF₃ (>5.0 equiv) was applied and that the reaction under air atmosphere did not proceed as efficiently as the one that proceeded under O₂.

Since the facile conversion of copper(I) to copper(II) under aerobic conditions is well-known, we suspected that Cu(OAc)₂ is the active oxidative species. To test this hypothesis, we conducted the reaction with a stoichiometric amount of Cu(OAc)₂ in the absence of O₂ and observed full recovery of the starting material (Table 2, entry 1). This result suggests that copper(II) unlikely acts alone as the oxidant. We further explored a range of oxidants, including iodine(III), iodine(V), silver(I), DDQ, and TBHP, all of which proved ineffective for this reaction (Table 2).

Under optimized conditions, we evaluated a series of symmetric and desymmetric phenanthrolines for the *ortho* trifluoromethylation (Table 3). Symmetric 4,7-diphenyl phenanthroline proves an excellent substrate (Table 3, entry 2, 92% yield); however, 4,7-dimethylphenanthroline has a lower reactivity with a decreased yield (Table 3, entry 3, 61% yield). Desymmetric 3-phenylphenanthroline is an acceptable substrate for the *ortho* trifluoromethylation at the C-2 position (Table 3, entry 4, 64% yield) and symmetric 3,8-diphenylphenanthroline provides the trifluoromethylation product in a moderate yield (Table 3, entry 5, 52%). The trifluoromethylation of 5-nitrophenanthroline provides two readily separable products with a decent combined yield (Table 3, entry 6).

Intrigued by the dominant *ortho* selectivity and the fact that the catalytic cycle turns over in the absence of any fluoride-based activator, we then carried out several control experiments to probe for a possible mechanism (Scheme 2). First, when a stoichiometric amount of TEMPO was applied to the standard conditions, the product was isolated with a good yield

(71% yield). At the same time, we did not detect any TEMPO–CF₃ adduct (Scheme 2,A). This result suggests that the CF₃ radical is unlikely to be involved in this reaction. Next, when we subjected **1** to catalytic conditions under an argon atmosphere, a trifluoromethyl-group addition product 1,2-dihydrophenanthroline **3** was isolated after 24 hours albeit with a low yield (Scheme 2,B). Subsequently, we tested the reactivity of an aniline-derived ketoimine **4** with an *ortho*-pyridyl moiety under an argon atmosphere and observed the imine trifluoromethylation product **5** with an acceptable yield (Scheme 2,C). These results suggest that an N,N-bidentate directing group is crucial for the copper-catalyzed CF₃-group transfer.

Based on the collective results from the control experiments, we propose the following mechanistic working hypothesis (Scheme 3). Since copper(II) is known to coordinate with **1** and generate the tetrahedral complex **6**,⁹ the acetate ligand may be activated by the substrate **1**. Subsequently, the acetate ligand may then activate the CF₃ group from TMSCF₃ through a hypervalent silicon species. From this anionic metathesis, a Cu(CF₃)(OAc) (phenanthroline) intermediate **7** can be generated. Since the C=N bond of a phenanthroline is prone to nucleophilic addition,¹⁰ the coordination of the substrate to the Lewis acidic copper(II) presumably activates the C=N bond to nucleophilic addition.¹¹ At the same time, the coordination of the Lewis basic substrate may cooperatively enhance the nucleophilicity of the CF₃ group. The addition product **8**, a 1,2-dihydrophenanthroline derivative can then undergo aerobic oxidation in the presence of O₂ to furnish **2**.^{12,13}

In summary, we have discovered a new copper-catalyzed aerobic C–H trifluoromethylation of phenanthrolines. Our preliminary mechanistic studies revealed that a Lewis acid–Lewis base cooperative activation mechanism may be involved and that the reaction does not proceed through a CF₃ radical species. Further exploration with this new trifluoromethylation mechanism is ongoing.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


This work was supported by Georgia State University and the National Institutes of Health (GM110382).

References and Notes


- (a) Müller K, Faeh C, Diederich F. *Science*. 2007; 317:1881. [PubMed: 17901324] (b) Purser S, Moore PR, Swallow S, Gouverneur V. *Chem. Soc. Rev.* 2008; 37:320. [PubMed: 18197348]
- For selected examples of transition-metal-catalyzed trifluoromethylation of aromatics and related compounds, see: Chen Q-Y, Wu S-W. *J. Chem. Soc. Chem. Commun.* 1989:705. Oishi M, Kondo H, Amii H. *Chem. Commun.* 2009:1909. Cho EJ, Senecal TD, Kinzel T, Zhang Y, Watson DA, Buchwald SL. *Science*. 2010; 328:1679. [PubMed: 20576888] Teverovskiy G, Surry DS, Buchwald SL. *Angew. Chem. Int. Ed.* 2011; 50:7312. Parsons AT, Senecal TD, Buchwald SL. *Angew. Chem. Int. Ed.* 2012; 51:2947. Dubinina GG, Furutachi H, Vicic DA. *J. Am. Chem. Soc.* 2008; 130:8600. [PubMed: 18543912] Zanardi A, Novikov MA, Martin E, Benet-Buchholz J, Grushin VV. *J. Am. Chem. Soc.* 2011; 133:20901. [PubMed: 22136628] Tomashenko OA, Escudero-Adán EC, Martínez Belmonte M, Grushin VV. *Angew. Chem. Int. Ed.* 2011; 50:7655. Morimoto H, Tsubogo

- T, Litvinas ND, Hartwig JF. *Angew. Chem. Int. Ed.* 2011; 50:3793. Kawai H, Furukawa T, Nomura Y, Tokunaga E, Shibata N. *Org. Lett.* 2011; 13:3596. [PubMed: 21667988] Knauber T, Arikian F, Röschenhaler G-V, Gooßen LJ. *Chem. Eur. J.* 2011; 17:2689. [PubMed: 21274956] Hafner A, Bräse S. *Angew. Chem. Int. Ed.* 2012; 51:3713. Qi Q, Shen Q, Lu L. *J. Am. Chem. Soc.* 2012; 134:6548. [PubMed: 22458339] Shimizu R, Egami H, Nagi T, Chae J, Hamashima Y, Sodeoka M. *Tetrahedron Lett.* 2010; 51:5947. He Z, Luo T, Hu M, Cao Y, Hu J. *Angew. Chem. Int. Ed.* 2012; 51:3944. Liu L-P, Xu B, Mashuta MS, Hammond GB. *J. Am. Chem. Soc.* 2008; 130:17642. [PubMed: 19055329] Morandi B, Carreira EM. *Angew. Chem. Int. Ed.* 2010; 49:938.
4. (a) Nagib DA, MacMillan DWC. *Nature (London, U.K.)*. 2011; 480:224. [PubMed: 22158245] (b) Ji Y, Brueckl T, Baxter RD, Fujiwara Y, Seiple IB, Su S, Blackmond DG, Baran PS. *Proc. Natl. Acad. Sci. U.S.A.* 2011; 108:14411. [PubMed: 21844378] (c) Fujiwara Y, Dixon JA, O'Hara F, Funder ED, Dixon DD, Rodriguez RA, Baxter RD, Herle B, Sach N, Collins MR, Ishihara Y, Baran PS. *Nature (London, U.K.)*. 2012; 492:95. [PubMed: 23201691] (d) Ye Y, Sanford MS. *J. Am. Chem. Soc.* 2012; 134:9034. [PubMed: 22624669]
5. (a) Wang X, Truesdale L, Yu J-Q. *J. Am. Chem. Soc.* 2010; 132:3648. [PubMed: 20184319] (b) Ball ND, Kampf JW, Sanford MS. *J. Am. Chem. Soc.* 2010; 132:2878. [PubMed: 20155920] (c) Ball ND, Gary JB, Ye Y, Sanford MS. *J. Am. Chem. Soc.* 2011; 133:7577. [PubMed: 21513271] (d) Mu X, Wu T, Wang H-Y, Guo Y-L, Liu G. *J. Am. Chem. Soc.* 2012; 134:878. [PubMed: 22191420]
6. Chu L, Qing F-L. *J. Am. Chem. Soc.* 2012; 134:1298. [PubMed: 22145831]
7. For a selected example of functionalized phenanthrolines as ligands in catalysis, see: Altman RA, Buchwald SL. *Org. Lett.* 2006; 8:2779. [PubMed: 16774255] a thorough search of REAXYS® database revealed that trifluoromethylated phenanthrolines have not been reported.
8. Amii, Qing, and Hartwig have each developed arene trifluoromethylation methods with the copper(I)/I complex. See ref. 3b,i, 6.
9. A closely related example is the commercially available dichloro(1,10-phenanthroline) copper(II).
10. For an example of nucleophilic addition to the C=N bond of 1,10-phenanthroline, see: Nishikawa Y, Yamamoto H. *J. Am. Chem. Soc.* 2011; 133:8432. [PubMed: 21553923]
11. For relevant studies in this group, see: Lu D-F, Zhu C-L, Xu H. *Chem. Sci.* 2013; 4:2478. Zhang Y-Q, Liu J-D, Xu H. *Org. Biomol. Chem.* 2013; 11:6242. [PubMed: 23942911]
12. Schönecker B, Zheldakova T, Liu Y, Kötteritzsch M, Günther W, Görls H. *Angew. Chem. Int. Ed.* 2003; 42:3240.
13. **Typical Experimental Procedure** To a mixture of 1,10-phenanthroline (**1**, 36 mg, 0.2 mmol), Cu(OAc)₂ (7.4 mg, 0.04 mmol), KOAc (9.8 mg, 0.1 mmol), and AcOH (6.0 μL, 0.1 mmol) in toluene (4.0 mL), TMSCF₃ (30 μL, 0.2 mmol) was added at room temperature under an O₂ atmosphere. The reaction mixture was stirred at 40 °C for 3 h. Another portion of TMSCF₃ (30 μL, 0.2 mmol) was added, and stirring was continued at 40 °C. The reaction was monitored by TLC. After the starting material was completely consumed, the reaction was quenched by H₂O and extracted with EtOAc (3 × 4 mL). The combined organic layer was washed by brine, dried over Na₂SO₄, and concentrated in vacuo. The product **2** was isolated through a silica gel flash column (40% EtOAc in hexanes) as a white foam (41 mg, 80% yield). **2-(Trifluoromethyl)-1,10-phenanthroline (2)** ¹H NMR (400 MHz, CDCl₃): δ = 9.24 (s, H_a), 8.38 (d, *J* = 8.4 Hz, H_f), 8.23 (d, *J* = 8.6 Hz, H_c), 7.93 (d, *J* = 8.4 Hz, H_g), 7.85, 7.78 (d, *J* = 8.8 Hz, H_d and H_e), 7.65 (m, H_b) ppm. ¹³C NMR (100 MHz CDCl₃): δ = 151.06, 147.7 (q, *J* = 35.0 Hz), 145.72, 145.56, 137.83, 136.09, 129.76, 129.10, 128.79, 125.70, 123.59, 124.30 (q, *J* = 273 Hz), 119.00 ppm. ¹⁹F NMR (377 MHz, CDCl₃): δ = -66.59 (s, 3F) ppm. IR (neat): ν_{max} = 3696, 2967, 2217, 1595, 1337, 1112, 851, 746 cm⁻¹. ESI-HRMS: *m/z* calcd for C₁₃H₈N₂F₃⁺ [M + H⁺]: 249.0640; found: 249.0642.

A) C–H trifluoromethylation of heteroarenes under strongly basic conditions reported by Qing




B) current work: C–H aerobic trifluoromethylation of phenanthrolines


**Scheme 1.**

Copper-catalyzed nonradical-based C–H trifluoromethylation of heteroarenes


A) a control experiment in the presence of TEMPO




B) CF_3 group addition to 1,10-phenanthroline under argon




C) imine trifluoromethylation without the quinolinyl directing group



Scheme 2.
Control experiments for mechanistic insights

**Scheme 3.**

Mechanistic working hypothesis for the copper-catalyzed aerobic C-H trifluoromethylation of phenanthroline


Table 1**Catalyst Discovery for Direct Phenanthroline Trifluoromethylation**

Entry ^a	Catalyst	Additive (equiv)	Conversion (%) ^b	Yield (%) ^c
1	none	KF (3.0)	<5	<5
2	CuI	KF (3.0)	<5	<5
3	CuBr ₂	KF (3.0)	<5	<5
4	CuOAc	KF (3.0)	>95	72
5	Cu(OAc) ₂	KF (3.0)	>95	75
6	Cu(TFA) ₂	KF (3.0)	78	56
7	Cu(OTf) ₂	KF (3.0)	<5	<5
8	Cu(SO ₄) ₂	KF (3.0)	<5	<5
9	Cu(OAc) ₂	none	100	58
10	Cu(OAc) ₂	KOAc (1.0)	100	78
11	Cu(OAc) ₂	KOAc (0.5) AcOH (0.5)	100	80

^aReactions were carried out under O₂ at 40 °C, unless stated otherwise.

^bConversions were determined by ¹H NMR analysis with 1,3,5-trimethoxybenzene as an internal standard.

^cIsolated yield.

Table 2Oxidant Screening for the Cu(OAc)₂-Catalyzed Phenanthroline Trifluoromethylation

Entry	Oxidant	Conversion (%)	Yield (%)
1	none	<5	<5
2	O ₂	>95	80
3	PhI(OAc) ₂	56	42
4	AgOAc	<5	<5
5	DDQ ^a	15	10
6	DMP ^b	35	22
7	TBHP ^c	<5	<5

^aDDQ: 2,3-dichloro-5,6-dicyano-1,4-benzoquinone.

^bDMP: Dess–Martin periodinane.

^cTBHP: *tert*-butyl hydroperoxide.

Table 3

Substrate Scope for the Copper-Catalyzed Trifluoromethylation of Phenanthrolines

Entry	Substrate	Product	Yield (%)
1			80
2			92
3			61
4			64
5			52
6			17
			54