

# Eleventh-century shift in timber procurement areas for the great houses of Chaco Canyon

Christopher H. Guiterman<sup>a,b,1</sup>, Thomas W. Swetnam<sup>a</sup>, and Jeffrey S. Dean<sup>a</sup>

<sup>a</sup>Laboratory of Tree-Ring Research, University of Arizona, Tucson, AZ 85721; and <sup>b</sup>School of Natural Resources and the Environment, University of Arizona, Tucson, AZ 85721

Edited by Stephen Plog, University of Virginia, Charlottesville, VA, and approved October 21, 2015 (received for review July 20, 2015)

**An enduring mystery from the great houses of Chaco Canyon is the origin of more than 240,000 construction timbers. We evaluate probable timber procurement areas for seven great houses by applying tree-ring width-based sourcing to a set of 170 timbers. To our knowledge, this is the first use of tree rings to assess timber origins in the southwestern United States. We found that the Chuska and Zuni Mountains (>75 km distant) were the most likely sources, accounting for 70% of timbers. Most notably, procurement areas changed through time. Before 1020 Common Era (CE) nearly all timbers originated from the Zunis (a previously unrecognized source), but by 1060 CE the Chuskas eclipsed the Zuni area in total wood imports. This shift occurred at the onset of Chaco florescence in the 11th century, a time with substantial expansion of existing great houses and the addition of seven new great houses in the Chaco Core area. It also coincides with the proliferation of Chuskan stone tools and pottery in the archaeological record of Chaco Canyon, further underscoring the link between land use and occupation in the Chuska area and the peak of great house construction. Our findings, based on the most temporally specific and replicated evidence of Chacoan resource procurement obtained to date, corroborate the long-standing but recently challenged interpretation that large numbers of timbers were harvested and transported from distant mountain ranges to build the great houses at Chaco Canyon.**

Ancestral Puebloans | archaeology | human–environment interactions | dendrochronology | timber origins

The high desert landscape of Chaco Canyon, New Mexico was the locale of a remarkable cultural development of Ancestral Puebloan peoples, including the construction of some of the largest pre-Columbian buildings in North America (1) (Fig. 1). The monumental “great houses” of Chaco Canyon reflect an elaborate socioecological system that spanned much of the 12,000-km<sup>2</sup> San Juan Basin from 850 to 1140 Common Era (CE) (2). These massive stone masonry structures required a wealth of resources to erect, including an estimated 240,000 trees incorporated as roof beams, door and window lintels, and other building elements (3). The incongruity of the great houses located in a nearly treeless landscape has led archaeologists and paleoecologists to investigate the origins of timbers used in construction (4–9). Beyond the simple curiosity driving this question, the answer has important implications for understanding the complexities of human–environmental interactions, the sociopolitical organization, and the economic structure of Chacoan society (10–12).

The first excavators of the great houses in the early 20th century speculated that construction timbers were harvested locally, perhaps resulting in deforestation of the surrounding landscape (13). Paleoecological studies conducted during the late 1970s and early 1980s, however, showed that ponderosa pine (*Pinus ponderosa*), the primary tree species used in construction, was not abundant enough at the relatively low elevations (1,800–2,000 m above sea level) of Chaco Canyon and nearby mesas to support timber demand (14–16). Spruce (*Picea* spp.) and fir (*Abies* spp.), which account for tens of thousands of construction beams, have been absent from Chaco Canyon for at least 12,000 y and could have only been logged from distant, higher-elevation sites (2,500–3,450 m

above sea level) (4). An inadequate supply of timbers in Chaco Canyon and its immediate surroundings during Puebloan occupation strongly suggests long-distance procurement from surrounding mountain ranges, where all three conifers now grow in abundance. This inference was corroborated by strontium isotope (<sup>87</sup>Sr/<sup>86</sup>Sr)-based sourcing. Through a comparison of <sup>87</sup>Sr/<sup>86</sup>Sr values from great-house timbers to <sup>87</sup>Sr/<sup>86</sup>Sr values from conifer stands growing today in mountains surrounding the San Juan Basin, two studies concluded that the Chuska Mountains (75 km west) and Mount Taylor (85 km southeast) were the most likely sources for spruce, fir, and ponderosa pine trees (6, 7). Recently, the explanation of long-distance timber transport and the related interpretations of <sup>87</sup>Sr/<sup>86</sup>Sr evidence have been challenged and an alternative has been proposed that most great-house timbers (particularly ponderosa pine) were just as likely to have originated from nearby and low-elevation sites within, east, and south of Chaco Canyon (8, 9).

We assessed probable timber origins independently from previous efforts by applying tree-ring width-based sourcing techniques to a set of 170 beams from our archives at the University of Arizona. These beams comprise six tree species from seven great-house structures (Table S1). Tree-ring-based sourcing uses correlation and Student's *t* tests between beams of unknown origin and site chronologies from likely timber harvesting areas (17) (Fig. S1). Each site chronology, as the average of 40–100 trees, represents tree-ring growth patterns peculiar to an individual landscape. This method of identifying the probable origin of timbers has been applied widely in Europe in the study of archaeological and nautical timbers and

## Significance

The iconic great houses of Chaco Canyon occupy a nearly treeless landscape and yet were some of the largest pre-Columbian structures in North America. This incongruity has sparked persistent debate over the origins of more than 240,000 trees used in construction. We used tree-ring methods for determining timber origins for the first time to our knowledge in the southwestern United States and show that 70% of timbers likely originated over 75 km from Chaco. We found that a previously unrecognized timber source, the Zuni Mountains, supplied construction beams as early as the 850s in the Common Era. Further, we elucidate shifting dynamics of procurement that highlight the importance of a single landscape, the Chuska Mountains, in the florescence of the Chacoan system.

Author contributions: C.H.G., T.W.S., and J.S.D. designed research; C.H.G. performed research; C.H.G., T.W.S., and J.S.D. analyzed data; and C.H.G., T.W.S., and J.S.D. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: All tree-ring data used here are available in Dataset S1. Tree-ring chronologies developed as part of this study are available on the International Tree-Ring Data Bank, [www.ncdc.noaa.gov/data-access/paleoclimatology-data/datasets/tree-ring](http://www.ncdc.noaa.gov/data-access/paleoclimatology-data/datasets/tree-ring) (accession nos. NMS588 and NMS589). The database of great house beams is available on the Chaco Research Archive, [www.chacoarchive.org/cra/chaco-resources/tree-ring-database](http://www.chacoarchive.org/cra/chaco-resources/tree-ring-database).

See Commentary on page 1118.

<sup>1</sup>To whom correspondence should be addressed. Email: [chguiterman@email.arizona.edu](mailto:chguiterman@email.arizona.edu).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1514272112/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1514272112/-DCSupplemental).



Fig. 1. Aerial view of Pueblo Bonito, the largest of the Chaco Canyon great houses. Image courtesy of Adriel Heisey.

artifacts, musical instruments, and paintings on oak panels (17–20). These techniques are underused in North America, but recent efforts in the northeastern United States have revealed distant, inland sources for 18th- and 19th-century nautical timbers (21, 22).

Tree-ring sourcing can only be applied where tree growth patterns are distinguishable between the potential locations of origin. In the southwestern United States tree growth primarily responds to regionally coherent winter precipitation (23, 24), and as a consequence trees across the region tend to share roughly half of their interannual variability (25). Differences between site chronologies are predominantly attributed to variations in topography and subregional-scale climate conditions (26).

We compared great-house beams to the site chronologies of eight potential harvesting areas surrounding the San Juan Basin. Chaco Canyon was not included as one of our sites because it lacked enough remnant wood from the Chaco era to build a local site chronology. To assess the efficacy and accuracy of the tree-ring sourcing method within the San Juan Basin, we tested whether tree-ring growth patterns could be distinguished between the various mountain ranges surrounding Chaco Canyon by applying sourcing methods to living trees of known origin (*SI Text* and *Table S2*). We found that within the San Juan Basin, despite a broadly coherent climate–tree growth signal, the ring-width growth patterns in individual trees can be used to correctly identify the specific mountain range from which they came (*Fig. S2*).

## Results and Discussion

We defined the source of each of the 170 great-house timbers as the tree growth location with the highest significant Student's

$t$  value (a combination of correlation and length of overlap, assessed at  $\alpha = 0.01$ ). It is worth noting that many beams had significant  $t$  values with multiple source areas, and tests of differences among these failed to separate the majority of them. Our use of maximum  $t$  values to designate probable origins is justified against the results of our evaluation of tree-ring sourcing in the San Juan Basin (*SI Text* and *Fig. S2*). The accuracy rate of this test is 90% across all four test sites but rises to 100% for the two test sites located within the same mountain range as a source-area chronology. Because the mountain ranges contain far more forest resources than the mesa locations of the other two test sites, it is plausible that much of the great-house construction wood came from areas represented by our eight source-area chronologies. Here, we describe a broad and compelling pattern of origins for great-house timbers that is both independent of and in accord with the sourcing of other Chacoan materials, including timbers.


Maximum  $t$  values for tree-ring sourcing of beams ranged from 2.86 to 16.01, with 75% greater than 5.58 and only four beams below the often-used yet arbitrary threshold for significance of  $t = 3.5$  (27) (*Dataset S1*). Average  $t$  values for the Chuska and Zuni Mountains were the highest (*Fig. S3A*). Differences in sourcing strength between species were negligible, although spruce and fir trees had the lowest mean  $t$  value (*Fig. S3B*). Using more conservative subsets of the sourcing results generated little change in our findings and no differences in interpretation (*SI Text* and *Table S3*).

We found that 70% of the timbers most likely originated from the Chuska and Zuni Mountains, each >75 km from Chaco Canyon (*Fig. 2*). No other potential location accounted for more than 9% ( $n \leq 16$ ) of the beams, fewer than would be expected by chance ( $\chi^2 = 204$ ,  $P < 0.001$ ). Sourcing patterns differed somewhat by species (*Fig. S4*). In the case of ponderosa pine, the Chuskas accounted for 50% and the Zunis for 29%. The Zuni Mountains were also important for piñon (*Pinus edulis*) and juniper (*Juniperus* spp.) trees, accounting for 58%. Results for spruce and fir were similar to those of English et al. (6), with Mount Taylor as the primary source (29%).

Our findings that identify the Chuska and Zuni Mountains as a significant source of great-house beams are consistent with previous interpretations of long-distance timber procurement (4–7, 10, 11). A pair of recent studies, however, challenge this idea based on similarities in the  $^{87}\text{Sr}/^{86}\text{Sr}$  values for trees in the Chuska Mountains and soil in Chaco Canyon (8, 9). They conclude, therefore, that low-elevation sites within and near the Canyon were just as likely sources for ponderosa beams as were the Chuskas, despite a lack of ponderosa pine in the paleoenvironmental record of the Canyon and surrounding areas since ~5,000 y ago (14–16, 28). If large numbers of local ponderosa pines were to exist in the canyon during the Chaco era and were used as construction timbers (8, 9), then we would expect the sourcing of the great-house beams to show a similar pattern to the “sourcing” of modern pines from low-elevation sites within Chaco Canyon (*Fig. S2F*). Modern (post-1300 CE) ponderosa pines growing in north-facing alcoves on the eastern end of the Chaco Core associate most strongly to the Jemez region on the eastern side of the San Juan Basin. The Chuska and Zuni Mountains, where the great-house timbers source, are instead to the west (*Fig. 2*), suggesting that it is unlikely that trees within or near Chaco Canyon were major sources of construction timbers.

Previously, the Zuni area had been identified as a logical procurement area for timber (11) and other resources (29, 30). Our study is the first to our knowledge to source great-house timbers to the Zuni Mountains. Owing to the general lack of large stands of spruce and fir in the present species composition of the mountain range, the area was not considered in the strontium sourcing studies (6, 7). Our results indicate that wood was imported from the Zunis as early as the 850s CE.

To evaluate changes in procurement areas through time we analyzed only beam specimens that have cutting or near-cutting




**Fig. 2.** Source locations for great-house timbers ( $n = 170$ ). The sizes of the green dots are proportional to the percent of beams sourcing to that location; values provide the percentages. Locations of outlier sites and prehistoric road segments come from Mills et al. (43) and Kantner and Kintigh (44), respectively.

outside dates spanning the temporal range of timber harvesting (Fig. S5). We found that before 1020 CE, nearly all beams sourced to the Zuni Mountains, and thereafter the Chuska Mountains, rose in importance, eclipsing the Zuni region by 1060 CE (Fig. 3 and Fig. S6). This apparent shift in timber procurement coincides with the proliferation of other Chuskan goods in the Chaco Canyon archaeological record (Fig. 4). After *ca.* 1020 CE, Narbona Pass Chert (unique to the Chuska Mountains) accounted


for >20% of the chipped stone tool assemblage in Chaco (31). Correspondingly, Chuskan pottery, usually made with temper from Narbona Pass, was extensively imported after *ca.* 1040 CE (32, 33).

At the same time the Chuska Mountains rose in importance during the middle 11th century CE, there was an onset of major architectural expansion in the Chaco Core area that is additionally defined by a change in masonry style (1). Expansion occurred at existing great houses (Pueblo Bonito, Peñasco Blanco, and Una Vida) and seven new great houses were built, comprising half of the great-house structures in the canyon. Whether this surge in construction and new design drove regional expansion of the Chaco system toward the Chuska Mountains or was in response to expansion in that area remains an open question. An explanation for the shift from mostly Zuni-area timbers in favor of Chuskan wood remains equally elusive but may relate to shorter travel distances between the Chuskas and Chaco that would have facilitated large-scale expansion in the Canyon, or a dwindling resource supply along the Zuni front after centuries of occupation. Improved dating and analyses of the Chaco outlier network, particularly on the eastern front of the Chuska Mountains, may provide answers to both questions (12).

The results of our study independently confirm the hypothesis that distant forested landscapes were the sources for hundreds of thousands of timbers used to build the great houses of Chaco Canyon. Our tree-ring-based results, in accordance with the strontium isotope studies, determine the Chuska Mountains as the primary procurement area. Our discovery of the Zuni Mountains as a source of timber has not only expanded the area from which beams are most likely to have been procured, but it also elucidates a temporal shift in timber procurement. The presence of a network of outlier sites and village clusters at the base of the Chuska and Zuni Mountains (Fig. 2) further supports our conclusions by suggesting a direct connection between these upland forests and the semiarid desert of the Chaco Core area. Given that our findings correspond with isotopic and archeological evidence of increasing use of the



**Fig. 3.** Time series of great-house timber origins. Both plots are on the same time scale. (A) The cumulative sum of beams sourcing to the Zuni (green) and Chuska (blue) Mountains. (B) Distribution of cutting dates by 5-y bins for each source location. The bars are plotted on the center year of each bin.



**Fig. 4.** Time series of the Chaco great-house archaeological record. Percentages for chipped stone and pottery represent the amount of Narbona Pass Chert (31) and Chuskan pottery (1) in those records. Curves for construction timbers are the same as in Fig. 3A. Highlighted time periods for construction and importation of pottery come from the Chaco Synthesis Project (1).

Chuskan landscape after 1020 CE (Fig. 4), a preponderance of evidence now indicates the importance of this particular landscape in the development and florescence of the Chacoan system.

The great houses of Chaco Canyon represent an enormous investment in materials, labor, and human ingenuity, requiring favorable environmental conditions and complex socioeconomic structures to construct and maintain. Although Chacoan exchange networks extended as far as Mesoamerica for prestige items (34, 35), the procurement of basic, labor-intensive resources from multiple distant landscapes, with shifting dynamics of use, is a prominent feature of Chacoan society. Testing for similar patterns at other pre-Columbian and historic sites in North America would be a fruitful endeavor and is possible now because of a dense network of long tree-ring chronologies and the rich collection of archaeological wood material housed at the University of Arizona and National Park Service archives.

## Materials and Methods

**San Juan Basin Tree-Ring Chronologies.** We used eight tree-ring chronologies in the San Juan Basin to represent the likely timber harvesting areas for Chaco Canyon great houses (Table 1 and Dataset S1). Each chronology is composed of one to four tree species, with greater numbers of species further back in time. In most cases, the chronologies are built from living tree samples dating back to ~1300 CE, and then extended back in time with archaeological wood from individual sites or site clusters. We assume that beams from these upland archaeological sites represent locally harvested trees because of the abundance of suitable trees near the sites, which is not the case for Chaco Canyon.

One chronology (Mesa Alta) contains no archaeological specimens, with the pre-1600 period consisting of subfossil logs found on the landscape. For this site, we combined two chronologies composed of different species to boost the strength of the local tree-ring growth signal by adding sample depth during the Chaco period. One of these chronologies (MEA) is composed of Douglas-fir (*Pseudotsuga menziesii*) and southwestern white pine (*Pinus strobiformis*) (36), and the other (CDP) is ponderosa pine that we collected

for this study. The Chuska Mountains chronology combines ponderosa pine and piñon subfossil wood from Narbona Pass (NPS) that we collected and archaeological wood excavated from 13<sup>th</sup>-century pueblos at the eastern foot of the mountain range (CHU). This combination was done to boost sample depth and increase the local common signal in the pre-1300s period, and to fill a time gap in our Narbona Pass chronology resulting from a lack of material dating between 1081 and 1140 CE.

**Great-House Timbers.** We selected Chaco Canyon great-house specimens from the archaeological collections of the Laboratory of Tree-Ring Research at the University of Arizona. To enable systematic selection from thousands of archived specimens, we first digitized the complete record of individual specimen catalog cards for the great houses. These data were assembled into a Microsoft Access database consisting of 6,421 records (the complete database is available from [www.chacoarchive.org/cra/chaco-resources/tree-ring-database/](http://www.chacoarchive.org/cra/chaco-resources/tree-ring-database/)). Of these, 2,497 beams have been assigned exact calendar dates. Our selection criteria for specimens to source consisted of conifers that had at least 30 measurable rings, were solid wood (i.e., not charcoal because of its fragility and associated difficulty to measure), came from a distribution of great houses, had cutting or near-cutting outside dates, and represented the temporal distribution of available specimens (Fig. S5). These criteria narrowed our available selection to 1,048 trees. This includes some duplicates where a single beam was sampled more than once by different excavators, resulting in more than one specimen ID or database record. In the process of measuring beams, we took care to avoid duplication by consulting laboratory technician notes, where this is often indicated. In selecting and measuring specimens, we gave preference to older trees, and thus more than 75% of our samples have more than 75 rings.

We measured ring widths to the nearest 0.001 mm on a Velmex system, recorded observations of injuries, ring anomalies, false rings, and frost rings, and photographed each sample. We used the COFECHA computer program (37) to assess the quality of measurements and crossdating (raw measurements are available in Dataset S1). Finally, biological growth trends of the measured beams were removed by division against 50-y cubic smoothing splines (38) in R using the dplR package (39, 40).

In all, we measured 174 specimens, but four were omitted from further analyses because they did not significantly match any of the San Juan Basin chronologies, leaving 170 beams in our final dataset of great-house timbers. These specimens include at least six tree species from seven great-house structures (Table S1). Because spruce and fir are difficult to differentiate (4) they have not historically been differentiated in the archaeological collection, nor has a systematic effort been made to differentiate *Abies lasiocarpa* from *Abies concolor*. Because of these uncertainties, and the relative consistency of spruce and fir as codominants in upper elevation forests, we grouped the two genera into a single category for analyses. We know, however, that our dataset contains both spruce and fir because seven of our beams were identified to their genus by scanning electron microscopy (4).

It is uncertain how well our sample of beams represents the population of great-house construction timbers, of which there were at least 240,000 harvested trees (3). The tree-ring archaeological record is limited to a very small portion of this total estimated population. Fewer than 10,000 beams have been excavated from the great houses, and dendrochronologists could date fewer than 2,500 of those (primarily because many lacked enough rings or adequate growth variability for confirmed dating). This record is all that is available for study, and we have therefore designed our test for origins on a sample of it. We chose a representative sample of the available record with respect to the component of cutting dates rather than other possible parameters or the population of timbers in its entirety. Although there is uncertainty about potential bias in our available sample, we would note that this problem is not unique to this study and exists in paleoecology and

**Table 1.** Source-area chronologies from the San Juan Basin

Location	Site code	Species*	Date range	Latitude	Longitude	Elevation, m above sea level	Source
Mesa Verde	MVER	1	550–1989	37.22	–108.48	2,263	41
Northern Jemez Mountains	MEA/CDP	1, 2, 3	620–2012	36.28	–106.63	2,525	36, this study
Southern Jemez Mountains	JEM	2, 4	598–1972	35.67	–106.50	2,011	41
Gobernador	GOB	2	623–1989	36.42	–107.19	2,230	41
San Juan Mountains	DUR	1, 4, 5	–319–2009	37.42	–107.86	2,213	41, 42
Zuni Mountains	CIB	1, 2, 4, 5	435–1972	35.33	–108.50	2,072	41
Chuska Mountains	CHU/NAR	1, 2, 4, 6	532–2012	36.09	–108.86	2,650	41, this study
Mount Taylor	CEB	4	680–1986	35.55	–107.50	2,072	41

\*Species: 1, *P. menziesii*; 2, *P. ponderosa*; 3, *P. strobiformis*; 4, *P. edulis*; 5, *Pinus flexilis*; 6, *Juniperus* spp.

archaeology in general. Animal or plant specimens and artifacts that survive long times, are discovered, sampled, and effectively dated and analyzed are commonly a very small subset of a much a larger population of unknown statistical distribution. Nevertheless, useful and accurate interpretations and information are often obtained, especially when combined with testing and other, independent lines of evidence, as we have discussed in the present case. It is hoped that further sampling and dating work in the future, and perhaps new methods of analysis, will further test our interpretations.

**Analyses.** Sourcing archaeological timbers consisted of performing correlation analyses between the measured beams and the eight San Juan Basin chronologies. Before calculating correlations, we removed autocorrelation from each series via autoregression modeling. Statistical significance of the correlations was assessed with one-tailed *t* tests ( $\alpha = 0.01$ ), whereby the correlation values (Pearson's *r*) were converted to *t* values using

$$t = \frac{r\sqrt{n-2}}{(1-r^2)},$$

where *n* is the length of overlap between the two series (27). Any non-significant correlations were excluded from analysis. We assigned the

origin of each beam to the location of the regional chronology with the highest significant *t* value.

To assess whether the number of trees sourcing to each location exceeded what might be expected by chance, we applied the  $\chi^2$  goodness-of-fit test (chisq.test) in the R stats package.

**ACKNOWLEDGMENTS.** We thank the Navajo Nation and the Navajo Forestry Department in particular for permission to sample on Narbona Pass. We were aided in the field and laboratory by E. Ahanonu, C. D. Allen, A. Arizpe, C. H. Baisan, E. Bigio, R. Brown, C. Dwyer, J. Farella, J. Johnston, A. Macalady, K. Miller, T. Murphy, and A. Penaloza. Archaeological tree-ring records were digitized and their associated specimens retrieved by P. P. Creasman and G. D. Flax. Edward Cook provided invaluable assistance with the archaeological chronologies. We thank A. Heisey for use of his Pueblo Bonito photo in Fig. 1, J. Betancourt for comments on an earlier draft of this manuscript, D. Meko for statistical advice, and D. Ford and B. Mills for their many insights and support. This study was supported by Western National Parks Association Grant 13-02 and National Park Service Desert Southwest Cooperative Ecosystem Studies Units Grant UAZDS-381. Additional support was provided by National Science Foundation Dynamics of Coupled Natural and Human Systems Award 1114898 (to T.W.S.) and by the Laboratory of Tree-Ring Research.

- Lekson SH, ed (2006) *The Archaeology of Chaco Canyon: An Eleventh-Century Pueblo Regional Center* (School of American Research Press, Santa Fe, NM).
- Doyel DE ed. (1992) Anasazi Regional Organization in the Chaco System. Anthropological Paper No. 5 (Maxwell Museum of Anthropology, Univ of New Mexico, Albuquerque).
- Dean JS, Warren RL (1983) Dendrochronology. *The Architecture and Dendrochronology of Chetro Keti, Chaco Canyon, New Mexico*, ed Lekson SH (US Department of the Interior National Park Service, Albuquerque, NM), pp 105–240.
- Betancourt JL, Dean JS, Hull HM (1986) Prehistoric long-distance transport of construction beams, Chaco Canyon, New Mexico. *Am Antiq* 51(2):370–375.
- Durand SR, Shelley PH, Antweiler RC (1999) Trees, chemistry, and prehistory in the American Southwest. *J Archaeol* 26(2):185–203.
- English NB, Betancourt JL, Dean JS, Quade J (2001) Strontium isotopes reveal distant sources of architectural timber in Chaco Canyon, New Mexico. *Proc Natl Acad Sci USA* 98(21):11891–11896.
- Reynolds AC, et al. (2005) 87Sr/86Sr sourcing of ponderosa pine used in Anasazi great house construction at Chaco Canyon, New Mexico. *J Archaeol Sci* 32(7):1061–1075.
- Drake BL, Wills WH, Hamilton MI, Dorshow W (2014) Strontium isotopes and the reconstruction of the Chaco regional system: evaluating uncertainty with Bayesian mixing models. *PLoS One* 9(5):e95580.
- Wills WH, Drake BL, Dorshow WB (2014) Prehistoric deforestation at Chaco Canyon? *Proc Natl Acad Sci USA* 111(32):11584–11591.
- Windes TC, Ford D (1996) The Chaco wood project: The chronometric reappraisal of Pueblo Bonito. *Am Antiq* 61(2):295–310.
- Windes TC, McKenna PJ (2001) Going against the grain: Wood production in Chacoan society. *Am Antiq* 66(1):119–140.
- Mills BJ (2002) Recent research on Chaco: Changing views on economy, ritual, and society. *J Archaeol Res* 10(1):65–117.
- Judd NM (1954) *The Material Culture of Pueblo Bonito* (Smithsonian Institution, Washington, DC).
- Betancourt JL, VAN Devender TR (1981) Holocene vegetation in Chaco Canyon, New Mexico. *Science* 214(4521):656–658.
- Betancourt JL, Van Devender TR, Martin PS (1983) Fossil packrat middens from Chaco Canyon, New Mexico: Cultural and ecological significance. *Chaco Canyon Country, A Field Guide to the Geomorphology, Quaternary Geology, Paleogeology, and Environmental Geology of Northwestern New Mexico*, eds Wells S, Love D, Gardner T (American Geomorphological Field Group, Albuquerque, NM), pp 207–217.
- Hall SA (1988) Prehistoric vegetation and environment at Chaco Canyon. *Am Antiq* 53(3):582–592.
- Bridge M (2012) Locating the origins of wood resources: A review of dendroprovenancing. *J Archaeol Sci* 39(8):2828–2834.
- Baillie M, Hillam J, Briffa K, Brown D (1985) Re-dating the English art-historical tree-ring chronologies. *Nature* 315:317–319.
- Wazny T (2002) Baltic timber in Western Europe—an exciting dendrochronological question. *Dendrochronologia* 3:313–320.
- Haneca K, Wazny T, Van Acker J, Beekman H (2005) Provenancing Baltic timber from art historical objects: success and limitations. *J Archaeol Sci* 32(2):261–271.
- Martin-Benito D, et al. (2014) Dendrochronological dating of the World Trade Center ship, Lower Manhattan, New York City. *Tree-Ring Res* 70(2):65–77.
- Creasman PP, Baisan CH, Guiterman CH (2015) Dendrochronological evaluation of ship timber from Charlestown Navy Yard (Boston, MA). *Dendrochronologia* 33:8–15.
- St. George S, Ault TR (2014) The imprint of climate within Northern Hemisphere trees. *Quat Sci Rev* 89:1–4.
- Sheppard P, Comrie A, Packin G, Angersbach K, Hughes M (2002) The climate of the US Southwest. *Clim Res* 21:219–238.
- Cropper JP, Fritts HC (1982) Density of tree-ring grids in western North America. *Tree Ring Bull* 42:3–9.
- Fritts HC (1974) Relationships of ring widths in arid-site conifers to variations in monthly temperature and precipitation. *Ecol Monogr* 44(4):411–440.
- Baillie MGL, Pilcher JR (1973) A simple crossdating program for tree-ring research. *Tree-Ring Bull* 33:7–14.
- Betancourt JL (1990) Late Quaternary biogeography of the Colorado Plateau. *Packrat Middens: The Last 40,000 Years of Biotic Change*, eds Betancourt JL, Van Devender TR, Martin PS (Univ of Arizona Press, Tucson, AZ), pp 259–292.
- Benson LV (2012) Development and application of methods used to source prehistoric Southwestern maize: a review. *J Archaeol Sci* 39(4):791–807.
- Grimstead DN, Reynolds AC, Hudson AM, Akins NJ, Betancourt JL (2014) Reduced population variance in strontium isotope ratios informs domesticated turkey use at Chaco Canyon, New Mexico, USA. *J Archaeol Method Theory* 21:1–23.
- Cameron CM (2001) Pink chert, projectile points, and the Chacoan Regional System. *Am Antiq* 66(1):70–101.
- Toll HW (1981) Ceramic comparisons concerning redistribution in Chaco Canyon, New Mexico. *Production and Distribution: A Ceramic Viewpoint*, eds Howard H, Morris EL (BAR International Series No. 120, Oxford), pp 83–121.
- Mills BJ, Carpenter AJ, Grimm W (1997) Sourcing Chuskan ceramic production: Petrographic and experimental analyses. *Kiva* 62(3):261–282.
- Crown PL, Hurst WJ (2009) Evidence of cacao use in the Prehispanic American Southwest. *Proc Natl Acad Sci USA* 106(7):2110–2113.
- Watson AS, et al. (2015) Early procurement of scarlet macaws and the emergence of social complexity in Chaco Canyon, NM. *Proc Natl Acad Sci USA* 112(27):8238–8243.
- Touchan R, Woodhouse CA, Meko DM, Allen CD (2011) Millennial precipitation reconstruction for the Jemez Mountains, New Mexico, reveals changing drought signal. *Int J Climatol* 31:896–906.
- Grissino-Mayer HD (2001) Evaluating crossdating accuracy: A manual and tutorial for the computer program COFECHA. *Tree-Ring Res* 57(2):205–221.
- Cook ER, Peters K (1981) The smoothing spline: A new approach to standardizing forest interior tree-ring width series for dendroclimatic studies. *Tree-Ring Bull* 41:45–53.
- R Core Team (2014) R: A language and environment for statistical computing. Available at [www.R-project.org/](http://www.R-project.org/).
- Bunn AG (2008) A dendrochronology program library in R (dplR). *Dendrochronologia* 26:115–124.
- Dean JS, Funkhouser G (1995) Dendroclimatic reconstructions for the southern Colorado Plateau. *Climate in the Four Corners and Adjacent Regions: Implications for Environmental Restoration and Land-Use Planning*, eds Waugh WJ, Peterson KL, Wigand PE, Louthan BD, Walker RD (US Department of Energy, Grand Junction Projects Office, Grand Junction, CO), pp 85–104.
- Bigio ER (2013) Late Holocene fire and climate history of the western San Juan Mountains, Colorado: Results from alluvial stratigraphy and tree-ring methods. PhD dissertation (Univ of Arizona, Tucson, AZ).
- Mills BJ, Clark JJ, Peeples MA (2014) Chaco Social Networks Site Database version 1.0 (Archeology Southwest, Tucson, AZ).
- Kantner JW, Kintigh KW (2006) The Chaco world. *The Archaeology of Chaco Canyon: An Eleventh-Century Pueblo Regional Center*, ed Lekson SH (School of American Research Press, Santa Fe, NM), pp 153–188.
- Daly A (2007) Timber, trade and tree-rings: A dendrochronological analysis of structural oak timber in northern Europe, C. AD 1000 to C. AD 1650. PhD dissertation (Univ of Southern Denmark, Denmark).
- Windes TC (2006) Early Puebloan occupation in the Chaco Region: Excavations and survey of Basketmaker III and Pueblo I sites (Anthropology Projects, Cultural Resources Management, Intermountain Region Support Office, National Park Service, Department of the Interior, Santa Fe, NM).
- Sass-Klaassen U, Vernimmen T, Baittinger C (2008) Dendrochronological dating and provenancing of timber used as foundation piles under historic buildings in The Netherlands. *Int Biodeterior Biodegradation* 61(1):96–105.
- Kampstra P (2008) Beanplot: A boxplot alternative for visual comparison of distributions. *J Stat Softw* 28:1–9.