

# Six *Pseudoalteromonas* Strains Isolated from Surface Waters of Kabeltonne, Offshore Helgoland, North Sea

Melissa B. Duhaime,<sup>a</sup> Antje Wichels,<sup>b</sup> Matthew B. Sullivan<sup>c,d</sup>

Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, USA<sup>a</sup>; Biologische Anstalt Helgoland, Alfred Wegener Institute, Helgoland, Germany<sup>b</sup>; Department of Microbiology, The Ohio State University, Columbus, Ohio<sup>c</sup>; and Department of Civil, Environmental and Geodetic Engineering, The Ohio State University, Columbus, Ohio, USA<sup>d</sup>

**Draft genomes are presented for 6 *Pseudoalteromonas* sp. strains isolated from surface waters at Kabeltonne, Helgoland, a long-term ecological research station in the North Sea. These strains contribute knowledge of the genomic underpinnings of a developing model system to study phage-host dynamics of a particle-associated ocean copiotroph.**

Received 9 December 2015 Accepted 21 December 2015 Published 11 February 2016

**Citation** Duhaime MB, Wichels A, Sullivan MB. 2016. Six *Pseudoalteromonas* strains isolated from surface waters of Kabeltonne, offshore Helgoland, North Sea. *Genome Announc* 4(1):e01697-15. doi:10.1128/genomeA.01697-15.

**Copyright** © 2016 Duhaime et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Melissa B. Duhaime, duhaimem@umich.edu.

Members of the genus *Pseudoalteromonas* (*Gammaproteobacteria*; *Alteromonadales*; *Alteromonadaceae*) are Gram-negative aerobic chemoorganoheterotrophs of marine microbial communities (1). They span habitats ranging from surface (2) to deep sea (3), polar waters (4) and sea ice (5), and temperate salt-erns (6). While comprising up to 6% of free-living seawater communities in polar regions (2), *Pseudoalteromonas* spp. are specialized in surface-associated habitats (7). *Pseudoalteromonas* spp. are prolific biofilm formers and are found among the dominant community members living on ocean particles (8) and on surfaces of marine eukaryotes (9, 10). They produce myriad bioactive compounds (11) that can prohibit settlement of fouling larvae and algae (9), presumably to give them a competitive advantage in biofilms. Their ability to utilize diverse and complex carbon substrates (12) likely enables their capacity to boom as r-strategist copiotrophs when potential carbon sources become available (13), for example, in the wake of the Deepwater Horizon oil spill (14), in the oil-polluted North Sea (15), or in decaying phytoplankton blooms (9). There are 49 established species in the genus, though 3,174 strains have not been described at the species level (January 2015, GenBank). Of these, 15 and 28 genomes have been sequenced, respectively (January 2015, GenBank). This *Pseudoalteromonas* sequencing project is a step in the development of a model system to study the ecology and evolution of these particle-associated ocean copiotrophs and their viruses (16, 17, M. B. Duhaime et al., submitted for publication).

DNA was extracted from these nonpigmented clonal isolates using the Qiagen DNeasy blood and tissue kit (Qiagen, USA). Indexed paired-end sequencing libraries were prepared with an Illumina TruSeq DNA sample preparation kit (Illumina, Inc., USA) and sequenced on an Illumina HiSeq2000 platform. The 100-bp reads were quality trimmed using sickle version 1.33 (18) with no 5' trimming, and all sequences with Ns or <Q20 were removed. The remaining singleton and paired reads were assembled using Velvet version 1/1/07 (19). Contigs from the velvet assemblies were used to generate *in silico* jumping libraries (1,000 kb) to be further assembled with the original reads using

ALLPATHS-LG (20). Proteins were predicted and annotated using NCBI's PGAP pipeline (December 2015; <http://www.ncbi.nlm.nih.gov/books/NBK174280>). Genome completeness was estimated with CheckM (21).

To further develop the Helgoland *Pseudoalteromonas* phage-host system, prophages were predicted with VirSorter and the RefSeq virus database (22). One prophage was predicted in strains 13-15 and H103 (15.7 and 55 kb, respectively) and two in strain H105 (87.3 and 15.6 kb). Functional annotations were screened for insights into potential substrates (12). All strains contain at least one outer membrane phospholipase and endoglucanase, which are involved, respectively, in hemolysis/bacteriocin release (23) and hydrolysis of cellulose, a major plant cell wall component (24). A glyoxalase of the beta-lactamase superfamily involved in antibiotic resistance is also found in all strains.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited in GenBank under the following accession numbers: [LODJ00000000](https://genbank.ncbi.nlm.nih.gov/GenBank/LODJ00000000) (*Pseudoalteromonas* sp. strain 13-15), [LODI00000000](https://genbank.ncbi.nlm.nih.gov/GenBank/LODI00000000) (strain H71), [LODK00000000](https://genbank.ncbi.nlm.nih.gov/GenBank/LODK00000000) (strain H100), [LOFG00000000](https://genbank.ncbi.nlm.nih.gov/GenBank/LOFG00000000) (strain H103), [LOFH00000000](https://genbank.ncbi.nlm.nih.gov/GenBank/LOFH00000000) (strain H105), and [LOFI00000000](https://genbank.ncbi.nlm.nih.gov/GenBank/LOFI00000000) (strain 10-33). This paper describes the first versions.

## ACKNOWLEDGMENTS

This study was supported in part by a University of Michigan ADVANCE Elizabeth Crosby award to M.B.D. and Gordon and Betty Moore Foundation grants #2631 and #3790 to M.B.S.

## FUNDING INFORMATION

University of Michigan ADVANCE Elizabeth Crosby Award provided funding to Melissa B. Duhaime. Gordon and Betty Moore Foundation provided funding to Matthew Sullivan under grant numbers 2631 and 3790.

## REFERENCES

- Gauthier G, Gauthier M, Christen R. 1995. Phylogenetic analysis of the genera *Alteromonas*, *Shewanella*, and *Moritella* using genes coding for small-subunit rRNA sequences and division of the genus *Alteromonas* into

- two genera, *Alteromonas* (emended) and *Pseudoalteromonas* gen. nov., and proposal of twelve new species combinations. *Int J Syst Bacteriol* 45:755–761. <http://dx.doi.org/10.1099/00207713-45-4-755>.
2. Wietz M, Gram L, Jørgensen B, Schramm A. 2010. Latitudinal patterns in the abundance of major marine bacterioplankton groups. *Aquat Microb Ecol* 61:179–189. <http://dx.doi.org/10.3354/ame01443>.
  3. Qin QL, Li Y, Zhang YJ, Zhou ZM, Zhang WX, Chen XL, Zhang XY, Zhou BC, Wang L, Zhang YZ. 2011. Comparative genomics reveals a deep-sea sediment-adapted life style of *Pseudoalteromonas* sp. SM9913. *ISME J* 5:274–284. <http://dx.doi.org/10.1038/ismej.2010.103>.
  4. Médigue C, Krin E, Pascal G, Barbe V, Bernsel A, Bertin PN, Cheung F, Cruveiller S, D'Amico S, Duillio A, Fang G, Feller G, Ho C, Mangenot S, Marino G, Nilsson J, Parrilli E, Rocha EP, Rouy Z, Sekowska A, Tutino ML, Vallenet D, von Heijne G, Danchin A. 2005. Coping with cold: the genome of the versatile marine Antarctica bacterium *Pseudoalteromonas haloplanktis* TAC125. *Genome Res* 15:1325–1335. <http://dx.doi.org/10.1101/gr.4126905>.
  5. Bian F, Xie BB, Qin QL, Shu YL, Zhang XY, Yu Y, Chen B, Chen XL, Zhou BC, Zhang YZ. 2012. Genome sequences of six *Pseudoalteromonas* strains isolated from Arctic sea ice. *J Bacteriol* 194:908–909. <http://dx.doi.org/10.1128/JB.06427-11>.
  6. De la Haba RR, Sánchez-Porro C, León MJ, Papke RT, Ventosa A. 2013. Draft genome sequence of the moderately halophilic bacterium *Pseudoalteromonas ruthenica* strain CP76. *Genome Announc* 1(3):e00268–13. <http://dx.doi.org/10.1128/genomeA.00268-13>.
  7. Thomas T, Evans FF, Schleheck D, Mai-Prochnow A, Burke C, Penevyan A, Dalisay DS, Stelzer-Braid S, Saunders N, Johnson J, Ferreira S, Kjelleberg S, Egan S. 2008. Analysis of the *Pseudoalteromonas tunicata* genome reveals properties of a surface-associated life style in the marine environment. *PLoS One* 3:e3252. <http://dx.doi.org/10.1371/journal.pone.0003252>.
  8. Fontanez KM, Eppley JM, Samo TJ, Karl DM, DeLong EF. 2015. Microbial community structure and function on sinking particles in the north Pacific subtropical gyre. *Front Microbiol* 6:469. <http://dx.doi.org/10.3389/fmicb.2015.00469>.
  9. Holmström C, Kjelleberg S. 1999. Marine *Pseudoalteromonas* species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiol Ecol* 30:285–293.
  10. Skovhus TL, Holmström C, Kjelleberg S, Dahllöf I. 2007. Molecular investigation of the distribution, abundance and diversity of the genus *Pseudoalteromonas* in marine samples. *FEMS Microbiol Ecol* 61:348–361. <http://dx.doi.org/10.1111/j.1574-6941.2007.00339.x>.
  11. Bowman JP. 2007. Bioactive compound synthetic capacity and ecological significance of marine bacterial genus *Pseudoalteromonas*. *Mar Drugs* 5:220–241. <http://dx.doi.org/10.3390/md504220>.
  12. Ivanova EP, Bakunina IY, Nedashkovskaya OI, Gorshkova NM, Alexeeva YV, Zelepuga EA, Zvyagintseva TN, Nicolau DV, Mikhailov VV. 2003. Ecophysiological variabilities in ectohydrolytic enzyme activities of some *Pseudoalteromonas* species, *P. Citrea*, *P. Issachenkonii*, and *P. nigri-faciens*. *Curr Microbiol* 46:6–10. <http://dx.doi.org/10.1007/s00284-002-3794-6>.
  13. Allers E, Gómez-Consarnau L, Pinhassi J, Gasol JM, Simek K, Pernthaler J. 2007. Response of *Alteromonadaceae* and *Rhodobacteriaceae* to glucose and phosphorus manipulation in marine mesocosms. *Environ Microbiol* 9:2417–2429. <http://dx.doi.org/10.1111/j.1462-2920.2007.01360.x>.
  14. Dubinsky EA, Conrad ME, Chakraborty R, Bill M, Borglin SE, Holli-baugh JT, Mason OU, Piceno YM, Reid FC, Stringfellow WT, Tom LM, Hazen TC, Andersen GL. 2013. Succession of hydrocarbon-degrading bacteria in the aftermath of the *Deepwater Horizon* oil spill in the Gulf of Mexico. *Environ Sci Technol* 47:10860–10867. <http://dx.doi.org/10.1021/es401676y>.
  15. Chronopoulou PM, Sanni GO, Silas-Olu DI, van der Meer JR, Timmis KN, Brussaard CP, McGenity TJ. 2015. Generalist hydrocarbon-degrading bacterial communities in the oil-polluted water column of the North Sea. *Microb Biotechnol* 8:434–447. <http://dx.doi.org/10.1111/1751-7915.12176>.
  16. Wichels A, Biel SS, Gelderblom HR, Brinkhoff T, Muyzer G, Schütt C. 1998. Bacteriophage diversity in the North Sea. *J Appl Environ Microbiol* 64:4128–4133.
  17. Duhaime MB, Wichels A, Waldmann J, Teeling H, Glöckner FO. 2011. Ecogenomics and genome landscapes of marine *Pseudoalteromonas* phage H105/1. *ISME J* 5:107–121. <http://dx.doi.org/10.1038/ismej.2010.94>.
  18. Joshi NA, Fass JN. 2011. Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. <https://github.com/najoshi/sickle>.
  19. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
  20. Ribeiro FJ, Przybylski D, Yin S, Sharpe T, Gnerre S, Abouelleil A, Berlin AM, Montmayeur A, Shea TP, Walker BJ. 2012. Finished bacterial genomes from shotgun sequence data. *Genome Res* 22:2270–2277. <http://dx.doi.org/10.1101/gr.141515.112>.
  21. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <http://dx.doi.org/10.1101/gr.186072.114>.
  22. Roux S, Enault F, Hurwitz BL, Sullivan MB. 2015. VirSorter: mining viral signal from microbial genomic data. *PEER J* 3:e985. <http://dx.doi.org/10.7717/peerj.985>.
  23. Istivan TS, Coloe PJ. 2006. Phospholipase A in Gram-negative bacteria and its role in pathogenesis. *Microbiology* 152:1263–1274. <http://dx.doi.org/10.1099/mic.0.28609-0>.
  24. Kumar R, Singh S, Singh OV. 2008. Bioconversion of lignocellulosic biomass: biochemical and molecular perspectives. *J Ind Microbiol Biotechnol* 35:377–391. <http://dx.doi.org/10.1007/s10295-008-0327-8>.