



REVIEW

Conservation in the face of climate change: recent developments [version 1; referees: 3 approved]

Joshua Lawler¹, James Watson^{2,3}, Edward Game^{4,5}

¹School of Environmental and Forest Sciences, University of Washington, Seattle, WA, USA

²School of Geography, Planning, and Environmental Management, University of Queensland, St. Lucia, Queensland, Australia

³Wildlife Conservation Society, Global Conservation Program, Bronx, NY, USA

⁴The Nature Conservancy, West End, Queensland, Australia

⁵School of Biological Sciences, University of Queensland, St. Lucia, Queensland, Australia

v1 First published: 28 Oct 2015, 4(F1000 Faculty Rev):1158 (doi: 10.12688/f1000research.6490.1)

Latest published: 28 Oct 2015, 4(F1000 Faculty Rev):1158 (doi: 10.12688/f1000research.6490.1)

Abstract

An increased understanding of the current and potential future impacts of climate change has significantly influenced conservation in practice in recent years. Climate change has necessitated a shift toward longer planning time horizons, moving baselines, and evolving conservation goals and targets. This shift has resulted in new perspectives on, and changes in, the basic approaches practitioners use to conserve biodiversity. Restoration, spatial planning and reserve selection, connectivity modelling, extinction risk assessment, and species translocations have all been reimagined in the face of climate change. Restoration is being conducted with a new acceptance of uncertainty and an understanding that goals will need to shift through time. New conservation targets, such as geophysical settings and climatic refugia, are being incorporated into conservation plans. Risk assessments have begun to consider the potentially synergistic impacts of climate change and other threats. Assisted colonization has gained acceptance in recent years as a viable and necessary conservation tool. This evolution has paralleled a larger trend in conservation—a shift toward conservation actions that benefit both people and nature. As we look forward, it is clear that more change is on the horizon. To protect biodiversity and essential ecosystem services, conservation will need to anticipate the human response to climate change and to focus not only on resistance and resilience but on transitions to new states and new ecosystems.



This article is included in the **F1000 Faculty Reviews** channel.

Open Peer Review

Referee Status:

	Invited Referees		
	1	2	3
version 1 published 28 Oct 2015			

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty. In order to make these reviews as comprehensive and accessible as possible, peer review takes place before publication; the referees are listed below, but their reports are not formally published.

- 1 **Paul Beier**, Northern Arizona University USA
- 2 **A Townsend Peterson**, The University of Kansas USA
- 3 **Bruce Stein**, National Wildlife Federation USA

Discuss this article

Comments (0)

Corresponding author: Joshua Lawler (jlawler@uw.edu)

How to cite this article: Lawler J, Watson J and Game E. **Conservation in the face of climate change: recent developments [version 1; referees: 3 approved]** *F1000Research* 2015, 4(F1000 Faculty Rev):1158 (doi: [10.12688/f1000research.6490.1](https://doi.org/10.12688/f1000research.6490.1))

Copyright: © 2015 Lawler J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Grant information: Joshua J. Lawler thanks the Wilburforce Foundation and the Denman endowment at the University of Washington for funding. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Competing interests: The authors declare that they have no competing interests.

First published: 28 Oct 2015, 4(F1000 Faculty Rev):1158 (doi: [10.12688/f1000research.6490.1](https://doi.org/10.12688/f1000research.6490.1))

Introduction

Climate change is one of the largest threats to biodiversity and to natural systems in general^{1,2}. Recent changes in climate are driving shifts in the timing of ecological events³, the distribution of species^{4,5}, and the functioning of ecosystems⁶. Models project even greater changes for the future^{7–10}.

These climate-driven changes challenge the way the planning and practice of conservation have traditionally been done. Most conservation has been predicated on the fact that the environment is relatively stable over the time frames of management planning (generally less than 50 years). However, projected changes in species distributions and ecosystem functions present obvious challenges to this assumption. Over the last decade, there has been a growing literature on what climate change means for biodiversity¹¹ and the implications this has for conservation planning and action¹². As a consequence, there have been some notable shifts in the way conservation is being conducted, from the goals and structure of conservation organizations to the planning and execution of conservation projects on the ground^{12,13}.

Climate change is increasingly integrated into the daily operations of conservation organizations. The anticipated impacts of climate change have driven planners and managers to consider longer time horizons and to anticipate the potentially synergistic effects of climate change and other threats and the need to address them quickly. Likewise, conservation biologists have begun to acknowledge the importance of planning for extreme weather events in addition to slow, long-term climatic change. These shifts have led to new perspectives on, and alterations to, approaches used to conserve biodiversity. Here, we describe some of the more prominent changes in the approaches conservation planners and practitioners have taken to help address the threat of climate change. Some of these approaches, such as restoration and the prioritization of species for conservation actions, have been around for quite some time but needed reframing as a consequence of the likely impacts of climate change. Others, such as assisted colonization, are new twists on old practices and have arisen directly out of the challenges posed by climate change.

Changing conservation approaches to address climate change

Rethinking restoration

Restoration is one of the basic tools of a conservation practitioner. It has traditionally involved returning a system to its state prior to some disturbance¹⁴. However, climate change challenges the very nature of restoration based on such a definition¹⁵. It brings into question the utility of historical benchmarks as restoration targets, the species or seed sources to be used, and the time frame for planning.

Rethinking how restoration efforts are applied in light of climate change has led to shifts in thinking as well as in practice. Most fundamentally, practicing restoration in a changing climate requires embracing uncertainty and accepting that the goals of a project may need to change over time¹⁶. Instead of relying on historical benchmarks, restoration efforts will likely need to look into the future and anticipate change—perhaps relying on a dynamic reference process that accounts for variability in reference ecosystems^{16,17}. Looking forward will also include rethinking the mix of species to

be planted and potentially focusing on ecosystem function rather than particular assemblages of species. Restoration efforts have begun to make use of the same niche modelling methods that have been used to assess potential species responses to climate change¹⁸, but there is clear recognition that better models are needed¹⁹.

Many recent climate-adaptation efforts have involved restoration. Of the projects funded by the Wildlife Conservation Society's Climate Adaptation Fund, at least 70% have involved restoration²⁰. These projects have included riparian restoration to enhance connectivity, coastal restoration to prevent storm damage, forest restoration to reduce fire risk, and prairie restorations to enhance watershed function. In addition to the projects that clearly engage in restoration, several projects are aimed at converting one type of ecosystem to another—not necessarily restoring a historical condition but preparing an ecosystem to function differently in a future climate. An example of this type of project involved converting forested areas to grasslands to facilitate marsh establishment in the face of sea-level rise. Whether such actions are classified as restoration could be debated, but clearly the lessons learned from decades of restoration will be essential to these new types of adaptation projects.

Changes in how conservation planning is undertaken

Systematic conservation planning²¹ has been used widely around the world to help prioritize conservation efforts, particularly the location of protected areas²². Most conservation planning has been based on “static” representations of biodiversity across a region, an approach that is clearly challenged by climate-driven changes in the distribution of species and communities²³. Consequently, substantial thought has been put into how best to incorporate climate change into the conservation-planning process.

The cornerstone of initial approaches to integrate climate change into conservation planning was the use of correlative niche models to predict future distributions of species and ensure that these were adequately represented by present-day conservation efforts. The more sophisticated niche-based planning efforts included the ability of species to track changing habitat conditions through space and time^{24,25} and consideration of uncertainties in predicted distributions^{26,27}. Climate, however, is only one of many factors determining the distributions of species, and the relationship is complex, uncertain, and in many cases evolving²⁸. The magnitude of these ecological uncertainties compounded by the uncertainties associated with climate predictions²⁹ led to calls to integrate climate change into planning using approaches that were more robust to uncertainty in predicted climate impacts^{30–32}.

An approach to planning that has gained some traction for conservation in a dynamic climate involves conserving the underlying geophysical variation in a region, also referred to as “conserving nature's stage”³³. The rationale for this approach is that, theoretically, there should be a strong relationship between species distributions and geophysical settings (for example, elevation and geology)³⁴ such that conserving representative examples of geophysical settings will protect representative ecological communities under both current and future climates³⁵. A similar approach that focuses on current and known patterns in a region emphasizes conserving connectivity between climatically diverse areas^{31,36,37}.

Increasing connectivity

Increasing the connectivity of landscapes to allow species to move in response to climate change is the most-often cited climate change adaptation strategy^{38–40}. Traditionally, connectivity planning has focused on connecting patches of habitat with what amount to linear strips or stepping-stones of more habitat. Although this approach may allow species to move through the landscape, it may do little to facilitate movement into what might become newly suitable habitat. Early attempts to model connectivity expressly for addressing climate change involved projecting shifts in species distributions through time and either identifying overlap of current and future distributions or mapping pathways that tracked those shifting distributions^{41,42}. A more mechanistic application of this approach involved mapping potential corridors through climate-driven shifts in suitable levels of snowpack for wolverines in the northwestern United States⁴³. Other studies have taken different approaches to identifying important areas for species movement in the face of climate change—approaches that are less reliant on projected future changes in climate and species distributions. Brost and Beier⁴⁴ and Beier⁴⁵ focused on the geophysical settings mentioned in the previous section, and charted routes across the landscape that either connected similar geophysical setting or linked a diversity of settings. Nuñez and colleagues⁴⁶ mapped routes through landscapes that connected slightly warmer patches of intact land to slightly cooler ones with routes that followed gentle temperature gradients and avoided human-impacted landscapes.

Although efforts continue to develop more relevant climate-connectivity methods, the critical question of whether corridors are really needed—or whether there might be other strategies and approaches that would achieve the same result—has been repeatedly raised³⁷. Some argue that the focus on corridors is misguided and that, alternatively, protecting large intact ecosystems should be prioritized^{47,48}. Others have argued that the benefits of increasing the size and number of corridors are fewer than those resulting from simply increasing the amount of protected land, which, if regularly distributed, would increase connectivity⁴⁹.

Assessing extinction risk through the climate change lens

Although a growing wealth of studies predict increased extinction risk for species because of climate change^{50,51}, many of these vary enormously in their estimations. It is also increasingly recognized that the predictions of extinction risk do not reflect the number of species that have become vulnerable (or extinct) to date, nor do they match the number identified as threatened because of climate change on the International Union for the Conservation of Nature (IUCN) Red List (only 10.5% of the 22,176 species)⁵². A fundamental challenge has involved integrating the projections of species niche models (the most-often used tool for assessing the impacts of climate change on species) into the processes of real-world extinction assessments.

Simple measures of population size, geographic range size, and other indicators of current status already used as IUCN Red List criteria are likely to be good predictors of climate change-associated extinction risk⁵³. Such IUCN Red List criteria have been used to predict the risk of extinction in the absence of conservation action

and the time lag between assessment and extinction^{54,55}. This time lag amounts to a warning period in which adaptation efforts can be taken to prevent extinction. Although there is a warning period, it is finite and thus delays in developing and implementing conservation plans after a species is identified as being threatened could be costly. Roughly half of listed species are likely to go extinct within 20 years of being listed as critically endangered⁵⁴.

An additional challenge is that most species risk assessments treat climate change as a problem driven by relatively slow, predictable, and continuous change in environmental conditions and fail to account for other important components of climate change, such as increasing extreme weather and climate events (for example, cyclones, floods, and drought)^{56,57}. It is increasingly recognized that the increases in frequencies and intensities of extreme events are critical determinants of patterns of biological diversity and will affect it differently from impacts resulting from steady climate change⁵⁸. A good example of this is how climate change will impact bat species: extreme maximum temperature is now considered a critical factor in the vulnerability of bats to climate change⁵⁹, but many studies (for example, 60) fail to consider it in projections of species distributions under climate change.

Even though our understanding of which extremes are most important and how they are shifting is limited⁶¹, there are good examples of assessments that do account for extremes. Recently, Ameca y Juárez and colleagues⁶² produced a comprehensive analysis of the impacts of cyclones and droughts on terrestrial mammals, one of the few large-scale studies to consider exposure to extreme events. They followed this exposure analysis with an assessment of terrestrial mammal sensitivity to extreme weather and climate events, identifying biological traits that make large terrestrial mammals more susceptible to climate-induced population declines.

Alongside species risk assessments, the assessment of the vulnerability of species to climate change—as well as the climate-related vulnerability of places and natural resources in general—has emerged as an important step in the adaptation process⁶³. As with extinction risk assessments, many different approaches to assessing vulnerability have been proposed (for example, 64), each evaluating some subset or combination of sensitivity, exposure, and adaptive capacity⁶⁵. These vulnerability assessments serve not only to determine which species are likely to be most vulnerable but also to identify the factors that make a species vulnerable and thus potential conservation actions—adaptation measures—that can be taken to reduce vulnerability.

Assisted colonization

One of the relatively new tools in the conservation toolbox is assisted colonization—broadly defined by the IUCN as the movement of an organism outside of its native range to avoid extinction of populations due to current or future threats⁶⁶. There are some species, particularly endemics with relatively specific habitat requirements and poor dispersal abilities, that will be unable to move to suitable climates. When these species are threatened with extinction—because of either climate change or some other factor—it may become necessary to move them to prevent their loss.

The question of whether assisted colonization should even be considered as an option spawned a lively debate^{67–72}. Those opposed to assisted colonization argue that the history of invasive species has taught us that the potential impacts on the ecosystems into which organisms would be moved could be too great⁶⁸. Those in favour of keeping the option of assisted colonization on the table argue that it would likely be necessary for preventing the extinction of certain species⁶⁹, that the potential impact of translocated species is likely overstated for several reasons^{69,71} (including that the traits of species that will need to be moved are not those traits generally associated with invasive species⁷³), and that the amount of change that systems will experience over the coming decades will likely overshadow the impacts of translocated individuals of a rare and declining species⁷⁴.

As the debate worked its way through the scientific literature, many researchers started to ask more productive questions—facing the reality that assisted colonization was already being used. There were, for example, several early efforts to develop frameworks for determining under what circumstances assisted colonization would be a viable conservation option^{75,76}. Other studies have highlighted the importance of the timing of assisted colonization efforts⁷⁷, developed advanced modelling approaches for identifying potential sites for translocations⁷⁸, and explored the situations in which invasions will be less likely and hence assisted colonization a less risky venture⁷⁹. Furthermore, calls for the development of policies to address assisted colonization⁸⁰ have begun to be met^{81,82}. Overall, it appears that, at least in the scientific literature, assisted colonization is gaining acceptance as a tool in the conservation toolbox and one that may not differ so much from other movements of species for conservation reasons^{83,84}.

Future trends

The practice, and to some degree the study, of conservation is currently undergoing a major shift—a shift from a focus on nature to a focus on nature and people. The idea that people are a part of ecosystems and that conservation needs to include the social sciences is not new and this is not the shift to which we are referring. This new shift is one from conserving nature for nature’s sake to conserving nature both for nature’s sake and for the use and enjoyment by people⁸⁵. This shift has resulted in an apparent change in the missions and the actions of several major non-governmental conservation organizations (for example, The Nature Conservancy, the World Wildlife Fund, and Conservation International). Like the subject of assisted colonization, however, this shift has not been well received by all in the conservation community and there remains a heated debate in conservation circles as well as in the literature about the degree to which conservation should focus on the needs of people^{85–88}.

The impact of this shift can be clearly seen in the way that conservation organizations are addressing climate change and is reflected in the application of all of the approaches mentioned above. With respect to restoration, conservation planning, and connectivity,

conservation practitioners have begun to target efforts that consider the roles that natural systems play in protecting people against the potential impacts of climate change^{89,90}. These ecosystem-based adaptation strategies may be more cost-effective than hard infrastructure-based solutions. One striking example of this approach is The Nature Conservancy’s “flood plains by design” strategy in which stretches of river are restored in places that will simultaneously reduce flooding of nearby communities and restore fish habitat. Another example is the active protection of coastal habitats, which has the potential to reduce the risks and the costs of sea-level rise, providing a critical service in the face of climate change⁹¹.

Because climate-driven changes are likely to be so large in some places, climate change is in part causing conservation practitioners to question their goals as well as the approaches they use⁷. These new goals are beginning to take people’s needs into account. For example, restoration efforts are now being refocused toward ecosystem function and ecosystem services instead of the specific set of species in a given ecosystem¹⁹. In addition, assisted colonization may be called on not just to preserve threatened species, but also to provide certain functions—and perhaps to allow ecosystems to provide certain services^{73,92}.

Given that our understanding of climate change impacts is still evolving, the theory and practice of conservation will likely continue to change at a relatively fast pace. One of the greatest future challenges to the conservation of biodiversity will likely come from how people respond to climate change⁹³. Sea-level rise is forcing human populations to consider radical adaptation action, including the construction of massive sea walls and the migration of coastal and island communities^{94,95}. Water shortages and crop failures will similarly result in human migrations, shifts in agriculture, and increased water withdrawals. There is increasing recognition that, in many places, human responses to climate change may further constrain options for biodiversity conservation, and therefore planning needs to simultaneously consider both human and biodiversity responses⁹³. The tools that conservation practitioners have to address climate change (for example, conservation planning, restoration, species risk assessments, and assisted colonization) will likely be most effective if their application takes human responses to climate change into account.

Continued rapid climate change will also necessitate a shift from discussions of resistance and resilience to more strategies that embrace change and foster transitions⁹⁶. Particularly if society hopes to continue to be the recipient of essential ecosystem services and to enjoy a diversity of plants and animals, conservation efforts will need to focus on smoothly transitioning ecosystems from one state to another. The enormity of that challenge necessitates policies and actions that reduce greenhouse-gas emissions and increase carbon sequestration. Unless adaptation is accompanied by meaningful mitigation efforts, it will be hard for conservation practitioners to accomplish even their shifting and evolving goals.

Abbreviation

IUCN, International Union for the Conservation of Nature.

Competing interests

The authors declare that they have no competing interests.

Grant information

Joshua J. Lawler thanks the Wilburforce Foundation and the Denman endowment at the University of Washington for funding.

I confirm that the funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References



- Sala OE, Chapin FS, Armesto JJ, *et al.*: **Global biodiversity scenarios for the year 2100**. *Science*. 2000; **287**(5459): 1770–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Millennium Ecosystem Assessment: **Ecosystems and Human Well-being Biodiversity Synthesis**. Washington DC, World Resources Institute. 2005.
[Reference Source](#)
- Gregory RD, Willis SG, Jiguet F, *et al.*: **An indicator of the impact of climatic change on European bird populations**. *PLoS One*. 2009; **4**(3): e4678.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- F** Chen IC, Hill JK, Ohlemüller R, *et al.*: **Rapid range shifts of species associated with high levels of climate warming**. *Science*. 2011; **333**(6045): 1024–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- F** Pinsky ML, Worm B, Fogarty MJ, *et al.*: **Marine taxa track local climate velocities**. *Science*. 2013; **341**(6151): 1239–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- F** Lane JE, Kruuk LE, Charmantier A, *et al.*: **Delayed phenology and reduced fitness associated with climate change in a wild hibernator**. *Nature*. 2012; **489**(7417): 554–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Watson JEM, Iwamura T, Butt N: **Mapping vulnerability and conservation adaptation strategies under climate change**. *Nat Clim Change*. 2013; **3**(11): 989–94.
[Publisher Full Text](#)
- Thuiller W, Lavorel S, Araujo MB, *et al.*: **Climate change threats to plant diversity in Europe**. *Proc Natl Acad Sci U S A*. 2005; **102**(23): 8245–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Peterson AT, Ortega-Huerta MA, Bartley J, *et al.*: **Future projections for Mexican faunas under global climate change scenarios**. *Nature*. 2002; **416**(6881): 626–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lawler JJ, Shafer SL, White D, *et al.*: **Projected climate-induced faunal change in the Western Hemisphere**. *Ecology*. 2009; **90**(3): 588–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chapman S, Mustin K, Renwick AR, *et al.*: **Publishing trends on climate change vulnerability in the conservation literature reveal a predominant focus on direct impacts and long time-scales**. *Divers Distrib*. 2014; **20**(10): 1221–8.
[Publisher Full Text](#)
- Schmitz OJ, Lawler JJ, Beier P, *et al.*: **Conserving Biodiversity: Practical Guidance about Climate Change Adaptation Approaches in Support of Land-use Planning**. *Nat Areas J*. 2015; **35**(1): 190–203.
[Publisher Full Text](#)
- Seimon A, Watson J, Dave R, *et al.*: **A Review of Climate Change Adaptation Initiatives within the Africa Biodiversity Collaborative Group Members**. *ABCG Arlingt USA*. 2011; 124.
[Reference Source](#)
- National Research Council: **Restoration of Aquatic Ecosystems: Science, Technology, and Public Policy**. Haworth Press. 1992.
[Reference Source](#)
- Harris JA, Hobbs RJ, Higgs E, *et al.*: **Ecological restoration and global climate change**. *Restor Ecol*. 2006; **14**(2): 170–6.
[Publisher Full Text](#)
- Wiens JA, Hobbs RJ: **Integrating Conservation and Restoration in a Changing World**. *BioScience*. 2015; **65**(3): 302–12.
[Publisher Full Text](#)
- Hiers JK, Mitchell RJ, Barnett A, *et al.*: **The dynamic reference concept: measuring restoration success in a rapidly changing no-analogue future**. *Ecol Restor*. 2012; **30**(1): 27–36.
[Publisher Full Text](#)
- Padonou EA, Tekka O, Bachmann Y, *et al.*: **Using species distribution models to select species resistant to climate change for ecological restoration of bowé in West Africa**. *Afr J Ecol*. 2015; **53**(1): 83–92.
[Publisher Full Text](#)
- Starzomski BM: **Novel ecosystems and climate change**. In: Hobbs RJ, Higgs ES, Hall CA, editors. *Novel Ecosystems: Intervening in the New Ecological World Order*. West Sussex UK, Wiley-Blackwell. 2013; 88–101.
[Publisher Full Text](#)
- Wildlife Conservation Society: **WCS Climate Adaptation Fund Grants List**. Bozeman Montana USA, Wildlife Conservation Society. 2014.
[Reference Source](#)
- Margules CR, Pressey RL: **Systematic conservation planning**. *Nature*. 2000; **405**(6783): 243–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Groves CR: **Drafting a Conservation Blueprint: A Practitioners Guide to Planning for Biodiversity**. Washington DC, Island Press. 2003; **22**(2): 147–148.
[Reference Source](#)
- Pressey RL, Cabeza M, Watts ME, *et al.*: **Conservation planning in a changing world**. *Trends Ecol Evol*. 2007; **22**(11): 583–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Williams P, Hannah L, Andelman S, *et al.*: **Planning for climate change: Identifying minimum-dispersal corridors for the Cape proteaceae**. *Conserv Biol*. 2005; **19**(4): 1063–74.
[Publisher Full Text](#)
- Carroll C: **Role of climatic niche models in focal-species-based conservation planning: Assessing potential effects of climate change on Northern Spotted Owl in the Pacific Northwest, USA**. *Biol Conserv*. 2010; **143**(6): 1432–7.
[Publisher Full Text](#)
- Carvalho SB, Brito JC, Crespo EG, *et al.*: **Conservation planning under climate change: Toward accounting for uncertainty in predicted species distributions to increase confidence in conservation investments in space and time**. *Biol Conserv*. 2011; **144**(7): 2020–30.
[Publisher Full Text](#)
- F** Kujala H, Moilanen A, Araujo MB, *et al.*: **Conservation planning with uncertain climate change projections**. *PLoS One*. 2013; **8**(2): e53315.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- F** Gillson L, Dawson TP, Jack S, *et al.*: **Accommodating climate change contingencies in conservation strategy**. *Trends Ecol Evol*. 2013; **28**(3): 135–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Lawler JJ, Tear TH, Pyke C, *et al.*: **Resource management in a changing and uncertain climate**. *Front Ecol Environ*. 2010; **8**(1): 35–43.
[Publisher Full Text](#)
- Millar CI, Stephenson NL, Stephens SL: **Climate change and forests of the future: managing in the face of uncertainty**. *Ecol Appl*. 2007; **17**(8): 2145–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Game ET, Lipsett-Moore G, Saxon E, *et al.*: **Incorporating climate change adaptation into national conservation assessments**. *Glob Change Biol*. 2011; **17**(10): 3150–60.
[Publisher Full Text](#)
- Groves CR, Game ET, Anderson MG, *et al.*: **Incorporating climate change into systematic conservation planning**. *Biodivers Conserv*. 2012; **21**(7): 1651–71.
[Publisher Full Text](#)
- Anderson MG, Ferree CE: **Conserving the stage: climate change and the geophysical underpinnings of species diversity**. *PLoS One*. 2010; **5**(7): e11554.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lawler JJ, Ackerly DD, Albano CM, *et al.*: **The theory behind, and the challenges of, conserving nature's stage in a time of rapid change**. *Conserv Biol*. 2015; **29**(3): 618–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Beier P, Brost B: **Use of land facets to plan for climate change: conserving the arenas, not the actors**. *Conserv Biol*. 2010; **24**(3): 701–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Ashcroft MB, Chisholm LA, French KO: **Climate change at the landscape scale: predicting fine-grained spatial heterogeneity in warming and potential refugia for vegetation**. *Glob Change Biol*. 2009; **15**(3): 656–67.
[Publisher Full Text](#)

37. Hodgson JA, Thomas CD, Wintle BA, *et al.*: **Climate change, connectivity and conservation decision making: back to basics.** *J Appl Ecol.* 2009; **46**(5): 964–9. [Publisher Full Text](#)
38. Mawdsley JR, O'Malley R, Ojima DS: **A review of climate-change adaptation strategies for wildlife management and biodiversity conservation.** *Conserv Biol.* 2009; **23**(5): 1080–9. [PubMed Abstract](#) | [Publisher Full Text](#)
39. Lawler JJ: **Climate change adaptation strategies for resource management and conservation planning.** *Ann N Y Acad Sci.* 2009; **1162**: 79–98. [PubMed Abstract](#) | [Publisher Full Text](#)
40. Heller NE, Zavaleta ES: **Biodiversity management in the face of climate change: A review of 22 years of recommendations.** *Biol Conserv.* 2009; **142**(1): 14–32. [Publisher Full Text](#)
41. Phillips SJ, Williams P, Midgley G, *et al.*: **Optimizing dispersal corridors for the Cape Proteaceae using network flow.** *Ecol Appl.* 2008; **18**(5): 1200–11. [PubMed Abstract](#) | [Publisher Full Text](#)
42. Vos CC, Berry P, Opdam P, *et al.*: **Adapting landscapes to climate change: examples of climate-proof ecosystem networks and priority adaptation zones.** *J Appl Ecol.* 2008; **45**(6): 1722–31. [Publisher Full Text](#)
43. McKelvey KS, Copeland JP, Schwartz MK, *et al.*: **Climate change predicted to shift wolverine distributions, connectivity, and dispersal corridors.** *Ecol Appl.* 2011; **21**(8): 2882–97. [Publisher Full Text](#)
44. Brost BM, Beier P: **Use of land facets to design linkages for climate change.** *Ecol Appl.* 2012; **22**(1): 87–103. [PubMed Abstract](#) | [Publisher Full Text](#)
45. Beier P: **Conceptualizing and designing corridors for climate change.** *Ecol Restor.* 2012; **30**(4): 312–9. [Publisher Full Text](#)
46. Nuñez TA, Lawler JJ, McRae BH, *et al.*: **Connectivity planning to address climate change.** *Conserv Biol.* 2013; **27**(2): 407–16. [PubMed Abstract](#) | [Publisher Full Text](#)
47. Mackey BG, Watson JEM, Hope G, *et al.*: **Climate change, biodiversity conservation, and the role of protected areas: An Australian perspective.** *Biodiversity.* 2008; **9**(3–4): 11–8. [Publisher Full Text](#)
48. Watson JEM, Fuller RA, Watson AWT, *et al.*: **Wilderness and future conservation priorities in Australia.** *Divers Distrib.* 2009; **15**(6): 1028–36. [Publisher Full Text](#)
49. Hodgson JA, Thomas CD, Corderby S, *et al.*: **Habitat re-creation strategies for promoting adaptation of species to climate change.** *Conserv Lett.* 2011; **4**(4): 289–97. [Publisher Full Text](#)
50. Thomas CD, Cameron A, Green RE, *et al.*: **Extinction risk from climate change.** *Nature.* 2004; **427**(6970): 145–8. [PubMed Abstract](#) | [Publisher Full Text](#)
51. **F** Foden WB, Butchart SHM, Stuart SN, *et al.*: **Identifying the world's most climate change vulnerable species: a systematic trait-based assessment of all birds, amphibians and corals.** *PLoS One.* 2013; **8**(6): e65427. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. Akçakaya HR, Butchart SHM, Watson JEM, *et al.*: **Preventing species extinctions resulting from climate change.** *Nat Clim Change.* 2014; **4**(12): 1048–9. [Publisher Full Text](#)
53. Pearson RG, Stanton JC, Shoemaker KT, *et al.*: **Life history and spatial traits predict extinction risk due to climate change.** *Nat Clim Change.* 2014; **4**(3): 217–21. [Publisher Full Text](#)
54. **F** Stanton JC, Shoemaker KT, Pearson RG, *et al.*: **Warning times for species extinctions due to climate change.** *Glob Chang Biol.* 2015; **21**(3): 1066–77. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
55. **F** Keith DA, Mahony M, Hines H, *et al.*: **Detecting extinction risk from climate change by IUCN Red List criteria.** *Conserv Biol.* 2014; **28**(3): 810–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. Kerr RA: **Climate change. Humans are driving extreme weather; time to prepare.** *Science.* 2011; **334**(6059): 1040. [PubMed Abstract](#) | [Publisher Full Text](#)
57. IPCC: **Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.** Cambridge, UK, Cambridge University Press. 2014. [Reference Source](#)
58. **F** Walther GR: **Community and ecosystem responses to recent climate change.** *Philos Trans R Soc Lond B Biol Sci.* 2010; **365**(1549): 2019–24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
59. Sherwin HA, Montgomery WI, Lundy MG: **The impact and implications of climate change for bats.** *Mammal Rev.* 2013; **43**(3): 171–82. [Publisher Full Text](#)
60. Rebelo H, Tarraso P, Jones G: **Predicted impact of climate change on European bats in relation to their biogeographic patterns.** *Glob Chang Biol.* 2010; **16**(2): 561–76. [Publisher Full Text](#)
61. Hegerl GC, Hanlon H, Beierkuhnlein C: **Climate science: Elusive extremes.** *Nat Geosci.* 2011; **4**(3): 142–3. [Publisher Full Text](#)
62. Ameca y Juárez EI, Mace GM, Cowlshaw G, *et al.*: **Assessing exposure to extreme climatic events for terrestrial mammals.** *Conserv Lett.* 2013; **6**(3): 145–53. [Publisher Full Text](#)
63. Lawler P, Stein BA, Edelson NA: **Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessment.** Washington DC, National Wildlife Federation. 2011. [Reference Source](#)
64. Lankford AJ, Svancara LK, Lawler JJ, *et al.*: **Comparison of climate change vulnerability assessments for wildlife.** *Wildl Soc Bull.* 2014; **38**(2): 386–94. [Publisher Full Text](#)
65. **F** Dawson TP, Jackson ST, House JL, *et al.*: **Beyond predictions: biodiversity conservation in a changing climate.** *Science.* 2011; **332**(6025): 53–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
66. IUCN S: **Guidelines for reintroductions and other conservation translocations.** Gland Switz Camb U K IUCN SSC Re-Intro Group. 2013. [Reference Source](#)
67. **F** McLachlan JS, Hellmann JJ, Schwartz MW: **A framework for debate of assisted migration in an era of climate change.** *Conserv Biol.* 2007; **21**(2): 297–302. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
68. **F** Ricciardi A, Simberloff D: **Assisted colonization is not a viable conservation strategy.** *Trends Ecol Evol.* 2009; **24**(5): 248–53. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
69. Schlaepfer MA, Helenbrook WD, Searing KB, *et al.*: **Assisted colonization: evaluating contrasting management actions (and values) in the face of uncertainty.** *Trends Ecol Evol.* 2009; **24**(9): 471–2. author reply 476–7. [PubMed Abstract](#) | [Publisher Full Text](#)
70. Seddon PJ, Armstrong DP, Soorae P, *et al.*: **The risks of assisted colonization.** *Conserv Biol.* 2009; **23**(4): 788–9. [PubMed Abstract](#) | [Publisher Full Text](#)
71. Vitt P, Havens K, Hoegh-Guldberg O: **Assisted migration: part of an integrated conservation strategy.** *Trends Ecol Evol.* 2009; **24**(9): 473–4. author reply 476–7. [PubMed Abstract](#) | [Publisher Full Text](#)
72. Hewitt N, Klenk N, Smith AL, *et al.*: **Taking stock of the assisted migration debate.** *Biol Conserv.* 2011; **144**(11): 2560–72. [Publisher Full Text](#)
73. Gallagher RV, Makinson RO, Hogbin PM, *et al.*: **Assisted colonization as a climate change adaptation tool.** *Austral Ecol.* 2015; **40**(1): 12–20. [Publisher Full Text](#)
74. Kostyack J, Lawler JJ, Goble DD, *et al.*: **Beyond reserves and corridors: policy solutions to facilitate the movement of plants and animals in a changing climate.** *Bioscience.* 2011; **61**(9): 713–9. [Publisher Full Text](#)
75. **F** Hoegh-Guldberg O, Hughes L, McIntyre S, *et al.*: **Ecology. Assisted colonization and rapid climate change.** *Science.* 2008; **321**(5887): 345–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
76. **F** Richardson DM, Hellmann JJ, McLachlan JS, *et al.*: **Multidimensional evaluation of managed relocation.** *Proc Natl Acad Sci U S A.* 2009; **106**(24): 9721–4. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
77. McDonald-Madden E, Runge MC, Possingham HP, *et al.*: **Optimal timing for managed relocation of species faced with climate change.** *Nat Clim Change.* 2011; **1**(5): 261–5. [Publisher Full Text](#)
78. **F** Fordham DA, Watts MJ, Delean S, *et al.*: **Managed relocation as an adaptation strategy for mitigating climate change threats to the persistence of an endangered lizard.** *Glob Chang Biol.* 2012; **18**(9): 2743–55. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
79. McIntyre S: **Ecological and anthropomorphic factors permitting low-risk assisted colonization in temperate grassy woodlands.** *Biol Conserv.* 2011; **144**(6): 1781–9. [Publisher Full Text](#)
80. Schwartz MW, Hellmann JJ, McLachlan JM, *et al.*: **Managed Relocation: Integrating the Scientific, Regulatory, and Ethical Challenges.** *BioScience.* 2012; **62**(8): 732–43. [Publisher Full Text](#)
81. Klenk NL: **The development of assisted migration policy in Canada: An analysis of the politics of composing future forests.** *Land Use Policy.* 2015; **44**: 101–9. [Publisher Full Text](#)
82. Klenk NL, Larson BMH: **The assisted migration of western larch in British Columbia: A signal of institutional change in forestry in Canada?** *Glob Environ Change.* 2015; **31**(0): 20–7. [Publisher Full Text](#)
83. **F** Thomas CD: **Translocation of species, climate change, and the end of trying to recreate past ecological communities.** *Trends Ecol Evol.* 2011; **26**(5): 216–21. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

84. Seddon PJ, Griffiths CJ, Soorae PS, *et al.*: **Reversing defaunation: restoring species in a changing world.** *Science*. 2014; **345**(6195): 406–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
85. Kareiva P, Marvier M: **What Is Conservation Science?** *BioScience*. 2012; **62**(11): 962–9.
[Publisher Full Text](#)
86. Soule M: **Also seeking common ground in conservation.** *Conserv Biol*. 2014; **28**(3): 637–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
87. Soulé M: **The “new conservation”.** *Conserv Biol*. 2013; **27**(5): 895–7.
[Publisher Full Text](#)
88. Tallis H, Lubchenco J: **Working together: A call for inclusive conservation.** *Nature*. 2014; **515**(7525): 27–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
89. Jones HP, Hole DG, Zavaleta ES: **Harnessing nature to help people adapt to climate change.** *Nat Clim Change*. 2012; **2**: 504–9.
[Publisher Full Text](#)
90. Hobbs RJ, Higgs E, Hall CM, *et al.*: **Managing the whole landscape: historical, hybrid, and novel ecosystems.** *Front Ecol Environ*. 2014; **12**(10): 557–64.
[Publisher Full Text](#)
91. Arkema KK, Guannel G, Verutes G, *et al.*: **Coastal habitats shield people and property from sea-level rise and storms.** *Nat Clim Change*. 2013; **3**(10): 913–8.
[Publisher Full Text](#)
92. Lunt ID, Byrne M, Hellmann JJ, *et al.*: **Using assisted colonisation to conserve biodiversity and restore ecosystem function under climate change.** *Biol Conserv*. 2013; **157**: 172–7.
[Publisher Full Text](#)
93. Watson JEM: **Human Responses to Climate Change will Seriously Impact Biodiversity Conservation: It’s Time We Start Planning for Them.** *Conserv Lett*. 2014; **7**(1): 1–2.
[Publisher Full Text](#)
94. Wyett K: **Escaping a Rising Tide: Sea Level Rise and Migration in Kiribati.** *Asia Pac Policy Stud*. 2014; **1**(1): 171–85.
[Publisher Full Text](#)
95. Birk T, Rasmussen K: **Migration from atolls as climate change adaptation: Current practices, barriers and options in Solomon Islands.** *Nat Resour Forum*. 2014; **38**(1): 1–13.
[Publisher Full Text](#)
96. Chornesky EA, Ackerly DD, Beier P, *et al.*: **Adapting California’s Ecosystems to a Changing Climate.** *BioScience*. 2015; **65**(3): 247–62.
[Publisher Full Text](#)

Open Peer Review

Current Referee Status:



Editorial Note on the Review Process

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty and are edited as a service to readers. In order to make these reviews as comprehensive and accessible as possible, the referees provide input before publication and only the final, revised version is published. The referees who approved the final version are listed with their names and affiliations but without their reports on earlier versions (any comments will already have been addressed in the published version).

The referees who approved this article are:

Version 1

- 1 **Bruce Stein**, Climate Adaptation and Resilience, National Wildlife Federation, Washington, DC, USA
Competing Interests: No competing interests were disclosed.
- 2 **A Townsend Peterson**, Biodiversity Institute and Department of Ecology and Evolutionary Biology, The University of Kansas, Lawrence, KS, USA
Competing Interests: No competing interests were disclosed.
- 3 **Paul Beier**, School of Forestry, Northern Arizona University, Flagstaff, AZ, USA
Competing Interests: No competing interests were disclosed.