



HHS Public Access

Author manuscript

Biochem Biophys Res Commun. Author manuscript; available in PMC 2017 February 19.

Published in final edited form as:

Biochem Biophys Res Commun. 2016 February 19; 470(4): 913–916. doi:10.1016/j.bbrc.2016.01.151.

Plant homologs of mammalian MBT-domain protein-regulated KDM1 histone lysine demethylases do not interact with plant Tudor/PWWP/MBT-domain proteins

Irfan Sadiq^{a,b}, Ido Keren^a, and Vitaly Citovsky^{a,*}

^a Department of Biochemistry and Cell Biology, State University of New York, Stony Brook, NY 11794-5215

^b Department of Biosciences, COMSATS Institute of Information Technology Islamabad, Park Road, Islamabad, 44000, Pakistan

Abstract

Histone lysine demethylases of the LSD1/KDM1 family play important roles in epigenetic regulation of eukaryotic chromatin, and they are conserved between plants and animals. Mammalian LSD1 is thought to be targeted to its substrates, i.e., methylated histones, by an MBT-domain protein SFMBT1 that represents a component of the LSD1-based repressor complex and binds methylated histones. Because MBT-domain proteins are conserved between different organisms, from animals to plants, we examined whether the KDM1-type histone lysine demethylases KDM1C and FLD of *Arabidopsis* interact with the *Arabidopsis* Tudor/PWWP/MBT-domain SFMBT1-like proteins SL1, SL2, SL3, and SL4. No such interaction was detected using the bimolecular fluorescence complementation assay in living plant cells. Thus, plants most likely direct their KDM1 chromatin-modifying enzymes to methylated histones of the target chromatin by a mechanism different from that employed by the mammalian cells.

Keywords

KDM1; histone lysine demethylases; Tudor/PWWP/MBT domain proteins; *Arabidopsis*

1. Introduction

Post-translational histone modifications, e.g., acetylation, methylation, and ubiquitination, play central role in gene regulation in all eukaryotic organisms, determining the active or inactive state of the chromatin. These modifications are effected by diverse histone-modifying enzymes, such as histone deacetylases, histone lysine demethylases, histone methyltransferases, and histone deubiquitinases. Among those, LSD1/KDM1-type histone

* Corresponding author. vitaly.citovsky@stonybrook.edu.
Irfan Sadiq: irfan_sadiq@comsats.edu.pk
Ido Keren: ido.keren@stonybrook.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

lysine demethylases [1] represent one of the most recently discovered [2], yet clearly central factors in controlling chromatin in different organisms, from animals to plants [3-5]. In animal cells, LSD1/KDM1A promotes demethylation of dimethylated lysine 4 (K4) of histone H3 [2] and often functions as negative regulator of gene expression [6]. LSD1/KDM1A usually acts in complex with CoREST that modulates the gene repression function of LSD1 *in vivo*, a histone methyltransferase (HMT; e.g., G9a), HDAC1/2 histone deacetylases, and DNA binding zinc finger proteins (e.g., either REST or ZNF217) [2, 7-17]. However, none of these components recognize methylated histones that represent the direct substrates of LSD1/KDM1A; recently, this role has been attributed to the SFMBT1 [Scm (Sex comb on midleg) with four MBT (malignant brain tumor) domains 1] protein, which functions as part of the LSD1/KDM1A-based repressor complex and is known to bind different forms of methylated histones [18, 19].

In plants, our knowledge of LSD1/KDM1 complexes is just emerging. For example, *Arabidopsis*, one of the major model plants, encodes four KDM1 proteins, KDM1A (FLD), KDM1B, KDM1C, and KDM1D [1]. KDM1C has been shown to interact with histone methyltransferase SURV5 [20-22] and histone deubiquitinase OTLD1 [23]. MBT domains are conserved within proteins from different organisms, from animals to insects to plants [24, 25]. Thus, we examined whether *Arabidopsis* LSD1/KDM1A-type histone lysine demethylases can recognize plant Tudor/PWWP/MBT-domain proteins [24, 25], i.e., the *Arabidopsis* SFMBT1-like proteins (SLs). Our data indicate no such interactions and suggest that plant LSD1/KDM1 histone lysine demethylases are directed to their substrates by a mechanism different from their mammalian counterparts.

2. Materials and methods

2.1. Plasmid construction

The coding sequences of *SL1*, *SL2*, *SL3*, and *SL4*, *KDM1C* and *FLD* were amplified from *Arabidopsis thaliana* cDNA library using primers detailed in Table S1. The *SL5* sequence failed to amplify and was not examined in this work. For transient expression in *N. benthamiana*, the amplified *SL1*, *SL2*, *SL3*, and *SL4* as well as *SURV5* [20] were cloned into the HindII-SalI, EcoRI-SalI, XhoI-SalI, SalI-SacII, and SalI sites of pSAT4-nYFP [26], respectively, resulting in pSAT4-SL1-nYFP, pSAT4-SL2-nYFP, pSAT4-SL3-nYFP, pSAT4-SL4-nYFP, and pSAT4-SURV5-nYFP. The amplified *KDM1C* and *KDM1A* (*FLD*) were cloned into the XhoI-SmaI and SacI-SalI sites of pSAT1-cYFP [26], respectively, resulting in pSAT1-cYFP-KDM1C and pSAT1-cYFP-FLD. Then, the expression cassettes from pSAT1-cYFP-KDM1C and pSAT1-cYFP-FLD were transferred into the AscI site of the binary vector for multigene expression pPZP-RCSII [27], resulting in pRCSII-KDM1C and pRCSII-FLD. Finally, the expression cassettes from pSAT4-SL1-nYFP, pSAT4-SL2-nYFP, pSAT4-SL3-nYFP, and pSAT4-SL4-nYFP were transferred into the I-SceI site of pRCSII-KDM1C and pRCSII-FLD, resulting in pRCSII-KDM1C-SL1, pRCSII-KDM1C-SL2, pRCSII-KDM1C-SL3, pRCSII-KDM1C-SL4, pRCSII-FLD-SL1, pRCSII-FLD-SL2, pRCSII-FLD-SL3, and pRCSII-FLD-SL4. For positive control experiments, the expression cassette from pSAT4-SURV5-nYFP was subcloned into the I-SceI site of pRCSII-KDM1C, resulting in pRCSII-KDM1C-SURV5.

2.2. BiFC assay

Nicotiana benthamiana plants were grown to a six-leaf stage at 25 °C with 16 h light (70-80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and 8 h dark. The tested proteins were transiently expressed in *N. benthamiana* plants by agroinfiltration as described [28] with modifications. Briefly, the binary constructs were transferred into *Agrobacterium tumefaciens* strain EHA105 by heat shock. The bacterial cells then were grown on LB medium supplemented with rifampicin (25 $\mu\text{g mL}^{-1}$) and spectinomycin (100 $\mu\text{g mL}^{-1}$) at 28°C for 48 h, resuspended in the infiltration buffer (10 mM MgCl_2 , 10 mM MES pH 5.6, 150 mM acetosyringone) to $A_{600} = 1.0$, and grown at room temperature for another 2-3 h. The abaxial side of mature leaves of 4-5-week-old *N. benthamiana* plants was vacuum-infiltrated with bacterial cell suspension harboring each tested construct. Plants were grown for additional 72 h, and the BiFC signal in the infiltrated areas was detected using a Zeiss LSM 5 Pascal confocal microscope. All experiments were repeated at least three times.

3. Results and Discussion

3.1. Arabidopsis SL proteins

Amino acid sequence analysis identified five relatively close homologs of the human SFMBT1 protein encoded by the Arabidopsis genome. These sequences were aligned with SFMBT1 by CLC Main Workbench 7.6.4 (<http://www.clcbio.com>) software, using default parameters (Fig. S1), and their phylogenetic tree was generated (Fig. 1). All of these five genes, i.e., *SL1* (At1g51745), *SL2* (At1g80810), *SL3* (At2g48160), *SL4* (At3g63070), and *SL5* (At5g08230,) belong to the Tudor domain “Royal family”, which includes proteins with Tudor, plant Agenet, Chromo, PWWP and MBT domains [24]. Protein products of the *SL3*, *SL4*, and *SL5* genes, known as members of the *HULK* gene family, localize in the plant cell nucleus and play important roles in regulating flowering timing [29]. The presence of Tudor/PWWP/MBT domains raised a possibility that these plant proteins, similarly to their mammalian homologs [18, 19], might act to direct KDM1 proteins to their sites of action on the target chromatin. In this scenario, the SL proteins should interact with KDM1 proteins. This idea was examined directly by monitoring potential KDM1-SL interactions in the living plant cells.

3.2. KDM1A and KDM1C do not recognize the SL1-4 proteins

Mammalian SFMBT1 is suggested to interact with LSD1/KDM1A through another component of the repressor complex CoREST [18]; however, plants do not encode CoREST homologs, suggesting that such interactions may happen directly, without the involvement of CoREST-like proteins. Thus, we assayed whether the SL1, SL2, or SL3 and SL4 proteins can interact with two best-studied representatives of the Arabidopsis KDM1 family, KDM1A (FLD) and KDM1C [1, 20, 30, 31]. The potential interactions were tested using bimolecular fluorescence complementation (BiFC) within living plant cells [26, 32]. Each of the tested KDM1 and SL proteins were tagged with carboxyl- and amino-terminal fragments of YFP, respectively, and transiently coexpressed in *N. benthamiana* leaves.

Fig. 2 shows that coexpression of cYFP-FLD with nYFP-SL1, nYFP-SL2, nYFP-SL3, or nYFP-SL4 did not produce a BiFC signal, indicating that FLD was unable to interact with

any of these SL1-4 proteins. In positive control experiments, the YFP fluorescence was reconstituted following coexpression of two known interactors KDM1C and SURV5 [20] tagged with cYFP and nYFP, respectively. As expected [20], the cYFP-KDM1C-nYFP-SURV5 complexes were observed in the cell nuclei (Fig. 2).

Coexpression of cYFP-KDM1C with nYFP-SL1, nYFP-SL2, nYFP-SL3, or nYFP-SL4 failed to regenerate the YFP signal, indicating no interaction between KDM1C and any of these SL proteins (Fig. 3). In this set of experiments, the positive control also detected the KDM1CSURV5 interaction following coexpression of cYFP-KDM1C and nYFP-SURV5, with the interacting proteins accumulating in the cell nucleus (Fig. 3). Collectively, these experiments indicate that neither FLD nor KDM1C bind any of the tested Tudor/PWWP/MBT-domain proteins SL1, SL2, SL3, or SL4.

These results also suggest that, similarly to KDM1C and FLD, other plant KDM1 histone lysine demethylases most likely do not recognize proteins containing the Tudor/PWWP/MBT domains. Potentially, plants may encode an as yet unidentified protein that mediates these interactions. Yet, in this scenario, plant cells should contain certain levels of this factor, which would be expected to support at least some degree of interactions of the coexpressed tested proteins. Thus, plants apparently do not direct their KDM1 chromatin-modifying enzymes to methylated histones of the target chromatin by the same mechanism that is employed by the mammalian cells.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

I.S. was supported in part by the Fulbright Fellowship. The work in the V.C. laboratory is supported by grants from NIH, NSF, USDA/NIFA, BARD, and BSF to V.C.

References

1. Zhou X, Ma H. Evolutionary history of histone demethylase families: distinct evolutionary patterns suggest functional divergence. *BMC Evol. Biol.* 2008; 8:294. [PubMed: 18950507]
2. Shi Y, Lan F, Matson C, Mulligan P, Whetstone JR, Cole PA, Casero RA, Shi Y. Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell.* 2004; 119:941–953. [PubMed: 15620353]
3. Mosammaparast N, Kim H, Laurent B, Zhao Y, Lim HJ, Majid MC, Dango S, Luo Y, Hempel K, Sowa ME, Gygi SP, Steen H, Harper JW, Yankner B, Shi Y. The histone demethylase LSD1/KDM1A promotes the DNA damage response. *J. Cell Biol.* 2013; 203:457–470. [PubMed: 24217620]
4. Prakash S, Singh R, Lodhi N. Histone demethylases and control of gene expression in plants. *Cell. Mol. Biol.* 2014; 60:97–105. [PubMed: 25535719]
5. Rudolph T, Beuch S, Reuter G. Lysine-specific histone demethylase LSD1 and the dynamic control of chromatin. *Biol. Chem.* 2013; 394:1019–1028. [PubMed: 23612539]
6. Jepsen K, Rosenfeld MG. Biological roles and mechanistic actions of co-repressor complexes. *J. Cell Sci.* 2002; 115:689–698. [PubMed: 11865025]
7. Ballas N, Battaglioli E, Atouf F, Andres ME, Chenoweth J, Anderson ME, Burger C, Moniwa M, Davie JR, Bowers WJ, Federoff HJ, Rose DW, Rosenfeld MG, Brehm P, Mandel G. Regulation of

- neuronal traits by a novel transcriptional complex. *Neuron*. 2001; 31:353–365. [PubMed: 11516394]
8. Shi YJ, Matson C, Lan F, Iwase S, Baba T, Shi Y. Regulation of LSD1 histone demethylase activity by its associated factors. *Mol. Cell*. 2005; 19:857–864. [PubMed: 16140033]
 9. Yang M, Gocke CB, Luo X, Borek D, Tomchick DR, Machius M, Otwinowski Z, Yu H. Structural basis for CoREST-dependent demethylation of nucleosomes by the human LSD1 histone demethylase. *Mol. Cell*. 2006; 23:377–387. [PubMed: 16885027]
 10. You A, Tong JK, Grozinger CM, Schreiber SL. CoREST is an integral component of the CoREST-human histone deacetylase complex. *Proc. Natl. Acad. Sci. USA*. 2001; 98:1454–1458. [PubMed: 11171972]
 11. Chong JA, Tapia-Ramirez J, Kim S, Toledo-Aral JJ, Zheng Y, Boutros MC, Altschuler YM, Frohman MA, Kraner SD, Mandel G. REST: a mammalian silencer protein that restricts sodium channel gene expression to neurons. *Cell*. 1995; 80:949–957. [PubMed: 7697725]
 12. Schoenherr CJ, Anderson DJ. The neuron-restrictive silencer factor (NRSF): a coordinate repressor of multiple neuron-specific genes. *Science*. 1995; 267:1360–1363. [PubMed: 7871435]
 13. Tachibana M, Sugimoto K, Fukushima T, Shinkai Y. Set domain-containing protein, G9a, is a novel lysine-preferring mammalian histone methyltransferase with hyperactivity and specific selectivity to lysines 9 and 27 of histone H3. *J. Biol. Chem*. 2001; 276:25309–25317. [PubMed: 11316813]
 14. Ropra A, Qazi R, Schoenike B, Daley TJ, Morrison JF. Localized domains of G9a-mediated histone methylation are required for silencing of neuronal genes. *Mol. Cell*. 2004; 14:727–738. [PubMed: 15200951]
 15. Ballas N, Mandel G. The many faces of REST oversee epigenetic programming of neuronal genes. *Curr. Opin. Neurobiol*. 2005; 15:500–506. [PubMed: 16150588]
 16. Grimes JA, Nielsen SJ, Battaglioli E, Miska EA, Speh JC, Berry DL, Atouf F, Holdener BC, Mandel G, Kouzarides T. The co-repressor mSin3A is a functional component of the REST-CoREST repressor complex. *J. Biol. Chem*. 2000; 275:9461–9467. [PubMed: 10734093]
 17. Wang J, Scully K, Zhu X, Cai L, Zhang J, Prefontaine GG, Krones A, Ohgi KA, Zhu P, Garcia-Bassets I, Liu F, Taylor H, Lozach J, Jayes FL, Korach KS, Glass CK, Fu XD, Rosenfeld MG. Opposing LSD1 complexes function in developmental gene activation and repression programmes. *Nature*. 2007; 446:882–887. [PubMed: 17392792]
 18. Tang M, Shen H, Jin Y, Lin T, Cai Q, Pinard MA, Biswas S, Tran Q, Li G, Shenoy AK, Tongdee E, Lin S, Gu Y, Law BK, Zhou L, McKenna R, Wu L, Lu J. The Malignant Brain Tumor (MBT) domain protein SFMBT1 is an integral histone reader subunit of the LSD1 demethylase complex for chromatin association and epithelial-to-mesenchymal transition. *J. Biol. Chem*. 2013; 288:27680–27691. [PubMed: 23928305]
 19. Zhang J, Bonasio R, Strino F, Kluger Y, Holloway JK, Modzelewski AJ, Cohen PE, Reinberg D. SFMBT1 functions with LSD1 to regulate expression of canonical histone genes and chromatin-related factors. *Genes Dev*. 2013; 27:749–766. [PubMed: 23592795]
 20. Krichevsky A, Gutgarts H, Kozlovsky SV, Tzfira T, Sutton A, Sternglanz R, Mandel G, Citovsky V. C2H2 zinc finger-SET histone methyltransferase is a plant-specific chromatin modifier. *Dev. Biol*. 2007; 303:259–269. [PubMed: 17224141]
 21. Krichevsky A, Kozlovsky SV, Gutgarts H, Citovsky V. Arabidopsis co-repressor complexes containing polyamine oxidase-like proteins and plant-specific histone methyltransferases. *Plant Signal. Behav*. 2007; 2:174–177. [PubMed: 19704688]
 22. Caro E, Stroud H, Greenberg MV, Bernatavichute YV, Feng S, Groth M, Vashisht AA, Wohlschlegel J, Jacobsen SE. The SET-domain protein SUV5 mediates H3K9me2 deposition and silencing at stimulus response genes in a DNA methylation-independent manner. *PLOS Genet*. 2012; 8:e1002995. [PubMed: 23071452]
 23. Krichevsky A, Lacroix B, Zaltsman A, Citovsky V. Involvement of KDM1C histone demethylase-OTLD1 otubain-like histone deubiquitinase complexes in plant gene repression. *Proc. Natl. Acad. Sci. USA*. 2011; 108:11157–11162. [PubMed: 21690391]

24. Maurer-Stroh S, Dickens NJ, Hughes-Davies L, Kouzarides T, Eisenhaber F, Ponting CP. The Tudor domain “Royal Family”: Tudor, plant Agenet, Chromo, PWWP and MBT domains. *Trends Biochem. Sci.* 2003; 28:69–74. [PubMed: 12575993]
25. Alvarez-Venegas R, Avramova Z. Evolution of the PWWP-domain encoding genes in the plant and animal lineages. *BMC Evol. Biol.* 2012; 12:101. [PubMed: 22734652]
26. Citovsky V, Lee LY, Vyas S, Glick E, Chen MH, Vainstein A, Gafni Y, Gelvin SB, Tzfira T. Subcellular localization of interacting proteins by bimolecular fluorescence complementation in planta. *J. Mol. Biol.* 2006; 362:1120–1131. [PubMed: 16949607]
27. Tzfira T, Tian GW, Lacroix B, Vyas S, Li J, Leitner-Dagan Y, Krichevsky A, Taylor T, Vainstein A, Citovsky V. pSAT vectors: a modular series of plasmids for fluorescent protein tagging and expression of multiple genes in plants. *Plant Mol. Biol.* 2005; 57:503–516. [PubMed: 15821977]
28. Magori S, Citovsky V. Agrobacterium counteracts host-induced degradation of its F-box protein effector. *Sci. Signal.* 2011; 4:ra69. [PubMed: 22009152]
29. Jali SS, Rosloski SM, Janakirama P, Steffen JG, Zhurov V, Berleth T, Clark RM, Grbic V. A plant-specific HUA2-LIKE (HULK) gene family in *Arabidopsis thaliana* is essential for development. *Plant J.* 2014; 80:242–254. [PubMed: 25070081]
30. Jiang D, Yang W, He Y, Amasino RM. Arabidopsis relatives of the human Lysine-Specific Demethylase1 repress the expression of FWA and FLOWERING LOCUS C and thus promote the floral transition. *Plant Cell.* 2007; 19:2975–2987. [PubMed: 17921315]
31. He Y, Michaels SD, Amasino RM. Regulation of flowering time by histone acetylation in *Arabidopsis*. *Science.* 2003; 302:1751–1754. [PubMed: 14593187]
32. Hu CD, Chinenov Y, Kerppola TK. Visualization of interactions among bZIP and Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol. Cell.* 2002; 9:789–798. [PubMed: 11983170]
33. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment tool. *J. Mol. Biol.* 1990; 215:403–410. [PubMed: 2231712]

Highlights

- Interaction of plant SFMBT1-like proteins with KDM1 histone demethylases was tested
- Four Arabidopsis homologs of the mammalian SFMBT1 were selected for study
- Two Arabidopsis KDM1 histone demethylases, KDM1C and FLD, were tested
- None of the Arabidopsis SFMBT1 homologs showed interaction with KDM1C or FLD
- Plants and animals may use different mechanisms to direct KDM1 to methyl-histones




Fig. 1. Phylogenetic tree of human SFMBT1 and its Arabidopsis SL1, SL2, SL3, SL4, and SL5 homologs. The SL proteins were identified by the BLASTA search [33]. GenBank accession numbers are shown in parentheses next to each protein name. The tree was built by the CLC Main Workbench 7.6.4 software (<http://www.clcbio.com>), using default parameters.




Fig. 2. The BiFC assay for possible interactions between SL1, SL2, SL3, or SL4 and FLD. Constructs encoding the tested proteins were coexpressed in agroinfiltrated *N. benthamiana* leaves. The interaction between KDM1C and SURV5 served as positive control. YFP signal is in green. Fluorescence images are single confocal sections. Scale bar = 20 μ m.




Fig. 3. The BiFC assay for possible interactions between SL1, SL2, SL3, or SL4 and KDM1C. Constructs encoding the tested proteins were coexpressed in agroinfiltrated *N. benthamiana* leaves. The interaction between KDM1C and SURV5 served as positive control. YFP signal is in green. Fluorescence images are single confocal sections. Scale bar = 20 μ m.