

Erratum to: Torniainen M, Mittendorfer-Rutz E, Tanskanen A, Björkenstam C, Suvisaari J, Alexanderson K, Tiihonen J. Antipsychotic treatment and mortality in schizophrenia. *Schizophr Bull.* 2015;41(3):656–663.

In **Table 1**, total figures of Females, High education level, Previous inpatient care due to psychosis and Receiving disability pension in 2005, as well as some of the percentages were incorrect. The correct figures are below in bold. In addition, in Table 3, the number of deaths among males with Moderate antipsychotic exposure should be 344 (instead of 433). The authors regret that the following errors occurred, yet the main findings of the study remain the same.

Table 1. Demographic and Clinical Characteristics of Patients

	Total		No antipsychotic exposure		Low antipsychotic exposure		Moderate antipsychotic exposure		High antipsychotic exposure	
<i>n</i> (%)	21 492		2 077	(10%)	4110	(19%)	8468	(39%)	6837	(32%)
Gender										
Female, <i>n</i> (%)	8442	(39%)	696	(34%)	1780	(43%)	3472	(41%)	2494	(36%)
Male, <i>n</i> (%)	13 050	(61%)	1381	(66%)	2330	(57%)	4996	(59%)	4343	(64%)
Age, M (SD)	45.5	(11.1)	45.3	(11.5)	46.0	(11.4)	45.4	(11.2)	45.2	(10.8)
Educational level, <i>n</i> (%)										
Low (≤ 9 years)	7457	(35%)	640	(31%)	1224	(30%)	2809	(33%)	2784	(41%)
Medium	11 257	(52%)	1090	(52%)	2220	(54%)	4551	(54%)	3396	(50%)
High (≥ 12 years)	2311	(11%)	284	(14%)	583	(14%)	963	(11%)	481	(7%)
Information missing	467	(2%)	63	(3%)	83	(2%)	145	(2%)	176	(3%)
Marital status, <i>n</i> (%)										
Married or living with a partner	2782	(13%)	349	(17%)	701	(17%)	1144	(14%)	588	(9%)
Previous treatment contact										
Outpatient care due to psychosis	3799	(18%)	347	(17%)	902	(22%)	1665	(20%)	885	(13%)
Previous inpatient care due to psychosis	11 295	(53%)	946	(46%)	2042	(50%)	4544	(54%)	3763	(55%)
In hospital during 2005 due to psychosis	4267	(20%)	176	(8%)	641	(16%)	1578	(19%)	1872	(27%)
Identified only through disability pension due to schizophrenia	2131	(10%)	608	(29%)	525	(13%)	681	(8%)	317	(5%)
Months since the first psychosis in the beginning of follow-up	70.3	(38.4)	72.8	(40.4)	64.4	(38.7)	68.3	(38.0)	75.5	(38.0)
Receiving disability pension in 2005	18 717	(87%)	1625	(78%)	3317	(81%)	7383	(87%)	6392	(93%)
First episode patients <i>n</i> (%)	1230		232	(19)	329	(27)	329	(27)	340	(28)
Gender										
Male, <i>n</i> (%)	806	(66%)	154	(66%)	213	(65%)	201	(61%)	238	(70%)
Female, <i>n</i> (%)	424	(34%)	78	(34%)	116	(35%)	128	(39%)	102	(30%)
Age, M (SD)	36.3	(11.7)	40.2	(12.1)	36.8	(11.5)	35.6	(11.4)	33.8	(11.3)