

**Ditengou FA, Teale WD, Kochersperger P, et al.** 2008. Mechanical induction of lateral root initiation in *Arabidopsis thaliana*. Proceedings of the National Academy of Sciences, USA **105**, 18818–18823.

**Dubrovsky JG, Sauer M, Napsucialy-Mendivil S, Ivanchenko MG, Friml J, Shishkova S, Celenza J, Benkova E.** 2008. Auxin acts as a local morphogenetic trigger to specify lateral root founder cells. Proceedings of the National Academy of Sciences, USA **105**, 8790–8794.

**Kircher S, Schopfer M.** 2016. Priming and positioning of lateral roots in *Arabidopsis*. An approach for an integrating concept. Journal of Experimental Botany **67**, 1411–1420.

**Laskowski M.** 2013. Lateral root initiation is a probabilistic event whose frequency is set by fluctuating levels of auxin response. Journal of Experimental Botany **64**, 2609–2617.

**Laskowski M, Grieneisen VA, Hoffhuis H, ten Hove CA, Hogeweg P, Maree AFM, Scheres B.** 2008. Root system architecture from coupling cell shape to auxin transport. Plos Biology **6**, 2721–2735.

**Lucas M, Godin C, Jay-Allemand C, Laplaze L.** 2008. Auxin fluxes in the root apex co-regulate gravitropism and lateral root initiation. Journal of Experimental Botany **59**, 55–66.

**Moreno-Risueno MA, Van Norman JM, Moreno A, Zhang JY, Ahnert SE, Benfey PN.** 2010. Oscillating gene expression determines competence for periodic *Arabidopsis* root branching. Science **329**, 1306–1311.

**Okushima Y, Fukaki H, Onoda M, Theologis A, Tasaka M.** 2007. ARF7 and ARF19 regulate lateral root formation via direct activation of LBD/ASL genes in *Arabidopsis*. The Plant Cell **19**, 118–130.

**Peret B, De Rybel B, Casimiro I, Benkova E, Swarup R, Laplaze L, Beckman T, Bennett MJ.** 2009. *Arabidopsis* lateral root development: an emerging story. Trends in Plant Science **14**, 399–408.

**Richter GL, Monshausen GB, Krol A, Gilroy S.** 2009. Mechanical stimuli modulate lateral root organogenesis. Plant Physiology **151**, 1855–1866.

**Van Norman JM, Xuan W, Beckman T, Benfey PN.** 2013. To branch or not to branch: the role of pre-patterning in lateral root formation. Development **140**, 4301–4310.

**Xuan W, Audenaert D, Parizot B, et al.** 2015. Root cap-derived auxin pre-patterns the longitudinal axis of the *Arabidopsis* root. Current Biology **25**, 1381–1388.

## Insight

# Flavonoid biosynthesis and *Arabidopsis* genetics: more good music

Anton R. Schäffner

Institute of Biochemical Plant Pathology, Helmholtz Zentrum München, German Research Center for Environmental Health (GmbH), Ingolstädter Landstr. 1, 85764 Neuherberg, Germany  
[schaeffner@helmholtz-muenchen.de](mailto:schaeffner@helmholtz-muenchen.de)

**In this issue of *Journal of Experimental Botany* (pages 1505–1517), Ishihara *et al.* report the identification of a gene responsible for the production of flavonol 3-*O*-gentiobioside 7-*O*-rhamnosides by elegantly tackling the ivories of *Arabidopsis* genetics and genetic resources combined with straightforward metabolite analysis: it is a model case of functional evaluation.**

Flavonoids are secondary metabolites derived from the phenylpropanoid biosynthetic pathway that occur in a huge number and variety in plants. They have been implicated in diverse processes, including pigmentation, redox and UV protection, plant–microbe interactions, development and regulation of auxin transport (Winkel-Shirley, 2006; Yonekura-Sakakibara *et al.*, 2008; Kuhn *et al.*, 2011; Peer *et al.*, 2011; Grunewald *et al.*, 2012; Buer *et al.*, 2013; Emiliani *et al.*, 2013; Saito *et al.*, 2013; Yin *et al.*, 2014; Ishihara *et al.*, 2016).

Genetics and flavonoid biosynthesis were already successfully engaged on the verge of *Arabidopsis* becoming the plant model organism. The *transparent testa* (*tt*) mutant loci, which affect the biosynthesis of flavonoids, defined easily scorable genetic markers due to the loss of seed coat pigmentation. Their molecular identification established many crucial steps in the biosynthesis of the flavonoid core structure (Koornneef *et al.*, 1983; Shirley *et al.*, 1995; Winkel-Shirley, 2006; Saito *et al.*, 2013). This core is formed by a phenyl ring condensed with an oxygen-containing heterocycle in different oxidation states which is further substituted at different positions with another phenyl side group. These variable cores constitute different flavonoid classes occurring

in plants and within a given plant species. However, only the decoration of these aglycones with various carbohydrate side chains and further chemical modification provides the full flavonoid range.

Again, genetics in combination with biochemistry, metabolite analyses and, in particular, gene co-expression patterns led to the identification of several UDP-carbohydrate-dependent glycosyltransferases (UGTs) conjugating flavonoids with different carbohydrates at different positions (Jones *et al.*, 2003; Yonekura-Sakakibara *et al.*, 2008, 2012). Nevertheless, there are still unresolved cases, one of them being the production of the *Arabidopsis* accession-specific flavonol 3-*O*-gentiobioside 7-*O*-rhamnosides (F3GG7Rs), comprising the flavonols kaempferol, quercetin or isorhamnetin with the specific carbohydrate decoration 3GG7R, which are most probably derived through glucosylation from flavonol 3-*O*-glucoside 7-*O*-rhamnoside (F3G7R) precursors.

## A novel flavonol glucosyltransferase

Ishihara *et al.* (2016) have now identified a gene responsible for this final step in the production of F3GG7Rs. Previous reports that the accessions *Ler* and Nö-0, but not Col-0 and Cvi, contain F3GG7Rs were extended to a collection of 81 accessions, of which just half were F3GG7R-producers. The *Ler* F3GG7R trait was inherited in a dominant manner in a cross with the F3GG7R-lacking Col-0. Linkage analysis using an F3GG7R-metabotyped, 95-member-sized *Ler* × Col recombinant inbred (RI) population as

well as recombination events within the originally identified interval using a further 200 additional RI lines eventually revealed the locus responsible in a small 87 kb region on chromosome 1. No obvious candidate, such as a *UGT* gene, was located in that interval; however, genome-wide association mapping of the F3GG7R metabotype of the 81 accessions confirmed the RI linkage mapping and eventually identified a single-nucleotide polymorphism leading to a premature stop codon in the Col-0 allele of *BETA GLUCOSIDASE 6 (BGLU6)*. In contrast, *Ler* contains a fully functional *BGLU6* gene and F3GG7R production was generally associated with functional *BGLU6* alleles.

Transcriptional co-expression analysis has been shown to be a valuable tool in the study of flavonol biosynthesis (see above); thus, the association of *BGLU6* with this pathway further supported its likely involvement in F3GG7R biosynthesis. Nevertheless, this finding is remarkable at first sight, since the underlying expression data were primarily derived from the F3GG7R-deficient Col-0 harbouring only the *BGLU6* pseudogene. However, the promoter sequences of the *Ler* and Col-0 alleles are highly similar and thus transcriptional co-regulation was not affected by either functional (*Ler*) or non-functional (Col) gene transcripts. On the other hand, [Ishihara et al. \(2016\)](#) point out based on published RNA-Seq data that the expression level of functional *BGLU6* alleles was about twofold higher than the transcription from *BGLU6* alleles leading to transcripts harbouring the premature stop codon. The reason for this negative impact on the abundance of the non-functional transcript (either transcription or stability of the mRNA) is not clear, but it may be an interesting future issue in relation to the pseudogenization of gene copies.

After this genetic free-form jazz, the scales were still completed successfully: luckily, *BGLU6*-targeting insertion lines in two F3GG7R-accumulating accessions, *Ler* and *Ws-4*, were available and both led to the loss of F3GG7R production. Conversely, the F3GG7R-deficient Col-0 gained the ability to synthesize F3GG7R after genetic transformation with a functional *Ler BGLU6* gene fragment.

## Genetics leading the way

Biochemical proof by an *in vitro* enzymatic activity test could not be provided by [Ishihara et al. \(2016\)](#), since expression of the recombinant *BGLU6* protein failed in several systems. However, genetics has provided overwhelming evidence that *BGLU6* is indeed responsible for F3GG7R formation. This not only adds another piece of information about the complex formation of flavonol glycosides in *Arabidopsis*, but also provides strong evidence that acyl-carbohydrates utilized as sugar donors by beta-glucosidases, such as the putative beta-glucosidase *BGLU6*, are involved in flavonol glycosylation in addition to the well-known UDP-carbohydrate donors used by UGTs. This extends recent reports on beta-glucosidases being involved in *Arabidopsis* anthocyanin glycosylation ([Miyahara et al., 2013](#)).

Nevertheless, the identification of this new molecular player being responsible for producing the accession-specific F3GG7R flavonol glycosides could not provide clues to a

specific physiological or ecological role. The same is mostly true for the plethora of specifically decorated flavonoids. Most probably, only genetics will be able to lead the way to unraveling such functional relations between specific flavonoid glycosides and particular processes and functions ([Yin et al., 2014](#)). More music expected.

Key words: *Arabidopsis thaliana*, flavonoid, flavonol glucosyltransferase, glycoside hydrolase-type, natural variation, whole-genome association mapping.

Journal of Experimental Botany, Vol. 67 No. 5 pp. 1203-1204, 2016 doi: 10.1093/jxb/erw050

## References

- Buer CS, Kordbacheh F, Truong TT, Hocart CH, Djordjevic MA.** 2013. Alteration of flavonoid accumulation patterns in *transparent testa* mutants disturbs auxin transport, gravity responses, and imparts long-term effects on root and shoot architecture. *Planta* **238**, 171–189.
- Emiliani J, Grotewold E, Falcone Ferreyra ML, Casati P.** 2013. Flavonols protect *Arabidopsis* plants against UV-B deleterious effects. *Molecular Plant* **6**, 1376–1379.
- Grunewald W, De Smet I, Lewis DR, et al.** 2012. Transcription factor WRKY23 assists auxin distribution patterns during *Arabidopsis* root development through local control on flavonol biosynthesis. *Proceedings of the National Academy of Sciences, USA* **109**, 1554–1559.
- Ishihara H, Tohge T, Viehöver P, Fernie AR, Weisshaar B, Stracke R.** 2016. Natural variation in flavonol accumulation in *Arabidopsis* is determined by the flavonol glucosyltransferase *BGLU6*. *Journal of Experimental Botany* **67**, 1505–1517.
- Jones P, Messner B, Nakajima J, Schäffner AR, Saito K.** 2003. UGT73C6 and UGT78D1, glucosyltransferases involved in flavonol glycoside biosynthesis in *Arabidopsis thaliana*. *The Journal of Biological Chemistry* **278**, 43910–43918.
- Koornneef M, van Eden J, Hanhart CJ, Stam P, Braaksma FJ, Feenstra WJ.** 1983. Linkage map of *Arabidopsis thaliana*. *Journal of Heredity* **74**, 265–272.
- Kuhn BM, Geisler M, Bigler L, Ringli C.** 2011. Flavonols accumulate asymmetrically and affect auxin transport in *Arabidopsis*. *Plant Physiology* **156**, 585–595.
- Miyahara T, Sakiyama R, Ozeki Y, Sasaki N.** 2013. Acyl-glucose-dependent glucosyltransferase catalyzes the final step of anthocyanin formation in *Arabidopsis*. *Journal of Plant Physiology* **170**, 619–624.
- Peer WA, Blakeslee JJ, Yang H, Murphy AS.** 2011. Seven things we think we know about auxin transport. *Molecular Plant* **4**, 487–504.
- Saito K, Yonekura-Sakakibara K, Nakabayashia R, Higashi Y, Yamazaki M, Tohge T, Fernie AR.** 2013. The flavonoid biosynthetic pathway in *Arabidopsis*: Structural and genetic diversity. *Plant Physiology and Biochemistry* **72**, 21–34.
- Shirley BW, Kubasek WL, Storz G, Bruggemann E, Koornneef M, Ausubel FM, Goodman HM.** 1995. Analysis of *Arabidopsis* mutants deficient in flavonoid biosynthesis. *The Plant Journal* **8**, 659–671.
- Winkel-Shirley B.** 2006. The biosynthesis of flavonoids. In: Grotewold E, ed. *The science of flavonoids*. Springer Science & Business Media, 71–95.
- Yin R, Han K, Heller W, Albert A, Dobrev PI, Zažímalová E, Schäffner AR.** 2014. Kaempferol 3-*O*-rhamnoside-7-*O*-rhamnoside is an endogenous flavonol inhibitor of polar auxin transport in *Arabidopsis* shoots. *New Phytologist* **201**, 466–475.
- Yonekura-Sakakibara K, Fukushima A, Nakabayashi R, et al.** 2012. Two glucosyltransferases involved in anthocyanin modification delineated by transcriptome independent component analysis in *Arabidopsis thaliana*. *The Plant Journal* **69**, 154–167.
- Yonekura-Sakakibara K, Tohge T, Matsuda F, Nakabayashi R, Takayama H, Niida R, Watanabe-Takahashi A, Inoue E, Saito K.** 2008. Comprehensive flavonol profiling and transcriptome coexpression analysis leading to decoding gene-metabolite correlations in *Arabidopsis*. *The Plant Cell* **20**, 2160–2176.