

Ranibizumab for the treatment of wet AMD: a summary of real-world studies

V Chong

Abstract

Data from real-world studies of ranibizumab in neovascular (wet) age-related macular degeneration suggest that outcomes in clinical practice fail to match those seen in clinical trials. These real-world studies follow treatment regimens that differ from the fixed dosing used in the pivotal clinical trial programme. To better understand the effectiveness of ranibizumab in clinical practice, we conducted a comprehensive evaluation of 12-month outcomes reported in peer-reviewed 'real-world' publications. Key measures included in our analysis were mean change in visual acuity (VA) and the proportion of patients gaining ≥ 15 letters or losing ≤ 15 letters. Twenty studies were eligible for inclusion in our study, with 18 358 eyes having sufficient data for analysis of 12-month outcomes. Mean baseline VA ranged from 48.8 to 61.6 Early Treatment Diabetic Retinopathy Study letters. Mean change in VA was between -2.0 and $+5.5$ letters, with a grand mean of $+2.9 \pm 3.2$, and a weighted mean (adjusted for the number of eyes in the study) of $+1.95$. Eleven studies reported that 19 ± 7.5 (mean value) of patients gained ≥ 15 letters, while in 12 studies the mean percentage of patient losing ≤ 15 letters was $89 \pm 6.5\%$. Our comprehensive analysis of real-world ranibizumab study data confirm that patient outcomes are considerably poorer than those reported in randomised control trials of both fixed and *pro re nata* regimens.

Eye (2016) 30, 270–286; doi:10.1038/eye.2015.217; published online 4 December 2015

Introduction

Wet age-related macular degeneration (AMD) is a chronic, progressive disease of the central retina and a major cause of irreversible vision loss worldwide.^{1,2} Central to the pathogenesis of the disease is the overexpression of vascular endothelial growth factor (VEGF), which

stimulates choroidal neovascularization and causes blood and fluid to leak into the macula. The last decade has seen the introduction of intravitreal anti-VEGF agents, which have revolutionized the treatment of wet AMD, offering patients previously unachievable improvements in vision.

Ranibizumab (Lucentis), a humanised monoclonal antibody fragment, was the first anti-VEGF agent shown to improve visual acuity (VA) in patients with wet AMD.^{3–5} Regulatory approval for the use of ranibizumab was granted on the basis of the MARINA^{4,6,7} and ANCHOR^{3,8} study data, which showed mean gains of 7–11 letters over 12 months with monthly dosing. Up to 40% of patients gained more than 15 letters during this time and few lost vision. Moreover, improvements were largely maintained at the 24-month follow-up.

The early clinical trial data clearly demonstrated the benefits that ranibizumab could offer; however, the requirement for monthly intravitreal injections places a high burden on patients and healthcare systems. Subsequent studies, therefore, investigated whether equivalent VA gains could be achieved with less frequent injections. In the PIER study, patients received three monthly loading doses followed by quarterly injections of 0.5 mg ranibizumab.⁹ However, there was a mean loss of 0.2 letters by month 12. Other studies used *pro re nata* (PRN) regimens, where the decision to administer the drug is dependent on the disease status assessed by the physician at regular monitoring visits (eg, vision worsening or increase in macula thickness). Outcomes of these PRN regimens are variable with mean gains of 2.3–9.3 letters.^{10–16} For example, in the SAILOR study, which utilised a PRN regimen based around quarterly monitoring visits following three initial monthly doses, there was a mean gain of only 2.3 letters.¹⁰ By contrast, the results from the PrONTo^{13,14} study (PRN regimen with

Oxford Eye Hospital, Oxford University Hospitals, Oxford, UK

Correspondence: V Chong, Oxford Eye Hospital, Oxford University Hospitals, Headley Way, Headington, Oxford OX3 9DU, England, UK
Tel: +44 (0)1865 234736;
Fax: +44 (0)1865 234515.
E-mail: victor@eretina.org

Received: 12 June 2015
Accepted in revised form: 21 September 2015
Published online: 4 December 2015

monthly monitoring visits following three initial monthly doses) were more promising with VA gains approaching MARINA and ANCHOR results (9.3 letters over 12 months).^{3,4,6–8} It seems that only regimens with frequent monitoring and strict retreatment criteria can achieve the visual outcomes anywhere near those seen with fixed monthly dosing.

In the meantime, ranibizumab became available for use in routine clinical practice and its effectiveness in real life started to be evaluated. The last few years saw several studies report that outcomes achieved with ranibizumab in clinical practice failed to match the efficacy observed in the early ranibizumab clinical trial programme upon which its license was granted. These include WAVE,¹⁷ HELIOS,¹⁸ LUMIERE,¹⁹ AURA,²⁰ and the MEDISOFT database.²¹ Interestingly, these studies showed VA gains of 3.8–6.7 letters during the most intensive treatment period (loading period), but unlike MARINA and ANCHOR, these initial visual outcomes were not maintained over time. In fact, most of these studies saw a change in mean VA at 12 months of –1 to 3.2 letters.^{17–21} These poor visual outcomes may stem from an inability to adhere to strict a PRN regimen (which is required for PRN to be effective) in routine clinical practice; this hypothesis is supported by the low injection frequency reported by these studies (a mean number of 4.3–5.1 injections over 12 months).^{17,19–21}

To better understand the overall picture concerning the real-world effectiveness of ranibizumab, a review of the literature was conducted to identify studies of ranibizumab in clinical practice. This review provides both a summary of the design and methodological quality of the studies and a basic analysis of VA outcomes from the studies.

Materials and methods

Search criteria

We conducted a systematic search of the PubMed database for English language peer-reviewed papers published before 1 December 2014. The search protocol and structure of the review was based on that used by the Cochrane Collaboration. Search terms were as follows: (1) (long-term OR real-life OR longitudinal OR cohort OR clinical experience OR open-label OR real-world OR database OR non-interventional OR non-interventional OR observational) NOT (randomised[ti] OR randomized [ti]); and (2) (wet AMD OR AMD OR exudative AMD OR neovascular AMD); and (3) (ranibizumab[ti] OR Lucentis[ti]).

Further studies to be considered for inclusion were identified by reviewing the references lists from the studies identified during the PubMed search and by assessing relevant papers already known to the authors.

Inclusion and exclusion criteria

Studies considered for inclusion were screened for eligibility in three sequential stages, by the review of: (1) the title; (2) the abstract, and (3) the full manuscript (Figure 1). At each stage, the rationale for retention or rejection of the study was recorded.

Inclusion criteria were as follows: non-randomised controlled studies where treatment was given according to the licensed posology (monthly injections until there are no signs of disease activity and/or maximum VA is achieved, followed by PRN), and where baseline VA and change in VA at 12 months were reported. There were no restrictions on geographic region, clinical characteristics, baseline VA or previous treatment.

Exclusion criteria were as follows: studies that were primarily focused on specific sub-types of wet AMD (eg, polypoidal choroidal vasculopathy or retinal angioma-tous proliferation), and studies specifically evaluating switch from other agents (bevacizumab and aflibercept). Follow-up and extensions to Phase III studies were not included as these were not considered real-world studies.

Data analysis

Key outcomes were mean change in VA from baseline and the proportion of patients gaining ≥ 15 letters or losing ≤ 15 letters at 12 months. Simple descriptive analyses of these data, including grand means (mean of the means of the several studies) and weighted means (mean weighted against number of eyes per study), were used to pool VA outcomes across the total population. Additional sub-analyses were used to compare visual outcomes according to patient previous treatment history (naive *vs* non-naive patients), type of study (prospective *vs* retrospective), and approach to missing data (last observation carried forward). For the purpose of our analysis conversion to Early Treatment Diabetic Retinopathy Study letters (ETDRS) letters was carried out in studies providing other measures of VA (eg, logMAR).

Results

Study selection

Figure 1 shows how the studies were selected for inclusion in our analysis. The PubMed search identified a total of 197 citations. Upon screening using our pre-defined inclusion criteria, 143 were discarded after review of the publication titles, 27 were discarded after the review of publication abstracts, and a further 8 were discarded after the review of full text. No additional studies were identified through the review of reference lists within the publications we identified; however, we were aware of an important study (AURA)²⁰ that was not




Figure 1 Study selection flow chart.

identified by the pre-defined search terms but was relevant to this review. Therefore, 20 studies in total were included in our evaluation.

Study characteristics

The characteristics of the 20 real-world studies of ranibizumab included in this review are summarised in Table 1. Most studies were conducted specifically to evaluate the effectiveness of intravitreal ranibizumab for the treatment of wet AMD in a real-world setting; however, the design and endpoints employed to achieve this aim were variable (Table 1 and Supplementary Table 1). Retrospective studies were the most common, with only six having a prospective design. The strengths and weaknesses of the individual studies are summarised in Table 2.

The size of the studies varied widely, with the number of eyes ranging from 54²² to 12 951.²¹ In total, 23 261 eyes were included, of which 18 358 had 12-month follow-up data. Approximately half of the studies considered only one eye per patient (generally the first treated eye).

Mean VA at baseline ranged from 48.8 to 61.6 letters, excluding the studies by Shona *et al*,³² Pushpoth *et al*,²⁶ and Nomura *et al*,²⁸ who reported VA outcomes for groups of patients stratified according to different subcategories (eg, baseline VA) but did not report a study mean. The grand mean (mean of the means of the several subsamples) was 54.6 ± 7.3, and when weighted according to the number of eyes in the study was 54.1.

Two of the 20 studies did not provide any information regarding previous treatment history.^{20,23} Eleven studies enrolled only treatment-naïve patients (Table 1), while

Table 1 Study characteristics: aims, design, and treatment protocols in real-world studies of ranibizumab for the treatment of wet AMD

Study	No. of patients	Duration	Country	Aim	Design	Treatment	Treatment-naïve?
Chavan <i>et al</i> ³³	123 eyes in 120 patients	36 months	UK	To describe bilateral visual outcomes with ranibizumab, and the effects of incomplete follow-up	Retrospective: data collected from consecutive patients commencing treatment between two defined time points	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Cohen <i>et al</i> (LUMIERE) ¹⁹	551 eyes in 551 patients (first eye treated)	12 months	France	To examine compliance with recommended treatment protocols in a real-world setting, and effects on treatment outcomes	Retrospective: ophthalmologists asked to provide historical data on their first 40–60 patients who had been treated with ranibizumab for ≥ 12 months	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Finger <i>et al</i> (WAVE) ¹⁷	3470 patients	12 months	Germany	To evaluate effectiveness, tolerability and safety of repeated injections of ranibizumab in a real-world setting	Prospective: all patients who were recommended for treatment with ranibizumab between two defined time points were included	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	75.1% (2605) patients were treatment-naïve; 11.7% (405) had received previous intravitreal injections
Frennesson and Nilsson ²⁵	312 eyes in 268 patients (44 patients bilaterally treated)	36 months	Sweden	To examine the effect of LOCF on visual outcomes in a 3-year study in a real-world setting	Retrospective: data collected from records of all patients who were followed up for 36 months	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	No (27 eyes had received previous PDT)
Gabai <i>et al</i> ³⁰	100 eyes in 92 patients	12 months	Italy	To evaluate the safety and efficacy of ranibizumab for the treatment of wet AMD in a real-world setting	Retrospective: examined records of all patients who began treatment with ranibizumab for newly diagnosed wet AMD, with ≥ 12 months' follow-up	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Hjelmqvist <i>et al</i> (Swedish Lucentis Quality Registry) ²³	471 patients (272 retrospectively and 199 prospectively)	12 months	Sweden	To evaluate effectiveness of ranibizumab in a real-world setting	Prospective and multicentre	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Not specified
Holz <i>et al</i> (AURA) ²⁰	2227 patients in the effectiveness analysis set (1695 patients in the first-year completers' set and	24 months	Europe, Canada and Venezuela	To assess the management of patients with wet AMD receiving anti-VEGF treatment in clinical practice	Retrospective, non-interventional and observational (consecutive screening of patients); multicentre	Treated with ranibizumab as prescribed by physician; 441 patients of the effectiveness set did not have a	Not specified

Table 1. (Continued)

Study	No. of patients	Duration	Country	Aim	Design	Treatment	Treatment-naïve?
	1184 in the second-year completers' set			between 2009 and 2011		ranibizumab loading dose and 1786 patients of the effectiveness set did have a ranibizumab loading dose	
Kumar <i>et al</i> ³⁴	81 eyes in 81 patients (data from first eye treated in 3 patients with bilateral disease)	12 months	UK	To examine effectiveness of ranibizumab in a real-world setting	Prospective: consecutive recruitment of patients commencing ranibizumab	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Matsumiya <i>et al</i> 2013 ²²	54 patients (24 with tAMD and 30 with PCV)	12 months	Japan	To compare the outcomes of intravitreal ranibizumab between two different phenotypes of wet AMD	Retrospective, interventional cohort study of consecutive case series	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	34 (62.9%) of patients were treatment-naïve
Muether <i>et al</i> ³⁵	102 patients (89 followed up for 12 months)	12 months	Germany	To examine the effects of latency between diagnosis of recurrence and retreatment on outcome	Prospective: consecutive enrolment of patients diagnosed between two defined time points	Loading dose of three injections (each 0.5 mg), followed by retreatment on PRN basis (German healthcare funding system introduced a delay of 23.5 ± 10.4 days between indication to treat and treatment)	Yes
Nomura <i>et al</i> ²⁸	123 patients (108 VMA-negative; 15 VMA-positive)	12 months	Japan	To investigate the effects of VMA on intravitreal ranibizumab treatment in patients with wet AMD	Retrospective comparative study of consecutive patients	Loading dose of three injections (each 0.5 mg), followed by retreatment at the discretion of the attending physician	Yes
Pagliarini <i>et al</i> (EPICOHORT) ²⁷	755 patients (133 patients received bilateral treatment)	24 months	Europe	To assess safety profile of ranibizumab in Europe-wide study in real-world setting	Prospective: enrolment in 54 European clinical centres; Phase IV observational study	For newly diagnosed patients, loading dose of three injections (each 0.5 mg), followed by retreatment on PRN basis; for patients with previous treatment, continue ranibizumab PRN	270 (35.8%) patients reported prior ocular treatment – ranibizumab most common (251; 33.2%)

Table 1. (Continued)

Study	No. of patients	Duration	Country	Aim	Design	Treatment	Treatment-naïve?
Piermarocchi <i>et al</i> ³⁶	94 eyes in 94 patients	12 months	Italy	To investigate whether genetic and non-genetic risk factors influence 12-month response to ranibizumab treatment for wet AMD	Prospective	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Pushpothi <i>et al</i> ²⁶	1086 eyes in 1017 patients	48 months	UK	To evaluate effectiveness of ranibizumab in a real-world setting	Retrospective: data collected from all patients who began treatment between two defined time points, and who completed ≥24 months' follow-up Prospective, observational, multicentre study	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	181 eyes had received previous treatment
Rakic <i>et al</i> (HELIOS) ¹⁸	309 eyes in 267 patients	24 months	Belgium	To evaluate effectiveness of ranibizumab in a real-world setting	Prospective, observational, multicentre study	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	74.9% treatment-naïve
Ross <i>et al</i> ²⁷	406 eyes in 406 patients	24 months	UK	To define which VA measurements are the best indicators of high-quality care	Retrospective analysis of data collected from treatment-naïve patients with ≥12 months' follow-up	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Shona <i>et al</i> ³²	87 eyes in 87 patients	12 months	UK	To evaluate effectiveness of ranibizumab in a real-world setting for patients with differing baseline VA	Retrospective: chart review of treatment-naïve patients who initiated ranibizumab; patients divided into three subgroups according to baseline VA: 24–34 letters 35–54 letters ≥55 letters	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Tufail <i>et al</i> ²¹	12 951 eyes in 11 135 patients	36 months	UK	To evaluate effectiveness of ranibizumab in a real-world setting and to benchmark standards of care	Retrospective: up to 5 years of routinely collected data were extracted remotely from 14 UK centres using an electronic medical records system; all patients receiving	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes (although prior use of bevacizumab was an exclusion criterion)

Table 1. (Continued)

Study	No. of patients	Duration	Country	Aim	Design	Treatment	Treatment-naïve?
Williams and Blyth ²⁹	615 eyes, including 88 eyes with baseline logMAR VA <0.30	12 months	UK	To assess the effect of baseline VA on outcome, including those with baseline VA <0.30 logMAR	ranibizumab for wet AMD were included Consecutive recruitment of treatment-naïve patients; only those with ≥12 months' follow-up included	Loading dose of three injections (each 0.5 mg), followed by retreatment on a PRN basis	Yes
Zhu <i>et al.</i> ²⁴	886 patients (208 eyes of 208 patients completed the study)	60 months	Australia	To assess the visual and anatomical outcomes and safety profile of intravitreal ranibizumab in treating wet AMD	Retrospective; consecutive patients treated	Loading dose of three injections (each 0.5 mg), followed by retreatment at the physician's discretion	71/208 patients were treatment-naïve; 137/208 received one or more wet AMD treatments

Abbreviations: AMD, age-related macular degeneration; LOCF, last observation carried forward; logMAR, logarithm of the minimal angle of resolution; PCV, polypoidal choroidal vasculopathy; PDT, photodynamic therapy; PRN, *pro re nata*; tAMD, typical age-related macular degeneration; VA, visual acuity; VEGF, vascular endothelial growth factor; VMA, vitreomacular adhesion. ^aPart of LUMINOUS (Holz and colleagues, 2013).

seven could have included patients who received previous treatment for wet AMD. Previous treatments included photo dynamic therapy (PDT) ($n = 3$)^{22,24,25} and anti-VEGF intravitreal injections ($n = 4$).^{17,18,26,27}

Most studies ($n = 18$) were conducted in a single country, of which 15 were European, 1 was Australian,²⁴ and 2 were Japanese.^{22,28} The country with the largest number of studies ($n = 7$) was the UK. The remaining two studies included in the analysis were multinational.

Change in VA

The grand mean (\pm SD) VA change from baseline to 12 months reported in the 20 studies was 2.9 ± 3.2 letters; however, when weighted according to the number of eyes evaluated, the change was 1.95 letters (Figure 2). Fourteen studies reported an improvement in VA, with a maximum mean gain in VA recorded in a single study of 5.5 letters.²⁹ Two studies reported a decline in visual outcomes with the largest decline in VA of 2.0 letters.³⁰ In the remaining four studies the VA at study end remained similar to baseline values (-0.8 to 0.97 letters).

VA gains in the 12 studies of anti-VEGF-naïve patients were slightly greater than that for the full study population, with a grand mean of 3.5 ± 3.9 letters and a weighted mean of 3.5 letters.

Retrospective studies found greater mean gains than prospective studies (grand mean of 3.5 ± 3.5 vs 1.3 ± 1.5 letters; weighted mean of 2.5 letters vs 0 letters). Studies that used a last observation carried forward (LOCF) approach to their data analysis ($n = 16$) reported lower VA gains on average than the full study population (grand mean gain at 12 months of 3.1 ± 3.5 letters; weighted mean 1.9 letters).

Patients gaining 15 or more letters at 12 months Eleven studies (including 3869 eyes with 12-month data) reported the percentage of patients gaining ≥ 15 letters. At 12 months, the means was $19 \pm 7.5\%$. Weighting for the number of eyes available for analysis had little effect (Figure 3a).

Similar results were observed for the studies of anti-VEGF-naïve patients. On average $19 \pm 8.4\%$ of these patients gained ≥ 15 letters (13–24%), with a weighted mean of 20%.

No noteworthy changes were seen in studies reporting LOCF, when compared with the overall analysis.

In retrospective studies, a greater percentage of patients gained ≥ 15 letters than in prospective studies (grand mean of $19 \pm 8.3\%$ and $16 \pm 1.4\%$). Weighting for the number of eyes had little effect with mean values changing to 20% and 15.5%, respectively.

Patients losing 15 letters or less at 12 months Twelve studies (including 12 467 eyes with 12-month data) reported the percentage of patients losing ≤ 15 letters. The

Table 2 Strengths, weaknesses and approaches to missing data in real-world studies of ranibizumab in wet AMD

Study	Strengths	Weaknesses	Approach regarding missing values	Attrition
Chavan <i>et al</i> ³³	Consecutive enrolment to reduce selection bias Separate analysis of data with/without LOCF for patients who discontinued	Relatively small sample size	Separate analysis of data with/without LOCF for patients who discontinued	30% over 3 years
Cohen <i>et al</i> (LUMIERE) ¹⁹	Consecutive enrolment to reduce selection bias	Some ophthalmologists used Snellen system; VA values had to be converted to ETDRS-equivalent values Poor compliance made it difficult to evaluate changes in VA (however, compliance was one of the study outcomes)	Only included patients with 12-month follow-up	NA
Finger <i>et al</i> (WAVE) ¹⁷	Relatively large sample size Sample should be broadly representative of those treated with ranibizumab (no additional selection criteria applied at enrolment)	Only about one-third of patients received further injections in the maintenance phase LOCF not applied	Excluded	25.5%
Frennesson and Nilsson ²⁵	Compared effects on VA outcomes of applying LOCF <i>vs</i> disregarding data from drop-outs	Substantial number of patients received bilateral treatment (not independent samples)	Separate analysis of data with/without LOCF for patients who discontinued	20.8%
Gabai <i>et al</i> ³⁰	—	—	Only included patients with 12-month follow-up Excluded	NA
Hjelmqvist <i>et al</i> (Swedish Lucentis Quality Registry) ²³	Both retrospective and prospective components	In 100 of the retrospective patients, Snellen testing was used at baseline and the data were converted to ETDRS values		370/471 patients completed 12 months and formed the 'on-treatment' population: 21.4% attrition
Holz <i>et al</i> (AURA) ²⁰	Large sample size Real-life assessment of anti-VEGF use Monitoring and visual outcomes in consecutively enrolled patients across multiple centres and countries Retrospective design (prevents investigator bias) Prospective study with monthly clinic attendance	Different disease management between countries Clinical centres included in AURA might not represent patient management in the entire country Study limited by the observational and uncontrolled nature of the design	To account for missing data, mean change in VA was assessed using LOCF	Overall 2609 patients: Effectiveness analysis set: 2227 First-year completers' set: 1695 Second-year completers' set: 1184
Kumar <i>et al</i> ³⁴	Prospective study with monthly clinic attendance	Presence of coexisting ocular pathology or time to treatment from first diagnosis not taken into consideration	Excluded	80/81 patients received 3 loading injections 2 died (although data available for months 3 and 6, respectively)

Table 2. (Continued)

Study	Strengths	Weaknesses	Approach regarding missing values	Attrition
Matsumiya <i>et al</i> ²²	Cohort study of consecutive case series Differential diagnosis of typical AMD and PCV carried out after inclusion To exclude possible influence of previous PDT treatments, visual outcomes were evaluated in sub-population of patients that were treatment-naive Time course of changes in VA tracked	Relatively small sample Possibility of under-treatment	NA	Further 3 patients lost to follow-up So 75/81 completed = attrition: 7.5% NA
Muether <i>et al</i> ³⁵	Cohort study of consecutive case series	As B-mode ultrasonography was not performed, some eyes with vitreous completely attached to the retina might have been categorised into the VMA (-) group in the analysis	Relatively small sample size	Excluded NA
Nomura <i>et al</i> ²⁸	Cohort study of consecutive case series	The patients were from a single institution (results do not present a general overview of exudative AMD in Japanese patients)	NA	NA
Pagliarini <i>et al</i> (EPICOHORT) ²⁷	Mean BCVA	Heterogeneous patient cohort: some patients had previously been treated with ranibizumab	States in methods that analysis was performed with and without LOCF, but LOCF efficacy results are not reported	22.8% over 24 months
Piermarocchi <i>et al</i> ³⁶	Mean change in BCVA at study end analysed based on patient's genetic characteristics associated with development of AMD	Relatively limited sample size Need for a more prolonged follow-up Clinical risk factors associated with worse prognosis after treatment were not considered	NA	All patients completed the 12-month follow-up
Pushpoth <i>et al</i> ²⁶	Long duration	High attrition in a long-duration study in an elderly patient group makes later results difficult to interpret	Excluded	No. of patients remaining under follow-up: 12 months: 897/1017 24 months: 730/1017 36 months: 468/730 48 months: 110/217

Table 2. (Continued)

Study	Strengths	Weaknesses	Approach regarding missing values	Attrition
Rakic <i>et al</i> (HELIOS) ¹⁸	Examined effects of no. of injections in maintenance phase	Data collection began soon after approval of ranibizumab in Belgium; thus, results may not reflect current levels of physician expertise	Excluded	255 patients evaluable at baseline 24 months: 184 patients Attrition = 27.9%
Ross <i>et al</i> ³⁷	Examined effects of baseline VA on outcomes Analysed data with and without LOCF but reported only VA data not change in VA data for LOCF analyses	High attrition	Only included patients with 12-month follow-up Analysed data with and without LOCF but reported only VA data not change in VA data for LOCF analyses	Data extracted on 700 eyes in 629 patients; 247 eyes in 176 patients did not meet eligibility criteria = 453 eyes 47 eyes in 47 patients excluded because no data available at 12 months 198 patients completed 24 months Attrition = 1 – (198/453) = 56.3%
Shona <i>et al</i> ³²	Examined effects of baseline VA, and no. of injections, on outcomes Very large sample size	—	Only included patients with 12-month follow-up	NA
Tufail <i>et al</i> ²¹		Missing data excluded	Missing data excluded in main analysis. However, data was also analysed separately for eyes that completed 168 weeks, follow-up (<i>n</i> = 1138), 78 eyes that received ≥1 injection at time 0 (<i>n</i> = 12 951) and <i>n</i> = 1138 at Week 168 Change in mean VA was similar for the two groups Only patients with 12 months' follow-up included	12 months: 8598 eyes 24 months: 4990 eyes 36 months: 2470 eyes
Williams and Blyth ²⁹	Examined effectiveness of ranibizumab in patients with relatively high baseline VA Strict inclusion and exclusion criteria Standardised retreatment criteria Long duration Retrospective study of consecutive patients treated for AMD All patients treated by a single physician	Difficulties reporting cataract progression due to non-standardised lens grading at each visit Use of multiple OCT devices ICGA screening for PCV was not performed regularly at baseline (PCV patients might have been included in the study)	NA	All patients included in the study (208) had 5-year VA assessment
Zhu <i>et al</i> ²⁴				

Abbreviations: AMD, age-related macular degeneration; BCVA, best corrected visual acuity; ETDRS, Early Treatment Diabetic Retinopathy Study; ICGA, indocyanine green angiography; NA, not applicable; OCT, optical coherence tomography; PDT, photodynamic therapy; VA, visual acuity; VEGF, vascular endothelial growth factor.




Figure 2 Mean change from baseline in visual acuity and number of injections in real-world studies and pivotal randomised controlled studies after 12 months of ranibizumab treatment in patients with wet AMD. *Represents average data for the 20 studies included in the current review (see Table 3).

means \pm SD at 12 months was $89 \pm 6.5\%$ (74.4–97.4%) and was similar to the weighted mean (Figure 3b).

On average, $91 \pm 5.9\%$ of anti-VEGF-naive patients lost ≤ 15 letters. Again, weighting for the number of eyes had little effect (89%).

No noteworthy results were observed in studies reporting LOCF.

Similar results were observed between retrospective and prospective studies regarding the percentage (grand mean) of patients losing ≤ 15 letters ($90 \pm 5.2\%$ vs $92 \pm 7.4\%$). No noteworthy differences were found when these changes were weighted for the number of eyes (at 12 months).

Number of injections over 12 months

The mean number of injections ranged from 4.2 to 7.5 (Table 3), with a grand mean (\pm SD) of 5.5 ± 0.8 . Anti-VEGF-naive patients received a similar number of injections to the full study population (grand mean of 5.4 ± 0.7). Weighting for the number of eyes at 12 months had little effect on the outcomes of these analyses.

Similar number of injections were received by patients in the retrospective and prospective studies (grand mean number of injections of 5.6 ± 0.7 and 5.1 ± 1 , respectively). In this analysis, weighting according to the number of eyes assessed at 12 months gave different results, showing that patients treated in prospective studies

received one less injection on average than those evaluated by retrospective studies (4.5 vs 5.7 injections).

Other outcomes

Ten studies reported mean changes in central retinal thickness, six reported safety data and three reported vision-related quality of life. A detailed overview of these and other outcomes is provided for reference in Supplementary Table 1.

Discussion

This review was performed in order to determine whether, in patients with wet AMD, real-world outcomes achieved with ranibizumab match the positive outcomes reported in the pivotal ranibizumab trials,^{3,4,6–9} and subsequent studies of 0.5 mg PRN regimens.^{10–16}

The grand mean (+2.9 letters) and weighted mean (+1.95 letters) calculated from the changes in VA reported at 12 months in the trials included in this review are considerably lower than those reported in the pivotal studies of ranibizumab (fixed monthly dosing),^{3,4,6–8} with the exception of the PIER study^{9,31} (fixed quarterly dosing) that showed a mean loss of 0.2 letters. Specifically, the recently published AURA study conducted in multiple countries and considered a true and current depiction of the real-world use of ranibizumab for the treatment of AMD, clearly

Table 3 Effects of ranibizumab on key visual acuity outcomes in real-world studies of wet AMD

Study	No. of patients	Duration	Country	No. of ranibizumab injections (mean ± SD unless otherwise stated)	Baseline visual acuity (mean ± SD unless otherwise stated, ETDRS letters unless otherwise stated)	Mean change in visual acuity from baseline to end of study (ETDRS letters unless otherwise stated)	Patients gaining ≥15 letters/3 lines (%)	Patients losing <15 letters/3 lines (%)
Chavan <i>et al</i> 2014 ³³	123 eyes in 120 patients	36 months	UK	Full analysis set: Yr 1: 5.9 (1–11) (mean, range) Yr 2: 4.1 (0–10) Yr 3: 4.2 (0–11) Per protocol set: Yr 1: 6.4 (1–11) (mean, range) Yr 2: 4.7 (0–10) Yr 3: 4.2 (0–11)	Full analysis set: 52.89 ± 14.63 Per protocol set: 53.75 ± 13.59	Full analysis set: –1.68 ± 17.76 Per protocol set: +1.47 ± 15.60	16.8% 19.1%	78.2% 84.3%
Cohen <i>et al</i> (LUMIERE) ¹⁹	551 eyes in 551 patients	12 months	France	5.1	LOCF set: 50.33 ± 17.10 At 12 months: 52.24 ± 14.69 (all treated eyes)	LOCF set: –11.03 ± 20.29 At 12 months: –1.34 ± 15.34	LOCF set: 10% At 12 months: 13%	LOCF set: 60% At 12 months: 84.6%
Finger <i>et al</i> (WAVE) ¹⁷	3 470 patients	12 months	Germany	4.3 ± 0.05 (SEM)	48.8 ± 18.7 from Holz and colleagues, 2013 (logMAR of Snellen values)	–0.8 from Holz and colleagues, 2013 (logMAR of Snellen values)	—	—
Frennesson and Nilsson ²⁵	312 eyes in 268 patients (44 patients bilaterally treated)	36 months	Sweden	VA of drop-outs disregarded: Yr 1: 5.4 (CI 5.2–5.7) Yr 2: 2.5 (CI 2.2–2.8) Yr 3: 2.3 (CI 1.7–2.9) LOCF: Yr 1: 5.3 (CI 5.1–5.6) Yr 2: 2.1 (CI 1.8–2.4) Yr 3: 1.9 (CI 1.4–2.4)	58.4 (CI 56.9–59.9)	VA of drop-outs disregarded: +0.1 [58.5 (CI 54.2–62.8)] LOCF: –4.1 [54.3 (CI 49.6–59.0)] At 12 months: drop-outs disregarded, +1.8; LOCF, +1.0	VA of drop-outs disregarded: 12.6% Not available for LOCF	VA of drop-outs disregarded: 77.2% Not available for LOCF
Gabai <i>et al</i> ³⁰	100 eyes in 92 patients	12 months	Italy	4.8	61.6 ± 14.8	–2.0 ± 17.6	16%	79%
Hjelmqvist <i>et al</i> (Swedish Lucentis Quality Registry) ²³	471 patients (272 retrospectively and 199 prospectively)	12 months	Sweden	4.7 ± 1.6 (on-treatment population, ie, excluding those who discontinued)	58.3 ± 12.2 (on-treatment population)	+1.0 ± 13.6	14.7%	74.4%
Holz <i>et al</i> (AURA) ²⁰	2227 patients in the effectiveness analysis set (1695 patients in the first-year completers' set and 1184 in the second-year completers' set)	24 months	Europe, Canada and Venezuela	Effectiveness set: Yr 1: mean of 5.0 Yr 2: mean of 2.2 First-year completers: Yr 1: mean of 5.5 Yr 2: mean of 2.9 Second-year completers: Yr 1: mean of 5.6 Yr 2: mean of 3.5	Effectiveness set: 55.4 ± 18.4 (n = 2147) First-year completers: 56.9 ± 17.8 (n = 1642) Second-year completers: 57.2 ± 17.9 (n = 1146)	Effectiveness set: Yr 1: +2.4 (LOCF) Yr 2: +0.6 (LOCF) First-year completers: Yr 1: +2.7 Yr 2: +0.3 Second-year completers: Yr 1: +2.7 Yr 2: +0.3	—	—
Kumar <i>et al</i> 2011 ³⁴ Matsumiya <i>et al</i> ²²	81 eyes in 81 patients 54 eyes from 54 patients	12 months 12 months	UK Japan	5.6 ± 2.3	49.5 ± 13.4 (logMAR values) 0.60 ± 0.28 for total	+3.7 ± 10.8 Total: 1 month: –0.03 3 months: –0.11	17.1%	97.4%

Table 3. (Continued)

Study	No. of patients	Duration	Country	No. of ranibizumab injections (mean ± SD unless otherwise stated)	Baseline visual acuity (mean ± SD unless otherwise stated, ETDRS letters unless otherwise stated)	Mean change in visual acuity from baseline to end of study (ETDRS letters unless otherwise stated)	Patients gaining ≥15 letters/3 lines (%)	Patients losing <15 letters/3 lines (%)
Muether <i>et al</i> ³⁵	102 patients [89 followed up for 12 months]	12 months	Germany	3.9 ± 0.9 in tAMD 4.2 ± 0.9 in PCV group	0.47 ± 0.26 for PCV 0.53 ± 0.28 for tAMD	6 months: -0.10 12 months: -0.09 PCV: 1 month: 0 3 months: -0.06 6 months: -0.06 12 months: -0.04 tAMD: 1 month: -0.06 3 months: -0.17 6 months: -0.15 12 months: -0.16	—	—
Nomura <i>et al</i> ²⁸	123 patients: 108 VMA (-); 15 VMA (+)	12 months	Japan	VMA (-): 5.2 VMA (+): 5.1	VMA (-): 0.41 ± 0.35 VMA (+): 0.42 ± 0.39	VMA (-): 6 letters VMA (+): 1.5 letters	—	—
Pagliari <i>et al</i> (EPICOHORT) ²⁷	755 patients (133 patients received bilateral treatment)	24 months	Europe	Yr 1: 4.4 Yr 2: 1.8	52.8 ± 20.3	-1.3 ± 0.72 (SEM) At 12 months: +1.5 ± 0.61 (SEM)	—	—
Piermarocchi <i>et al</i> ³⁶ Pushpoth <i>et al</i> ²⁶	94 eyes of 94 patients 1086 eyes in 1017 patients	12 months 48 months	Italy UK	4.2 ± 1.2 Pre-treatment: Yr 1: 6.2 ± 2.6 Yr 2: 8.3 ± 3.0 Yr 3: 9.7 ± 4.7 Yr 4: 12.9 ± 7.2 (cumulative totals) Treatment-naive: Yr 1: 5.2 ± 2.7 Yr 2: 8.3 ± 3.7 Yr 3: 10.8 ± 5.8 Yr 4: 12.8 ± 7.8 (cumulative totals)	Month 48: 50.43 ± 15.58 (pre-treatment; n = 180) 54.09 ± 15.25 (treatment-naive; n = 906)	0.97 ± 9.1 Month 48: 53.82 ± 15.58 (pre-treatment; n = 40) [- +3.39] 58.73 ± 17.19 (treatment-naive; n = 75) [+4.64] At 12 months: pre-treatment, +2.67; treatment-naive, +3.82	Pre-treatment: Month 48: 92.5% (n = 37) Treatment-naive: Month 48: 81.3% (n = 61)	Pre-treatment: Month 48: 0.1% (n = 4) Treatment-naive: Month 48: 21.3% (n = 16) At 12 months: pre-treatment, 20.5%; treatment-naive, 20.1%
Rakic <i>et al</i> (HELIOS) ¹⁸	309 eyes in 267 patients	24 months	Belgium	Yr 1: 5.0 ± 2.1 Yr 2: 3.7 ± 2.1	56.3 ± 14.3	-2.4 ± 17.4 At 12 months: +1.6 ± 15.6	14.1% At 12 months: 15.1%	81.5% At 12 months: 86.9%
Ross <i>et al</i> ³⁷	406 eyes in 406 patients	24 months	UK	Yr 1: 5.9 (3-13) Yr 2: 3.6 (0-10)	54.4 ± 14.2	+1.6 ± 17.6 At 12 months: +4.1 ± 14.2	20.7% At 12 months: 20.9%	86.7% At 12 months: 90.1%
Shona <i>et al</i> ³²	87 eyes in 87 patients	12 months	UK	Baseline VA: Good: 5.7 Medium: 6.1 Poor: 5.3	Good: 61.67 ± 6.21 Medium: 43.79 ± 4.67 Poor: 28.93 ± 4.7	Good: +2.85 Medium: +7.10 Poor: +14.00	Good: 10% Medium: 14% Poor: 41%	Good: 95% Medium: 92% Poor: 100%
Tufail <i>et al</i> ²¹	12 951 eyes in 11 135 patients	36 months	UK	Yr 1: 5.7 (1-13) Yr 2: 3.7 (0-13) Yr 3: 3.7 (0-12)	For eyes with ≥ 36 months' follow-up: 55 letters	For eyes with ≥ 36 months' follow-up: -2 letters at Week 156 At 12 months: +2.0	—	82% At 12 months: 90%

Table 3. (Continued)

Study	No. of patients	Duration	Country	No. of ranibizumab injections (mean ± SD unless otherwise stated)	Baseline visual acuity (mean ± SD unless otherwise stated, ETDRS letters unless otherwise stated)	Mean change in visual acuity from baseline to end of study (ETDRS letters unless otherwise stated)	Patients gaining ≥15 letters/3 lines (%)	Patients losing <15 letters/3 lines (%)
Williams and Blyth ²⁹	615 eyes, including 88 eyes with baseline VA <0.30	12 months	UK	All: 5.6 Baseline VA: <0.30: 5.4 0.30–0.59: 5.6 0.60–0.99: 5.8 1.00–1.20: 5.2	All: 0.60 (logMAR) Baseline VA (logMAR): <0.30 (mean 0.19) 0.30–0.59 (mean 0.42) 0.60–0.99 (mean 0.73) 1.00–1.20 (mean 1.06) Baseline VA (ETDRS) <0.30 (mean 75.48) 0.30–0.59 (mean 63.84) 0.60–0.99 (mean 48.58) 1.00–1.20 (mean 31.97)	All: 0.49 (logMAR) Baseline VA (logMAR): <0.30: 0.20 0.30–0.59: 0.37 0.60–0.99: 0.60 1.00–1.20: 0.76 Baseline VA (ETDRS) <0.30: –0.69 0.30–0.59: +2.61 0.60–0.99: +6.32 1.00–1.20: +15.05	All: 24% Baseline VA: <0.30: 1% 0.30–0.59: 16% 0.60–0.99: 33% 1.00–1.20: 46%	All: 92% Baseline VA: <0.30: 93% 0.30–0.59: 88% 0.60–0.99: 92% 1.00–1.20: 100%
Zhu <i>et al</i> ²⁴	886 patient (208 eyes of 208 patients completed the study)	60 months	Australia	Yr 1: 7.5±2.6 Yr 2: 5.8±3.9 Yr 3: 6.4±4.5 Yr 4: 5.6±4.5 Yr 5: 5.8±4.7	53.6 ± 19.3 (208 patients)	Month 6: +3.2 ± 10.4 Yr 1: +1.9 Yr 5: -2.4	—	—

Abbreviations: AMD, age-related macular degeneration; CI, confidence interval; ETDRS, Early Treatment Diabetic Retinopathy Study; LOCF, last observation carried forward; logMAR, logarithm of the minimal angle of resolution; PCV, polypoidal choroidal vasculopathy; tAMD, typical age-related macular degeneration; VA, visual acuity; VEGF, vascular endothelial growth factor; Yr, year.

demonstrated that the VA outcomes are far worse than the ones reported in the landmark trials (MARINA and ANCHOR).^{3,4,6–8} Rather, they are more in line with those observed in PIER^{9,31} and SAILOR,¹⁰ which used less intensive treatment schedules (fixed quarterly dosing and PRN dosing based on a quarterly monitoring schedule, respectively), and compare unfavourably with MARINA^{4,6,7} and ANCHOR.^{3,8}

What are the factors that underlie the reduced effectiveness of ranibizumab in clinical practice? A key consideration may be the reduced number of injections that patients receive in clinical practice. Across the 20 real-world studies included in the present review, the mean number of injections ranged from 4.2–7.5; the grand mean at 12 months was 5.5 ± 0.8 injections and the weighted mean was 5.4 injections, notably less than the 12 injections received in ANCHOR/MARINA^{3,4,6–8} (fixed monthly dosing). A reduced number of injections in the real-world setting is not unexpected as adherence to treatment regimens tends to be harder to maintain owing to the challenges of implementing intensive treatment regimens outside of the clinical trial setting. However, the reduced efficacy of ranibizumab in real-world practice observed in our study cannot be explained simply by the reduced number of injections, as the pooled mean values were: (1) less than the 0.5 mg PRN arms of CATT^{11,12} and HARBOUR (6.9 and 7.7 injections, respectively),^{15,16} which saw greater increases in VA; (2) similar to PrONTO (5.6 injections),^{13,14} which demonstrated a greater increase in VA; and (3) greater than in SAILOR (4.6 injections),¹⁰ which showed a similar increase in VA as observed in the present analysis (see Figure 2 for change in VA from baseline).

Besides the reduced number of injections, the schedule on which they are given as well as the monitoring frequency and retreatment criteria might also be a reason for the limited efficacy of ranibizumab in the clinic. The studies in this and previous analyses typically employed PRN regimens as a means of appropriately managing treatment burden. By definition, a PRN regimen only treats patient with symptomatic disease; hence, it is possible that recurring fluid may cause progressive damage. Results from CATT^{11,12} and HARBOUR^{15,16} show that it is possible to achieve good visual outcomes with a PRN regimen when frequent regimented monitoring visits and strict retreatment criteria are in place. However, with less intensive monitoring, as seen in the SAILOR study (quarterly monitoring schedule)¹⁰ and in routine clinical practice as per the ranibizumab label, visual outcomes using PRN are poor.

Recently, the treat-and-extend regimen has emerged as a potential alternative method of delivering the best-possible visual outcomes, with the appropriate injection frequency, while minimising the treatment burden.




Figure 3 (a) Percentage of patients gaining ≥ 15 letters. *Represents average data for the 11 studies included in the current review which reported this outcome (Table 3). (b) Percentage of patients losing < 15 letters. *Represents average data for the 12 studies included in the current review, which reported this outcome (Table 3).

With treat-and-extend, the physician can identify the appropriate injection frequency for each patient, proactively injecting at each visit and deciding when to treat next rather than whether to treat at that time. Thus, treatment remains ahead of the disease. In the future, it will be interesting to evaluate whether this newer treatment approach can improve outcomes in the real-life situation.

Baseline VA is an additional determinant of outcome in terms of VA gains (ceiling effect), with patients with poorer vision tending to show more gain. However, the mean baseline acuity of the studies included in the present analysis is comparable to that reported in the clinical trials (respective ranges 48.8–61.6 vs 47.1–61.5) with the grand mean (54.6) and weighted mean (54.1)

lying in the centre of these values. Therefore, it is difficult to ascribe the poorer outcomes observed in the present analysis to baseline acuity alone.

As with any review of this type, it is important to consider the limitations of the current analysis. By their nature, real-world studies (non-randomised and not controlled studies) are not as rigorously designed as clinical trials and therefore require careful interpretation. In particular, 13 out of the 20 included studies, including the largest study by Tufail *et al.*²¹ were retrospective. Data may be incomplete, with patients being lost to follow-up or not having outcomes recorded at appropriate time points. Some studies employed an LOCF method to account for these missing data; although a standard and valid technique, LOCF analysis may skew study results particularly in conditions that change over time, for example, wet AMD. Moreover, differences in VA measurements in real-life studies *vs* clinical trials (Snellen *vs* ETDRS) could also be considered as a limitation.

In addition, the inclusion criteria vary across studies included in the present analysis, with many including patients irrespective of their treatment history or baseline VA. Such variability can affect outcomes. For example, patients with good baseline acuity demonstrated a smaller improvement in response to ranibizumab than patients with poor baseline acuity,^{29,32} while treatment-naïve patients showed a better response than previously treated individuals.²⁶ Differences such as these may obscure the effects of treatment if the numbers of patients from a particular group are unrepresentative of the population as a whole. However, with nearly 20 000 patients analysed, the real-life data reviewed here could be considered as representative of current practice.

From an analytical perspective, the validity of data pooling across studies with different designs is inherently limited. The strengths and weaknesses of each of the 20 real-world studies included in the review, along with a description of the approaches adopted to missing values in the studies, are summarised in Table 2.

Finally, only peer-reviewed studies indexed on PubMed were included. It is possible that additional relevant studies not listed on PubMed are not captured here.

In conclusion, the current review strongly suggests that, in patients with wet AMD, VA outcomes achieved with ranibizumab in clinical practice do not match those reported in pivotal studies. Although factors such as reduced injection frequency and patient heterogeneity are likely to have a part in this phenomenon, further research is required to determine the reasons for this discrepancy. Nonetheless, clinicians should be aware of this discrepancy when consenting patients for receiving ranibizumab for wet AMD.

Conflict of interest

VC is a consultant for Allergan, Bayer, Novartis, Quantel Medical, Pfenex.

Acknowledgements

I take full responsibility for the scope, direction, and the content of the manuscript, and has approved the submitted manuscript. Medical writing assistance was provided by Ana Tadeu of Porterhouse Medical Ltd., and was funded by Bayer HealthCare Pharmaceuticals.

References

- 1 Stewart MW. Clinical and differential utility of VEGF inhibitors in wet age-related macular degeneration: focus on aflibercept. *Clin Ophthalmol* 2012; **6**: 1175–1186.
- 2 Kawasaki R, Yasuda M, Song SJ, Chen SJ, Jonas JB, Wang JJ *et al.* The prevalence of age-related macular degeneration in Asians: a systematic review and meta-analysis. *Ophthalmology* 2010; **117**(5): 921–927.
- 3 Brown DM, Kaiser PK, Michels M, Soubrane G, Heier JS, Kim RY *et al.* Ranibizumab *vs* verteporfin for neovascular age-related macular degeneration. *N Engl J Med* 2006; **355**(14): 1432–1444.
- 4 Rosenfeld PJ, Brown DM, Heier JS, Boyer DS, Kaiser PK, Chung CY *et al.* Ranibizumab for neovascular age-related macular degeneration. *N Engl J Med* 2006; **355**(14): 1419–1431.
- 5 Blick SK, Keating GM, Wagstaff AJ. Ranibizumab. *Drugs* 2007; **67**(8): 1199–1206 discussion 1207–1199.
- 6 Chang TS, Bressler NM, Fine JT, Dolan CM, Ward J, Klesert TR. Improved vision-related function after ranibizumab treatment of neovascular age-related macular degeneration: results of a randomized clinical trial. *Arch Ophthalmol* 2007; **125**(11): 1460–1469.
- 7 Kaiser PK, Blodi BA, Shapiro H, Acharya NR. Angiographic and optical coherence tomographic results of the MARINA study of ranibizumab in neovascular age-related macular degeneration. *Ophthalmology* 2007; **114**(10): 1868–1875.
- 8 Brown DM, Michels M, Kaiser PK, Heier JS, Sy JP, Ianchulev T. Ranibizumab *vs* verteporfin photodynamic therapy for neovascular age-related macular degeneration: Two-year results of the ANCHOR study. *Ophthalmology* 2009; **116**(1): 57–65 e55.
- 9 Regillo CD, Brown DM, Abraham P, Yue H, Ianchulev T, Schneider S *et al.* Randomized, double-masked, sham-controlled trial of ranibizumab for neovascular age-related macular degeneration: PIER Study year 1. *Am J Ophthalmol* 2008; **145**(2): 239–248.
- 10 Boyer DS, Heier JS, Brown DM, Francom SF, Ianchulev T, Rubio RG. A Phase IIIb study to evaluate the safety of ranibizumab in subjects with neovascular age-related macular degeneration. *Ophthalmology* 2009; **116**(9): 1731–1739.
- 11 Martin DF, Maguire MG, Ying GS, Grunwald JE, Fine SL, Jaffe GJ. Ranibizumab and bevacizumab for neovascular age-related macular degeneration. *N Engl J Med* 2011; **364**(20): 1897–1908.
- 12 Martin D, Maguire M, Fine SL, Ying GS, Jaffe GJ, Grunwald JE *et al.* Ranibizumab and bevacizumab for treatment of

- neovascular age-related macular degeneration: two-year results. *Ophthalmology* 2012; **119**: 1388–1398.
- 13 Lalwani GA, Rosenfeld PJ, Fung AE, Dubovy SR, Michels S, Feuer W *et al.* A variable-dosing regimen with intravitreal ranibizumab for neovascular age-related macular degeneration: year 2 of the PrONTO Study. *Am J Ophthalmol* 2009; **148**(1): 43–58 e41.
 - 14 Fung AE, Lalwani GA, Rosenfeld PJ, Dubovy SR, Michels S, Feuer WJ *et al.* An optical coherence tomography-guided, variable dosing regimen with intravitreal ranibizumab (Lucentis) for neovascular age-related macular degeneration. *Am J Ophthalmol* 2007; **143**(4): 566–583.
 - 15 Busbee BG, Ho AC, Brown DM, Heier JS, Suner IJ, Li Z *et al.* Twelve-month efficacy and safety of 0.5mg or 2.0mg ranibizumab in patients with subfoveal neovascular age-related macular degeneration. *Ophthalmology* 2013; **120**(5): 1046–1056.
 - 16 Ho AC, Busbee BG, Regillo CD, Wieland MR, Van Everen SA, Li Z *et al.* Twenty-four-month efficacy and safety of 0.5mg or 2.0mg ranibizumab in patients with subfoveal neovascular age-related macular degeneration. *Ophthalmology* 2014; **121**(11): 2181–2192.
 - 17 Finger RP, Wiedemann P, Blumhagen F, Pohl K, Holz FG. Treatment patterns, visual acuity and quality-of-life outcomes of the WAVE study—A noninterventional study of ranibizumab treatment for neovascular age-related macular degeneration in Germany. *Acta Ophthalmol* 2013; **91**(6): 540–546.
 - 18 Rakic JM, Leys A, Brie H, Denhaerynck K, Pacheco C, Vancayzeele S *et al.* Real-world variability in ranibizumab treatment and associated clinical, quality of life, and safety outcomes over 24 months in patients with neovascular age-related macular degeneration: the HELIOS study. *Clin Ophthalmol* 2013; **7**: 1849–1858.
 - 19 Cohen SY, Mimoun G, Oubraham H, Zourdani A, Malbrel C, Quere S *et al.* Changes in visual acuity in patients with wet age-related macular degeneration treated with intravitreal ranibizumab in daily clinical practice: the LUMIERE study. *Retina* 2013; **33**(3): 474–481.
 - 20 Holz FG, Tadayoni R, Beatty S, Berger A, Cereda MG, Cortez R *et al.* Multi-country real-life experience of anti-vascular endothelial growth factor therapy for wet age-related macular degeneration. *Br J Ophthalmol* 2014; **99**: 220–226.
 - 21 Tufail A XW, Johnston R, Akerele T, McKibbin M, Downey L, Natha S, Chakravarthy U, Bailey C, Khan R, Antcliff R, Armstrong S, Varma A, Kumar V, Tsaloumas M, Mandal K, Bunce C. The neovascular age-related macular degeneration database: multicenter study of 92 976 ranibizumab injections: report I: visual acuity. *Ophthalmology* 2014; **121**(5): 1092–1101.
 - 22 Matsumiya W, Honda S, Kusuhara S, Tsukahara Y, Negi A. Effectiveness of intravitreal ranibizumab in exudative age-related macular degeneration (AMD): comparison between typical neovascular AMD and polypoidal choroidal vasculopathy over a 1 year follow-up. *BMC ophthalmology* 2013; **13**: 10.
 - 23 Hjelmqvist L, Lindberg C, Kanulf P, Dahlgren H, Johansson I, Siewert A. One-year outcomes using ranibizumab for neovascular age-related macular degeneration: results of a prospective and retrospective observational multicentre study. *Journal of ophthalmology* 2011; **2011**: 405724.
 - 24 Zhu M, Chew JK, Broadhead GK, Luo K, Joachim N, Hong T *et al.* Intravitreal Ranibizumab for neovascular Age-related macular degeneration in clinical practice: five-year treatment outcomes. *Graefes Arch Clin Exp Ophthalmol* 2015; **253**(8): 1217–1225.
 - 25 Frennesson CI, Nilsson SE. A three-year follow-up of ranibizumab treatment of exudative AMD: impact on the outcome of carrying forward the last acuity observation in drop-outs. *Acta Ophthalmol* 2014; **92**(3): 216–220.
 - 26 Pushpoth S, Sykakis E, Merchant K, Browning AC, Gupta R, Talks SJ. Measuring the benefit of 4 years of intravitreal ranibizumab treatment for neovascular age-related macular degeneration. *Br J Ophthalmol* 2012; **96**(12): 1469–1473.
 - 27 Pagliarini S, Beatty S, Lipkova B, Perez-Salvador Garcia E, Reynders S, Gekkieva M *et al.* A 2-year, phase IV, multicentre, observational study of ranibizumab 0.5mg in patients with neovascular age-related macular degeneration in routine clinical practice: The EPICOHORT Study. *J Ophthalmol* 2014; **2014**: 857148.
 - 28 Nomura Y, Takahashi H, Tan X, Fujimura S, Obata R, Yanagi Y. Effects of vitreomacular adhesion on ranibizumab treatment in Japanese patients with age-related macular degeneration. *Jpn J Ophthalmol* 2014; **58**(5): 443–447.
 - 29 Williams TA, Blyth CP. Outcome of ranibizumab treatment in neovascular age related macula degeneration in eyes with baseline visual acuity better than 6/12. *Eye (Lond)* 2011; **25**(12): 1617–1621.
 - 30 Gabai A, Veritti D, Lanzetta P. One-year outcome of ranibizumab for neovascular age-related macular degeneration: a thorough analysis in a real-world clinical setting. *Eur J Ophthalmol* 2014; **24**(3): 396–401.
 - 31 Abraham P, Yue H, Wilson L. Randomized, double-masked, sham-controlled trial of ranibizumab for neovascular age-related macular degeneration: PIER study year 2. *Am J Ophthalmol* 2010; **150**(3): 315–324 e311.
 - 32 Shona O, Gupta B, Vemala R, Sivaprasad S. Visual acuity outcomes in ranibizumab-treated neovascular age-related macular degeneration; stratified by baseline vision. *Clin Experiment Ophthalmol* 2011; **39**(1): 5–8.
 - 33 Chavan R, Panneerselvam S, Adhana P, Narendran N, Yang Y. Bilateral visual outcomes and service utilization of patients treated for 3 years with ranibizumab for neovascular age-related macular degeneration. *Clin Ophthalmol* 2014; **8**: 717–723.
 - 34 Kumar A, Sahni JN, Stangos AN, Campa C, Harding SP. Effectiveness of ranibizumab for neovascular age-related macular degeneration using clinician-determined retreatment strategy. *Br J Ophthalmol* 2011; **95**(4): 530–533.
 - 35 Muether PS, Hoerster R, Hermann MM, Kirchhof B, Fauser S. Long-term effects of ranibizumab treatment delay in neovascular age-related macular degeneration. *Graefes Arch Clin Exp Ophthalmol* 2013; **251**(2): 453–458.
 - 36 Piermarocchi S, Miotto S, Colavito D, Leon A, Segato T. Combined effects of genetic and non-genetic risk factors affect response to ranibizumab in exudative age-related macular degeneration. *Acta Ophthalmol* 2015; **93**(6): e451–e457.
 - 37 Ross AH, Donachie PH, Sallam A, Stratton IM, Mohamed Q, Scanlon PH *et al.* Which visual acuity measurements define high-quality care for patients with neovascular age-related macular degeneration treated with ranibizumab? *Eye (Lond)* 2013; **27**(1): 56–64.

Supplementary Information accompanies this paper on Eye website (<http://www.nature.com/eye>)