

Research article

Open Access

Differential gene expression by integrin $\beta 7^+$ and $\beta 7^-$ memory T helper cells

Madeleine W Rodriguez¹, Agnès C Paquet¹, Yee Hwa Yang¹ and David J Erle*^{1,2}

Address: ¹Department of Medicine, University of California, San Francisco, CA 94143-0854 USA and ²Program in Immunology, University of California, San Francisco, CA 94143-0854 USA

Email: Madeleine W Rodriguez - mwillkom@yahoo.com; Agnès C Paquet - apaquet@medsfgh.ucsf.edu; Yee Hwa Yang - jean@biostat.ucsf.edu; David J Erle* - erle@itsa.ucsf.edu

* Corresponding author

Published: 05 July 2004

Received: 12 February 2004

BMC Immunology 2004, 5:13 doi:10.1186/1471-2172-5-13

Accepted: 05 July 2004

This article is available from: <http://www.biomedcentral.com/1471-2172/5/13>

© 2004 Rodriguez et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The cell adhesion molecule integrin $\alpha 4\beta 7$ helps direct the migration of blood lymphocytes to the intestine and associated lymphoid tissues. We hypothesized that $\beta 7^+$ and $\beta 7^-$ blood memory T helper cells differ in their expression of genes that play a role in the adhesion or migration of T cells.

Results: RNA was prepared from $\beta 7^+$ and $\beta 7^-$ CD4⁺ CD45RA⁻ blood T cells from nine normal human subjects and analyzed using oligonucleotide microarrays. Of 21357 genes represented on the arrays, 16 were more highly expressed in $\beta 7^+$ cells and 18 were more highly expressed in $\beta 7^-$ cells (≥ 1.5 fold difference and adjusted $P < 0.05$). Several of the differentially expressed transcripts encode proteins with established or putative roles in lymphocyte adhesion and chemotaxis, including the chemokine receptors CCR9 and CCR10, the integrin $\alpha 4$ subunit, L-selectin, KLRB1 (CD161), NT5E (CD73), LGALS1 and LGALS2 (galectin-1 and -2), and RGS1. Flow cytometry was used to determine whether differences in levels of transcripts encoding cell surface proteins were associated with differential expression of those proteins. Using this approach, we found that surface expression of KLRB1, LAIR1, and NT5E proteins was higher on $\beta 7^+$ memory/effector T cells than on $\beta 7^-$ cells.

Conclusions: Memory/effector T cells that express integrin $\beta 7$ have a distinct pattern of expression of a set of gene transcripts. Several of these molecules can affect cell adhesion or chemotaxis and are therefore likely to modulate the complex multistep process that regulates trafficking of CD4⁺ memory T cell subsets with different homing behaviors.

Background

Lymphocyte migration is a multistep process that involves a complex interplay between adhesion molecules and chemokines and their G protein-coupled receptors [1,2]. Naïve T cells express the adhesion molecule L-selectin, the chemokine receptor CCR7 and other molecules that allow

these cells to migrate preferentially to secondary lymphoid organs where they can encounter antigen-presenting cells. When presented with appropriate antigens, these T cells can differentiate into memory T cells. Some memory cells continue to express L-selectin and/or CCR7 and to migrate efficiently to secondary lymphoid organs,

whereas others lose expression of these molecules and instead express other molecules that direct migration (or "homing") to other organs [1,2].

Extensive investigation has helped to define the role of some adhesion molecules and chemokine receptors in CD4⁺ memory T cell homing to the skin and the gut. The adhesion molecule cutaneous lymphocyte antigen (CLA) and the cutaneous T cell-attracting chemokine receptor CCR10 help control T cell homing to the skin [3-8]. The adhesion molecule integrin $\alpha 4\beta 7$ and the chemokine receptor CCR9 play key roles in homing of lymphocytes to the intestine and Peyer's patches [9-13]. Integrin $\alpha 4\beta 7$ is a receptor for mucosal addressin cell adhesion molecule-1 (MAdCAM-1), a glycoprotein that is expressed by gut endothelium. CCR9 is a receptor for the chemokine TECK (CCL25), which is expressed by endothelial cells and other cells in the small intestine [14,15]. The adhesion molecule and chemokine receptor expression pattern of memory/effector CD4⁺ T cells is strongly influenced by whether initial T cell activation takes place in cutaneous or intestinal lymph nodes [16-18]. Recent evidence suggests that T cell homing receptor expression patterns are "imprinted" by dendritic cells during antigen presentation [19-21]. Although many other molecules have been shown to help control lymphocyte adhesion or migration, it is not clear which if any of these are selectively expressed in gut homing memory CD4⁺ T cells.

We used DNA microarrays to systematically compare RNA transcript expression in human blood $\beta 7^+$ and $\beta 7^-$ CD4⁺ memory T cells. We identified a substantial number of differentially expressed genes, many of which have been previously shown to have effects on cell adhesion and migration. In addition, we showed that some of these transcript expression differences were reflected in differences in surface expression of the encoded proteins.

Results

Microarray analysis of differential gene expression by $\beta 7^+$ and $\beta 7^-$ CD4⁺ memory T cells

As previously reported [22], human blood CD4⁺ T cells could be divided into three distinct subsets based upon their surface expression of integrin $\beta 7$ and the naïve cell marker CD45RA (Fig. 1A). CD45RA⁺ naïve cells expressed low levels of $\beta 7$. CD45RA⁻ memory T cells were divided into a larger population of $\beta 7^-$ cells and a smaller population of $\beta 7^+$ cells. The integrin $\beta 7$ subunit can combine with either integrin $\alpha 4$ (to form the gut homing receptor $\alpha 4\beta 7$) or αE . Integrin $\alpha E\beta 7$ is highly expressed on gut intraepithelial lymphocytes but is rarely expressed by blood T cells and its possible role in homing is less clear [23-25]. Consistent with previous reports [22,23,26], staining with antibodies specific for the $\alpha 4\beta 7$ and $\alpha E\beta 7$ complexes (Act-1 and HML-1 respectively) showed that

the virtually all $\beta 7$ -expressing CD4⁺ T cells in the blood were $\alpha 4\beta 7^+$, and that only $0.5 \pm 0.3\%$ (mean \pm SD) of blood CD4⁺ T cells expressed $\alpha E\beta 7$. We were able to obtain highly purified populations of $\beta 7^+$ and $\beta 7^-$ CD4⁺ CD45RA⁻ cells by flow cytometric sorting (Fig. 1B and 1C).

We used DNA microarrays to compare RNA transcript expression by $\beta 7^+$ and $\beta 7^-$ cells from 9 healthy adult human subjects. Using this approach, we identified 16 genes that were expressed at higher levels in $\beta 7^+$ cells (at least 1.5-fold higher expression and $P < 0.05$ when adjusted for multiple comparisons). These are listed in Table 1. Our estimates of fold difference are likely to be conservative since the use of amplified RNA (as opposed to unamplified cDNA) tends to underestimate fold differences [27]. The most highly differentially expressed transcript encodes the chemokine receptor CCR9. CCR9 protein has previously been shown to be preferentially expressed on $\alpha 4\beta 7^+$ blood CD4⁺ T cells [13]. The ITGA4 transcript encodes the α subunit of integrin $\alpha 4\beta 7$, and was expressed at 1.7-fold higher levels on $\beta 7^+$ cells. Other genes listed in Table 1 include genes encoding cell surface receptors (LRRN3, CD1C, KLRB1, LAIR1, IL18RAP, KLRG1), transcription factors (RAM2, SREBF1) and a transcriptional repressor known to play a role in T cell differentiation (GFI1). A complete listing of all gene expression results is provided in Appendix 2 [see additional file 2].

We also identified 18 gene transcripts that were expressed at significantly lower levels in $\beta 7^+$ cells compared with $\beta 7^-$ cells (at least 1.5 times lower and adjusted $P < 0.05$). These are listed in Table 2. GPR2, the most highly differentially expressed transcript, encodes CCR10, the receptor for the cutaneous T cell attracting chemokine (CTACK/CCL27). Hudak et al. [8] found that CCR10 protein was present on a subset of CD4⁺ CD45RA⁻ T cells that express the skin homing receptor CLA but do not express integrin $\alpha 4\beta 7$. We also found that $\beta 7^-$ cells expressed more transcript for SELL, which encodes L-selectin (CD62L), a non-integrin cell adhesion molecule that mediates lymphocyte adhesion to peripheral lymph node high endothelial venules. None of the other genes listed in Table 2 have established roles in organ-specific homing, although both RGS1 and LGALS1 have been reported to modulate lymphocyte adhesion (see Discussion). Given the importance of G protein-coupled receptors in regulating leukocyte migration, we note that in addition to GPR2/CCR10, one other G protein-coupled receptor, P2RY5, was expressed at higher levels on $\beta 7^-$ cells. Although classified as a purinergic receptor, expression of P2RY5 did not result in nucleotide-evoked signaling responses, and the function of this receptor remains unknown [28].




Figure 1
Integrin $\beta 7$ expression on blood T helper cells and on purified $\beta 7^-$ and $\beta 7^+$ memory T helper cell subsets. Dot density plots depicting expression of the naïve cell marker CD45RA and integrin $\beta 7$ on CD4⁺ blood T cells (A) and on purified $\beta 7^+$ (B) and $\beta 7^-$ (C) memory cells in arbitrary fluorescence units. Sorted populations were ~99% pure. Quadrant markers indicate settings used to sort populations based on CD45RA and $\beta 7$ expression. Colors indicate the relative frequency of cells (from lowest to highest density: green, purple, light blue, orange, dark blue, and yellow).

Table 1: Gene transcripts with higher expression in $\beta 7^+$ versus $\beta 7^-$ CD4⁺ CD45RA⁻ T helper cells*

Symbol	Name	Accession	Fold Difference	P value
CCR9	chemokine (C-C motif) receptor 9	NM_031200	+3.0	< 0.01
CCL5	chemokine (C-C motif) ligand 5	NM_002985	+2.4	< 0.01
RAM2	transcription factor RAM2	NM_018719	+2.2	< 0.01
LRRN3	leucine rich repeat neuronal 3	AL442092	+2.1	< 0.01
GFI1	growth factor independent 1	NM_005263	+1.8	< 0.01
ITGA4	integrin, alpha 4 (CD49D)	NM_000885	+1.7	< 0.01
CD1C	CD1C antigen, c polypeptide	NM_001765	+1.7	< 0.01
KLRB1	killer cell lectin-like receptor subfamily B, member 1	NM_002258	+1.7	< 0.01
LAIR1	leukocyte-associated Ig-like receptor 1	NM_002287	+1.7	< 0.01
RRM2	ribonucleotide reductase M2 polypeptide	NM_001034	+1.6	< 0.01
--	Homo sapiens cDNA FLJ32290 fis, clone PROST2000463	AK056852	+1.6	< 0.01
HHL	expressed in hematopoietic cells, heart, liver	NM_014857	+1.6	0.02
IL18RAP	interleukin 18 receptor accessory protein	NM_003853	+1.6	< 0.01
SREBF1	sterol regulatory element binding transcription factor 1	NM_004176	+1.6	< 0.01
KLRG1	killer cell lectin-like receptor subfamily G, member 1	NM_005810	+1.5	< 0.01
LGALS2	lectin, galactoside-binding, soluble, 2 (galectin 2)	NM_006498	+1.5	0.01

* Includes all transcripts with fold difference ≥ 1.5 and adjusted P < 0.05. Positive fold difference values indicate higher expression on $\beta 7^+$ cells.

Comparison of surface protein expression on $\beta 7^+$ and $\beta 7^-$ CD4⁺ memory T cells

We wished to determine whether differences in expression of selected transcripts were reflected in differences in expression of the cell surface proteins encoded by those transcripts. As previously discussed, the products of the two most highly differentially expressed genes (CCR9 and GPR2/CCR10) have already been shown to be highly

selectively expressed on $\alpha 4\beta 7^+$ and $\alpha 4\beta 7^-$ CD4⁺ T cells respectively [8,13]. We selected four other cell surface proteins encoded by transcripts that were expressed more highly on $\beta 7^+$ cells for further analysis based upon the availability of suitable antibodies. Three of these (LAIR1, KLRB1, and KLRG1) are listed in Table 1. A fourth, NT5E (ecto-5'-nucleotidase, also known as CD73), was also significantly higher in $\beta 7^+$ cells (adjusted P < 0.01),

Table 2: Gene transcripts with higher expression in $\beta7^-$ versus $\beta7^+ CD4^+ CD45RA^-$ T helper cells*

Symbol	Name	Accession	Fold Difference	P value
GPR2	G protein-coupled receptor 2 (CCR10)	NM_016602	-3.2	< 0.01
RGS1	regulator of G-protein signalling 1	NM_002922	-2.0	< 0.01
TRIM2	tripartite motif-containing 2	NM_015271	-1.9	< 0.01
C1orf24	chromosome 1 open reading frame 24	NM_052966	-1.9	< 0.01
CAMTA1	calmodulin binding transcription activator 1	AB020640	-1.9	< 0.01
LMNA	lamin A/C	NM_005572	-1.9	< 0.01
LGALS1	lectin, galactoside-binding, soluble, 1 (galectin 1)	NM_002305	-1.9	< 0.01
P2RY5	purinergic receptor P2Y, G-protein coupled, 5	NM_005767	-1.8	< 0.01
SELL	selectin L (lymphocyte adhesion molecule 1)	NM_000655	-1.8	0.02
--	similar to RIKEN cDNA 1700007B22	AK021437	-1.8	< 0.01
ChGn	chondroitin beta1,4 N-acetylgalactosaminyltransferase	NM_018371	-1.8	< 0.01
--	similar to BcDNA:GH11415 gene product	AK056276	-1.7	< 0.01
PTPLA	protein tyrosine phosphatase-like, member a	NM_014241	-1.7	< 0.01
MRC2	mannose receptor, C type 2	NM_006039	-1.6	< 0.01
PRNP	prion protein (p27-30)	NM_000311	-1.6	< 0.01
PHLDA1	pleckstrin homology-like domain, family A, member 1	AK026181	-1.6	< 0.01
KRT1	keratin 1	NM_006121	-1.5	< 0.01
SEMA5A	semaphorin 5A	NM_003966	-1.5	< 0.01

* Includes all transcripts with fold difference ≤ -1.5 and adjusted $P < 0.05$. Negative fold difference values indicate higher expression on $\beta7^-$ cells.

Table 3: Surface expression of five gene products on $\beta7^+$ versus $\beta7^- CD4^+$ T cells


Gene	Fold difference: Transcript**	Fold difference: Surface protein***	Mean surface protein expression*		
			$CD45RA^- \beta7^+$ cells	$CD45RA^- \beta7^-$ cells	$CD45RA^+$ cells
LAIR1	+1.7	+2.5 (+1.6 to +3.9)	34.6	15.3	37.3
KLRB1	+1.7	+2.5 (+2.2 to +3.6)	26.1	10.7	1.1
KLRG1	+1.5	1.0 (-3.2 to +2.1)	17.7	15.6	9.1
NT5E	+1.3	+1.7 (+1.2 to +3.2)	20.3	11.7	6.8
SELL	-1.8	-1.8 (-2.1 to -1.5)	22.8	40.2	50.7

* Mean fluorescence intensity (MFI, in arbitrary units) was measured for 5–8 subjects. ** As determined using DNA microarrays. *** Values represent mean (range) of the fold difference in MFI between $CD45RA^- \beta7^+$ cells and $CD45RA^- \beta7^-$ cells for all subjects. Positive fold difference values indicate higher expression on $\beta7^+$ cells and negative fold difference values indicate higher expression on $\beta7^-$ cells.

although the estimated magnitude of the difference was smaller (1.3-fold). We used flow cytometry to measure expression of each of these four molecules on $CD4^+$ T cell subsets from human blood. For comparison, we also analyzed surface expression of one protein (SELL/CD62L) encoded by a transcript that was expressed at higher levels on $\beta7^-$ cells. We used a different population of study subjects from that used in the microarray studies to ensure that our findings could be validated in an independent sample. For four of the five proteins examined, the extent of differential surface expression was at least as great as predicted from the microarray data (Table 3). The other

protein, KLRG1, was not consistently differentially expressed.

We used flow cytometry to conduct a more detailed analysis of the expression of KLRB1, NT5E and LAIR1 on T helper cell subsets with distinct homing properties. A small subset of memory cells expressed the skin homing receptor CLA (~4% of $CD4^+ CD45RA^-$ T cells). These CLA⁺ cells had low expression of KLRB1, NT5E and LAIR1 (similar to levels found in other $\beta7^- CD45RA^-$ cells, data not shown). The adhesion molecule L-selectin (SELL, CD62L) and the chemokine receptor CCR7 both play roles in homing to secondary lymphoid organs and each is

**Figure 2**

A $\beta 7^{\text{high}}$ subset of memory/effector T helper cells lacks CD62L and CCR7. Dot density plots depicting expression of integrin $\beta 7$ and (A) SELL (CD62L) or (B) CCR7 on $\text{CD4}^+ \text{CD45RA}^-$ T cells. Quadrant lines were drawn based on staining obtained with control antibodies.

expressed on subsets of memory/effector T cells. Memory/effector cells that lacked L-selectin and CCR7 included a subset with particularly high $\beta 7$ expression and a larger subset with no $\beta 7$ expression (Fig. 2). CD45RA^+ cells that expressed L-selectin and CCR7 (central memory cells) also included $\beta 7^+$ and $\beta 7^-$ subsets. We found that KLRB1, a member of the killer cell lectin-like receptor superfamily also known as CD161 or NKR1A [29], was highly expressed on $\beta 7^+$ CCR7 $^-$ memory effector T helper cells (Fig. 3). There was less expression on $\beta 7^-$ CCR7 $^-$ memory effector cells and on CCR7 $^+$ central memory cells. KLRB1 was essentially undetectable on CD45RA^+ naïve cells. The expression pattern of NT5E/CD73 was similar, although NT5E expression was limited to a smaller fraction of cells than KLRB1 expression (Fig. 4). In contrast, LAIR1 had a different pattern of expression (Fig. 5). LAIR1 was most highly expressed on naïve cells, and was also highly expressed on $\beta 7^+$ CD45RA^- cells (both CCR7 $^-$ and CCR7 $^+$ subsets). LAIR1 was less highly expressed on $\beta 7^-$ CD45RA^- cells (especially the CCR7 $^-$ subset). LAIR1 contains two ITIM domains and can transmit signals following ligation by antibody, however the nature of the LAIR1 ligand(s) and the biological function of LAIR1 on CD4^+ T cells remains unknown [30].

Discussion

We used DNA microarrays to identify gene transcripts that were differentially expressed on the subset of CD4^+ memory/effector T cells that express integrin $\beta 7$. Since the overwhelming majority of $\beta 7^+$ cells in blood express the gut homing receptor integrin $\alpha 4\beta 7$, this approach was designed to identify genes that might play a role in organ-specific homing. We identified 16 transcripts that were more highly expressed in $\beta 7^+$ cells and 18 that were more highly expressed on $\beta 7^-$ cells. We can relate transcript expression to expression of protein in seven cases (Table 3 and refs. [8,13]). In six of these cases (CCR9, GPR2/CCR10, KLRB1, LAIR1, NT5E, and SELL), differential transcript expression was accompanied by differential expression of the corresponding protein on the surface of $\beta 7^+$ versus $\beta 7^-$ cells. In one case (KLRG1), we did not find evidence for a consistent difference in surface protein expression (Table 3). These data suggest that most of the differentially expressed transcripts correspond to differentially expressed proteins, although it will be important to confirm this directly when suitable reagents become available.

Some of the differentially expressed transcripts encode proteins with known roles in organ-specific homing. The




Figure 3

Expression of KLRB1 (CD161) on T helper cell subsets. Histograms depict expression of KLRB1 (CD161) on (A) CD45RA⁺ naïve cells and (B) four subsets of CD45RA⁻ memory/effector cells. These subsets were distinguished by their expression of $\beta 7$ and CCR7, a marker for central memory cells, as shown in Fig. 2B. Black lines indicate KLRB1 staining and filled red lines indicate staining obtained with an isotype control antibody.

two most highly differentially expressed genes we detected in our analysis encode chemokine receptors already shown to have important roles in gut and non-gut (skin) homing. These are CCR9, the TECK/CCL25 receptor which is expressed preferentially on $\alpha 4\beta 7^+$ gut homing cells [13], and GPR2/CCR10, the CTACK/CCL27 receptor which is expressed preferentially on CLA⁺ skin homing cells [8]. We found that transcripts encoding the α subunit (ITGA4) of integrin $\alpha 4\beta 7$, was more highly expressed in $\beta 7^+$ cells. The microarray experiments indicated that the transcript encoding the $\beta 7$ subunit itself (ITGB7) was not differentially expressed, consistent with previous results obtained using quantitative RT-PCR (M.W.R and D.J.E, unpublished results). This suggests that surface expression of the $\alpha 4\beta 7$ heterodimer on CD4⁺ memory T cells may be determined primarily by the amount of $\alpha 4$ subunit available and not by the amount of $\beta 7$ produced. The other differentially expressed transcript with a known role in homing is SELL, which encodes the peripheral lymph node homing receptor L-selectin (CD62L). We found that expression of both L-selectin transcript and surface pro-

tein was higher in $\beta 7^-$ than $\beta 7^+$ memory/effector T cells. Although many $\beta 7$ -expressing memory cells also expressed L-selectin, cells expressing the highest levels of $\beta 7$ did not express L-selectin (Fig. 2). This presumably contributes to the development of CD4⁺ T cell subsets with distinct homing properties.

We also found evidence that $\beta 7^+$ and $\beta 7^-$ cells differ in their expression of other molecules that have been reported to modulate cell adhesion and migration. KLRB1 transcript and surface protein expression was higher on $\beta 7^+$ memory effector T cells. Overexpression of KLRB1 (also known as CD161 or NKR1A) was previously found to increase transendothelial migration of CD4⁺ T cells, perhaps by modulating integrin activity [31]. The finding that KLRB1 was preferentially expressed on T cells from human intestine led to a recent suggestion that KLRB1 may help direct intestinal homing [32]. This hypothesis remains to be tested, but would be consistent with our finding of preferential expression of KLRB1 on $\beta 7^+$ blood memory CD4⁺ T cells. We also found higher expression of




Figure 4

Expression of NT5E (CD73) on T helper cell subsets. Histograms depict expression of NT5E (CD73) on (A) CD45RA⁺ naïve cells and (B) four subsets of CD45RA⁻ memory/effector cells. These subsets were distinguished by their expression of $\beta 7$ and CCR7, a marker for central memory cells, as shown in Fig. 2B. Black lines indicate NT5E staining and filled red lines indicate staining obtained with an isotype control antibody.

NT5E (CD73) transcripts and surface protein on $\beta 7^+$ memory effector T cells. NT5E is an ecto 5'-nucleotidase expressed on lymphocytes and on endothelial cells and has been reported to mediate lymphocyte-endothelial adhesion and transmigration by various mechanisms, including promotion of integrin-mediated binding via effects on integrin clustering [33-36]. Transcripts encoding two members of the galectin family were differentially expressed on $\beta 7^+$ versus $\beta 7^-$ cells. Both LGALS1/galectin-1 (expressed at higher levels in $\beta 7^-$ cells) and LGALS2/galectin-2 (expressed at higher levels in $\beta 7^+$ cells) have been shown to have several effects on cell adhesion in lymphocytes and other cells [37]. Transcripts for RGS1 were expressed at higher levels in $\beta 7^-$ cells. RGS1 is a GTPase activating protein that regulates G α i-stimulated pathways. Expression of RGS1 inhibits chemokine-induced cell migration and integrin-mediated adhesion in lymphocytes [38]. Our data suggest that there may be less RGS1-mediated inhibition of chemokine signaling in $\beta 7^+$ cells. We speculate that the differential expression of KLRB1, NT5E, LGALS1, LGALS2, and RGS1 on $\beta 7^+$ and $\beta 7^-$

T helper cell subsets may contribute to homing specificity by modulating the complex multistep process of cell adhesion and migration.

The primary activation of T cells leads to changes in the expression of integrin $\alpha 4\beta 7$ and other molecules that control homing. It has recently been shown that upregulation of integrin $\alpha 4\beta 7$ expression and increased responsiveness to the CCR9 ligand TECK occur within two days after activation of murine naïve CD4⁺ T cells in intestinal, but not cutaneous, lymph nodes [18] and that activation by dendritic cells from Peyer's patches but not peripheral lymph node or spleen promotes murine CD8⁺ T cell $\alpha 4\beta 7$ expression and TECK responsiveness [19]. We found that both KLRB1 and NT5E were virtually absent on CD45RA⁺ naïve CD4⁺ T cells and were expressed at low levels on CD45RA⁻ CCR7⁺ central memory cells and on $\beta 7^-$ CD45RA⁻ non-gut homing memory effector cells but were expressed at higher levels in the $\beta 7^{\text{high}}$ CCR7⁻ CD45RA⁻ memory effector subset. In contrast, LAIR1 was highly expressed on naïve T cells and on both CCR7⁻ and CCR7⁺ $\beta 7^+$ CD45RA⁻




Figure 5

Expression of LAIR1 on T helper cell subsets. Histograms depict expression of LAIR1 on (A) CD45RA⁺ naive cells and (B) four subsets of CD45RA⁻ memory/effector cells. These subsets were distinguished by their expression of $\beta 7$ and CCR7, a marker for central memory cells, as shown in Fig. 2B. Black lines indicate LAIR1 staining and filled red lines indicate staining obtained with an isotype control antibody.

cells but expression was lower on $\beta 7^-$ CD45RA⁻ cells. These results suggest that expression of $\alpha 4\beta 7$ and expression of KLRB1, NT5E and LAIR1 may be coordinately regulated at the time of human T cell activation, as is apparently the case for $\alpha 4\beta 7$ and CCR9 in mouse systems. Alternatively, it is possible that differences in expression arise from changes in gene expression that occur after the transition from naive to memory effector cells or from the preferential retention of survival of particular memory effector cell subsets. Although we showed that expression of KLRB1, NT5E and LAIR1 was higher for the $\beta 7^+$ subset as a whole, many $\beta 7^+$ cells did not express these proteins and some $\beta 7^-$ cells did. This suggests that the homing properties of individual T cells might be "fine tuned" by complex combinatorial regulation of molecules known to have major roles in determining organ-specific recruitment (e.g., integrin $\alpha 4\beta 7$ and CCR9) and other molecules that can modulate adhesion and migration (e.g., KLRB1 and NT5E).

Conclusions

We used DNA microarrays to identify a set of gene transcripts that were differentially expressed on $\beta 7^+$ versus $\beta 7^-$ blood memory CD4⁺ T cells. Some of these encode adhesion receptors (integrin $\alpha 4$ and L-selectin) and chemokine receptors (CCR9 and GPR2/CCR10) already known to be involved in organ-specific lymphocyte migration. Other differentially expressed transcripts encode additional molecules previously reported to affect cell adhesion and migration, including KLRB1, NT5E/CD73, LGALS1 and LGALS2 (galectin-1 and -2) and RGS1. As predicted from the transcript expression data, we found that KLRB1, NT5E and LAIR1 proteins were all preferentially expressed on the surface of $\beta 7^+$ versus $\beta 7^-$ memory CD4⁺ T cells. Our results suggest that expression of several molecules that can modify T cell adhesion, endothelial transmigration and chemotaxis are also selectively regulated on CD4⁺ T cell subsets, perhaps during the "imprinting" process that takes place during initial T cell activation. These molecules are likely to contribute to the complex multistep process that regulates trafficking of

CD4⁺ memory T cell subsets with different homing behaviors.

Methods

Isolation of $\beta 7^+$ and $\beta 7^-$ lymphocytes from human blood

This protocol was approved by the Committee on Human Research at the University of California, San Francisco and informed consent was obtained from all subjects. Peripheral blood mononuclear cells (PBMC) were isolated from nine normal adult donors (six females and three males, ages 24–42 years) using sodium heparin coated Vacutainer Cell Preparation Tubes (Becton Dickinson) according to the manufacturer's instructions. PBMC were stained with CD4-FITC, CD45RA-Cy5 and anti-integrin $\beta 7$ -PE (all from BD Pharmingen). Anti-FITC microbeads and a MACS bead sorter (Miltenyi Biotec) were used to enrich for the CD4⁺ cells prior to flow cytometry. CD4⁺ memory (CD45RA⁻) $\beta 7^+$ and $\beta 7^-$ T cells were then purified using a FACS Vantage (Becton Dickinson).

Microarray analysis of transcript expression

Total RNA from $\beta 7^+$ and $\beta 7^-$ CD4⁺ memory T cells was purified using the Mini RNA Isolation Kit (Zymo Research) and DNase treated using the DNA-free RNA Kit (Zymo Research) according to the manufacturer's instructions. RNA quality was evaluated using an Agilent Bioanalyzer. RNA amplification, labeling, and hybridization were performed as previously described with slight modifications [27]. In brief, two rounds of amplification were performed using T7 RNA polymerase (Ambion MessageAmp aRNA Kit). Cy3 or Cy5 was incorporated into the amplified RNA products using amino allyl-modified nucleotides. Fluorescently-labeled amplified RNAs were fragmented using Ambion RNA Fragmentation Reagents and hybridized to DNA microarrays using Ambion Slide-Hyb Glass Array Hybridization Buffer #1. Each hybridization involved $\beta 7^+$ cell RNA from a single subject (labeled with one dye) and $\beta 7^-$ cell RNA from the same subject (labeled with the other dye). A total of 27 arrays were used to analyze samples from the 9 subjects (at least 2 arrays per subject, including a dye swap).

Arrays were produced in our microarray facility using the Operon version 2 set of 70-mer oligonucleotide probes supplemented with some custom-designed 70-mers as described [27]. Arrays included probes for 21357 genes plus control probes. After hybridization, arrays were scanned using an Axon GenePix 4000B scanner and images were processed using GenePix 5.0 software. The "print-tip loess" normalization was used to correct for within-array dye and spatial effects [39] and single channel quantile normalization was used to facilitate comparison between arrays [40]. We used functions in the library *marrayNorm* [41] of the R / Bioconductor package [42] to perform these normalizations. After normalization we

determined a log ratio, $\log_2(\beta 7^+$ sample intensity/ $\beta 7^-$ sample intensity), for each probe on each array. No background subtraction was performed. Complete information about the array platform and data from each of the individual arrays is available from GEO [43] (GEO accession GSE1039). Appendix 1 [see additional file 1] is a MIAME-compliant description of the array experiments.

Identification of differentially expressed gene transcripts

A fixed effects linear model was used to estimate differences in transcript expression between $\beta 7^+$ and $\beta 7^-$ CD4⁺ T cells. For each individual gene (probe) on the array, the model can be described as

$$Y_{ij} = \mu + A_i + \epsilon_{ij},$$

where Y_{ij} represents the normalized log ratio determined on a single array, i represents the subject number and j represents the array number. The parameter μ represents the actual log ratio of gene expression between $\beta 7^+$ and $\beta 7^-$ cells, which we wished to estimate. A_i represents the difference between the log ratio for subject i and the average log ratio for all subjects and ϵ denotes the experimental error. We fit the model using a zero sum constraint ($\sum A_i = 0$). Fold difference was calculated from the estimated log ratio. We computed moderated t-statistics [44], log-odds ratios of differential expression (based on empirical Bayes shrinkage of the standard errors towards a common value [45]), and adjusted p-values (obtained using the Bonferroni correction) using functions in the *limma* library of the Bioconductor software package [41,42]. Similar results were obtained under a random effects model (not shown).

Analysis of surface protein expression on blood CD4⁺ T cell subsets

Whole blood was collected from nine normal adult subjects (seven females and two males, ages 23–38 years, none of whom were included in the microarray experiments described above). Blood was stained with various antibody combinations and analyzed on a FACScalibur equipped for six color analysis. Antibodies recognizing CD4 (SK3), integrin $\beta 7$ (FIB504), CD62L (DREG-56), CD73 (AD2), and CD161 (DX12) and isotype control antibodies were all from BD Biosciences. Other antibodies used were LAIR1 (DX26, generous gift of L. Lanier), KLRG1 (13A2, generous gift of D. Voehringer and H. Pircher), CD45RA (Immunotech), the anti- $\alpha 4\beta 7$ mAb Act-1 [46], the anti- $\alpha E\beta 7$ antibody HML-1 (Beckman Coulter), CCR7 (R&D Systems). Mean fluorescence intensity (MFI) was corrected by subtracting the MFI obtained with isotype control antibodies. The percentage of positive cells was determined by setting the threshold so that $\leq 1\%$ of cells were considered positive after staining with control antibody.

Authors' contributions

MWR recruited the subjects, performed the experiments and participated in data analysis and drafting of the manuscript. ACP performed the statistical analysis of the microarray data. YHY supervised the statistical analysis of the microarray data. DJE conceived of the study and participated in design, analysis, and preparation of the manuscript. All authors read and approved the final manuscript.

Additional material

Additional File 2

Tab-delimited text file with calculated gene expression ratios for each probe on each array. Values represent normalized \log_2 (C_{γ5} intensity/C_{γ3} intensity).

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2172-5-13-S2.txt>]

Additional File 1

MIAME-compliant description of the array experiments in rich text format.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2172-5-13-S1.rtf>]

Acknowledgements

Microarray studies were supported by Andrea Barczak, Michael Salazar, and the UCSF/San Francisco General Hospital General Clinical Research Center Core Genomics Laboratory (supported by grant M01-RR00083 from the National Center for Research Resources of the NIH). Flow cytometry was performed in the Flow Cytometry Core Laboratory of the J. David Gladstone Institutes. We thank Gordon Smyth and Mark Segal for helpful discussions regarding the microarray data analysis. The work was supported by NIH grant DK/AI54212.

References

- von Andrian UH, Mackay CR: **T-cell function and migration. Two sides of the same coin.** *N Engl J Med* 2000, **343**:1020-1034.
- Kunkel EJ, Butcher EC: **Chemokines and the tissue-specific migration of lymphocytes.** *Immunity* 2002, **16**:1-4.
- Berg EL, Yoshino T, Rott LS, Robinson MK, Warnock RA, Kishimoto TK, Picker LJ, Butcher EC: **The cutaneous lymphocyte antigen is a skin lymphocyte homing receptor for the vascular lectin endothelial cell-leukocyte adhesion molecule 1.** *J Exp Med* 1991, **174**:1461-1466.
- Tietz W, Allemand Y, Borges E, von Laer D, Hallmann R, Vestweber D, Hamann A: **CD4+ T cells migrate into inflamed skin only if they express ligands for E- and P-selectin.** *J Immunol* 1998, **161**:963-970.
- Homey B, Wang W, Soto H, Buchanan ME, Wiesenborn A, Catron D, Muller A, McClanahan TK, Dieu-Nosjean MC, Orozco R, Ruzicka T, Lehmann P, Oldham E, Zlotnik A: **Cutting edge: the orphan chemokine receptor G protein-coupled receptor-2 (GPR-2, CCR10) binds the skin-associated chemokine CCL27 (CTACK/ALP/ILC).** *J Immunol* 2000, **164**:3465-3470.
- Reiss Y, Proudfoot AE, Power CA, Campbell JJ, Butcher EC: **CC chemokine receptor (CCR)4 and the CCR10 ligand cutaneous T cell-attracting chemokine (CTACK) in lymphocyte trafficking to inflamed skin.** *J Exp Med* 2001, **194**:1541-1547.
- Homey B, Alenius H, Muller A, Soto H, Bowman EP, Yuan W, McEvoy L, Lauerma AI, Assmann T, Bunemann E, Lehto M, Wolff H, Yen D, Marxhausen H, To W, Sedgwick J, Ruzicka T, Lehmann P, Zlotnik A: **CCL27-CCR10 interactions regulate T cell-mediated skin inflammation.** *Nat Med* 2002, **8**:157-165.
- Hudak S, Hagen M, Liu Y, Catron D, Oldham E, McEvoy LM, Bowman EP: **Immune surveillance and effector functions of CCR10(+) skin homing T cells.** *J Immunol* 2002, **169**:1189-1196.
- Berlin C, Berg EL, Briskin MJ, Andrew DP, Kilshaw PJ, Holzmann B, Weissman IL, Hamann A, Butcher EC: **Alpha 4 beta 7 integrin mediates lymphocyte binding to the mucosal vascular addressin MAdCAM-1.** *Cell* 1993, **74**:185-185.
- Hamann A, Andrew DP, Jablonski-Westrich D, Holzmann B, Butcher EC: **Role of alpha 4-integrins in lymphocyte homing to mucosal tissues in vivo.** *J Immunol* 1994, **152**:3282-3293.
- Abitorabi MA, Mackay CR, Jerome EH, Osorio O, Butcher EC, Erle DJ: **Differential expression of homing molecules on recirculating lymphocytes from sheep gut, peripheral, and lung lymph.** *J Immunol* 1996, **156**:3111-3117.
- Williams MB, Butcher EC: **Homing of naive and memory T lymphocyte subsets to Peyer's patches, lymph nodes, and spleen.** *J Immunol* 1997, **159**:1746-1752.
- Zabel BA, Agace WW, Campbell JJ, Heath HM, Parent D, Roberts AI, Ebert EC, Kassam N, Qin S, Zovko M, LaRosa GJ, Yang LL, Soler D, Butcher EC, Ponath PD, Parker CM, Andrew DP: **Human G protein-coupled receptor GPR-9-6/CC chemokine receptor 9 is selectively expressed on intestinal homing T lymphocytes, mucosal lymphocytes, and thymocytes and is required for thymus-expressed chemokine-mediated chemotaxis.** *J Exp Med* 1999, **190**:1241-1256.
- Kunkel EJ, Campbell JJ, Haraldsen G, Pan J, Boisvert J, Roberts AI, Ebert EC, Vierra MA, Goodman SB, Genovese MC, Wardlaw AJ, Greenberg HB, Parker CM, Butcher EC, Andrew DP, Agace WW: **Lymphocyte CC chemokine receptor 9 and epithelial thymus-expressed chemokine (TECK) expression distinguish the small intestinal immune compartment: Epithelial expression of tissue-specific chemokines as an organizing principle in regional immunity.** *J Exp Med* 2000, **192**:761-768.
- Papadakis KA, Landers C, Prehn J, Kouroumalis EA, Moreno ST, Gutierrez-Ramos JC, Hodge MR, Targan SR: **CC chemokine receptor 9 expression defines a subset of peripheral blood lymphocytes with mucosal T cell phenotype and Th1 or T-regulatory 1 cytokine profile.** *J Immunol* 2003, **171**:159-165.
- Picker LJ, Treer JR, Ferguson-Darnell B, Collins PA, Buck D, Terstappen LV: **Control of lymphocyte recirculation in man. I. Differential regulation of the peripheral lymph node homing receptor L-selectin on T cells during the virgin to memory cell transition.** *J Immunol* 1993, **150**:1105-1121.
- Picker LJ, Treer JR, Ferguson-Darnell B, Collins PA, Bergstresser PR, Terstappen LV: **Control of lymphocyte recirculation in man. II. Differential regulation of the cutaneous lymphocyte-associated antigen, a tissue-selective homing receptor for skin-homing T cells.** *J Immunol* 1993, **150**:1122-1136.
- Campbell DJ, Butcher EC: **Rapid acquisition of tissue-specific homing phenotypes by CD4(+) T cells activated in cutaneous or mucosal lymphoid tissues.** *J Exp Med* 2002, **195**:135-141.
- Mora JR, Bono MR, Manjunath N, Weninger W, Cavanagh LL, Roseblatt M, Von Andrian UH: **Selective imprinting of gut-homing T cells by Peyer's patch dendritic cells.** *Nature* 2003, **424**:88-93.
- Stagg AJ, Kamm MA, Knight SC: **Intestinal dendritic cells increase T cell expression of alpha4beta7 integrin.** *Eur J Immunol* 2002, **32**:1445-1454.
- Dudda JC, Simon JC, Martin S: **Dendritic cell immunization route determines CD8+ T cell trafficking to inflamed skin: role for tissue microenvironment and dendritic cells in establishment of T cell-homing subsets.** *J Immunol* 2004, **172**:857-863.
- Erle DJ, Briskin MJ, Butcher EC, Garcia-Pardo A, Lazarovits AI, Tidswell M: **Expression and function of the MAdCAM-1 receptor, integrin alpha 4 beta 7, on human leukocytes.** *J Immunol* 1994, **153**:517-528.
- Cerf-Bensussan N, Jarry A, Brousse N, Lisowska-Groszpierska B, Guy-Grand D, Griscelli C: **A monoclonal antibody (HML-1) defining a novel membrane molecule present on human intestinal lymphocytes.** *Eur J Immunol* 1987, **17**:1279-1285.

24. Cerf-Bensussan N, Begue B, Gagnon J, Meo T: **The human intraepithelial lymphocyte marker HML-1 is an integrin consisting of a beta 7 subunit associated with a distinctive alpha chain.** *Eur J Immunol* 1992, **22**:273-277.
25. Parker CM, Cepek KL, Russell GJ, Shaw SK, Posnett DN, Schwarting R, Brenner MB: **A family of beta 7 integrins on human mucosal lymphocytes.** *Proc Natl Acad Sci U S A* 1992, **89**:1924-1928.
26. Erle DJ, Brown T, Christian D, Aris R: **Lung epithelial lining fluid T cell subsets defined by distinct patterns of beta 7 and beta 1 integrin expression.** *Am J Respir Cell Mol Biol* 1994, **10**:237-244.
27. Barczak A, Rodriguez MW, Hanspers K, Koth LL, Tai YC, Bolstad BM, Speed TP, Erle DJ: **Spotted long oligonucleotide arrays for human gene expression analysis.** *Genome Res* 2003, **13**:1775-1785.
28. Ralevic V, Burnstock G: **Receptors for purines and pyrimidines.** *Pharmacol Rev* 1998, **50**:413-492.
29. Lanier LL, Chang C, Phillips JH: **Human NKR-PIA. A disulfide-linked homodimer of the C-type lectin superfamily expressed by a subset of NK and T lymphocytes.** *J Immunol* 1994, **153**:2417-2428.
30. Meyaard L, Adema GJ, Chang C, Woollatt E, Sutherland GR, Lanier LL, Phillips JH: **LAIR-1, a novel inhibitory receptor expressed on human mononuclear leukocytes.** *Immunity* 1997, **7**:283-290.
31. Poggi A, Costa P, Zocchi MR, Moretta L: **NKRPIA molecule is involved in transendothelial migration of CD4+ human T lymphocytes.** *Immunol Lett* 1997, **57**:121-123.
32. Iiai T, Watanabe H, Suda T, Okamoto H, Abo T, Hatakeyama K: **CD161+ T (NT) cells exist predominantly in human intestinal epithelium as well as in liver.** *Clin Exp Immunol* 2002, **129**:92-98.
33. Henttinen T, Jalkanen S, Yegutkin GG: **Adherent leukocytes prevent adenosine formation and impair endothelial barrier function by Ecto-5'-nucleotidase/CD73-dependent mechanism.** *J Biol Chem* 2003, **278**:24888-24895.
34. Airas L, Niemela J, Jalkanen S: **CD73 engagement promotes lymphocyte binding to endothelial cells via a lymphocyte function-associated antigen-1-dependent mechanism.** *J Immunol* 2000, **165**:5411-5417.
35. Airas L, Hellman J, Salmi M, Bono P, Puurunen T, Smith DJ, Jalkanen S: **CD73 is involved in lymphocyte binding to the endothelium: characterization of lymphocyte-vascular adhesion protein 2 identifies it as CD73.** *J Exp Med* 1995, **182**:1603-1608.
36. Arvilommi AM, Salmi M, Airas L, Kalimo K, Jalkanen S: **CD73 mediates lymphocyte binding to vascular endothelium in inflamed human skin.** *Eur J Immunol* 1997, **27**:248-254.
37. Hughes RC: **Galectins as modulators of cell adhesion.** *Biochimie* 2001, **83**:667-676.
38. Bowman EP, Campbell JJ, Druey KM, Scheschonka A, Kehrl JH, Butcher EC: **Regulation of chemotactic and proadhesive responses to chemoattractant receptors by RGS (regulator of G-protein signaling) family members.** *J Biol Chem* 1998, **273**:28040-28048.
39. Yang YH, Dudoit S, Luu P, Lin DM, Peng V, Ngai J, Speed TP: **Normalization for cDNA microarray data: a robust composite method addressing single and multiple slide systematic variation.** *Nucleic Acids Res* 2002, **30**:e15.
40. Yang YH, Thorne N: **Normalization for Two-color cDNA Microarray Data.** *Science and Statistics: A Festschrift for Terry Speed* 2003, **40**:403-418.
41. Dudoit S, Yang YH: **Bioconductor R packages for exploratory analysis and normalization of cDNA microarray data.** *The Analysis of Gene Expression Data: Methods and Software* Edited by: Parmigiani G, Garrett ES, Irizarry RA and Zeger SL. New York, Springer; 2002.
42. Ihaka R, Gentleman R: **R: A language for data analysis and graphics.** *Journal of Computational and Graphical Statistics* 1996, **5**:299-314.
43. **Gene Expression Omnibus** [<http://www.ncbi.nlm.nih.gov/geo/>]
44. Smyth GK: **Linear models and empirical Bayes methods for assessing differential expression in microarray experiments.** *Statistical Applications in Genetics and Molecular Biology* 2004, **3**:3.
45. Lönstedt I, Speed TP: **Replicated microarray data.** *Statistica Sinica* 2002, **12**:31-46.
46. Schweighoffer T, Tanaka Y, Tidswell M, Erle DJ, Horgan KJ, Luce GE, Lazarovits AI, Buck D, Shaw S: **Selective expression of integrin**

alpha 4 beta 7 on a subset of human CD4+ memory T cells with Hallmarks of gut-trophism. *J Immunol* 1993, **151**:717-729.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

