

Minireview

MicroRNA Pharmacoepigenetics: Posttranscriptional Regulation Mechanisms behind Variable Drug Disposition and Strategy to Develop More Effective Therapy

Ai-Ming Yu, Ye Tian, Mei-Juan Tu, Pui Yan Ho, and Joseph L. Jilek

Department of Biochemistry & Molecular Medicine, University of California Davis School of Medicine, Sacramento, California

Received September 30, 2015; accepted November 12, 2015

ABSTRACT

Knowledge of drug absorption, distribution, metabolism, and excretion (ADME) or pharmacokinetics properties is essential for drug development and safe use of medicine. Varied or altered ADME may lead to a loss of efficacy or adverse drug effects. Understanding the causes of variations in drug disposition and response has proven critical for the practice of personalized or precision medicine. The rise of noncoding microRNA (miRNA) pharmacoepigenetics and pharmacoepigenomics has come with accumulating evidence supporting the role of miRNAs in the modulation of ADME gene expression and then drug disposition and response. In this article,

we review the advances in miRNA pharmacoepigenetics including the mechanistic actions of miRNAs in the modulation of Phase I and II drug-metabolizing enzymes, efflux and uptake transporters, and xenobiotic receptors or transcription factors after briefly introducing the characteristics of miRNA-mediated posttranscriptional gene regulation. Consequently, miRNAs may have significant influence on drug disposition and response. Therefore, research on miRNA pharmacoepigenetics shall not only improve mechanistic understanding of variations in pharmacotherapy but also provide novel insights into developing more effective therapeutic strategies.

Introduction

The utility of a drug depends on its efficacy and safety profiles. Upon entering the body, the drug is subjected to absorption, distribution, metabolism, and excretion (ADME) processes before acting on its molecular target to exert pharmacological or toxicological effects. Change in ADME may lead to variable levels of drug for target binding and consequently have significant impact on drug efficacy and safety profiles, which could cause a reduction/loss of pharmacological effects or adverse events (Lu, 1998; Haga et al., 2006; Giacomini et al., 2010; Yu and Pan, 2012). Therefore, research on ADME processes and causes of variation is essential for developing better drugs and ensuring the safe use of approved medications.

ADME processes are mechanistically controlled by drug-metabolizing enzymes and transporters expressed in various tissues including small intestine, liver, and kidney. Drug-metabolizing enzymes

consist of Phase I [e.g., cytochrome P450 (CYP or P450)] and Phase II [e.g., uridine 5'-diphospho-glucuronosyltransferase (UGT)] enzymes that are able to convert the drug to a more hydrophilic and polar metabolite and determine hepatic drug clearance. Transporters including ATP binding cassette (ABC) and solute carrier (SLC) proteins mediate the transport (e.g., efflux or uptake) of many drugs and may have significant effects on drug absorption, distribution, and excretion processes. Therefore, change in drug-metabolizing enzyme and transporter gene expression or protein activity would ultimately alter ADME or pharmacokinetics properties and subsequently affect therapeutic outcomes.

Many mechanisms behind variable ADME have been discovered, which may help to develop more rational and improved therapeutics. For instance, genetic variations can have significant impact on the expression or function of drug-metabolizing enzymes (e.g., CYP2D6 and UGT1A1) and transporters (e.g., ABCB1) and consequently alter drug disposition and response. Therefore, doses may be adjusted or an alternative drug may be prescribed for patients with particular high-risk genotypes or phenotypes, namely personalized or precision

A.-M. Yu was supported by the National Institutes of Health National Cancer Institute [Grant 1U01CA17531].

dx.doi.org/10.1124/dmd.115.067470

ABBREVIATIONS: ABC, ATP-binding cassette; ADME, absorption, distribution, metabolism, and excretion; AHR, aryl hydrocarbon receptor; BCRP or ABCG2, breast cancer resistance protein; CAR or NR1I3, constitutive androstane receptor; CDDP, cisplatin; CYP or P450, cytochrome P450; DGCR8, DiGeorge syndrome critical region 8; ER α or NR3A1, estrogen receptor alpha; GST, glutathione S-transferase; HCPT, hydroxycamptothecin; HNF4 α or NR2A1, hepatocyte nuclear factor 4 alpha; MDR, multidrug resistance; miR or miRNA, microRNA; miRISC, miRNA-induced silencing complex; MRE, miRNA response element; MRP or ABCC, multidrug resistance-associated protein; ncRNA, noncoding RNA; NQO, NAD(P)H:quinone oxidoreductase; NR, nuclear receptor; Nrf2 or NFE2L2, nuclear factor (erythroid-derived 2)-like 2; PEPT, peptide transporter; P-gp or ABCB1 or MDR1, P-glycoprotein; PPAR α or NR1C1, peroxisome proliferator-activated receptor alpha; pre-miRNA, precursor miRNA; pri-miRNA, primary miRNA; PXR or NR1I2, pregnane X receptor; RISC, miRNA-induced silencing complex; RXR α or NR2B1, 9-cis retinoic acid receptor alpha; SLC, solute carrier; SULT, sulfotransferase; TRBP, transactivation-response RNA-binding protein; UGT, uridine 5'-diphospho-glucuronosyltransferase; 3'UTR, 3'-untranslated region; NR1I1, vitamin D receptor.

medicine, to achieve the desired efficacy and prevent adverse effects. Furthermore, drug-metabolizing enzyme and transporter gene expression is regulated by nuclear receptors [NRs; e.g., pregnane X receptor (PXR or NR1I2)] and transcription factors, and modulated through signal transduction, posttranslational modification, membrane trafficking and subcellular organization pathways (for reviews, see Correia and Liao, 2007; Morgan, 2009; Gu and Manautou, 2010; Klaassen and Aleksunes, 2010; Tolson and Wang, 2010)]. Activation or suppression of such regulatory factors or pathways would cause significant change in enzyme/transporter levels and activities and lead to multidrug resistance (MDR), loss of efficacy, or adverse drug effects. Knowledge of these mechanisms has proven helpful for the prediction and prevention of possible toxicity risks and the development of more effective and safer treatments.

Increased research on pharmacoeigenetics and pharmacoeigenomics has demonstrated the role of epigenetic factors in controlling ADME gene expression, in particular, by noncoding microRNAs (miRNAs or miR), DNA methylation proteins, and histone modification proteins (see recent reviews, Ivanov et al., 2012; Ingelman-Sundberg et al., 2013; Zhong and Leeder, 2013). A number of studies have shown that methylation of cytosine-phosphate-guanine sites located in the promoter regions of ADME genes or acetylation of histones may alter ADME gene expression in cells. In addition, there is growing evidence that miRNAs may modulate cellular ADME processes through posttranscriptional regulation of ADME gene expression (Fig. 1). In this review, we briefly introduce the general characteristics of noncoding miRNAs and focus on the discussion of miRNA-controlled epigenetic regulation of drug metabolism and disposition.

MicroRNAs in Posttranscriptional Gene Regulation

MicroRNAs are short, genome-derived, noncoding RNAs (ncRNAs) that exhibit cell specific expression profiles during development and in response to xenobiotics or other environmental factors. Since the discovery of the first miRNA *lin-4* in *Caenorhabditis elegans* in 1993, more than 1800 miRNA sequences have been identified in humans and predicted to regulate thousands of protein-coding genes essentially modulating all cellular processes.




Fig. 1. Role of miRNAs in the regulation of drug metabolism and transport. MicroRNAs may modulate cellular drug metabolism and transport capacity through targeting of drug-metabolizing enzymes, transporters, xenobiotic receptors, and/or other regulatory factors whereas the biogenesis of miRNAs may be regulated by xenobiotic receptors or other regulatory factors.

Biogenesis of miRNAs. Mature miRNAs are produced within cells via canonical or alternative biogenesis pathways. The canonical pathway is dependent upon both DiGeorge syndrome critical region 8 (Drosha-DGCR8) and Dicer-TRBP (transactivation-response RNA-binding protein) complexes (Fig. 2), whereas the noncanonical pathway is not (Yang and Lai, 2011). Canonical biogenesis starts from the transcription of miRNA-coding sequences by RNA polymerase II to long primary miRNA (pri-miRNA) transcripts. Drosha-DGCR8 then cleaves pri-miRNAs to shorter hairpin precursor miRNAs (pre-miRNAs) within the nucleus. After being transported to the cytoplasm by Exportin-5, pre-miRNAs are cleaved by the Dicer-TRBP complex into double-stranded miRNA duplexes, which upon unwinding produce single-stranded mature miRNAs (Fig. 2). By contrast, noncanonical miRNAs are generated from short hairpin introns (mirtrons) by splicing machinery and lariat-debranching enzyme (Curtis et al., 2012) or derived from particular hairpin ncRNA species (e.g., small nucleolar RNAs or transfer RNAs) by other RNases (Cole et al., 2009; Babiarz et al., 2011), rather than Drosha-DGCR8 and Dicer-TRBP.

Mechanisms of miRNA-Controlled Posttranscriptional Gene Expression. Once loaded into the argonaute 2-containing miRNA-induced silencing complexes (miRISCs) in the cytoplasm, miRNAs act on target transcripts via complementary Watson-Crick base pairing to corresponding miRNA response elements (MREs; Fig. 2), which are usually present within the 3'-untranslated regions (3'UTRs) of target genes. Upon binding to MRE sites, miRNAs reduce protein outcome from target transcripts, which may involve translational repression and/or mRNA deadenylation and decay mechanisms (Fig. 2). Although there had been some confusion about the contribution and timing of miRNA-induced translational inhibition and mRNA degradation effects, two recent *in vivo* studies clearly demonstrated that miRNAs largely inhibit translation (e.g., translation initiation) before causing mRNA decay (Bazzini et al., 2012; Djuranovic et al., 2012), emphasizing the importance of evaluating protein abundance rather than only examining mRNA levels for understanding miRNA-controlled posttranscriptional gene regulation.

Identification and Evaluation of microRNA Target Genes. Based on the general principles of miRNA-target interactions, such as the strength of Watson-Crick complementarity, accessibility of MRE sites, and evolutionary conservation, a number of computational algorithms have been established for *in silico* prediction of putative MRE sites and targeted genes (for reviews, see Yu, 2007, 2009). Although computational prediction is helpful for the identification of possible miRNA targets for further investigation, there are unsurprisingly high levels of false positives, whereas other true targets are overlooked. In any case, biologic experiments are essential for the identification and verification of miRNA targets as well as delineation of the significance of miRNA-controlled gene regulation.

One experimental approach to screen miRNA targets is to examine the transcriptome of cells with lost or gained miRNA expression/function. Given the translational repression mechanism of miRNA-controlled gene regulation, many true miRNA targets may likely be overlooked by this method. Furthermore, other transcripts might be presented as false-positive targets that were actually altered as a result of transcriptional regulation or signal transduction caused by true miRNA targets. Another biologic approach is to directly pull down transcripts using the miRNA of interest or isolate argonaute 2 or miRISC complex-associated transcripts through immunoprecipitation and thus analyze those isolated mRNAs (and miRNAs) using various techniques including deep sequencing. This "gold standard" method relies largely on the stringency of experimental conditions, specificity of antibody for immunoprecipitation, and selectivity of other reagents. Given the fact that gene expression within live cells is very dynamic, experimental




Fig. 2. The canonical pathway of miRNA biogenesis and the mechanisms of miRNA controlled posttranscriptional gene regulation. The miRNA coding sequences are transcribed by RNA polymerase II to pri-miRNAs in the nucleus, which are then processed by the Drosha-DGCR8 complex to pre-miRNAs. Exportin 5 transports pre-miRNAs into cytoplasm, where pre-miRNAs are further processed by the Dicer-TRBP complex to double-stranded miRNAs. The miRNA duplexes are subsequently unwound, and the single-stranded mature miRNAs are loaded into argonaute 2-containing miRISCs. As discussed in the main text, some miRNAs may be generated through a number of different noncanonical pathways. By pairing to the MREs that usually occur within the 3'UTRs of target mRNAs, miRNAs mediate translational repression or mRNA degradation, which both may lead to lower levels of protein expression.

results may differ with experimental timing. Indeed this method presents many possible “targets”; however, this cannot be confirmed by other assays and the actual effects on protein outcome remain unknown. For this reason, proteomic approaches have become increasingly popular for the identification of possible miRNA targets in cells after the perturbation of miRNA expression or function (Vinther et al., 2006; Baek et al., 2008; Selbach et al., 2008; Leivonen et al., 2009a; Yang et al., 2009). Although proteomic approaches are able to directly define the effects of miRNAs on protein outcome and provide insight into miRNA targets, it is still unknown whether those altered proteins are direct or downstream effects of miRNA-controlled gene regulation.

Many methodologies have been developed for the validation of computationally predicted or experimentally identified miRNA target genes to delineate the significance of specific miRNAs in the control of cellular processes. Approaches include the evaluation of miRNA-MRE interactions (e.g., luciferase reporter assays with wild-type and mutant MRE sites, RNA electrophoresis mobility shift assay, etc.), assessment of relationship between miRNA and mRNA target expression (protein and/or mRNA levels), and examination of the impact of miRNA on target gene mRNA and protein expression as well as biologic functions. While one study (e.g., reporter assay) may offer some helpful information, the use of a combination of approaches is necessary to provide a more comprehensive understanding of miRNA-governed regulation and biologic significance. Furthermore, these studies often involve the use of both loss- and gain-of-miRNA-function strategies as well as in vitro and in vivo model systems (Yu, 2009; Yu and Pan, 2012). Gain-of-function approaches are commonly used in a system with minimal or no expression of miRNA, whereas loss-of-function approaches may be used in a model with excessive or regular levels of miRNA expression. Specifically, miRNA expression/function may be perturbed in cells or animals with miRNA mimics, precursors, antagomirs/inhibitors, sponges, and siRNAs as well as corresponding expression plasmids/viral vectors. In addition, some genetically modified animal models (Lu et al., 2007; Thai et al., 2007; Xiao

et al., 2007; Han et al., 2015) have been developed and used for the examination of specific miRNA targets and functions at the whole organism level. Through the actions of target genes, miRNAs are thus able to control many cellular processes, including ADME, which may subsequently alter drug efficacy and safety profiles (Yu, 2009; Ivanov et al., 2012; Yu and Pan, 2012; Ingelman-Sundberg et al., 2013; Yokoi and Nakajima, 2013; Zhong and Leeder, 2013).

MicroRNAs Modulate Cellular Drug Metabolism Capacity through the Regulation of Phase I and II Enzyme Expression. In an effort to address the gap in understanding posttranscriptional gene regulation mechanisms underlying variable drug metabolism, many studies have demonstrated that some *CYP* genes may be regulated by miRNAs in human cell line model systems (Table 1). Among them, a well-conserved miRNA miR-27b was shown to directly act on multiple CYPs including CYP1B1 and CYP3A4 (Tsuchiya et al., 2006; Pan et al., 2009a; Li et al., 2014b). Luciferase reporter assays were first employed to define the interactions of miR-27b with corresponding MRE sites within the 3'UTRs of CYP1B1 in MCF-7 and CYP3A4 in HEK-293 cells, which were predicted by bioinformatic algorithms. Other biologic experiments were conducted to elucidate the effects of miR-27b on CYP1B1 and CYP3A4 mRNA and protein expression as well as drug-metabolizing capacity (e.g., midazolam 1'-hydroxylation by endogenous CYP3A4) in various cell lines. The inverse relationship between miR-27b and CYP1B1 protein expression levels in human breast cancerous tissues (Tsuchiya et al., 2006) further demonstrated the significance of miR-27b-controlled regulation underlying variable CYP1B1 expression in cancerous tissues. Furthermore, the existence of miRNA-controlled posttranscriptional regulation mechanisms may help to understand developmental and sexual differences in *CYP* gene expression in humans and animals (for review, see Zhong and Leeder, 2013). Indeed the increase of hepatic miR-27b and murine mmu-miR-298 levels in mice after puberty (Pan et al., 2009a) may provide insight into the decreased or lost expression of CYP3A4 found in *CYP3A4*-transgenic mice (Granvil et al., 2003; Yu et al., 2005; Felmler et al., 2008). It is also noteworthy that miR-27b/a was

also shown to act on a number of ABC transporters (e.g., ABCB1; Table 2), xenobiotic receptors and transcription factors, e.g., vitamin D receptor (*NR1H1*); Table 3 (see the following sections), suggesting that miR-27 may be an important miRNA related to cellular defense against xenobiotics.

There are also many other experimentally evaluated *CYP* gene regulation by miRNAs (Table 1), including *CYP1A1* by miR-892a (Choi et al., 2012), *CYP2C8* by miR-103 and miR-107 (Zhang et al., 2012), *CYP2C9* by miR-128-3p and miR-130b (Rieger et al., 2015; Yu et al., 2015a), *CYP2E1* by miR-378 and miR-570 (Mohri et al., 2010; Nakano et al., 2015), *CYP2J2* by let-7b (Chen et al., 2012a), and *CYP24A1* by miR-125b (Komagata et al., 2009). The effects of miR-130b on endogenous *CYP2C9* enzymatic activity (e.g., tolbutamide hydroxylation) in HepaRG cells were demonstrated by selective liquid chromatography-coupled tandem mass spectrometry assay (Rieger et al., 2015). Reporter assays further confirmed the direct action of miR-130b on *CYP2C9* 3'UTR. Interestingly, miR-130b also reduced the activities of all other measured CYPs by more than 30%, which was associated with significantly lower mRNA levels of constitutive androstane receptor (*CAR/NR1H3*), farnesoid X receptor, *CYP1A1*, *CYP1A2*, *CYP2A6*, *CYP2C8*, *CYP2C9*, *CYP2C19*, and glutathione *S*-transferase alpha 2 (*GSTA2*). Because miR-130b levels are highly elevated in cholestatic liver and during inflammation (Rieger et al., 2013), these findings should advance molecular understanding of altered drug metabolism under pathophysiologic conditions.

Some Phase II drug-metabolizing enzymes, e.g., *UGT1A1* and sulfotransferase 1A1 (*SULT1A1*), are posttranscriptionally regulated by specific miRNAs (Table 1). Computational analysis identified a putative miR-491-3p MRE site within the shared 3'UTR common to all *UGT1A* genes, which encode *UGT1A* enzymes responsible for the addition of glucuronosyl groups from uridine diphosphate glucuronic acids onto drugs to form glucuronide metabolites. This miR-491-3p MRE was first verified using a luciferase reporter assay and mutagenesis

studies (Dluzen et al., 2014). Furthermore, gain and loss of miR-491-3p function in Huh-7 cells led to a significant suppression and elevation of *UGT1A1* mRNA levels and raloxifene glucuronidation activities, respectively. In addition, there was an inverse correlation between miR-491-3p and *UGT1A3* as well as *UGT1A6* mRNA levels among a set of normal human liver specimens, indicating that miR-491-3p may be an important epigenetic regulator in the control of hepatic *UGT1A* expression. In another study, *SULT1A1*, the most abundant hepatic *SULT* involved in the sulfation of various drugs, was revealed to be targeted by miR-631 in an allele specific manner (Yu et al., 2010), suggesting the importance of miR-631 pharmacoeigenetics among populations with particular *SULT1A1* genotypes.

MicroRNAs Control Cellular Drug Transport Capacity via the Regulation of Transporter Expression. Transporters mediate the translocation of many drugs (and endogenous compounds) across the cell membrane and consequently influence ADME processes as well as pharmacokinetic and dynamic properties. Some ABC transporters [e.g., P-glycoprotein (P-gp, *MDR1* or *ABCB1*), breast cancer resistance protein (*BCRP* or *ABCG2*), and multidrug resistance-associated proteins (*MRPs* or *ABCCs*)] mainly function as efflux transporters and overexpression of such efflux transporter in *MDR* or diseased cells may confer *MDR* to chemotherapy and antibiotics. By contrast, many *SLC* family proteins [e.g., organic cation transporters, organic anion transporters, and peptide transporters (*PEPTs*)] may act as uptake transporters. With the understanding of pharmacoeigenetics, there is increasing evidence to support a critical role for miRNAs in posttranscriptional regulation of efflux and uptake transporter genes (Table 2), which may subsequently modulate intracellular drug accumulation and drug response.

P-gp/*MDR1/ABCB1* was first shown to be regulated by miR-27a and miR-451 (Kovalchuk et al., 2008; Zhu et al., 2008). Compared with drug-sensitive parental cell lines, there was an upregulation of *ABCB1* as well as dysregulation of a number of miRNAs including miR-451

TABLE 1
MicroRNAs are shown to modulate the expression of Phase I and II drug-metabolizing enzymes

Enzyme	miRNA	Experimental Methods	References
<i>CYP1A1</i>	miR-892a	Reporter assay, mRNA and protein expression, and functional assay	Choi et al., 2012
<i>CYP1B1</i>	miR-27b	Correlation, reporter assay, protein expression, and functional assay	Tsuchiya et al., 2006
<i>CYP2C8</i>	miR-103, miR-107	Protein and mRNA expression Reporter assay, mRNA and protein expression, and correlation analysis	Chuturgoon et al., 2014 Zhang et al., 2012
<i>CYP2C9</i>	miR-128-3p	RNA electrophoresis mobility shift assay, reporter assay, mRNA & protein expression and correlation	Yu et al., 2015a
<i>CYP2C19</i>	miR-130b miR-103 & miR-107 miR-103 & miR-107 miR-29a-3p	Reporter assay, protein expression, and function Protein expression Protein expression RNA electrophoresis mobility shift assay, mRNA & protein expression, correlation	Rieger et al., 2015 Zhang et al., 2012 Zhang et al., 2012 Yu et al., 2015c
<i>CYP2E1</i>	miR-378 miR-570	Reporter assay, mRNA & protein expression, correlation, and functional assay Reporter assay, mRNA & protein expression, and correlation study	Mohri et al., 2010 Nakano et al., 2015
<i>CYP2J2</i> <i>CYP3A4</i>	let-7b miR-27b	Reporter assay, protein expression, and functional assay Reporter assay, mRNA and protein expression, and functional assay	Chen et al., 2012a Pan et al., 2009a; Li et al., 2014b
<i>CYP24A1</i>	miR-34a miR-577, -1, -532-3p, -627 miR-125b	Correlation Correlation, reporter assay Reporter assay, mRNA & protein expression, and functional assay	Lamba et al., 2014 Wei et al., 2014b Komagata et al., 2009
<i>UGT1A1/1A3/1A6</i> <i>UGT2B15</i> & <i>2B17</i> <i>GSTπ</i> <i>SULT1A1</i>	miR-491-3p miR-376c miR-130b miR-631	Reporter assay, mRNA expression, correlation and/or activity Reporter assay, mRNA expression and enzymatic activity mRNA and protein expression Reporter assay, mRNA and protein expression	Dluzen et al., 2014 Wijayakumara et al., 2015 Zong et al., 2014 Yu et al., 2010

TABLE 2
MicroRNAs are revealed to control the expression of efflux and uptake drug transporters

Transporter	miRNA	Experimental Methods	References
ABCB1/P-gp	let-7g	mRNA and protein expression, functional assay, and correlation	Boyerinas et al., 2012
	miR-9	Protein expression and functions (increase)	Munoz et al., 2013
	miR-19a/b	mRNA and protein expression, functional assay (signal transduction)	Wang et al., 2013
	miR-21	Protein expression and functions (increase)	Xie et al., 2013; Dong et al., 2015
	miR-27a	mRNA and protein expression, and functional assay (increase)	Zhu et al., 2008; Li et al., 2010c
		Reporter assay, mRNA/protein expression, and function (increase; promoter?)	Zhang et al., 2010
		Reporter assay, protein expression, and function	Feng et al., 2011
		mRNA/protein expression, and function (decrease; via Frizzled Class Receptor 7/ β -catenin)	Chen et al., 2013
	miR-129-5p	Reporter assay, protein expression, and function	Wu et al., 2014
	miR-130a	mRNA and protein expression, and function (increase)	Yang et al., 2012
	miR-130b	mRNA and protein expression, and function (increase)	Zong et al., 2014
	miR-137	Protein expression (probably via Y-box binding protein-1)	Zhu et al., 2013
	miR-138	mRNA and protein expression and function (indirect effects?)	Zhao et al., 2010
	miR-145	Reporter assay, protein expression and function	Ikemura et al., 2013; Zhu et al., 2014
	miR-200c	mRNA and protein expression, and function	Chen et al., 2012c
		mRNA and protein expression (via Polycomb complex protein BMI-1)	Liu et al., 2012
		Reporter assay, mRNA/protein expression, and functional assay	Yang et al., 2013
	miR-223	Reporter assay, mRNA/protein expression, and function (increase; promoter?)	Hong et al., 2010
miR-296	Reporter assay, protein expression and function	Bao et al., 2012	
miR-298	Reporter assay, protein expression, and function	Feng et al., 2011	
miR-331-5p	Reporter assay, protein expression, and function	Xu et al., 2013	
miR-381 & -495	Reporter assay, mRNA and protein expression, and function	Kovalchuk et al., 2008	
miR-451	mRNA and protein expression, and functional assay (increase)	Zhu et al., 2008	
	mRNA and protein expression, and function	Bitarte et al., 2011	
miR-508-5p	Reporter assay, mRNA and protein expression, and functional assays	Shang et al., 2014	
ABCB9	miR-31	Reporter assay; mRNA expression; protein expression and function	Dong et al., 2014
ABCG2/BCRP	miR-142-3p	Reporter assay, mRNA and protein expression, and functional assays	Shen et al., 2013
	miR-145	Reporter assay, mRNA and protein expression, and functional assays	Shi et al., 2014c
	miR-181a	Reporter assay, mRNA and protein expression, and functional assays	Jiao et al., 2013
	miR-200c	mRNA and protein expression (via Polycomb complex protein BMI-1)	Liu et al., 2012
	miR-212	Reporter assay, protein expression	Turrini et al., 2012
	miR-328	Reporter assay, mRNA and protein expression, and functional assays	Pan et al., 2009g; Li et al., 2011; Xu et al., 2012c
	miR-487a	Reporter assay, mRNA and protein expression, and functional assays	Ma et al., 2013
	miR-519c	Reporter assay, mRNA and protein expression, and functional assays	To et al., 2008, 2009; Li et al., 2011
ABCC1/MRP1	miR-520h	Reporter assay, mRNA and protein expression	Liao et al., 2008; Wang et al., 2010; Li et al., 2011
	miR-7	Reporter assay, mRNA and protein expression, and association study	Liu et al., 2015
	miR-133a	Reporter assay, mRNA and protein expression, and functional assays	Ma et al., 2015
	miR-134	mRNA and protein expression	Guo et al., 2010
	miR-326	Reporter assay, mRNA and protein expression, and functional assays	Liang et al., 2010; Ma et al., 2015
miR-1291	Reporter assay, mRNA and protein expression, and functional assays	Pan et al., 2013; Li et al., 2015	
ABCC2/MRP2	let-7c	Reporter assay, mRNA and protein expression, and functional assay	Zhan et al., 2013
	miR-297	Reporter assay, mRNA and protein expression, and functional assay	Xu et al., 2012a
	miR-379	Reporter assay, mRNA and protein expression, and functional assays	Haenisch et al., 2011; Werk et al., 2014
ABCC4/MRP4	miR-124a & miR-506	Reporter assays, mRNA and protein expression, functional assay, and correlation study	Markova and Kroetz, 2014
ABCC5/MRP5	miR-128	Reporter assay, protein expression, and functional assay	Zhu et al., 2011
			Li et al., 2014a
ABCE1	miR-129-5p	Reporter assay, mRNA & protein expression, and function	Wu et al., 2014
	miR-203	Reporter assay and protein expression	Furuta et al., 2010

(continued)

TABLE 2—Continued

Transporter	miRNA	Experimental Methods	References
SLC15A1/PEPT1	miR-92b	Reporter assay, mRNA and protein expression, and functional assays	Dalmaso et al., 2011
	miR-193-3p	Correlation study, mRNA and protein expression, and functional assays	Dai et al., 2015
SLC16A1/MCT1	miR-124	Reporter assay, protein expression and function	Li et al., 2009
	miR-29a/b & -124	Reporter assay, mRNA and protein expression	Pullen et al., 2011

within drug-resistant cells. The actions of miR-451 in the regulation of ABCB1 were then demonstrated by computational analysis, luciferase reporter assay, and Western blot studies. Furthermore, treatment with miR-451 sensitized the ABCB1-overexpressing breast cancer cells to doxorubicin, suggesting that miRNAs may be employed to improve chemotherapy or combat MDR. Later, the effectiveness of combined use of miR-451 and an existing anticancer drug (e.g., irinotecan) in treating stem-like colon cancer cells was also demonstrated, which also involved targeting of the ABCB1 efflux transporter (Bitarte et al., 2011). However, some studies (Zhu et al., 2008; Li et al., 2010c; Zhang

et al., 2010) showed an upregulation of ABCB1 by miR-27a, whereas others (Chen et al., 2013) reported a downregulation. Among documented ABCB1 regulatory miRNAs (Table 2), miR-508-5p was shown to reverse MDR not only in gastric cancer cells in vitro but also in xenograft tumor mouse models in vivo (Shang et al., 2014), supporting the strategy to develop miRNA-based combination therapy (Fig. 3).

BCRP/ABCG2, another important efflux transporter in ADME and MDR, is also regulated by a number of miRNAs via direct or indirect mechanisms (Table 2). The MDR of S1 colon carcinoma cell line is

TABLE 3
MicroRNAs are shown to regulate the expression of xenobiotic receptors or transcription factors

Nuclear receptor	miRNA	Experimental Methods	References
AHR NR1I1/	miR-124	Reporter assay, mRNA and protein expression, association study	Huang et al., 2011
	miR-27b	Reporter assay, mRNA and protein expression	Pan et al., 2009a
	miR-125b	Reporter assay, mRNA and protein expression	Mohri et al., 2009
	miR-346	Reporter assay, protein expression	Chen et al., 2014b
	miR-148a	Reporter assay, mRNA and protein expression	Takagi et al., 2008
	miR-137	Association, reporter assay, mRNA/protein expression, function	Takwi et al., 2014)
	miR-10b	Reporter assay, mRNA and protein expression	Zheng et al., 2010
	miR-506	Reporter assay, mRNA and protein expression, and function	Tong et al., 2011
	miR-21	Reporter assay, mRNA and protein expression, and function	Zhou et al., 2011
	miR-21 & -27b	Reporter assay, mRNA/protein expression, function, association	Kida et al., 2011
NR1H3/LXR α	miR-141	Reporter assay, mRNA and protein expression	Hu et al., 2012
	miR-613	Reporter assay, mRNA/protein expression, function	Ou et al., 2011
	miR-1/-206	Reporter assay, mRNA/protein expression, function	Zhong et al., 2013; Vinod et al., 2014
NR2A1/HNF4 α	miR-21	Luciferase reporter assay	Wirsing et al., 2011
	miR-24	Reporter assay, mRNA/protein expression, function	Takagi et al., 2010
	miR-34/-449	Reporter assay, mRNA/protein expression, function	Takagi et al., 2010; Wirsing et al., 2011; Ramamoorthy et al., 2012; Wang and Burke, 2013
	miR-27a/b	Reporter assay, mRNA & protein expression	Ji et al., 2009
	miR-128-2	Reporter assay, mRNA and protein expression	Adlakha et al., 2013
NR3A1/ER α	miR-34a	Reporter assay, mRNA and protein expression, and association	Oda et al., 2014
	miR-221/-222	Reporter assay, mRNA/protein expression, association and function	Zhao et al., 2008; Guttilla et al., 2012; Wei et al., 2014a
	Let-7	Reporter assay, mRNA/protein expression, and function study	Zhao et al., 2011
	miR-18a	Reporter assay, protein expression, association and function study	Liu et al., 2009
	miR-18a/b, -19b, -20b, -193b, -206	Reporter assay, protein expression	Castellano et al., 2009; Leivonen et al., 2009b
	miR-145	Reporter assay, mRNA and protein expression, and function study	Spizzo et al., 2010
	miR-22	Reporter assay, protein expression, and function study	Pandey and Picard, 2009; Xiong et al., 2010
	miR-206	Reporter assay, protein expression, and function study	Adams et al., 2007; Kondo et al., 2008
	miR-27a	via zinc finger and BTB domain containing 10 (ZBTB10)	Li et al., 2010a
	miR-219-5p	Reporter assay, mRNA/protein expression, association and function	Huang et al., 2015
	miR-135b	Reporter assay, mRNA and protein expression	Aakula et al., 2015
	miR-144	Reporter assay, mRNA and protein expression, and function study	Sangokoya et al., 2010
	miR-28	Reporter assay, mRNA and protein expression	Yang et al., 2011
	miR-200a	(Activation) via Kelch-like ECH-associated protein 1 (Keap1)	Eades et al., 2011; Yang et al., 2014
	miR-153, -27a, -142-5p, -144	Reporter assay, mRNA and protein expression, and function study	Narasimhan et al., 2012, 2014
miR-93	Protein expression, and function study	Singh et al., 2013	
miR-507, -634, -450, -129-5p	Reporter assay, mRNA and protein expression, and function study	Yamamoto et al., 2014	
miR-34a	mRNA and protein expression	Huang et al., 2014	
miR-340	Reporter assay, mRNA and protein expression, and function study	Shi et al., 2014a	
miR-141	via Kelch-like ECH-associated protein 1 (Keap1)	Shi et al., 2015	

attributable to the overexpression of ABCG2. miR-519c was found to suppress ABCG2 expression in drug-sensitive parental S1 colon cancer cells via targeting of a distal miR-519c MRE site, whereas drug-resistant S1 colon cancer cells were found to consist of truncated ABCG2 transcripts lacking the distal miR-519c MRE site; together this provides a mechanistic understanding of ABCG2-mediated MDR in these cells (To et al., 2008, 2009). miR-328 was also revealed to regulate ABCG2 and consequently improve the chemosensitivity of ABCG2-overexpressing drug-resistant cells (Pan et al., 2009g). Another proximal MRE site for miR-519c (and miR-520h) were further identified and validated through luciferase report assay and mutagenesis (Li et al., 2011). In addition, the efficiency of miR-328, -519c, and -520h in the regulation of ABCG2 in human breast cancer cell lines was compared side by side (Li et al., 2011). Studies using gain- and loss-of-miRNA function approaches revealed that miR-519c and miR-328 were more effective than miR-520h in the modulation of ABCG2 protein outcome in MCF-7 cells (Li et al., 2011). As a result, both miR-519c and -328 had significant impact on intracellular accumulation of mitoxantrone, an ABCG2 substrate (Li et al., 2011). Therefore, understanding miRNA-controlled regulation not only advances the role of miRNAs in MDR but also offers clue to develop more effective therapeutic strategy, e.g., to use miRNAs to improve chemosensitivity (Fig. 3).

Many MRP/ABCC transporters are regulated by particular miRNAs (Table 2). Compared with wild-type MCF-7 cells, overexpression of ABCC1 in drug-resistant MCF-7 cells was associated with an altered miRNA expression profile, in which miR-326 was downregulated over 12-fold (Liang et al., 2010). Further studies demonstrated that miR-326 directly acted on the 3'UTR of ABCC1, reduced ABCC1 mRNA and protein expression, and sensitized drug-resistant MCF-7 cells to

chemotherapy (Liang et al., 2010). Other ABCC1 regulatory miRNAs evaluated by biologic experiments include miR-134 and miR-7 in small cell lung cancer cells (Guo et al., 2010; Liu et al., 2015), miR-1291 in pancreatic cancer cells (Pan et al., 2013), and miR-133a in hepatocarcinoma cells (Ma et al., 2015). Likewise, restoring the expression or function of such ABCC1 regulatory miRNAs increased intracellular drug accumulation and chemosensitivity, supporting the feasibility of miRNA-based strategies to improve therapy (Fig. 3). In addition, some other miRNAs might modulate the expression of ABCC1 and other transporters and be implicated in the progression of hepatocarcinoma tissues (Borel et al., 2012).

The reduction of ABCC2 protein expression by rifampicin in HepG2 cells was associated with an increased ABCC2 mRNA level and altered miRNA expression profile (Haenisch et al., 2011). Among these altered miRNAs, miR-379 was elevated and might directly target ABCC2 mRNA. Indeed pre-miR-379 showed a dose-dependent suppression of ABCC2 protein expression in HepG2 cells, and miR-379 antagomir was able to rescue rifampicin-reduced ABCC2 protein expression, demonstrating a mechanistic role for miR-379 in rifampicin-altered ABCC2 protein expression at the posttranscriptional level (Haenisch et al., 2011). Another study showed that miR-297 was able to modulate ABCC2 expression in drug-resistant HCT116 cells and thus increased the efficacy of anticancer drugs (Xu et al., 2012a).

By using multiple approaches, a very recent study demonstrated that miR-124a and miR-506 was effective in modulating the expression of ABCC4 in HEK293T/17 cells, likely through a translation inhibition mechanism (Markova and Kroetz, 2014). The reduction of ABCC4 protein outcome by miR-124a and miR-506 was translated into a significantly lower level of efflux of ABCC4 substrate drug, bimaneglutathione. Although there are six major ABCC4 3'UTR haplotypes at




Fig. 3. Coadministration of miRNAs with chemotherapeutic agents may be more effective to control tumor progression. There is growing evidence supporting the feasibility of combination therapy. MicroRNAs downregulate the expression of efflux transporters in tumor cells, resulting in a higher level of intracellular drug accumulation and thus an improved efficacy (Zhu et al., 2008; Pan et al., 2009g, 2013; Liang et al., 2010; Shang et al., 2014; Li et al., 2015). In addition, miRNAs may directly target (proto-) oncogenes and thus additively or synergistically suppress tumor growth (Deng et al., 2014; Zhao et al., 2015).

a frequency greater than 5% among Caucasians, African Americans, and Asians, none of the inferred polymorphisms were located within or in proximity to the MRE sites for miR-124a and miR-506 and thus showed any significant effects on miR-124a- and miR-506-mediated regulation of ABCC4. In addition, miR-124a and miR-506 levels were found to be inversely correlated with ABCC4 levels in 26 human kidney samples (Markova and Kroetz, 2014), illustrating the clinical significance of miRNA-controlled regulation of ABC transporters.

There are also some reports on posttranscriptional gene regulation of SLC transporters (Table 2). The increase of human peptide transporter 1 (PEPT1/SLC15A1) protein expression during differentiation of intestinal Caco2-BBE cells was associated with three- to four-fold decrease of mature miR-92b levels (Dalmaso et al., 2011). miR-92b was then shown to directly target the 3'UTR of SLC15A1, reduce SLC15A1 mRNA and protein expression, and subsequently inhibit cellular uptake of a tripeptide (lysine-proline-valine) substrate and suppress bacterial peptide-induced proinflammatory responses in cells (Dalmaso et al., 2011). Another uptake transporter, monocarboxylate transporter 1, was also found to be posttranscriptionally regulated by several miRNAs including miR-124, miR-29a, and miR-29b (Li et al., 2009; Pullen et al., 2011).

MicroRNAs Regulate the Expression of Xenobiotic Receptors or Transcription Factors to Modulate Drug Metabolism and Disposition. Transcriptional regulation of drug-metabolizing enzyme and transporter gene expression is primarily controlled by xenobiotic receptors or transcription factors [e.g., aryl hydrocarbon receptor (AHR), PXR/NR1I2, CAR/NR1I4, peroxisome proliferator-activated receptor α (PPAR α or NR1C1), /NR1I1, nuclear factor (erythroid-derived 2)-like 2 (Nrf2 or NFE2L2), estrogen receptor alpha (ER α or NR3A1), and 9-cis retinoic acid receptor alpha (RXR α or NR2B1), etc.]. Upon activation by corresponding ligands, xenobiotic receptors and transcription factors bind to target promoter elements and lead to the upregulation of drug-metabolizing enzymes (e.g., CYP3A4 and UGT1A1) and transporters (e.g., ABCB1 and ABCG2). Therefore, miRNAs may alter the expression of enzymes and transporters through posttranscriptional regulation of xenobiotic receptor or transcription factor genes (Table 3) and thus modulate drug disposition and response (Fig. 1).

The actions of miR-27b/a illustrate miRNA-NR-enzyme/transporter signaling in the modulation of cellular drug metabolism and disposition. In addition to direct targeting of CYP3A4 (Pan et al., 2009a), CYP1B1 (Tsuchiya et al., 2006), and ABCB1 (Zhu et al., 2008), miR-27b/a was able to act on a number of xenobiotic receptors and transcription factors including NR1I1/ (Pan et al., 2009a), NR1C1/PPAR α (Kida et al., 2011), ER α /NR3A1 (Li et al., 2010a), NFE2L2/Nrf2 (Narasimhan et al., 2012), and RXR α /NR2B1 (Ji et al., 2009). Therefore, alteration of enzyme/transporter protein outcome by miR-27b/a could be an accumulative effect from direct targeting of the enzyme/transporter and "indirect" targeting of regulatory factors including xenobiotic receptors and transcription factors (Yu, 2009; Yu and Pan, 2012). It should be noted that the miRNA-NR-enzyme/transporter signaling inevitably affects the mRNA levels of an enzyme/transporter, which might complicate the interpretation of miRNA regulatory mechanisms if overlooked.

PXR/NR1I2 controls transcriptional expression of many important ADME genes including CYP3A4, UGT1A, and ABCB1. miR-148a was shown to target NR1I2 and then influence CYP3A4 expression (Takagi et al., 2008). The miR-148a MRE site within NR1I2 3'UTR was defined by luciferase reporter assay. Further studies demonstrated that miR-148a precursor and antagomir were able to decrease and elevate, respectively, NR1I2 protein levels in HepG2 cells. In addition,

transfection of LS-180 cells with miR-148a precursor attenuated NR1I2-dependent induction of CYP3A4 mRNA expression, suggesting a miR-148a-NR1I2-CYP3A4 axis.

CAR/NR1I3 is another important xenobiotic receptor that transcriptionally regulates the expression of various enzymes (e.g., CYP2B6 and UGT1A1) and transporters (e.g., ABCB1 and ABCG2). NR1I3 was found to be directly targeted by miR-137, which not only suppressed the protein expression of ABCB1 and CYP2B6 in drug-resistant cells but also sensitized the cells to doxorubicin in vitro as well as neuroblastoma xenografts to doxorubicin in vivo (Takwi et al., 2014). A separate study also demonstrated the actions of miR-137 in the regulation of murine Car/Nr1i3 (Chen et al., 2014a). Although miR-137 and mouse Nr1i3 expression levels were inversely related in mice during development, knocking down miR-137 led to higher rate of bilirubin clearance, validating the miR-137-Nr1i3-Ugt1a1 link in a whole body system. In addition, ectopic expression of miR-137 suppressed Nr1i3 protein levels in primary mouse hepatocytes, which could be attenuated by miR-137 antagomir.

Upon ligand binding, transcription factor AHR activates the expression of many enzymes including CYP1A1, CYP1A2, and CYP1B1. There is also evidence suggesting that CYP enzymes might be regulated by miRNA through AHR pathways (Table 3). For instance, in vivo administration of locked nucleic acid-modified antisense miR-29 antagomir to mice altered the expression of many hepatic proteins, including the upregulation of Ahr, Cyp1a1, Cyp2b10, Cyp3a11, and Cyp4a12 (Kurtz et al., 2015). A conserved miR-29 MRE site within the 3'UTR of human AHR and mouse Ahr was then validated by luciferase reporter assay, demonstrating that AHR is a direct target of miR-29. Another study showed that AHR was modulated by miR-124 in neuroblastoma SK-N-SH cells (Huang et al., 2011). In addition, miR-24 was shown to regulate the AHR dimer partner, aryl hydrocarbon receptor nuclear translocator, and thus interfere with expression and induction of CYP1A1 in Huh-7 and HepG2 cells (Oda et al., 2012).

Hepatocyte nuclear factor 4 α (HNF4 α /NR2A1) is an important nuclear transcription factor that binds to target DNA segment as a homodimer and serves as a master regulator of ADME gene expression. For instance, HNF4 α is essential for PXR- and CAR-mediated transcriptional activation of CYP3A4 (Tirona et al., 2003). Some miRNAs were shown to directly regulate HNF4 α expression and thus affect cellular ADME capacity (Table 3). Both endogenous and exogenous HNF4 α protein outcome was reduced by miR-24 and miR-34a in cell line models, which likely acted through direct targeting of corresponding MRE sites (Takagi et al., 2010). As a result, introduction of miR-24 and miR-34a into HepG2 cells was able to suppress the mRNA levels of several HNF4 α target genes, including CYP7A1, CYP8B1, and CYP27A1, which was reasoned to affect bile acid homeostasis (Takagi et al., 2010).

PPAR α /NR1C1 is a major transcription factor in the control of lipid metabolism, which may be posttranscriptionally regulated by a number of miRNAs (Table 3). Both miR-21 and miR-27b were shown to modulate PPAR α protein expression levels but not mRNA levels in Huh7 cells (Kida et al., 2011). The miR-21 MRE site within PPAR α 3'UTR was further assessed by luciferase reporter assay, and the significance of miR-21 in the control of PPAR α protein expression was manifested by the inverse correlation between miR-21 and PPAR α protein levels in human liver samples (Kida et al., 2011). In addition, miR-506 was revealed to directly act on the 3'UTR of PPAR α and modulate the protein levels of PPAR α in cells (Tong et al., 2011). Although miR-506 levels were sharply higher in hydroxycamptothecin (HCPT)-resistant human colon cancer SW1116/HCPT cells, miR-506 antagomir was effective to sensitize the cells to HCPT

(Tong et al., 2011), supporting the development of miRNA-based strategy to combat MDR and improve cancer therapy (Fig. 3).

The redox-sensitive transcription factor Nrf2 plays a pivotal role in cellular response to oxidative and electrophilic stress via the induction of many drug-metabolizing enzymes [e.g., NAD(P)H:quinone oxidoreductase 1 and 2 (NQO1 and NQO2), glucose-6-phosphate dehydrogenase, GSTs, and UGTs] and efflux transporters (e.g., ABCG2, ABCC3, and ABCC4). Although inherently cytoprotective, recent evidence has demonstrated that Nrf2 overexpression may actually impart MDR by regulation of xenobiotic disposition in certain cancer cells (for review, see Leinonen et al., 2014). Additionally, evidence for miRNA crosstalk within Nrf2 signaling pathway has been reported (Table 3). Among them, miR-144 was demonstrated to act directly on Nrf2 3'UTR and then modulate the expression of Nrf2 downstream targets and cellular glutathione levels (Sangkoyka et al., 2010). Another study (Shi et al., 2014a) revealed that, compared with parental HepG2 cells, miR-340 was downregulated and Nrf2 was upregulated in cisplatin (CDDP)-resistant HepG2/CDDP cells. The effects of miR-340 mimics on Nrf2 and target gene (e.g., NQO1) expression were also demonstrated, which could be attenuated by miR-340 antagomir. Furthermore, administration of miR-340 mimics improved the sensitivity of HepG2/CDDP cells to cisplatin, whereas miR-340 antagomir could restore drug resistance, suggesting an important role for miR-340 in the development of cisplatin resistance in HepG2/CDDP cells and supporting the use of miRNA-based strategy to combat MDR (Fig. 3).

Conclusions

The focus on miRNA pharmacoeigenetics and pharmacoeigenomics is rising because a growing number of studies demonstrate the roles of noncoding miRNAs in the control of drug disposition and response. Although one miRNA (e.g., miR-27b) can regulate various ADME genes (e.g., *CYP3A4*) via direct and/or indirect targeting, the same ADME gene (e.g., *ABCG2*) may be modulated by multiple miRNAs (e.g., miR-328, miR-519c, and miR-520h) (for review, see Yu and Pan, 2012). As a result, miRNA signaling may have significant effects on ADME processes and drug response. Nevertheless, current understandings are mainly obtained from cell model systems and ex vivo tissue samples. Better physiologic models and in vivo studies are highly warranted to provide more direct evidence to support the importance of miRNAs in the regulation of pharmacokinetics and dynamics.

The knowledge of miRNA pharmacoeigenetics/epigenomics not only provides novel insights into interindividual variability in drug disposition and response (Yu, 2009; Yu and Pan, 2012; Ingelman-Sundberg et al., 2013; Zhong and Leeder, 2013) but also offers new clues to developing more effective treatments (Fig. 3). MicroRNAs are able to suppress the expression of some efflux transporters and enzymes and lead to a greater degree of intracellular drug accumulation, resulting in higher efficacy (Zhu et al., 2008; Pan et al., 2009g; Liang et al., 2010; Pan et al., 2013; Shang et al., 2014; Li et al., 2015). Meanwhile, some miRNAs may directly reduce the protein outcome of pharmacological targets [e.g., (proto)oncogenes] and thus control disease progression (e.g., tumor growth and metastasis) (Deng et al., 2014; Zhao et al., 2015). However, research on miRNA pharmacoeigenetics and development of miRNA-based therapy may be limited by the utilization of synthetic miRNA agents consisting of excessive artificial modifications on the phosphate linkages and/or ribose rings and thus exhibit different physicochemical and biologic properties or toxicities. By contrast, recombinant RNA technology has been successfully employed to cost-effectively produce large quantities of miRNA agents (Li et al., 2014b; Chen et al., 2015; Li et al., 2015; Wang et al., 2015), which are comprised of no or posttranscriptional modifications on the nucleobases

and may better capture the structures, functions, and safety properties of natural RNAs. These natural ncRNA agents are produced in cells and should represent a novel class of miRNA agents for research and development.

Authorship Contributions

Performed data analysis: Yu, Tian, Tu, Ho, and Jilek.

Wrote or contributed to the writing of the manuscript: Yu, Tian, Tu, Ho, and Jilek.

References

- Aakula A, Leivonen SK, Hintsanen P, Aittokallio T, Ceder Y, Børresen-Dale AL, Perälä M, Östling P, and Kallioniemi O (2015) MicroRNA-135b regulates ER α , AR and HIF1AN and affects breast and prostate cancer cell growth. *Mol Oncol* **9**:1287–1300.
- Adams BD, Furneaux H, and White BA (2007) The micro-ribonucleic acid (miRNA) miR-206 targets the human estrogen receptor-alpha (ERalpha) and represses ERalpha messenger RNA and protein expression in breast cancer cell lines. *Mol Endocrinol* **21**:1132–1147.
- Adlakha YK, Khanna S, Singh R, Singh VP, Agrawal A, and Saini N (2013) Pro-apoptotic miRNA-128-2 modulates ABCA1, ABCG1 and RXR α expression and cholesterol homeostasis. *Cell Death Dis* **4**:e780.
- Babiarz JE, Hsu R, Melton C, Thomas M, Ullian EM, and Blalock R (2011) A role for non-canonical microRNAs in the mammalian brain revealed by phenotypic differences in Dgcr8 versus Dicer1 knockouts and small RNA sequencing. *RNA* **17**:1489–1501.
- Baek D, Villén J, Shin C, Camargo FD, Gygi SP, and Bartel DP (2008) The impact of microRNAs on protein output. *Nature* **455**:64–71.
- Bao L, Hazari S, Mehra S, Kaushal D, Moroz K, and Dash S (2012) Increased expression of P-glycoprotein and doxorubicin chemoresistance of metastatic breast cancer is regulated by miR-298. *Am J Pathol* **180**:2490–2503.
- Bazzini AA, Lee MT, and Giraldez AJ (2012) Ribosome profiling shows that miR-430 reduces translation before causing mRNA decay in zebrafish. *Science* **336**:233–237.
- Bitarte N, Bandres E, Boni V, Zarate R, Rodriguez J, Gonzalez-Huarriz M, Lopez I, Javier Sola J, Alonso MM, and Fortes P, et al. (2011) MicroRNA-451 is involved in the self-renewal, tumorigenicity, and chemoresistance of colorectal cancer stem cells. *Stem Cells* **29**:1661–1671.
- Borel F, Han R, Visser A, Petry H, van Deventer SJ, Jansen PL, and Konstantinova P; Réseau Centre de Ressources Biologiques Foie (French Liver Biobanks Network), France (2012) Adenosine triphosphate-binding cassette transporter genes up-regulation in untreated hepatocellular carcinoma is mediated by cellular microRNAs. *Hepatology* **55**:821–832.
- Boyerinas B, Park SM, Murmann AE, Gwin K, Montag AG, Zillhardt M, Hua YJ, Lengyel E, and Peter ME (2012) Let-7 modulates acquired resistance of ovarian cancer to Taxanes via IMP-1-mediated stabilization of multidrug resistance 1. *Int J Cancer* **130**:1787–1797.
- Castellano L, Giamas G, Jacob J, Coombes RC, Lucchesi W, Thiruchelvam P, Barton G, Jiao LR, Wait R, and Waxman J, et al. (2009) The estrogen receptor-alpha-induced microRNA signature regulates itself and its transcriptional response. *Proc Natl Acad Sci USA* **106**:15732–15737.
- Chen F, Chen C, Yang S, Gong W, Wang Y, Cianflone K, Tang J, and Wang DW (2012a) Let-7b inhibits human cancer phenotype by targeting cytochrome P450 epoxigenase 2J2. *PLoS One* **7**:e39197.
- Chen J, Tian W, Cai H, He H, and Deng Y (2012c) Down-regulation of microRNA-200c is associated with drug resistance in human breast cancer. *Med Oncol* **29**:2527–2534.
- Chen QX, Wang WP, Zeng S, Urayama S, and Yu AM (2015) A general approach to high-yield biosynthesis of chimeric RNAs bearing various types of functional small RNAs for broad applications. *Nucleic Acids Res* **43**:3857–3869.
- Chen S, He N, Yu J, Li L, Hu Y, Deng R, Zhong S, and Shen L (2014a) Post-transcriptional regulation by miR-137 underlies the low abundance of CAR and low rate of bilirubin clearance in neonatal mice. *Life Sci* **107**:8–13.
- Chen Y, Du J, Zhang Z, Liu T, Shi Y, Ge X, and Li YC (2014b) MicroRNA-346 mediates tumor necrosis factor α -induced downregulation of gut epithelial vitamin D receptor in inflammatory bowel diseases. *Inflamm Bowel Dis* **20**:1910–1918.
- Chen Z, Ma T, Huang C, Zhang L, Lv X, Xu T, Hu T, and Li J (2013) MiR-27a modulates the MDR1/P-glycoprotein expression by inhibiting FZD7/ β -catenin pathway in hepatocellular carcinoma cells. *Cell Signal* **25**:2693–2701.
- Choi YM, An S, Lee EM, Kim K, Choi SJ, Kim JS, Jiang HH, An IS, and Bae S (2012) CYP1A1 is a target of miR-892a-mediated post-transcriptional repression. *Int J Oncol* **41**:331–336.
- Chaturgoon AA, Phulkdaree A, and Moodley D (2014) Fumonisin B $_1$ modulates expression of human cytochrome P450 1b1 in human hepatoma (HepG2) cells by repressing Mir-27b. *Toxicol Lett* **227**:50–55.
- Cole C, Sobala A, Lu C, Thatcher SR, Bowman A, Brown JW, Green PJ, Barton GJ, and Hutvagner G (2009) Filtering of deep sequencing data reveals the existence of abundant Dicer-dependent small RNAs derived from tRNAs. *RNA* **15**:2147–2160.
- Correia MA and Liao M (2007) Cellular proteolytic systems in P450 degradation: evolutionary conservation from *Saccharomyces cerevisiae* to mammalian liver. *Expert Opin Drug Metab Toxicol* **3**:33–49.
- Curtis HJ, Sibley CR, and Wood MJ (2012) Mirtrons, an emerging class of atypical miRNA. *Wiley Interdiscip Rev RNA* **3**:617–632.
- Dai X, Chen X, Chen Q, Shi L, Liang H, Zhou Z, Liu Q, Pang W, Hou D, and Wang C, et al. (2015) MicroRNA-193a-3p Reduces Intestinal Inflammation in Response to Microbiota via Down-regulation of Colonic PepT1. *J Biol Chem* **290**:16099–16115.
- Dalmasco G, Nguyen HT, Yan Y, Laroui H, Charania MA, Obertone TS, Sitaraman SV, and Merlin D (2011) MicroRNA-92b regulates expression of the oligopeptide transporter PepT1 in intestinal epithelial cells. *Am J Physiol Gastrointest Liver Physiol* **300**:G52–G59.
- Deng X, Cao M, Zhang J, Hu K, Yin Z, Zhou Z, Xiao X, Yang Y, Sheng W, and Wu Y, et al. (2014) Hyaluronic acid-chitosan nanoparticles for co-delivery of MiR-34a and doxorubicin in therapy against triple negative breast cancer. *Biomaterials* **35**:4333–4344.
- Djuranovic S, Nahvi A, and Green R (2012) miRNA-mediated gene silencing by translational repression followed by mRNA deadenylation and decay. *Science* **336**:237–240.

- Dluzen DF, Sun D, Salzberg AC, Jones N, Bushey RT, Robertson GP, and Lazarus P (2014) Regulation of UDP-glucuronosyltransferase 1A1 expression and activity by microRNA 491-3p. *J Pharmacol Exp Ther* **348**:465–477.
- Dong Z, Ren L, Lin L, Li J, Huang Y, and Li J (2015) Effect of microRNA-21 on multidrug resistance reversal in A549/DDP human lung cancer cells. *Mol Med Rep* **11**:682–690.
- Dong Z, Zhong Z, Yang L, Wang S, and Gong Z (2014) MicroRNA-31 inhibits cisplatin-induced apoptosis in non-small cell lung cancer cells by regulating the drug transporter ABCB9. *Cancer Lett* **343**:249–257.
- Eades G, Yang M, Yao Y, Zhang Y, and Zhou Q (2011) miR-200a regulates Nrf2 activation by targeting Keap1 mRNA in breast cancer cells. *J Biol Chem* **286**:40725–40733.
- Felmllee MA, Lon HK, Gonzalez FJ, and Yu AM (2008) Cytochrome P450 expression and regulation in CYP3A4/CYP2D6 double transgenic humanized mice. *Drug Metab Dispos* **36**:435–441.
- Feng DD, Zhang H, Zhang P, Zheng YS, Zhang XJ, Han BW, Luo XQ, Xu L, Zhou H, and Qu LH, et al. (2011) Down-regulated miR-331-5p and miR-27a are associated with chemotherapy resistance and relapse in leukaemia. *J Cell Mol Med* **15**:2164–2175.
- Furuta M, Kozaki KI, Tanaka S, Arii S, Imoto I, and Inazawa J (2010) miR-124 and miR-203 are epigenetically silenced tumor-suppressive microRNAs in hepatocellular carcinoma. *Carcinogenesis* **31**:766–776.
- Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, Dahlin A, Evers R, Fischer V, and Hillgren KM, et al.; International Transporter Consortium (2010) Membrane transporters in drug development. *Nat Rev Drug Discov* **9**:215–236.
- Granvil CP, Yu AM, Elizondo G, Akiyama TE, Cheung C, Feigenbaum L, Krausz KW, and Gonzalez FJ (2003) Expression of the human CYP3A4 gene in the small intestine of transgenic mice: in vitro metabolism and pharmacokinetics of midazolam. *Drug Metab Dispos* **31**:548–558.
- Gu X and Manautou JE (2010) Regulation of hepatic ABC transporters by xenobiotics and in disease states. *Drug Metab Rev* **42**:482–538.
- Guo L, Liu Y, Bai Y, Sun Y, Xiao F, and Guo Y (2010) Gene expression profiling of drug-resistant small cell lung cancer cells by combining microRNA and cDNA expression analysis. *Eur J Cancer* **46**:1692–1702.
- Guttilla IK, Phoenix KN, Hong X, Tirnauer JS, Claffey KP, and White BA (2012) Prolonged mammosphere culture of MCF-7 cells induces an EMT and repression of the estrogen receptor by microRNAs. *Breast Cancer Res Treat* **132**:75–85.
- Haenisch S, Laechelt S, Bruckmueller H, Werk A, Noack A, Bruhn O, Remmler C, and Cascorbi I (2011) Down-regulation of ATP-binding cassette C2 protein expression in HepG2 cells after rifampicin treatment is mediated by microRNA-379. *Mol Pharmacol* **80**:314–320.
- Haga SB, Thummel KE, and Burke W (2006) Adding pharmacogenetics information to drug labels: lessons learned. *Pharmacogenomics* **16**:847–854.
- Han YC, Vidigal JA, Mu P, Yao E, Singh I, González AJ, Concepcion CP, Bonetti C, Ogradowski P, and Carver B, et al. (2015) An allelic series of miR-17 ~ 92-mutant mice uncovers functional specialization and cooperation among members of a microRNA polycistron. *Nat Genet* **47**:766–775.
- Hong L, Han Y, Zhang H, Li M, Gong T, Sun L, Wu K, Zhao Q, and Fan D (2010) The prognostic and chemotherapeutic value of miR-296 in esophageal squamous cell carcinoma. *Ann Surg* **251**:1056–1063.
- Hu W, Wang X, Ding X, Li Y, Zhang X, Xie P, Yang J, and Wang S (2012) MicroRNA-141 represses HBV replication by targeting PPARα. *PLoS One* **7**:e34165.
- Huang C, Cai Z, Huang M, Mao C, Zhang Q, Lin Y, Zhang X, Tang B, Chen Y, and Wang X, et al. (2015) miR-219-5p modulates cell growth of papillary thyroid carcinoma by targeting estrogen receptor α. *J Clin Endocrinol Metab* **100**:E204–E213.
- Huang TC, Chang HY, Chen CY, Wu PY, Lee H, Liao YF, Hsu WM, Huang HC, and Juan HF (2011) Silencing of miR-124 induces neuroblastoma SK-N-SH cell differentiation, cell cycle arrest and apoptosis through promoting AHR. *FEBS Lett* **585**:3582–3586.
- Huang X, Gao Y, Qin J, and Lu S (2014) The role of miR-34a in the hepatoprotective effect of hydrogen sulfide on ischemia/reperfusion injury in young and old rats. *PLoS One* **9**:e113305.
- Ikemura K, Yamamoto M, Miyazaki S, Mizutani H, Iwamoto T, and Okuda M (2013) MicroRNA-145 post-transcriptionally regulates the expression and function of P-glycoprotein in intestinal epithelial cells. *Mol Pharmacol* **83**:399–405.
- Ingelman-Sundberg M, Zhong XB, Hankinson O, Beednagar S, Yu AM, Peng L, and Osawa Y (2013) Potential role of epigenetic mechanisms in the regulation of drug metabolism and transport. *Drug Metab Dispos* **41**:1725–1731.
- Ivanov M, Kacevska M, and Ingelman-Sundberg M (2012) Epigenomics and interindividual differences in drug response. *Clin Pharmacol Ther* **92**:727–736.
- Ji J, Zhang J, Huang G, Qian J, Wang X, and Mei S (2009) Over-expressed microRNA-27a and 27b influence fat accumulation and cell proliferation during rat hepatic stellate cell activation. *FEBS Lett* **583**:759–766.
- Jiao X, Zhao L, Ma M, Bai X, He M, Yan Y, Wang Y, Chen Q, Zhao X, and Zhou M, et al. (2013) MiR-181a enhances drug sensitivity in mitoxantone-resistant breast cancer cells by targeting breast cancer resistance protein (BCRP/ABCG2). *Breast Cancer Res Treat* **139**:717–730.
- Kida K, Nakajima M, Mohri T, Oda Y, Takagi S, Fukami T, and Yokoi T (2011) PPARα is regulated by miR-21 and miR-27b in human liver. *Pharm Res* **28**:2467–2476.
- Klaassen CD and Aleksunes LM (2010) Xenobiotic, bile acid, and cholesterol transporters: function and regulation. *Pharmacol Rev* **62**:1–96.
- Komagata S, Nakajima M, Takagi S, Mohri T, Taniya T, and Yokoi T (2009) Human CYP24 catalyzing the inactivation of calcitriol is post-transcriptionally regulated by miR-125b. *Mol Pharmacol* **76**:702–709.
- Kondo N, Toyama T, Sugiura H, Fujii Y, and Yamashita H (2008) miR-206 Expression is down-regulated in estrogen receptor alpha-positive human breast cancer. *Cancer Res* **68**:5004–5008.
- Kovalchuk O, Filkowski J, Meservy J, Ilynskyy Y, Tryndyak VP, Chekhun VF, and Pogribny IP (2008) Involvement of microRNA-451 in resistance of the MCF-7 breast cancer cells to chemotherapeutic drug doxorubicin. *Mol Cancer Ther* **7**:2152–2159.
- Kurtz CL, Fannin EE, Toth CL, Pearson DS, Vickers KC, and Sethupathy P (2015) Inhibition of miR-29 has a significant lipid-lowering benefit through suppression of lipogenic programs in liver. *Sci Rep* **5**:12911.
- Lamba V, Ghodke Y, Guan W, and Tracy TS (2014) microRNA-34a is associated with expression of key hepatic transcription factors and cytochromes P450. *Biochem Biophys Res Commun* **445**:404–411.
- Leinonen HM, Kansanen E, Pöhlönen P, Heinäniemi M, and Levenon AL (2014) Role of the Keap1-Nrf2 pathway in cancer. *Adv Cancer Res* **122**:281–320.
- Leivonen SK, Mäkelä R, Ostling P, Kohonen P, Haapa-Painanen S, Kleivi K, Eerley E, Aakula A, Hellström K, and Sahlborg N, et al. (2009a) Protein lysate microarray analysis to identify microRNAs regulating estrogen receptor signaling in breast cancer cell lines. *Oncogene* **28**:3926–3936.
- Li B, Chen H, Wu N, Zhang WJ, and Shang LX (2014a) Downregulation of miR-128 in ovarian cancer promotes cisplatin resistance. *Int J Gynecol Cancer* **24**:1381–1388.
- Li KK, Pang JC, Ching AK, Wong CK, Kong X, Wang Y, Zhou L, Chen Z, and Ng HK (2009) miR-124 is frequently down-regulated in medulloblastoma and is a negative regulator of SLC16A1. *Hum Pathol* **40**:1234–1243.
- Li MM, Addepalli B, Tu MJ, Chen QX, Wang WP, Limbach PA, LaSalle JM, Zeng S, Huang M, and Yu AM (2015) Chimeric miR-1291 biosynthesized efficiently in *E. coli* is effective to reduce target gene expression in human carcinoma cells and improve chemosensitivity. *Drug Metab Dispos* **43**:1129–1136.
- Li MM, Wang WP, Wu WJ, Huang M, and Yu AM (2014b) Rapid production of novel pre-microRNA agent hsa-mir-27b in *Escherichia coli* using recombinant RNA technology for functional studies in mammalian cells. *Drug Metab Dispos* **42**:1791–1795.
- Li X, Mertens-Talcott SU, Zhang S, Kim K, Ball J, and Safe S (2010a) MicroRNA-27a Indirectly Regulates Estrogen Receptor alpha Expression and Hormone Responsiveness in MCF-7 Breast Cancer Cells. *Endocrinology* **151**:2462–2473.
- Li X, Pan YZ, Seigel GM, Hu ZH, Huang M, and Yu AM (2011) Breast cancer resistance protein BCRP/ABCG2 regulatory microRNAs (hsa-miR-328, -519c and -520h) and their differential expression in stem-like ABCG2+ cancer cells. *Biochem Pharmacol* **81**:783–792.
- Li Z, Hu S, Wang J, Cai J, Xiao L, Yu L, and Wang Z (2010c) MiR-27a modulates MDR1/P-glycoprotein expression by targeting HIPK2 in human ovarian cancer cells. *Gynecol Oncol* **119**:125–130.
- Liang Z, Wu H, Xia J, Li Y, Zhang Y, Huang K, Wagar N, Yoon Y, Cho HT, and Scala S, et al. (2010) Involvement of miR-326 in chemotherapy resistance of breast cancer through modulating expression of multidrug resistance-associated protein 1. *Biochem Pharmacol* **79**:817–824.
- Liao R, Sun J, Zhang L, Lou G, Chen M, Zhou D, Chen Z, and Zhang S (2008) MicroRNAs play a role in the development of human hematopoietic stem cells. *J Cell Biochem* **104**:805–817.
- Liu H, Wu X, Huang J, Peng J, and Guo L (2015) miR-7 modulates chemoresistance of small cell lung cancer by repressing MRP1/ABCC1. *Int J Exp Pathol* **96**:240–247.
- Liu S, Tetzlaff MT, Cui R, and Xu X (2012) miR-200c inhibits melanoma progression and drug resistance through down-regulation of BMI-1. *Am J Pathol* **181**:1823–1835.
- Liu WH, Yeh SH, Lu CC, Yu SL, Chen HY, Lin CY, Chen DS, and Chen PJ (2009) MicroRNA-18a prevents estrogen receptor-alpha expression, promoting proliferation of hepatocellular carcinoma cells. *Gastroenterology* **136**:683–693.
- Lu AY (1998) Drug-metabolism research challenges in the new millennium: individual variability in drug therapy and drug safety. *Drug Metab Dispos* **26**:1217–1222.
- Lu Y, Thomson JM, Wong HY, Hammond SM, and Hogan BL (2007) Transgenic over-expression of the microRNA miR-17-92 cluster promotes proliferation and inhibits differentiation of lung epithelial progenitor cells. *Dev Biol* **310**:442–453.
- Ma J, Wang T, Guo R, Yang X, Yin J, Yu J, Xiang Q, Pan X, Tang H, and Lei X (2015) Involvement of miR-133a and miR-326 in ADM resistance of HepG2 through modulating expression of ABCG1. *J Drug Target* **23**:519–524.
- Ma MT, He M, Wang Y, Jiao XY, Zhao L, Bai XF, Yu ZJ, Wu HZ, Sun ML, and Song ZG, et al. (2013) MiR-487a sensitizes mitoxantone (MX)-resistant breast cancer cells (MCF-7/MX) to MX by targeting breast cancer resistance protein (BCRP/ABCG2). *Cancer Lett* **339**:107–115.
- Markova SM and Kroetz DL (2014) ABCG4 is regulated by microRNA-124a and microRNA-506. *Biochem Pharmacol* **87**:515–522.
- Mohri T, Nakajima M, Fukami T, Takamiya M, Aoki Y, and Yokoi T (2010) Human CYP2E1 is regulated by miR-378. *Biochem Pharmacol* **79**:1045–1052.
- Mohri T, Nakajima M, Takagi S, Komagata S, and Yokoi T (2009) MicroRNA regulates human vitamin D receptor. *Int J Cancer* **125**:1328–1333.
- Morgan ET (2009) Impact of infectious and inflammatory disease on cytochrome P450-mediated drug metabolism and pharmacokinetics. *Clin Pharmacol Ther* **85**:434–438.
- Munoz JL, Bliss SA, Greco SJ, Ramkissoon SH, Ligon KL, and Rameshwar P (2013) Delivery of functional anti-miR-9 by mesenchymal stem cell-derived exosomes to glioblastoma multiforme cells conferred chemosensitivity. *Mol Ther Nucleic Acids* **2**:e126.
- Nakano M, Mohri T, Fukami T, Takamiya M, Aoki Y, McLeod HL, and Nakajima M (2015) SNPs in CYP2E1 3'-UTR affect the regulation of CYP2E1 by miR-570. *Drug Metab Dispos* **43**:1450–1457.
- Narasimhan M, Patel D, Vedpathak D, Rathinam N, Henderson G, and Mahimainathan L (2012) Identification of novel microRNAs in post-transcriptional control of Nrf2 expression and redox homeostasis in neuronal, SH-SY5Y cells. *PLoS One* **7**:e31111.
- Narasimhan M, Riar AK, Rathinam ML, Vedpathak D, Henderson G, and Mahimainathan L (2014) Hydrogen peroxide responsive miR153 targets Nrf2/ARE cytoprotection in paraquat induced dopaminergic neurotoxicity. *Toxicol Lett* **228**:179–191.
- Oda Y, Nakajima M, Mohri T, Takamiya M, Aoki Y, Fukami T, and Yokoi T (2012) Aryl hydrocarbon receptor nuclear translocator in human liver is regulated by miR-24. *Toxicol Appl Pharmacol* **260**:222–231.
- Oda Y, Nakajima M, Tsuneyama K, Takamiya M, Aoki Y, Fukami T, and Yokoi T (2014) Retinoid X receptor α in human liver is regulated by miR-34a. *Biochem Pharmacol* **90**:179–187.
- Ou Z, Wada T, Gramignoli R, Li S, Strom SC, Huang M, and Xie W (2011) MicroRNA hsa-miR-613 targets the human LXRα gene and mediates a feedback loop of LXRα autoregulation. *Mol Endocrinol* **25**:584–596.
- Pan YZ, Gao W, and Yu AM (2009a) MicroRNAs regulate CYP3A4 expression via direct and indirect targeting. *Drug Metab Dispos* **37**:2112–2117.
- Pan YZ, Morris ME, and Yu AM (2009b) MicroRNA-328 negatively regulates the expression of breast cancer resistance protein (BCRP/ABCG2) in human cancer cells. *Mol Pharmacol* **75**:1374–1379.
- Pan YZ, Zhou A, Hu Z, and Yu AM (2013) Small nucleolar RNA-derived microRNA hsa-miR-1291 modulates cellular drug disposition through direct targeting of ABC transporter ABCG1. *Drug Metab Dispos* **41**:1744–1751.
- Pandey DP and Picard D (2009) miR-22 inhibits estrogen signaling by directly targeting the estrogen receptor alpha mRNA. *Mol Cell Biol* **29**:3783–3790.

- Pullen TJ, da Silva Xavier G, Kelsey G, and Rutter GA (2011) miR-29a and miR-29b contribute to pancreatic beta-cell-specific silencing of monocarboxylate transporter 1 (Mct1). *Mol Cell Biol* **31**:3182–3194.
- Ramamoorthy A, Li L, Gaedigk A, Bradford LD, Benson EA, Flockhart DA, and Skaar TC (2012) In silico and in vitro identification of microRNAs that regulate hepatic nuclear factor 4 α expression. *Drug Metab Dispos* **40**:726–733.
- Rieger JK, Klein K, Winter S, and Zanger UM (2013) Expression variability of absorption, distribution, metabolism, excretion-related microRNAs in human liver: influence of nongenetic factors and association with gene expression. *Drug Metab Dispos* **41**:1752–1762.
- Rieger JK, Reutter S, Hofmann U, Schwab M, and Zanger UM (2015) Inflammation-associated microRNA-130b down-regulates cytochrome P450 activities and directly targets CYP2C9. *Drug Metab Dispos* **43**:884–888.
- Sangokoya C, Telen MJ, and Chi JT (2010) microRNA miR-144 modulates oxidative stress tolerance and associates with anemia severity in sickle cell disease. *Blood* **116**:4338–4348.
- Selbach M, Schwanhäusser B, Thierfelder N, Fang Z, Khanin R, and Rajewsky N (2008) Widespread changes in protein synthesis induced by microRNAs. *Nature* **455**:58–63.
- Shang Y, Zhang Z, Liu Z, Feng B, Ren G, Li K, Zhou L, Sun Y, Li M, and Zhou J, et al. (2014) miR-508-5p regulates multidrug resistance of gastric cancer by targeting ABCB1 and ZNRD1. *Oncogene* **33**:3267–3276.
- Shen WW, Zeng Z, Zhu WX, and Fu GH (2013) MiR-142-3p functions as a tumor suppressor by targeting CD133, ABCG2, and Lgr5 in colon cancer cells. *J Mol Med (Berl)* **91**:989–1000.
- Shi L, Chen ZG, Wu LL, Zheng JJ, Yang JR, Chen XF, Chen ZQ, Liu CL, Chi SY, and Zheng JY, et al. (2014a) miR-340 reverses cisplatin resistance of hepatocellular carcinoma cell lines by targeting Nrf2-dependent antioxidant pathway. *Asian Pac J Cancer Prev* **15**:10439–10444.
- Shi L, Wang Z, Sun G, Wan Y, Guo J, and Fu X (2014c) miR-145 inhibits migration and invasion of glioma stem cells by targeting ABCG2. *Neuromolecular Med* **16**:517–528.
- Shi L, Wu L, Chen Z, Yang J, Chen X, Yu F, Zheng F, and Lin X (2015) MiR-141 Activates Nrf2-Dependent Antioxidant Pathway via Down-Regulating the Expression of Keap1 Confering the Resistance of Hepatocellular Carcinoma Cells to 5-Fluorouracil. *Cell Physiol Biochem* **35**:2333–2348.
- Singh B, Ronghe AM, Chatterjee A, Bhat NK, and Bhat HK (2013) MicroRNA-93 regulates NRF2 expression and is associated with breast carcinogenesis. *Carcinogenesis* **34**:1165–1172.
- Spizzo R, Nicoloso MS, Lupini L, Lu Y, Fogarty J, Rossi S, Zagatti B, Fabbri M, Veronese A, and Liu X, et al. (2010) miR-145 targets with TP53 in a death-promoting regulatory loop and targets estrogen receptor- α in human breast cancer cells. *Cell Death Differ* **17**:246–254.
- Takagi S, Nakajima M, Kida K, Yamaura Y, Fukami T, and Yokoi T (2010) MicroRNAs regulate human hepatocyte nuclear factor 4 α , modulating the expression of metabolic enzymes and cell cycle. *J Biol Chem* **285**:4415–4422.
- Takagi S, Nakajima M, Mohri T, and Yokoi T (2008) Post-transcriptional regulation of human pregnane X receptor by micro-RNA affects the expression of cytochrome P450 3A4. *J Biol Chem* **283**:9674–9680.
- Takwi AA, Wang YM, Wu J, Michaelis M, Cinatti J, and Chen T (2014) miR-137 regulates the constitutive androstane receptor and modulates doxorubicin sensitivity in parental and doxorubicin-resistant neuroblastoma cells. *Oncogene* **33**:3717–3729.
- Thai TH, Calado DP, Casola S, Ansel KM, Xiao C, Xue Y, Murphy A, Frendewey D, Valenzuela D, and Kutok JL, et al. (2007) Regulation of the germinal center response by microRNA-155. *Science* **316**:604–608.
- Tirona RG, Lee W, Leake BF, Lan LB, Cline CB, Lamba V, Parviz F, Duncan SA, Inoue Y, and Gonzalez FJ, et al. (2003) The orphan nuclear receptor HNF4 α determines PXR- and CAR-mediated xenobiotic induction of CYP3A4. *Nat Med* **9**:220–224.
- To KK, Robey RW, Knutsen T, Zhan Z, Ried T, and Bates SE (2009) Escape from hsa-miR-519c enables drug-resistant cells to maintain high expression of ABCG2. *Mol Cancer Ther* **8**:2959–2968.
- To KK, Zhan Z, Litman T, and Bates SE (2008) Regulation of ABCG2 expression at the 3' untranslated region of its mRNA through modulation of transcript stability and protein translation by a putative microRNA in the S1 colon cancer cell line. *Mol Cell Biol* **28**:5147–5161.
- Tolson AH and Wang H (2010) Regulation of drug-metabolizing enzymes by xenobiotic receptors: PXR and CAR. *Adv Drug Deliv Rev* **62**:1238–1249.
- Tong JL, Zhang CP, Nie F, Xu XT, Zhu MM, Xiao SD, and Ran ZH (2011) MicroRNA 506 regulates expression of PPAR α in hydroxycamptothecin-resistant human colon cancer cells. *FEBS Lett* **585**:3560–3568.
- Tsuchiya Y, Nakajima M, Takagi S, Taniya T, and Yokoi T (2006) MicroRNA regulates the expression of human cytochrome P450 1B1. *Cancer Res* **66**:9090–9098.
- Turrini E, Haenisch S, Laechelt S, Dieuwok T, Bruhn O, and Cascorbi I (2012) MicroRNA profiling in K-562 cells under imatinib treatment: influence of miR-212 and miR-328 on ABCG2 expression. *Pharmacogenet Genomics* **22**:198–205.
- Vinod M, Chennamsetty I, Colin S, Belloy L, De Paoli F, Schaidt H, Graier WF, Frank S, Kratky D, and Staels B, et al. (2014) miR-206 controls LXR α expression and promotes LXR-mediated cholesterol efflux in macrophages. *Biochim Biophys Acta* **1841**:827–835.
- Vinther J, Hedegaard MM, Gardner PP, Andersen JS, and Arntander P (2006) Identification of mRNA targets with stable isotope labeling by amino acids in cell culture. *Nucleic Acids Res* **34**:e107.
- Wang F, Li T, Zhang B, Li H, Wu Q, Yang L, Nie Y, Wu K, Shi Y, and Fan D (2013) MicroRNA-19a/b regulates multidrug resistance in human gastric cancer cells by targeting PTEN. *Biochem Biophys Res Commun* **434**:688–694.
- Wang F, Xue X, Wei J, An Y, Yao J, Cai H, Wu J, Dai C, Qian Z, and Xu Z, et al. (2010) hsa-miR-520h downregulates ABCG2 in pancreatic cancer cells to inhibit migration, invasion, and side populations. *Br J Cancer* **103**:567–574.
- Wang WP, Ho PY, Chen QX, Addepalli B, Limbach PA, Li MM, Wu WJ, Jilek JL, Qiu JX, and Zhang HJ, et al. (2015) Bioengineering novel chimeric microRNA-34a for prodrug cancer therapy: high-yield expression and purification, and structural and functional characterization. *J Pharmacol Exp Ther* **354**:131–141.
- Wang Z and Burke PA (2013) The role of microRNAs in hepatocyte nuclear factor-4 α expression and transactivation. *Biochim Biophys Acta* **1829**:436–442.
- Wei Y, Lai X, Yu S, Chen S, Ma Y, Zhang Y, Li H, Zhu X, Yao L, and Zhang J (2014a) Exosomal miR-221/222 enhances tamoxifen resistance in recipient ER-positive breast cancer cells. *Breast Cancer Res Treat* **147**:423–431.
- Wei Z, Jiang S, Zhang Y, Wang X, Peng X, Meng C, Liu Y, Wang H, Guo L, and Qin S, et al. (2014b) The effect of microRNAs in the regulation of human CYP3A4: a systematic study using a mathematical model. *Sci Rep* **4**:4283.
- Werk AN, Bruckmueller H, Haenisch S, and Cascorbi I (2014) Genetic variants may play an important role in mRNA-miRNA interaction: evidence for haplotype-dependent down-regulation of ABCG2 (MRP2) by miRNA-379. *Pharmacogenet Genomics* **24**:283–291.
- Wijayakumara DD, Hu DG, Meech R, McKinnon RA, and Mackenzie PI (2015) Regulation of Human UGT2B15 and UGT2B17 by miR-376c in Prostate Cancer Cell Lines. *J Pharmacol Exp Ther* **354**:417–425.
- Wirsing A, Senkel S, Klein-Hitpass L, and Ryffel GU (2011) A systematic analysis of the 3'UTR of HNF4A mRNA reveals an interplay of regulatory elements including miRNA target sites. *PLoS One* **6**:e27438.
- Wu Q, Yang Z, Xia L, Nie Y, Wu K, Shi Y, and Fan D (2014) Methylation of miR-129-5p CpG island modulates multi-drug resistance in gastric cancer by targeting ABC transporters. *Oncotarget* **5**:11552–11563.
- Xiao C, Calado DP, Galler G, Thai TH, Patterson HC, Wang J, Rajewsky N, Bender TP, and Rajewsky K (2007) MiR-150 controls B cell differentiation by targeting the transcription factor c-Myb. *Cell* **131**:146–159.
- Xie Z, Cao L, and Zhang J (2013) miR-21 modulates paclitaxel sensitivity and hypoxia-inducible factor-1 α expression in human ovarian cancer cells. *Oncol Lett* **6**:795–800.
- Xiong J, Yu D, Wei N, Fu H, Cai T, Huang Y, Wu C, Zheng X, Du Q, and Lin D, et al. (2010) An estrogen receptor α suppressor, microRNA-22, is downregulated in estrogen receptor α -positive human breast cancer cell lines and clinical samples. *FEBS J* **277**:1684–1694.
- Xu K, Liang X, Shen K, Cui D, Zheng Y, Xu J, Fan Z, Qiu Y, Li Q, and Ni L, et al. (2012a) miR-297 modulates multidrug resistance in human colorectal carcinoma by down-regulating MRP-2. *Biochem J* **446**:291–300.
- Xu XT, Xu Q, Tong JL, Zhu MM, Nie F, Chen X, Xiao SD, and Ran ZH (2012c) MicroRNA expression profiling identifies miR-328 regulates cancer stem cell-like SP cells in colorectal cancer. *Br J Cancer* **106**:1320–1330.
- Xu Y, Ohms SJ, Li Z, Wang Q, Gong G, Hu Y, Mao Z, Shannon MF, and Fan JY (2013) Changes in the expression of miR-381 and miR-495 are inversely associated with the expression of the MDR1 gene and development of multi-drug resistance. *PLoS One* **8**:e82062.
- Yamamoto S, Inoue J, Kawano T, Kozaki K, Omura K, and Inazawa J (2014) The impact of miRNA-based molecular diagnostics and treatment of NRF2-stabilized tumors. *Mol Cancer Res* **12**:58–68.
- Yang JJ, Tao H, Hu W, Liu LP, Shi KH, Deng ZY, and Li J (2014) MicroRNA-200a controls Nrf2 activation by target Keap1 in hepatic stellate cell proliferation and fibrosis. *Cell Signal* **26**:2381–2389.
- Yang JS and Lai EC (2011) Alternative miRNA biogenesis pathways and the interpretation of core miRNA pathway mutants. *Mol Cell* **43**:892–903.
- Yang L, Li N, Wang H, Xia X, Wang X, and Luo J (2012) Altered microRNA expression in cisplatin-resistant ovarian cancer cells and upregulation of miR-130a associated with MDR1/P-glycoprotein-mediated drug resistance. *Oncol Rep* **28**:592–600.
- Yang M, Yao Y, Eades G, Zhang Y, and Zhou Q (2011) MiR-28 regulates Nrf2 expression through a Keap1-independent mechanism. *Breast Cancer Res Treat* **129**:983–991.
- Yang T, Zheng ZM, Li XN, Li ZF, Wang Y, Geng YF, Bai L, and Zhang XB (2013) MiR-223 modulates multidrug resistance via downregulation of ABCB1 in hepatocellular carcinoma cells. *Exp Biol Med (Maywood)* **238**:1024–1032.
- Yang Y, Chaerkady R, Beer MA, Mendell JT, and Pandey A (2009) Identification of miR-21 targets in breast cancer cells using a quantitative proteomic approach. *Proteomics* **9**:1374–1384.
- Yokoi T and Nakajima M (2013) microRNAs as mediators of drug toxicity. *Annu Rev Pharmacol Toxicol* **53**:377–400.
- Yu AM (2007) Small interfering RNA in drug metabolism and transport. *Curr Drug Metab* **8**:700–708.
- Yu AM (2009) Role of microRNAs in the regulation of drug metabolism and disposition. *Expert Opin Drug Metab Toxicol* **5**:1513–1528.
- Yu AM, Fukumachi K, Krausz KW, Cheung C, and Gonzalez FJ (2005) Potential role for human cytochrome P450 3A4 in estradiol homeostasis. *Endocrinology* **146**:2911–2919.
- Yu AM and Pan YZ (2012) Noncoding microRNAs: Small RNAs play a big role in regulation of ADME? *Acta Pharm Sin B* **2**:93–101.
- Yu D, Green B, Marrone A, Guo Y, Kadlubar S, Lin D, Fuscoe J, Pogribny I, and Ning B (2015a) Suppression of CYP2C9 by microRNA hsa-miR-128-3p in human liver cells and association with hepatocellular carcinoma. *Sci Rep* **5**:8534.
- Yu D, Green B, Tolleson WH, Jin Y, Mei N, Guo Y, Deng H, Pogribny I, and Ning B (2015c) MicroRNA hsa-miR-29a-3p modulates CYP2C19 in human liver cells. *Biochem Pharmacol* **98**:215–223.
- Yu X, Dhakal IB, Beggs M, Edavana VK, Williams S, Zhang X, Mercer K, Ning B, Lang NP, and Kadlubar FF, et al. (2010) Functional genetic variants in the 3'-untranslated region of sulfotransferase isoform 1A1 (SULT1A1) and their effect on enzymatic activity. *Toxicol Sci* **118**:391–403.
- Zhan M, Qu Q, Wang G, and Zhou H (2013) Let-7c sensitizes acquired cisplatin-resistant A549 cells by targeting ABCG2 and Bcl-XL. *Pharmacazie* **68**:955–961.
- Zhang H, Li M, Han Y, Hong L, Gong T, Sun L, and Zheng X (2010) Down-regulation of miR-27a might reverse multidrug resistance of esophageal squamous cell carcinoma. *Dig Dis Sci* **55**:2545–2551.
- Zhang S-Y, Surapreddi S, Coulter S, Ferguson SS, and Goldstein JA (2012) Human CYP2C8 is post-transcriptionally regulated by microRNAs 103 and 107 in human liver. *Mol Pharmacol* **82**:529–540.
- Zhao JJ, Lin J, Yang H, Kong W, He L, Ma X, Coppola D, and Cheng JQ (2008) MicroRNA-221/222 negatively regulates estrogen receptor α and is associated with tamoxifen resistance in breast cancer. *J Biol Chem* **283**:31079–31086.
- Zhao X, Yang L, Hu J, and Ruan J (2010) miR-138 might reverse multidrug resistance of leukemia cells. *Leuk Res* **34**:1078–1082.
- Zhao Y, Deng C, Lu W, Xiao J, Ma D, Guo M, Recker RR, Gatalica Z, Wang Z, and Xiao GG (2011) let-7 microRNAs induce tamoxifen sensitivity by downregulation of estrogen receptor α signaling in breast cancer. *Mol Med* **17**:1233–1241.
- Zhao Y, Tu MJ, Yu YF, Wang WP, Chen QX, Qiu JX, Yu AX, and Yu AM (2015). Combination therapy with bioengineered miR-34a prodrug and doxorubicin synergistically suppresses osteosarcoma growth. *Biochem Pharmacol*, **98**:602–613.
- Zheng L, Lv GC, Sheng J, and Yang YD (2010) Effect of miRNA-10b in regulating cellular steatosis level by targeting PPAR- α expression, a novel mechanism for the pathogenesis of NAFLD. *J Gastroenterol Hepatol* **25**:156–163.

- Zhong D, Huang G, Zhang Y, Zeng Y, Xu Z, Zhao Y, He X, and He F (2013) MicroRNA-1 and microRNA-206 suppress LXR α -induced lipogenesis in hepatocytes. *Cell Signal* **25**: 1429–1437.
- Zhong XB and Leeder JS (2013) Epigenetic regulation of ADME-related genes: focus on drug metabolism and transport. *Drug Metab Dispos* **41**:1721–1724.
- Zhou J, Wang KC, Wu W, Subramaniam S, Shyy JY, Chiu JJ, Li JY, and Chien S (2011) MicroRNA-21 targets peroxisome proliferators-activated receptor- α in an autoregulatory loop to modulate flow-induced endothelial inflammation. *Proc Natl Acad Sci USA* **108**: 10355–10360.
- Zhu H, Wu H, Liu X, Evans BR, Medina DJ, Liu CG, and Yang JM (2008) Role of MicroRNA miR-27a and miR-451 in the regulation of MDR1/P-glycoprotein expression in human cancer cells. *Biochem Pharmacol* **76**:582–588.
- Zhu X, Li Y, Shen H, Li H, Long L, Hui L, and Xu W (2013) miR-137 restoration sensitizes multidrug-resistant MCF-7/ADM cells to anticancer agents by targeting YB-1. *Acta Biochim Biophys Sin (Shanghai)* **45**:80–86.
- Zhu X, Li Y, Xie C, Yin X, Liu Y, Cao Y, Fang Y, Lin X, Xu Y, and Xu W, et al. (2014) miR-145 sensitizes ovarian cancer cells to paclitaxel by targeting Sp1 and Cdk6. *Int J Cancer* **135**: 1286–1296.
- Zhu Y, Yu F, Jiao Y, Feng J, Tang W, Yao H, Gong C, Chen J, Su F, and Zhang Y, et al. (2011) Reduced miR-128 in breast tumor-initiating cells induces chemotherapeutic resistance via Bmi-1 and ABCC5. *Clin Cancer Res* **17**:7105–7115.
- Zong C, Wang J, and Shi TM (2014) MicroRNA 130b enhances drug resistance in human ovarian cancer cells. *Tumour Biol* **35**:12151–12156.

Address correspondence to: Ai-Ming Yu, Department of Biochemistry & Molecular Medicine, UC Davis School of Medicine, 2700 Stockton Blvd., Suite 2132, Sacramento, CA 95817. E-mail: aimyu@ucdavis.edu.
