



HHS Public Access

Author manuscript

Cancer Epidemiol Biomarkers Prev. Author manuscript; available in PMC 2017 February 01.

Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2016 February ; 25(2): 417–424. doi:
10.1158/1055-9965.EPI-15-0594.

Dietary intake of one-carbon metabolism-related nutrients and pancreatic cancer risk: The Singapore Chinese Health Study

Joyce Y. Huang^{1,2}, Lesley M. Butler^{1,2}, Renwei Wang¹, Aizhen Jin³, Woon-Puay Koh^{4,5}, and Jian-Min Yuan^{1,2}

¹Cancer Control and Population Sciences, The University of Pittsburgh Cancer Institute

²Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

³National Registry of Diseases Office, Health Promotion Board, Singapore

⁴Duke-NUS Graduate Medical School Singapore, Singapore

⁵Saw Swee Hock School of Public Health, National University of Singapore, Singapore

Abstract

Background—Nutrients involved in one-carbon metabolism are hypothesized to protect against pancreatic cancer development.

Methods—The Singapore Chinese Health Study database was used to prospectively examine the association between intake of one-carbon metabolism-related nutrients and pancreatic cancer risk. Between 1993 and 1998, 63,257 men and women aged 45–74 years were enrolled into the cohort. The daily intakes of the following one-carbon metabolism-related nutrients were assessed at enrollment using a 165-item food frequency questionnaire: betaine, choline, folate, and vitamins B₂, B₆, and B₁₂. Multivariable hazard ratios (HRs) and 95% confidence intervals (CIs) for pancreatic cancer risk associated with dietary intakes of one-carbon metabolism-related nutrients were calculated.

Results—As of December 2013, 271 incident pancreatic cancer cases were identified during an average of 16.3 years of follow-up. Higher intake of vitamin B₆ and choline were associated with statistically significant decreases in the risk of developing pancreatic cancer. Compared with the lowest quartile, HRs (95% CIs) for the highest quartiles of vitamin B₆ and choline were 0.52 (0.36, 0.74) (*P* trend = 0.001) and 0.67 (0.48, 0.93) (*P* trend = 0.04), respectively. There were no clear associations between the other one-carbon metabolism-related nutrients and pancreatic cancer risk.

Conclusion—Our study suggests that higher intake of vitamin B₆ and choline may lower the risk of pancreatic cancer.

Corresponding Author: Dr. Lesley M. Butler, University of Pittsburgh Cancer Institute, UPMC Cancer Pavilion, 5150 Centre Avenue – Suite 4C, Pittsburgh, PA 15232. Tel: 412-623-3386; Fax: 412-864-7838; butlerl3@upmc.edu.

Conflicts of interest: none to declare

Impact—Our prospective findings are consistent with the *in vivo* evidence for protective roles of vitamin B₆ and choline on pancreatic cancer development.

Keywords

cohort studies; diet; one-carbon metabolism; pancreatic cancer; vitamin B₆

Introduction

Pancreatic cancer is among the most deadly malignancies in the world. In the U.S., pancreatic cancer is the fourth leading cause of cancer-related death in both men and women (1). In 2014, it is estimated that 20,170 men and 19,420 women died from pancreatic cancer (1). The median survival of pancreatic cancer is only three to six months, due in part to the lack of effective treatments (2). Cigarette smoking and obesity are the only established modifiable risk factors for pancreatic cancer (3, 4). The identification of novel primary prevention targets is a viable approach for reducing the burden of pancreatic cancer.

One-carbon metabolism is a set of interconnected pathways that supply methyl groups for DNA synthesis, repair, and methylation (5). Adequate DNA methylation maintains chromosome stability and prevents gene disruption (6). Studies using a global methylation profiling approach showed that a substantial number of genes were aberrantly methylated not just in advanced pancreatic cancers (7), but also in precancerous lesions (8), indicating a potential time window when chemoprevention agents that target DNA methylation pathways could have a beneficial impact. Diet is a major source for key substrates and cofactors involved in one-carbon metabolism, such as vitamin B₆, choline, and folate (9). Vitamin B₆ is a cofactor for multiple enzymes in the one-carbon metabolism pathway, including serine hydroxymethyltransferase for nucleotide synthesis (10) and remethylation of homocysteine (11), and cystathionine β-synthase and cystathionine γ-lyase for generation of glutathione (12). In rodents, diets deficient in methyl donors (i.e., choline, methionine, and/or folate) resulted in global hypomethylation (13–15) and increased incidence of neoplastic lesions induced by carcinogens in the pancreas (16–18). Therefore, low dietary intake of these nutrients may interfere with the normal function of one-carbon metabolism pathways (19, 20), and thus increase the risk of developing pancreatic cancer.

A majority of epidemiologic studies that have evaluated one-carbon metabolism-related nutrients in relation to pancreatic cancer risk have focused on the potential role of folate. A recent pooled analysis of 14 prospective cohort studies conducted mostly in U.S. populations, reported that dietary folate was not associated with pancreatic cancer risk (21). Similarly, there was no trend with increasing blood folate levels and pancreatic cancer risk in the European Investigation of Cancer (EPIC) cohort study (22) or in a pooled analysis of U.S. cohorts (23).

Few prospective studies of pancreatic cancer have considered the potential role of one-carbon metabolism-related nutrients other than folate. Dietary methionine was strongly associated with lower risk of pancreatic cancer in a Swedish study (24), but not in a cohort of Finnish male smokers (25). While neither cohort supported dietary vitamin B₆ as a possible protective factor for pancreatic cancer (24, 25), a recent report showed a

statistically inverse association for plasma pyridoxal phosphate (PLP), the bioactive form of vitamin B₆, in European women (22). A large population-based case-control study in the U.S. reported a positive association and trend with increasing pancreatic cancer risk for dietary vitamin B₁₂ (26). There have been no epidemiologic studies of other one-carbon related nutrients, such as choline or betaine and pancreatic cancer risk.

Due to the involvement of multiple nutrients and the complexity of one-carbon metabolism pathways, a comprehensive assessment of the nutrients involved and their associations with pancreatic cancer risk is needed. Therefore, it is worthwhile to investigate these associations in a prospective cohort with well-characterized dietary intake. We analyzed data from a prospective cohort of Singapore Chinese to test the hypothesis that higher intake of one-carbon metabolism-related nutrients are inversely associated with the risk of developing pancreatic cancer.

Materials and Methods

Study design and population

The design of Singapore Chinese Health Study has been previously described in detail (27). Briefly, between 1993 and 1998, 63,257 Chinese men and women between ages of 45–74 years living in Singapore were enrolled into the cohort study. The cohort was drawn from residents of government-built housing estates (86% of all Singaporeans during the recruitment time period). Study subjects were restricted to two major dialect groups of Chinese in Singapore (Hokkien and Cantonese). The study was approved by the Institutional Review Boards of the National University of Singapore and the University of Pittsburgh.

Assessment of one-carbon metabolism-related nutrients

At the time of recruitment, each participant completed an in-person interview with a structured questionnaire asking for information on demographics, uses of tobacco and alcohol, medical history, and physical activity. To assess usual dietary intake, a 165-item semi-quantitative food frequency questionnaire (FFQ) was used. The FFQ was developed for the target study population and validated using a series of 24-hour recall interviews and repeated administration of the FFQ in a subpopulation of the cohort (28).

Average daily intake of one-carbon metabolism-related nutrients was calculated by multiplying the usual frequency and portion size of each food item by the nutrient content using the food composition values from the Singapore Food Composition Database. The original Singapore Food Composition Database contained the levels of 98 nutritive/nonnutritive food components, including folate, and vitamins B₂, B₆ and B₁₂, per 100 g of edible food for each food item included in the FFQ (28). The nutrient content information of betaine (29–34), choline (30, 32, 34), and methionine (32, 34) has recently become available and has been added into the Singapore Food Composition Database based on the data published by US Department of Agriculture and the University of Minnesota's Nutrition Coordinating Center Food and Nutrient Database.

Our team has previously reported the correlation coefficients for total calorie intake and selected nutrients (including folate) from the FFQ versus the 24-hour recalls information

collected from 858 randomly chosen cohort members for the purpose of validating the FFQ (28). The correlation coefficient ranged from 0.31 to 0.53 for total calories and 0.50 to 0.69 for calorie-adjusted folate by residual method (35) across sex and dialect groups (Cantonese and Hokkien) (28).

Ascertainment of pancreatic cancer cases

Incident pancreatic cancer cases (ICD-Oncology codes, 2nd edition, C25) were identified through record linkage with the databases of the nationwide Singapore Cancer Registry and the Singapore Registry of Births and Deaths. The national cancer registry has been in place since 1968 and has been shown to be comprehensive in recording cancer cases among the entire population (36). As of December 31, 2013, only 57 participants from this cohort were known to be lost to follow-up because of migration out of Singapore (n=30) or for other reasons (n=27). Among those under follow-up, 276 cohort members who were free of cancer at baseline developed pancreatic cancer.

Statistical analysis

Person-years of follow-up time was counted from the date of baseline interview to the diagnosis of pancreatic cancer, lost to follow-up, death of any cause, or December 31, 2013, whichever occurred first. Among the cohort participants, 1,936 had a history of cancer before enrollment, and thus were excluded from the present analysis. Another 459 men and 564 women were excluded due to extreme values of total calorie intake (men: <700 or >3700 kcal/day, women: <600 or >3000 kcal/day). In total, 60,298 subjects including 271 pancreatic cancer cases were included in the analysis.

Nutrient variables were presented as the values adjusted for total calorie intake by using the residual method (35). Cox proportional hazards regression method (37) was used to estimate the hazard ratios (HRs) and 95% confidence intervals (CI) for the associations between one-carbon metabolism-related nutrients and pancreatic cancer risk. Study subjects were grouped into quartiles based on the distribution of nutrient intake in the entire cohort. The nutrients were coded as ordinal values (1, 2, 3, 4) of quartile variables to assess the linear trends of the nutrient-pancreatic cancer association. We did not identify any violation of the proportional hazard assumption.

On the basis of previous analyses in the cohort, we controlled for age at baseline (years), sex, year of baseline interview (1993–1995, 1996–1998), dialect group (Cantonese, Hokkien), and the level of education (no formal schooling, primary school, and secondary school or above) in all multivariable models. We further adjusted for body mass index (BMI) (<18.5, 18.5–21.4, 21.5–24.4, 24.5–27.4, 27.5 kg/m²) (38, 39), smoking status (never smokers, former smokers, current smokers), alcohol drinking (nondrinker, drinker), weekly use of vitamin/mineral supplement (no, yes), self-reported physician diagnosed diabetes (no, yes), any weekly physical activity (no, yes), and total daily caloric intake (tertiles). Covariates were included in the final multivariable regression models, because including the variable in the base model resulted in change of 10% or more in the hazard ratio for pancreatic cancer and at least one of the one-carbon metabolism-related nutrients,

or the variable had been previously reported to be associated with pancreatic cancer in the present study population or other populations.

We further performed stratified analyses by sex. To rule out potential effects of subclinical pancreatic cancer on intake of one-carbon metabolism-related nutrients we performed secondary analyses after excluding pancreatic cancer cases and person-years during the first two years of follow-up post-enrollment. Statistical analyses were conducted using SAS version 9.3 (SAS Institute, Inc., Cary, NC). All *P* values were two-sided and considered statistically significant if less than 0.05.

Results

After an average 16.3 years of follow-up, 271 cohort members (138 males and 133 females) developed pancreatic cancer. The mean age at cancer diagnosis was 72.0 years for males, and 71.6 years for females. The mean time from baseline interview to cancer diagnosis was 10.5 years (range, 3 months to 20.2 years).

Baseline characteristics were evaluated by highest and lowest quartile intake levels of vitamin B₆ and choline, two of the one-carbon metabolism-related nutrients with *a priori* hypothesis for associations with pancreatic cancer risk (Table 1). Overall, men (vs. women) were more likely to be ever smokers (57.7% vs. 8.7%) and alcohol drinkers (31.4% vs. 9.0%). Men and women in the highest quartile of vitamin B₆ intake were more likely to have a higher BMI and to be a never smoker than those in the lowest quartile. Among men, alcohol drinking and diabetes were more prevalent in the highest quartile of vitamin B₆. Men in the highest quartile of choline intake were more likely to be smokers and alcohol drinkers compared to those in the lowest quartile.

Positive correlations were observed between a majority of the one-carbon metabolism-related nutrients (Supplementary Table S1). The strongest correlations were observed between choline and other nutrients, including methionine [Spearman's correlation coefficient (*r*) = 0.70], vitamin B₂ (*r*=0.60) and B₁₂ (*r*=0.70). The weakest correlations were observed between methionine and other nutrients, including betaine (*r*=0.00) and folate (*r*=0.05). The Spearman's correlation coefficient of vitamin B₆ and choline was 0.50.

Risk of pancreatic cancer in females was 30% lower than that in males (Table 2). Compared with never smokers, no association was observed with former smokers, regardless of year since quitting (data not shown). Current smokers who smoked for 30 or more years had an increased pancreatic cancer risk (HR=1.61; 95% CI: 1.19–2.16) that was similar to those who smoked for less than 30 years (HR=1.55; 95% CI: 0.84–2.88), compared with never smokers. Among women, current smokers experienced a statistically significant 93% increased risk of pancreatic cancer compared with never smokers, and alcohol consumption of one or more drinks per week was associated with statistically non-significant 68% increased risk of pancreatic cancer, compared with nondrinkers. There were no significant associations between smoking or drinking and risk of pancreatic cancer in men.

Higher intakes of choline and vitamin B₆ were associated with decreased risk of pancreatic cancer in a dose-dependent manner (Table 3). The inverse associations between dietary

choline and vitamin B₆ and pancreatic cancer risk remained after excluding the first two years of follow-up. Comparing the highest versus the lowest quartile, the HRs (95% CIs) for choline and vitamin B₆ were 0.64 (0.45, 0.90) and 0.54 (0.37, 0.78), respectively (both *P*s for trend = 0.02). There was no association for pancreatic cancer risk with the intake of betaine, folate, methionine, vitamins B₂, or B₁₂ (Table 3). For comparison with a previous reports (24, 26), we evaluated potential joint effects between dietary folate and methionine on pancreatic cancer risk. Although no clear pattern emerged, higher dietary methionine was associated with reduced risk when folate was low (HR=0.64; 95% CI: 0.42–0.99; comparing third to first tertile) (Supplementary Table S2). There was no evidence for joint effects on pancreatic cancer risk with folate and vitamin B₆ or choline, or with vitamin B₆ and choline (all *P*s for interaction > 0.6).

The association between dietary choline or vitamin B₆ and pancreatic cancer risk was more apparent in men than in women (Table 3). However, we did not detect a statistically significant interaction between gender and either choline or vitamin B₆ on pancreatic risk (both *P*s = 0.15). There were also dose-dependent inverse associations for pancreatic cancer risk only in men for methionine (*P* for trend=0.02) and vitamin B₁₂ (*P* for trend=0.047). The gender- nutrient interaction on pancreatic cancer risk was statistically significant for vitamin B₁₂ (*P* for interaction=0.01), but not for methionine (*P* for interaction=0.11).

Discussion

We prospectively evaluated whether one-carbon metabolism-related nutrients were associated with pancreatic cancer risk and found that dietary intake of choline and vitamin B₆ demonstrated inverse, statistically significant trends. Highest quartiles, as compared to the lowest quartiles, of dietary vitamin B₆ and choline were associated with a 48% and 33% decrease in pancreatic cancer risk, respectively. We reported no association with the other one-carbon metabolism related-nutrients, including betaine, folate, methionine, vitamins B₂ and B₁₂.

Our finding of a dietary vitamin B₆-pancreatic cancer risk association differs from the null results observed in prospective cohorts in Sweden (24) and Finland (25), as well as a large population-based case-control study in the U.S. (26). Possible explanations for the discrepancy include the relatively low intake of vitamin B₆ in our cohort, as well as differences in the major food sources of vitamin B₆ in a Chinese versus Western diet. Less than 15% of our cohort met the U.S. Recommended Daily Allowance of 1.7mg and 1.5mg for men and women, respectively (9). In contrast, 75% of the Swedish cohort and 80% of the U.S. control population had consumed more than 1.7 mg/day from food only (24, 26). The major food sources of vitamin B₆ in our cohort were rice (25%) and fish (16%), compared with meat (29%) and cereals (17%) in Sweden (40), and read-to-eat cereal (13%), poultry (9.0%) and beef (8.7%) in the U.S. (41). The suggestive evidence that red meat intake is associated with an increased risk of pancreatic cancer (42) may partially explain the null results with vitamin B₆ that were observed in Western study populations. It is also possible that once the daily requirement is met there is no additional benefit with higher intake of vitamin B₆ on the risk of pancreatic cancer.

Our dietary vitamin B₆-pancreatic cancer finding supports the statistically significant inverse associations with higher circulating pyridoxal-5'-phosphate (PLP), the bioactive form of vitamin B₆, and pancreatic cancer risk that were reported in two European studies (22, 43), but not the null finding from a pooled analysis of four U.S. cohorts (23). We have previously reported a modest statistically significant correlation with plasma PLP and dietary vitamin B₆ in a healthy subset of our study population ($r = 0.17$, $P=0.0003$) (44). In addition to the lower intake and different major food sources of vitamin B₆ in our study population, perhaps our FFQ also captured the internal dose more accurately, compared with the studies that reported no association with dietary vitamin B₆ and pancreatic cancer risk.

A protective effect of vitamin B₆ on pancreatic cancer development is biologically plausible given vitamin B₆'s role as a cofactor for enzymes involved in the DNA synthesis and methylation pathways of one-carbon metabolism. As a cofactor for serine hydroxymethyltransferase, a diet low in vitamin B₆ results in a decreased production of the methyl donor, methylene-THF (45–47). A decrease in the methylene-THF pool may overload the DNA repair system by increasing uracil incorporation into DNA, and eventually lead to chromosome breaks (47, 48). Global DNA hypomethylation has been linked with genomic instability (49) and tumorigenesis (50, 51). The level of methylation of long interspersed nuclear element-1 (LINE-1) DNA sequences from peripheral lymphocytes is used as a biomarker for genomic DNA methylation status (52), and lower levels are associated with increased risk of some cancers (53). The level of LINE-1 methylation was measured in a healthy subset of our Singapore Chinese cohort (54), and in a secondary analysis, we found that dietary vitamin B₆ had a weak positive correlation with LINE-1 ($r=0.12$, $P=0.007$). In summary, it is biologically plausible that adequate intake of vitamin B₆ may reduce the risk of pancreatic cancer through its beneficial effects on DNA synthesis and methylation status.

Vitamin B₆ may play a role in preventing DNA from oxidative damage. In rats fed a vitamin B₆-deficient diet, decreased activity of pancreatic glutathione reductase, an enzyme that maintains the cellular glutathione level was reported (55). Glutathione is an antioxidant that is required for maintenance of the cellular redox state and detoxification of carcinogens, and low glutathione may impair the antioxidant defense system (12). Therefore, increasing oxidative stress may represent a mechanistic pathway by which low intake of vitamin B₆ may lead to increased pancreatic cancer risk.

To our knowledge, no epidemiological study has studied the relationship between dietary choline and pancreatic cancer risk. Our observed inverse association between choline intake and pancreatic cancer risk is consistent with the experimental evidence that dietary deficiencies of methyl donors, such as choline, led to aberrant differentiation and function of the exocrine pancreas and contributed to pancreatic carcinogenesis (56). A long-term choline-deficient ethionine (an antagonist of methionine)-supplemented diet in mice with induced chronic pancreatitis resulted in an increase in the expression of key molecules in the pancreatic carcinogenesis process, such as epidermal growth factor receptor (EGFR), K-Ras, and transforming growth factor α (TGF α) (57). Furthermore, a choline-deficient diet was able to shorten the induction period and increase the incidence of carcinogen-induced pancreatic carcinomas in hamsters (16, 58). In summary, our observed inverse association

between dietary choline and pancreatic cancer risk is biologically plausible, but it is not clear whether the role of choline is independent of the potential effects of methionine and/or vitamin B₁₂, given that intake of these three nutrients are strongly correlated with each other.

We did not observe statistically significant associations with dietary betaine, folate, methionine, vitamin B₂ or B₁₂ for pancreatic cancer risk. Dietary intake of betaine or vitamin B₂ has not been previously evaluated in relation to pancreatic cancer risk. Our finding for no association with folate was consistent with a pooled analysis of 14 prospective cohort studies (21). Our finding for no association with vitamin B₁₂ was consistent with results from the only other prospective study to evaluate an association with pancreatic cancer risk (25). Our finding for methionine was similar to two (25, 59), but not all (24) prospective studies that evaluated methionine-pancreatic cancer risk associations. In our sex-stratified results, a vitamin B₁₂-pancreatic cancer inverse association was only present in men, and vitamin B₁₂ was the only nutrient with a statistically significant interaction with sex. Vitamin B₁₂ functions as a cofactor for methionine synthase, an enzyme that converts homocysteine to methionine (9). In our data, vitamin B₁₂ was strongly correlated with methionine and choline (Supplementary Table S1), making it difficult to tease out the individual effects of these three nutrients on pancreatic cancer risk. The inverse associations with vitamin B₁₂, methionine, and choline intake and pancreatic cancer risk among men in our study suggests that compared to women, men may be more susceptible to low intake of these one-carbon metabolism-related nutrients (60). Our sex-specific findings, however, should be interpreted with caution, as they may be due to chance given the small number of cases available in the stratified analyses.

The strengths of our study include a prospective design, long duration of follow-up, and a comprehensive assessment of one-carbon metabolism-related nutrients. There are also limitations of our study. Due to the nature of an observational study and the one-time assessment of diet, our results may be influenced by misclassification of usual diet during the long follow-up period. However, given the prospective design, the potential for misclassification is unlikely to be different in cases and non-case participants; the non-differential misclassification could bias our results towards the null.

In summary, this prospective cohort study demonstrated statistically significant, inverse associations between dietary vitamin B₆ and choline, and pancreatic cancer risk. These novel findings support the hypotheses that vitamin B₆ and choline are relevant in pancreatic carcinogenesis. Future studies are needed to study the underlying mechanisms how vitamin B₆ and choline, as well as other correlated one-carbon metabolism-related nutrients, may protect against the development of pancreatic cancer.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: All authors received funding from National Institutes of Health/National Cancer Institute grants: R01 CA144034, U01 CA182876.

We thank Siew-Hong Low of the National University of Singapore for supervising the field work of the Singapore Chinese Health Study, and the Singapore Cancer Registry for assistance with the identification of cancer outcomes.

References

1. American Cancer Society. Cancer Facts & Figures 2014. American Cancer Society; Atlanta: 2014.
2. Gong Z, Holly EA, Bracci PM. Survival in population-based pancreatic cancer patients: San Francisco Bay area, 1995–1999. *Am J Epidemiol.* 2011; 174:1373–81. [PubMed: 22047824]
3. Iodice S, Gandini S, Maisonneuve P, Lowenfels AB. Tobacco and the risk of pancreatic cancer: a review and meta-analysis. *Langenbecks Arch Surg.* 2008; 393:535–45. [PubMed: 18193270]
4. Aune D, Greenwood DC, Chan DS, Vieira R, Vieira AR, Navarro Rosenblatt DA, et al. Body mass index, abdominal fatness and pancreatic cancer risk: a systematic review and non-linear dose-response meta-analysis of prospective studies. *Ann Oncol.* 2012; 23:843–52. [PubMed: 21890910]
5. Selhub J. Folate, vitamin B12 and vitamin B6 and one carbon metabolism. *J Nutr Health Aging.* 2002; 6:39–42. [PubMed: 11813080]
6. Lomberk G, Mathison AJ, Grzenda A, Urrutia R. The sunset of somatic genetics and the dawn of epigenetics: a new frontier in pancreatic cancer research. *Curr Opin Gastroenterol.* 2008; 24:597–602. [PubMed: 19122501]
7. Tan AC, Jimeno A, Lin SH, Wheelhouse J, Chan F, Solomon A, et al. Characterizing DNA methylation patterns in pancreatic cancer genome. *Mol Oncol.* 2009; 3:425–38. [PubMed: 19497796]
8. Sato N, Fukushima N, Maitra A, Matsubayashi H, Yeo CJ, Cameron JL, et al. Discovery of novel targets for aberrant methylation in pancreatic carcinoma using high-throughput microarrays. *Cancer Res.* 2003; 63:3735–42. [PubMed: 12839967]
9. Institute of Medicine (US). Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. Washington, D.C: National Academies Press (US); 1998. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes and its Panel on Folate OBV, and Choline.
10. Shane B. Folylpolyglutamate synthesis and role in the regulation of one-carbon metabolism. *Vitam Horm.* 1989; 45:263–335. [PubMed: 2688305]
11. Townsend JH, Davis SR, Mackey AD, Gregory JF 3rd. Folate deprivation reduces homocysteine remethylation in a human intestinal epithelial cell culture model: role of serine in one-carbon donation. *Am J Physiol Gastrointest Liver Physiol.* 2004; 286:G588–95. [PubMed: 14615285]
12. Choi SW, Friso S. Vitamins B6 and cancer. *Subcell Biochem.* 2012; 56:247–64. [PubMed: 22116703]
13. Jimenez-Chillaron JC, Diaz R, Martinez D, Pentinat T, Ramon-Krauel M, Ribo S, et al. The role of nutrition on epigenetic modifications and their implications on health. *Biochimie.* 2012; 94:2242–63. [PubMed: 22771843]
14. Pogribny IP, Tryndyak VP, Bagnyukova TV, Melnyk S, Montgomery B, Ross SA, et al. Hepatic epigenetic phenotype predetermines individual susceptibility to hepatic steatosis in mice fed a lipogenic methyl-deficient diet. *J Hepatol.* 2009; 51:176–86. [PubMed: 19450891]
15. Pogribny IP, Karpf AR, James SR, Melnyk S, Han T, Tryndyak VP. Epigenetic alterations in the brains of Fisher 344 rats induced by long-term administration of folate/methyl-deficient diet. *Brain Res.* 2008; 1237:25–34. [PubMed: 18694733]
16. Mizumoto K, Tsutsumi M, Denda A, Konishi Y. Rapid production of pancreatic carcinoma by initiation with N-nitroso-bis(2-oxopropyl)amine and repeated augmentation pressure in hamsters. *J Natl Cancer Inst.* 1988; 80:1564–7. [PubMed: 3193471]

17. Andry CD, Kupchik HZ, Rogers AE. L-azaserine induced preneoplasia in the rat pancreas. A morphometric study of dietary manipulation (lipotrope deficiency) and ultrastructural differentiation. *Toxicol Pathol.* 1990; 18:10–7. [PubMed: 2362985]
18. Longnecker DS, Chandar N, Sheahan DG, Janosky JE, Lombardi B. Preneoplastic and neoplastic lesions in the pancreas of rats fed choline-devoid or choline-supplemented diets. *Toxicol Pathol.* 1991; 19:59–65. [PubMed: 2047708]
19. Rios-Avila L, Coats B, Chi YY, Midttun O, Ueland PM, Stacpoole PW, et al. Metabolite profile analysis reveals association of vitamin B-6 with metabolites related to one-carbon metabolism and tryptophan catabolism but not with biomarkers of inflammation in oral contraceptive users and reveals the effects of oral contraceptives on these processes. *J Nutr.* 2015; 145:87–95. [PubMed: 25527663]
20. da Silva VR, Rios-Avila L, Lamers Y, Ralat MA, Midttun O, Quinlivan EP, et al. Metabolite profile analysis reveals functional effects of 28-day vitamin B-6 restriction on one-carbon metabolism and tryptophan catabolic pathways in healthy men and women. *J Nutr.* 2013; 143:1719–27. [PubMed: 23966327]
21. Bao Y, Michaud DS, Spiegelman D, Albanes D, Anderson KE, Bernstein L, et al. Folate intake and risk of pancreatic cancer: pooled analysis of prospective cohort studies. *J Natl Cancer Inst.* 2011; 103:1840–50. [PubMed: 22034634]
22. Chuang SC, Stolzenberg-Solomon R, Ueland PM, Vollset SE, Midttun O, Olsen A, et al. A U-shaped relationship between plasma folate and pancreatic cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Eur J Cancer.* 2011; 47:1808–16. [PubMed: 21411310]
23. Schernhammer E, Wolpin B, Rifai N, Cochrane B, Manson JA, Ma J, et al. Plasma folate, vitamin B6, vitamin B12, and homocysteine and pancreatic cancer risk in four large cohorts. *Cancer Res.* 2007; 67:5553–60. [PubMed: 17545639]
24. Larsson SC, Giovannucci E, Wolk A. Methionine and vitamin B6 intake and risk of pancreatic cancer: a prospective study of Swedish women and men. *Gastroenterology.* 2007; 132:113–8. [PubMed: 17241865]
25. Stolzenberg-Solomon RZ, Pietinen P, Barrett MJ, Taylor PR, Virtamo J, Albanes D. Dietary and other methyl-group availability factors and pancreatic cancer risk in a cohort of male smokers. *Am J Epidemiol.* 2001; 153:680–7. [PubMed: 11282796]
26. Gong Z, Holly EA, Bracci PM. Intake of folate, vitamins B6, B12 and methionine and risk of pancreatic cancer in a large population-based case-control study. *Cancer Causes Control.* 2009; 20:1317–25. [PubMed: 19415507]
27. Yuan JM, Stram DO, Arakawa K, Lee HP, Yu MC. Dietary cryptoxanthin and reduced risk of lung cancer: the Singapore Chinese Health Study. *Cancer Epidemiol Biomarkers Prev.* 2003; 12:890–8. [PubMed: 14504200]
28. Hankin JH, Stram DO, Arakawa K, Park S, Low SH, Lee HP, et al. Singapore Chinese Health Study: development, validation, and calibration of the quantitative food frequency questionnaire. *Nutr Cancer.* 2001; 39:187–95. [PubMed: 11759279]
29. Howe, JC.; Williams, JR.; Holden, JM.; Zeisel, SH.; Mar, MH. USDA Database for the Choline Content of Common Foods, Release One. 2004.
30. Patterson, KY.; Bhagwat, SA.; Williams, JR.; Howe, JC.; Holden, JM.; Zeisel, SH., et al. USDA Database for the Choline Content of Common Foods, Release Two. 2008.
31. U.S. Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Release 19. 2006.
32. U.S. Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Release 20. 2007.
33. U.S. Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Release 21. 2008.
34. Schakel SF, Sievert YA, Buzzard IM. Sources of data for developing and maintaining a nutrient database. *J Am Diet Assoc.* 1988; 88:1268–71. [PubMed: 3171020]
35. Willett, W. *Nutritional Epidemiology.* Oxford University Press; 1998.
36. Parkin DM, Whelan SI, Ferlay J, Teppo L, Thomas D. *Cancer Incidence in Five Continents.* 2002; VII

37. Cox DR. Regression Models and Life-Tables. *Journal of the Royal Statistical Society Series B (Methodological)*. 1972; 34:187–220.
38. Untawale S, Odegaard AO, Koh WP, Jin AZ, Yuan JM, Anderson KE. Body mass index and risk of pancreatic cancer in a Chinese population. *PLoS One*. 2014; 9:e85149. [PubMed: 24454807]
39. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2004; 363:157–63. [PubMed: 14726171]
40. Olsen A, Halkjaer J, van Gils CH, Buijsse B, Verhagen H, Jenab M, et al. Dietary intake of the water-soluble vitamins B1, B2, B6, B12 and C in 10 countries in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr*. 2009; 63(Suppl 4):S122–49. [PubMed: 19888270]
41. Cotton PA, Subar AF, Friday JE, Cook A. Dietary sources of nutrients among US adults, 1994 to 1996. *J Am Diet Assoc*. 2004; 104:921–30. [PubMed: 15175590]
42. World Cancer Research Fund, American Institute for Cancer Research. Food, nutrition, physical activity, and the prevention of cancer: A global perspective. AICR; Washington, DC: 2007.
43. Stolzenberg-Solomon RZ, Albanes D, Nieto FJ, Hartman TJ, Tangrea JA, Rautalahti M, et al. Pancreatic cancer risk and nutrition-related methyl-group availability indicators in male smokers. *J Natl Cancer Inst*. 1999; 91:535–41. [PubMed: 10088624]
44. Inoue-Choi M, Nelson HH, Robien K, Arning E, Bottiglieri T, Koh WP, et al. One-carbon metabolism nutrient status and plasma S-adenosylmethionine concentrations in middle-aged and older Chinese in Singapore. *Int J Mol Epidemiol Genet*. 2012; 3:160–73. [PubMed: 22724053]
45. Perry C, Yu S, Chen J, Matharu KS, Stover PJ. Effect of vitamin B6 availability on serine hydroxymethyltransferase in MCF-7 cells. *Arch Biochem Biophys*. 2007; 462:21–7. [PubMed: 17482557]
46. Stabler SP, Sampson DA, Wang LP, Allen RH. Elevations of serum cystathionine and total homocysteine in pyridoxine-, folate-, and cobalamin-deficient rats. *The Journal of Nutritional Biochemistry*. 1997; 8:279–89.
47. Ames BN. DNA damage from micronutrient deficiencies is likely to be a major cause of cancer. *Mutat Res*. 2001; 475:7–20. [PubMed: 11295149]
48. Huang YC, Chen W, Evans MA, Mitchell ME, Shultz TD. Vitamin B-6 requirement and status assessment of young women fed a high-protein diet with various levels of vitamin B-6. *Am J Clin Nutr*. 1998; 67:208–20. [PubMed: 9459368]
49. Chen RZ, Pettersson U, Beard C, Jackson-Grusby L, Jaenisch R. DNA hypomethylation leads to elevated mutation rates. *Nature*. 1998; 395:89–93. [PubMed: 9738504]
50. Gaudet F, Hodgson JG, Eden A, Jackson-Grusby L, Dausman J, Gray JW, et al. Induction of tumors in mice by genomic hypomethylation. *Science*. 2003; 300:489–92. [PubMed: 12702876]
51. Timp W, Bravo HC, McDonald OG, Goggins M, Umbricht C, Zeiger M, et al. Large hypomethylated blocks as a universal defining epigenetic alteration in human solid tumors. *Genome Med*. 2014; 6:61. [PubMed: 25191524]
52. Weisenberger DJ, Campan M, Long TI, Kim M, Woods C, Fiala E, et al. Analysis of repetitive element DNA methylation by MethyLight. *Nucleic Acids Res*. 2005; 33:6823–36. [PubMed: 16326863]
53. Woo HD, Kim J. Global DNA hypomethylation in peripheral blood leukocytes as a biomarker for cancer risk: a meta-analysis. *PLoS One*. 2012; 7:e34615. [PubMed: 22509334]
54. Inoue-Choi M, Nelson HH, Robien K, Arning E, Bottiglieri T, Koh WP, et al. Plasma S-adenosylmethionine, DNMT polymorphisms, and peripheral blood LINE-1 methylation among healthy Chinese adults in Singapore. *BMC Cancer*. 2013; 13:389. [PubMed: 23957506]
55. Dubick MA, Gretz D, Majumdar APN. Overt vitamin B-6 deficiency affects rat pancreatic digestive enzyme and glutathione reductase activities. *J Nutr*. 1995; 125:20–5. [PubMed: 7529303]
56. Longnecker DS. Abnormal methyl metabolism in pancreatic toxicity and diabetes. *J Nutr*. 2002; 132:2373S–6S. [PubMed: 12163695]
57. Ida S, Ohmuraya M, Hirota M, Ozaki N, Hiramatsu S, Uehara H, et al. Chronic pancreatitis in mice by treatment with choline-deficient ethionine-supplemented diet. *Exp Anim*. 2010; 59:421–9. [PubMed: 20660988]

58. Mizumoto K, Tsutsumi M, Kitazawa S, Denda A, Konishi Y. Usefulness of rapid production model for pancreatic carcinoma in male hamsters. *Cancer Lett.* 1990; 49:211–5. [PubMed: 2317782]
59. Skinner HG, Michaud DS, Giovannucci EL, Rimm EB, Stampfer MJ, Willett WC, et al. A prospective study of folate intake and the risk of pancreatic cancer in men and women. *Am J Epidemiol.* 2004; 160:248–58. [PubMed: 15257998]
60. Fischer LM, daCosta KA, Kwock L, Stewart PW, Lu TS, Stabler SP, et al. Sex and menopausal status influence human dietary requirements for the nutrient choline. *Am J Clin Nutr.* 2007; 85:1275–85. [PubMed: 17490963]

Table 1
Participant Characteristics According to Intake of Vitamin B₆ (mg) at Baseline, the Singapore Chinese Health Study, 1993–2013

Characteristics	Men				Women			
	Vitamin B ₆ intake ^a		Choline intake ^a		Vitamin B ₆ intake ^a		Choline intake ^a	
	Q1	Q4	Q1	Q4	Q1	Q4	Q1	Q4
N	8244	6953	8512	6680	6956	8130	15001	15012
Mean age, y	56.8	56.2	56.6	55.9	57.7	54.8	57.7	54.6
Body mass index (kg/m ²), %								
<18.5	7.8	5.5	6.4	7.3	6.8	5.2	5.9	5.8
18.5–21.4	24.7	21.8	23.2	23.1	22.2	22.0	22.1	22.3
21.5–24.4	45.0	42.8	45.4	41.8	47.8	44.3	48.2	43.2
24.5–27.4	16.0	20.7	17.6	18.5	15.0	18.2	15.5	18.5
27.5	6.6	9.3	7.4	9.3	8.2	10.2	8.2	10.2
Education, %								
No formal education	12.7	8.1	11.4	9.1	48.8	30.0	48.1	31.1
Primary school	55.0	46.5	53.4	48.7	38.2	40.0	38.0	40.2
Secondary school	32.3	45.4	35.2	42.2	13.0	30.1	13.9	28.7
Smoking status, %								
Never smokers	39.2	44.7	44.6	37.5	88.2	93.9	90.3	91.6
Former smokers	21.1	22.2	22.3	20.2	3.2	2.1	3.0	2.4
Current smokers	39.7	33.1	33.1	42.3	8.6	4.0	6.8	6.1
Alcohol drinking, %								
Nondrinkers	74.5	61.0	74.6	57.9	90.6	90.1	92.1	88.4
< 7 drinks/week	20.5	23.1	21.1	24.2	7.7	8.6	6.7	10.1
7 drinks/week	5.0	15.9	4.4	17.9	1.7	1.4	1.2	1.6
Weekly vitamin / minerals use (% Yes)	3.9	6.5	4.1	5.9	5.7	10.5	5.6	9.5
Weekly physical activity (% Yes)	42.1	46.9	44.6	41.7	21.3	30.1	22.6	27.5
Diabetes (% Yes)	6.9	9.5	6.8	10.3	8.4	8.2	8.0	10.1
Mean calorie intake, kcal/day	1879.8	1809.2	1878.6	1827.3	1540.1	1486.2	1415.8	1645.8
Mean nutrient intake ^b								
Betaine, mg/day	63.4	81.4	62.8	79.3	71.1	76.6	69.5	74.9

Characteristics	Men				Women			
	Vitamin B ₆ intake ^a		Choline intake ^a		Vitamin B ₆ intake ^a		Choline intake ^a	
	Q1	Q4	Q1	Q4	Q1	Q4	Q1	Q4
Choline, mg/day	188.7	259.8	160.4	306.0	199.7	260.8	174.3	292.4
Folate, µg/day	119.2	185.3	127.8	169.2	129.6	194.2	138.0	178.8
Methionine, mg/day	1171.2	1400.6	1098.0	1540.6	1208.5	1471.7	1129.8	1597.5
Vitamin B ₂ , mg/day	0.8	1.0	0.7	1.1	0.9	1.1	0.8	1.1
Vitamin B ₆ , mg/day	0.8	1.4	1.0	1.2	0.9	1.4	1.0	1.2
Vitamin B ₁₂ , µg/day	1.8	2.6	1.5	3.1	2.0	2.7	1.6	3.1

Abbreviations: Q1: 1st quartile; Q4, 4th quartile.

^a Nutrient intake was adjusted for daily total calorie intake by residual method. Quartiles of vitamin B₆ and choline were based on the distribution among the entire cohort.

^b Nutrient intake was adjusted for daily total calorie intake by residual method.

Associations between potential risk factors and pancreatic cancer risk overall, and stratified by sex, the Singapore Chinese Health Study, 1993–2013

Table 2

Characteristics	Total subjects			Men			Women		
	Cases, N	HR (95%CI) ^a	Cases, N	HR (95%CI) ^a	Cases, N	HR (95%CI) ^a	Cases, N	HR (95%CI) ^a	
Age	271	1.09 (1.08–1.11)	138	1.10 (1.07–1.12)	133	1.09 (1.06–1.12)			
Female vs. Male	271	0.70 (0.55–0.89)	138		133				
Body mass index, kg/m ²									
<18.5	22	1.44 (0.91–2.27)	13	1.53 (0.84–2.79)	9	1.33 (0.66–2.69)			
18.5–21.4	62	1.18 (0.86–1.60)	37	1.27 (0.84–1.92)	25	1.06 (0.66–1.70)			
21.5–24.4	114	1.00 (ref)	57	1.00 (ref)	57	1.00 (ref)			
24.5–27.4	55	1.33 (0.97–1.84)	26	1.20 (0.75–1.90)	29	1.48 (0.95–2.32)			
27.5	18	0.93 (0.56–1.52)	5	0.58 (0.23–1.45)	13	1.21 (0.66–2.21)			
Education									
No formal education	82	1.00 (ref)	14	1.00 (ref)	68	1.00 (ref)			
Primary school	121	1.07 (0.78–1.45)	75	1.30 (0.73–2.32)	46	1.00 (0.68–1.49)			
Secondary school	68	1.16 (0.81–1.68)	49	1.45 (0.79–2.67)	19	1.02 (0.60–1.76)			
Smoking									
Never smokers	166	1.00 (ref)	54	1.00 (ref)	112	1.00 (ref)			
Former smokers	34	0.96 (0.64–1.45)	31	0.92 (0.59–1.44)	3	0.76 (0.24–2.41)			
Current smokers	71	1.40 (1.02–1.92)	53	1.24 (0.84–1.81)	18	1.93 (1.17–3.20)			
Alcohol drinking									
Nondrinkers	222	1.00 (ref)	105	1.00 (ref)	117	1.00 (ref)			
< 7 drinks/week	39	1.00 (0.71–1.42)	24	0.77 (0.49–1.20)	15	1.68 (0.98–2.89) ^b			
7 drinks/week	10	0.80 (0.42–1.52)	9	0.79 (0.40–1.56)	1	--			
Diabetes									
No	241	1.00 (ref)	122	1.00 (ref)	119	1.00 (ref)			
Yes	30	1.30 (0.88–1.90)	16	1.49 (0.88–2.52)	14	1.19 (0.64–1.95)			
Weekly vitamin/mineral supplement use									
No	256	1.00 (ref)	132	1.00 (ref)	124	1.00 (ref)			
Yes	15	0.89 (0.53–1.51)	6	0.80 (0.35–1.82)	9	0.99 (0.50–1.94)			

^aHR and 95% CI were adjusted for age (years), sex, father's dialect group (Cantonese, Hokkien), and year of interview (1993–1995, 1996–1998).

^bHR and 95% CI were calculated for drinkers vs. nondrinkers. Only one case was in women < 7 drinks/week, and thus < 7 drinks/week was combined with 7 drinks/week.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Multivariate analysis of one-carbon metabolism dietary factors and pancreatic cancer incidence by sex, Singapore Chinese Health Study 1993–2013

Table 3

Nutrients	Median intake ^d	All		Men		Women	
		Cases, N	HR (95%CI) ^b	Cases, N	HR (95%CI) ^b	Cases, N	HR (95%CI) ^b
Betaine, mg/day							
Q1	41.56	68	1.00 (ref)	43	1.00 (ref)	25	1.00 (ref)
Q2	59.43	75	1.12 (0.81–1.57)	40	1.21 (0.78–1.87)	35	1.10 (0.65–1.83)
Q3	70.84	71	1.03 (0.73–1.44)	29	0.82 (0.51–1.33)	42	1.30 (0.79–2.14)
Q4	105.62	57	0.80 (0.56–1.14)	26	0.66 (0.40–1.08)	31	1.05 (0.62–1.79)
P-trend			0.19		0.051		0.71
Choline, mg/day							
Q1	176.30	97	1.00 (ref)	62	1.00 (ref)	35	1.00 (ref)
Q2	217.25	54	0.57 (0.40–0.80)	22	0.48 (0.29–0.78)	32	0.70 (0.43–1.13)
Q3	245.27	64	0.72 (0.52–1.00)	29	0.71 (0.45–1.11)	35	0.79 (0.49–1.27)
Q4	286.79	56	0.67 (0.48–0.93)	25	0.55 (0.34–0.88)	31	0.86 (0.53–1.41)
P-trend			0.04		0.02		0.69
Folate, µg/day							
Q1	108.24	74	1.00 (ref)	45	1.00 (ref)	29	1.00 (ref)
Q2	137.56	66	0.94 (0.67–1.32)	33	1.00 (0.63–1.58)	33	0.92 (0.55–1.52)
Q3	162.90	74	1.12 (0.81–1.56)	30	0.94 (0.58–1.50)	44	1.35 (0.83–2.17)
Q4	207.21	57	0.89 (0.63–1.28)	30	0.90 (0.56–1.44)	27	0.94 (0.54–1.61)
P-trend			0.82		0.62		0.73
Methionine, mg/day							
Q1	1073.17	72	1.00 (ref)	47	1.00 (ref)	35	1.00 (ref)
Q2	1268.95	88	1.26 (0.92–1.73)	48	1.35 (0.90–2.04)	40	1.17 (0.71–1.92)
Q3	1414.90	58	0.88 (0.62–1.25)	20	0.62 (0.37–1.05)	38	1.14 (0.69–1.89)
Q4	1625.25	53	0.82 (0.57–1.17)	23	0.67 (0.41–1.10)	30	1.02 (0.60–1.72)
P-trend			0.09		0.02		0.98
Vitamin B ₂ , mg/day							
Q1	0.71	71	1.00 (ref)	48	1.00 (ref)	23	1.00 (ref)
Q2	0.86	70	1.03 (0.74–1.44)	34	0.96 (0.61–1.51)	36	1.20 (0.71–2.04)

Nutrients	Median intake ^d	All		Men		Women	
		Cases, N	HR (95%CI) ^b	Cases, N	HR (95%CI) ^b	Cases, N	HR (95%CI) ^b
Q3	0.98	61	0.92 (0.65–1.30)	31	0.86 (0.54–1.36)	30	1.08 (0.63–1.87)
Q4	1.20	69	1.01 (0.72–1.42)	25	0.68 (0.42–1.11)	44	1.55 (0.93–2.59)
P-trend			0.88		0.12		0.13
Vitamin B ₆ , mg/day							
Q1	0.88	95	1.00 (ref)	62	1.00 (ref)	33	1.00 (ref)
Q2	1.02	64	0.71 (0.51–0.98)	22	0.48 (0.29–0.80)	42	1.03 (0.64–1.64)
Q3	1.13	68	0.80 (0.58–1.11)	31	0.73 (0.47–1.14)	37	0.98 (0.60–1.58)
Q4	1.33	44	0.52 (0.36–0.74)	23	0.45 (0.28–0.74)	21	0.66 (0.38–1.15)
P-trend			0.001		0.004		0.16
Vitamin B ₁₂ , µg/day							
Q1	1.43	74	1.00 (ref)	54	1.00 (ref)	20	1.00 (ref)
Q2	2.05	69	0.97 (0.69–1.35)	29	0.71 (0.45–1.12)	40	1.57 (0.91–2.70)
Q3	2.52	68	0.99 (0.70–1.38)	34	0.91 (0.59–1.41)	34	1.31 (0.75–2.28)
Q4	3.26	60	0.88 (0.62–1.24)	21	0.54 (0.33–0.90)	39	1.60 (0.93–2.74)
P-trend			0.51		0.047		0.21

^aNutrient intake levels were adjusted for daily total calorie intake using residual method.

^bAdjusted for age (continuous, year), sex (male, female), year of interview (1993–1995, 1996–1998), dialect group (Cantonese, Hokkien), education (no formal education, primary school, and secondary school or higher), BMI (<18.5, 18.5–21.4, 21.5–24.4, 24.5–27.4, 27.5 m/kg²), smoking status (never, former, current), diabetes (no, yes), alcohol drinking (no, yes), and weekly vitamin use (no, yes). Stratified analyses were not adjusted by sex. The cutoff values for one-carbon metabolism-related nutrients were the same for men and women.